

Physics of Partially Ionized Plasmas

Plasma is considered to be one of the four states of matter. The other three being solid, liquid and gas. It is formed by heating a gas, which ionizes its molecules or atoms. It consists of charged particles, namely positive ions and negative electrons. A partially ionized plasma has a large number of neutral particles too. Molecular clouds, diffuse interstellar gas, solar atmosphere, the earth's ionosphere and laboratory plasmas including fusion plasmas constitute partially ionized plasmas. The topic is an active area of study in plasma physics.

This book provides a comprehensive account of the various aspects of partially ionized plasmas including multifluid description, equilibrium, and waves. It includes a detailed discussion on the reionization phase of the universe along with a brief description of high discharge plasmas, tokomak plasmas and laser plasmas. It also discusses various elastic and inelastic collisions amongst the three particle species.

Using many examples, it demonstrates the novelties of partially ionized plasmas: for instance, in a partially ionized plasma, the magnetic induction is subjected to the ambipolar diffusion and the Hall effect in addition to the usual resistive dissipation. A discussion of kinematic dynamo in partially ionized plasma is also given in the book.

Vinod Krishan was Senior Professor at the Indian Institute of Astrophysics, Bangalore. She held visiting positions, at the Indian Institute of Science, the Lawrence Berkeley Laboratory, the National Institute for Space Research (Brazil), the University of Tokyo and the Raman Research Institute. During her long teaching and research career, she taught courses on plasma astrophysics, quantum mechanics and statistical physics at undergraduate and postgraduate levels. She has published more than a hundred research papers. Her major research interests are in the areas of space and astrophysical plasmas.



Cambridge Atmospheric and Space Science Series

Editors

Alexander J. Dessler John T. Houghton Michael J. Rycroft

Titles in Print in this Series

M. H. Rees

Physics and chemistry of the upper atmosphere

R. Dalev

Atmosphere data analysis

J. K. Hargreaves

The solar-terrestrial environment

J. R. Garratt

The atmosphere boundary layer

S. Sazhin

Whistler-mode waves in a hot plasma

S. P. Gary

Theory of space plasma microinstabilities

I. N. James

Introduction to circulating atmospheres

T. I. Gombosi

Gaskinetic theory

M. Walt

Introduction to geomagnetically trapped radiation

B. A. Kagan

Ocean-atmosphere interaction and climate modelling

D. Hastings and H. Garrett

Spacecraft-environment interactions

J. C. King and J. Turner

Antarctic meteorology and climatology

T. E. Cravens

Physics of solar system plasmas

J. F. Lemaire and K. I. Gringauz

The earth's plasmasphere

T. I. Gombosi

Physics of space environment

J. Green

Atmospheric dynamics

G. E. Thomas and K. Stamnes

Radiative transfer in the atmosphere and ocean

I. G. Enting

Inverse problems in atmospheric constituent transport

R. D. Hunsucker and J. K. Hargreaves

The high-latitude ionosphere and its effects on radio propagation

N. Meyer-Vernet

Basics of the solar wind

V. Y. Trakhtengerts and M. J. Rycroft

Whistler and Alfvén mode cyclotron masers in space

R. W. Schunk and A. F. Nagy

Ionospheres physics, plasma physics, and chemistry Second Edition

Mark C. Serreze and Roger G. Barry

The arctic climate system Second Edition

David Burgess and Manfred Scholer

Collisionless shocks in space plasmas: structure and accelerated particles



Physics of Partially Ionized Plasmas

Vinod Krishan





CAMBRIDGEUNIVERSITY PRESS

4843/24, 2nd Floor, Ansari Road, Daryaganj, Delhi - 110002, India

Cambridge University Press is part of the University of Cambridge.

It furthers the University's mission by disseminating knowledge in the pursuit of education, learning and research at the highest international levels of excellence.

www.cambridge.org

Information on this title: www.cambridge.org/9781107117396

© Vinod Krishan 2016

This publication is in copyright. Subject to statutory exception and to the provisions of relevant collective licensing agreements, no reproduction of any part may take place without the written permission of Cambridge University Press.

First published 2016

Printed in India

A catalogue record for this publication is available from the British Library

Library of Congress Cataloging-in-Publication Data

Names: Krishan, V. (Vinod)

Title: Physics of partially ionized plasmas / Vinod Krishan.

Description: Delhi, India : Cambridge University Press, 2016. \mid Includes

bibliographical references and index.

Identifiers: LCCN 2016001616 | ISBN 9781107117396 (hardback)

Subjects: LCSH: Space plasmas. | Plasma (Ionized gases)

Classification: LCC QC809.P5 K77 2016 | DDC 530.4/4-dc23 LC record available at

http://lccn.loc.gov/2016001616

ISBN 978-1-107-11739-6 Hardback

Cambridge University Press has no responsibility for the persistence or accuracy of URLs for external or third-party internet websites referred to in this publication, and does not guarantee that any content on such websites is, or will remain, accurate or appropriate.



> In loving memory of my father Shri Om Prakash Pabbi who taught me how to think



Contents

| | Figures | | |
|----------|---|---|------|
| | Prefa | ace | xiii |
| 1 | Par | tially Ionized Plasmas Here and Everywhere | 1 |
| | 1.1 | Plasma, a Matter of State | 1 |
| | 1.2 | Partial Ionization in the Early Universe | 1 |
| | 1.3 | Reionization Phase of the Universe | 6 |
| | 1.4 | Partially Ionized Plasma on Cosmic Scales | 7 |
| | 1.5 | Molecular Clouds | 7 |
| | 1.6 | Accretion Disks | 8 |
| | 1.7 | Interstellar Partially Ionized Plasma | 9 |
| | 1.8 | Solar Atmosphere | 9 |
| | | Cometary Tails | 14 |
| | | The Ionosphere | 14 |
| | | Partially Ionized Plasmas in Laboratory | 14 |
| | | Cold Plasmas in Medicine | 16 |
| | | Summary | 17 |
| | 1.14 | Problems | 17 |
| 2 | Multifluid Description of Partially Ionized Plasmas | | 18 |
| | 2.1 | Statistical Description | 18 |
| | | 2.1.1 The Maxwellian velocity distribution function | 19 |
| | 2.2 | Local Maxwell-Boltzmann Distribution Function | 25 |
| | | 2.2.1 The Boltzmann equation | 25 |
| | | 2.2.2 The fluid formalism | 28 |
| | | 2.2.3 The stress tensor | 33 |
| | | 2.2.4 Validity of the fluid picture | 36 |
| | | 2.2.5 The Boltzmann collision integral | 37 |
| | 2.3 | Three-fluid Description of a Partially Ionized Plasma | 37 |
| | 2.4 | Two-fluid Description of a Partially Ionized Plasma | 41 |
| | | 2.4.1 The MHD fluid | 41 |
| | | 2.4.2 Restrictions on the MHD fluid | 46 |
| | | 2.4.3 The neutral fluid | 47 |
| | | 2.4.4 The two-fluid system | 47 |



viii Contents

| | 2.5 | Single Partially Ionized Fluid—the SPIF | 48 |
|---|------|---|-----|
| | 2.6 | Weakly Ionized Plasma | 54 |
| | 2.7 | Summary | 56 |
| | 2.8 | Problems | 57 |
| 3 | Equ | iilibrium of Partially Ionized Plasmas | 59 |
| | 3.1 | The Equilibrium | 59 |
| | 3.2 | The Equilibrium Velocity Distribution Function | 59 |
| | 3.3 | Collisional Forces | 61 |
| | 3.4 | Collision Frequencies | 64 |
| | | 3.4.1 The electron-ion collision frequency | 64 |
| | | 3.4.2 The electron-electron collision frequency | 66 |
| | | 3.4.3 The electron-neutral collision frequency | 66 |
| | | 3.4.4 The ion-neutral collision frequency | 66 |
| | 3.5 | Equilibrium of the Three-fluid System | 67 |
| | 3.6 | Hydrostatic Equilibrium | 67 |
| | 3.7 | Hydrodynamic Equilibrium | 69 |
| | 3.8 | Equilibrium with Collisional Forces | 73 |
| | 3.9 | Equilibrium of Collisionally Dominated Fluids | 76 |
| | 3.10 | Three-fluid Equilibrium with Magnetic Field | 78 |
| | 3.11 | Three-fluid Equilibrium with Magnetic and Collisional Forces | 79 |
| | 3.12 | Equilibrium in the Two-fluid Description of a Partially Ionized | |
| | | Plasma | 83 |
| | | 3.12.1 Hydrostatic equilibrium | 83 |
| | | 3.12.2 Hydrodynamic equilibrium | 84 |
| | 3.13 | Equilibrium in Collisionally Dominated Two-fluids | 86 |
| | 3.14 | Two-fluid Equilibrium with Magnetic Field | 88 |
| | | 3.14.1 With current density $\vec{J} = J_{\theta}$ | 92 |
| | | 3.14.2 Current density $ec{J} \parallel ec{B}$ | 93 |
| | 3.15 | Two-fluid Equilibrium with Collisional Forces | 94 |
| | 3.16 | Equilibrium of the SPIF | 95 |
| | 3.17 | Equilibrium of the Weakly Ionized Plasma | 95 |
| | 3.18 | Hall Equilibrium of the Weakly Ionized Plasmas | 97 |
| | 3.19 | Summary | 99 |
| | 3.20 | Problems | 100 |
| 4 | Wa | ves in Partially Ionized Plasmas | 102 |
| | 4.1 | Waves in a Weakly Ionized Plasma | 102 |
| | | 4.1.1 Transverse waves in weakly ionized plasmas—the | |
| | | TWIP waves | 107 |



| | | | • | Contents | ix |
|---|---------------|--------|---|----------|-----|
| | | 4.1.2 | Polarization of TWIP waves | | 113 |
| | | 4.1.3 | Energy partition in TWIP waves | | 114 |
| | | 4.1.4 | | | 115 |
| | | 4.1.5 | Longitudinal waves in weakly ionized plasmas—the | 3 | |
| | | | LWIP waves | | 117 |
| | | | LWIP waves with $\vec{k} \perp \vec{B}_0$ | | 118 |
| | | | Polarization of LWIP waves with $\vec{k} \perp \vec{B}_0$ | | 119 |
| | | | 07 1 | | 119 |
| | | 4.1.9 | Dissipation of LWIP waves for $\vec{k} \perp \vec{B}_0$ | | 120 |
| | | | Oblique LWIP waves | | 121 |
| | | | in Single Partially Ionized Fluid—the SPIF | | 124 |
| | 4.3 | | in Two-fluid Picture | | 126 |
| | | | Incompressible wave-motions | | 126 |
| | | | Compressible wave-motions | | 135 |
| | | | ostatic Waves in Three-fluid Picture | | 140 |
| | | | ostatic Waves in Magnetized Three-fluids | | 149 |
| | 4.6 | | omagnetic Waves in Three-fluids | | 161 |
| | 4 = | | Electromagnetic waves in unmagnetized three-fluids | 3 | 163 |
| | 4.7 | | omagnetic Waves in Magnetized Three-fluids | | 168 |
| | | | Electromagnetic waves with $ec{k}\perpec{B}_0$ | | 170 |
| | | 4.7.2 | Electromagnetic waves with $ec{k} \parallel ec{B}_0$ | | 176 |
| | | Summ | • | | 181 |
| | 4.9 | Proble | ms | | 181 |
| 5 | \mathbf{Ad} | vance | d Topics in Partially Ionized Plasmas | | 184 |
| | 5.1 | - | ibrium of Partially Ionized Structures in the Solar | | |
| | | | sphere | | 184 |
| | 5.2 | _ | ibrium Plasma Structures with Hall Effect | | 186 |
| | 5.3 | | Equilibrium of the Weakly Ionized Plasmas | | 187 |
| | 5.4 | | Equilibrium of the Static Weakly Ionized Plasmas | | 188 |
| | 5.5 | • | mical Hall Equilibrium | | 189 |
| | 5.6 | | Equilibrium with Ambipolar and Resistive Effects | | 193 |
| | 5.7 | | r Relaxation in Weakly Ionized Plasmas | | 194 |
| | | | Taylor relaxation in a single-fluid | | 194 |
| | | | Taylor relaxation in an ideal single-fluid with flow | | 197 |
| | | | Taylor relaxation in a two-fluid system | | 197 |
| | F 0 | | Taylor relaxation in a weakly ionized plasma | | 199 |
| | 5.8 | | Naves in Rotating Weakly Ionized Plasmas | | 201 |
| | 5.9 | Exact | Non-linear Dispersion Relation | | 206 |



X Contents

| 5.10 | Kolmog | gorov Dissipation Scales in Weakly Ionized Plasmas | 207 |
|--------|----------|--|-----|
| 5.11 | Genera | tion of Magnetic Fields in Partially Ionized Plasmas | 213 |
| | 5.11.1 | Mean-field dynamo | 215 |
| | 5.11.2 | Mean-field dynamo in partially ionized plasma | 221 |
| | 5.11.3 | Hall dynamo | 222 |
| | 5.11.4 | Mean-field dynamo with ambipolar diffusion | 225 |
| | 5.11.5 | Hall and ambipolar dynamo | 228 |
| | 5.11.6 | Differential rotation-Hall-ambipolar dynamo | 229 |
| | 5.11.7 | Magnetic transport via ambipolar diffusion | 230 |
| | 5.11.8 | Stability analysis of ambipolar diffusion | 232 |
| | 5.11.9 | Sweet-Parker current slab in a weakly ionized plasma | 235 |
| | 5.11.10 | Parker-Sweet reconnection mechanism | 235 |
| | 5.11.11 | Parker-Sweet current slab | 237 |
| 5.12 | Summa | ry | 241 |
| 5.13 | Problen | ns | 242 |
| 6 Res | earch I | Problems in Partially Ionized Plasmas | 244 |
| 6.1 | The Thr | ee-fluid Description | 244 |
| 6.2 | The Two | p-fluid Description | 245 |
| 6.3 | The Sing | gle-fluid Description | 245 |
| 6.4 | Weakly 1 | Ionized Plasma | 245 |
| 6.5 | The Equ | ilibrium | 245 |
| 6.6 | Waves | | 246 |
| 6.7 | Continu | ation of Studies in Chapter 5 | 246 |
| Supple | ementary | Matter | 249 |
| Index | | | 259 |



Figures

| 1.1 | Time line of the universe. | 2 |
|------------|---|----------|
| 1.2 | Schematic variation of the ionization fraction, $f=n_e/n_H$, of hydrogen with redshift. | 6 |
| 1.3 | Multi-wavelength image of a cluster of galaxies containing a massive bright elliptical galaxy (large glow) that is ejecting jets of material into the cluster. New results from Chandra find that the ejected material significantly enriches the cluster with iron and other elements. | 7 |
| 1.4 | Image of M16 molecular cloud. | 8 |
| 1.5 | Accretion disk in a binary system, an illustration. | 9 |
| 1.6 | Schematic temperature variation with the radius (scaled to the radius of the sun) from the core to the photosphere. | 10 |
| 1.7 | Temperature variation of the sun from the photosphere to the corona. | 10 |
| 1.8 | Umbra and penumbra in a sunspot. | 11 |
| 1.9 | Hot and cold regions in a solar prominence. | 12 |
| 1.10 | Spicules, the cold and hot jets of plasma seen in the chromosphere. | 13 |
| 1.11 | This photograph of Comet West was taken by amateur astronomer John Laborde on March 9, 1976. The picture shows the two distinct tails. The thin tail is made up of gases, while the broad white tail is made up of tiny dust particles. | 13 |
| 1.12 | Various layers of the ionosphere and their predominant ion populations are listed at their respective heights above ground. | 15 |
| 1 12 | The density in the ionosphere varies considerably. | 16 |
| 2.1 | Cold plasma flame does not burn hands. Plot of the function g vs v at two different temperatures. | 23 |
| 2.2 | The three speeds, v_p , v_m and v_{rms} are schematically marked on a plot of the function g vs v at $T=300$ K. | 23 |
| 2.3 | The particles from a point P_1 move to a point P_2 in the phase space (x, v_x) such that the phase space particle density $f(x, v_x)$ remains | 26 |
| 2.1 | a constant. Deflection of an electron in the Coulomb field of an ion. | 26 64 |
| 3.1 | Simulation of an accretion disk. | 85 |
| 3.2 4.1 | (a) The equilibrium form of the uniform magnetic field frozen to all the three fluids. (b) The perturbed sinusoidal form of the | 63 |
| | magnetic field not frozen to any of the fluids. | 108 |



xii Figures

| 4.2 | Dispersion curves for the two TWIP waves including the Hall effect (a) Equation (4.46), (b) Equation (4.48) | 112 |
|-----|--|-----|
| 4.3 | The phase velocity curves for the two TWIP waves, Eq. (4.49). | |
| 1.0 | (a) $\left(\frac{V_{\text{PT}}}{V_{nA}}\right)$ (+) vs $(k_z\lambda_{nH})$, (b) $\left(\frac{V_{\text{PT}}}{V_{nA}}\right)$ (-) vs $(k_z\lambda_{nH})$ | 112 |
| 4.4 | The group velocity curves for the two TWIP waves, Eq. (4.52). | |
| | (a) $\left(\frac{V_{\text{GT}}}{V_{nA}}\right)$ (+) vs $(k_z\lambda_{nH})$, (b) $\left(\frac{V_{\text{GT}}}{V_{nA}}\right)$ (-) vs $(k_z\lambda_{nH})$ | 113 |
| 4.5 | The damping curves for the two TWIP waves, Eq. (4.70). (a) $[\omega_I(+)/(\alpha_0\omega_{\rm ic}D_T)]$ vs $(k_z\lambda_{nH})$ (b) $[\omega_I(-)/(\alpha_0\omega_{\rm ic}D_T)]$ vs $(k_z\lambda_{nH})$ | 116 |
| 4.6 | The positive charge Q is surrounded by negative charges forming the Debye cloud while the charged particles are also undergoing collisions with the neutrals. | 143 |
| 4.7 | Variation of the damping rate, ω_I , Eq. (4.245) with temperature $(T=T_e=T_i=T_n)$ for the sound wave of frequency $\omega_R=kC_{\rm ns}$ | 147 |
| 4.8 | Variation of the damping rate, ω_I , for the electron plasma wave of frequency $\omega_R = \omega_{\rm ep}$, Eq. (4.248) and the collision frequencies $\nu_{\rm ei0}$, Eq. (3.38) and $\nu_{\rm en0}$, Eq. (3.41) with temperature $(T=T_e=T_i=T_n)$ | 148 |
| 4.9 | Variation of the magnitude of the damping rate, $\omega_I \approx (\nu_{\rm in0}/4)$, with temperature $(T=T_e=T_i=T_n)$ for the ion acoustic wave | |
| | of frequency $\omega_R = \left(\omega_{ m ip}^2 + k^2 C_{ m is}^2\right)^{1/2}$ | 150 |
| 5.1 | Variation of the the ratio of the mass densities ρ_i/ρ_n as a function of the height on the solar atmosphere. | 185 |
| 5.2 | Cascading process in a 3-dimensional homogeneous turbulent fluid. There is a transfer of kinetic energy from large spatial scales to small spatial scales. The kinetic energy begins to dissipate into heat at a scale at which the Reynolds number becomes unity. | 208 |
| 5.3 | Magnetic induction $\pm B_y(x,z)$ being advected into the Parker-Sweet current sheet with speed $V_{\rm nz}$ in a weakly ionized | 200 |
| | plasma. | 236 |
| | | |



Preface

Retirement is a time to indulge oneself, determine your own deadlines and to meet them at your own pace. The cooking breaks are no brakes! This is how I came to write this book, my second, after my superannuation; the first one is called *Plasmas; The First State of Matter.*

I was introduced to the topic of partially ionized plasmas by Professor Kumar Chitre who handed me the A. Brandenburg and E. G. Zweibel paper (1995, Ap. J., 448, 734) during my visit to the Tata Institute of Fundamental Research sometime in the year 2000. This resulted in our paper "Ambipolar diffusion in the solar atmosphere'."

Partially ionized plasmas again came into my line of sight in the year 2005 when I visited the University of Tokyo campus near Edogawadai to work with Professor Yoshida-sansei. Since then, I have been studying plasma-typical problems in partially ionized plasmas. Around the same time I was also collaborating with Professor Swadesh Mahajan on the role of the Hall effect in diverse circumstances. It turned out that in the weakly ionized plasma model of a partially ionized plasma, the ion inertial scale, a hallmark of the Hall effect, gets multiplied by the inverse of the ionization fraction. As a result, the effective ion inertial scale acquires a much larger value than its counterpart in a fully ionized plasma. This was reported in "Equilibrium structures in partially ionized rotating plasmas within Hall magnetohydrodynamics"². During my visit to Professor S. Masuda's group in Nagoya University, Professor K. Shibata and I discussed the possibility of observing this new inertial scale phenomena on the solar atmosphere with a future solar mission. Additionally, during my visits to the Kyoto University and the National Astronomical Observatory of Japan, Mitaka, Tokyo, I had discussions on the mean field dynamo in partially ionized plasmas with Professor S. Tsuneta's group. The highly positive response from my peers galvanized me into pursuing the area further, and my own desire to present partially ionized plasmas as a subject in its own right resulted in this book.

The decade and a half's work could not have been done without the support of colleagues and collaborators, some of whom I have mentioned above. I wish to place on record my appreciation for Professor R. T. Gangadhara (my first student) for his continued collaboration on problems of our common interest. I have also enjoyed working with Professor B. P. Pandey, Dr Baba Varghese, Professor Nikhil Chakraborty, Professor H. Isobe, Professor Y. Hiraki, Dr K. A. P. Singh, and Dr A. Bhowmik.

¹Chitre, S. M., and V. Krishan. 2001. Mon. Not. R. Astron. 323: L23-L25

²Krishan, V., and Z. Yoshida. 2006. Physics of Plasmas. 13: 092303



xiv Preface

I thank Dr Ebenezer for helping me with the figures.

I am one amongst the many who learnt the hard lesson from B. K. Shivani that one is responsible for all the good and the not-so-good that happens in one's life; a lesson in ultimate freedom.

I thank my housekeeper, Nagaratna for relieving me from some of the daily household chores.

I am blessed with a very supportive extended family.

My husband, Professor Som Krishan, suffice it to say, has tolerated me for forty-seven years. The remaining, I hope, are easy, for him. My daughter, Dr Monika Krishan, a cognitive scientist and a spiritual healer maintains the flow of positive energy in our lives.

Most of all, I am indebted to Professor P. Sreekumar, Director, the Indian Institute of Astrophysics, for generously extending me institute facilities which enabled me to continue my work in a smooth manner.

I hope this book will serve the young and the not so young plasma researchers.