

Index

- accumulation, 71, 110, 111
 - non-ideal, 81
- agarose, 33, 35, 44
 - pore size, 59
- alkalinity, 35, 176, 183
- aluminium, 44, 67, 176, 238
- ammonia, 35, 176
- ammonium persulphate, 33, 266
- analytical techniques, 285
 - colorimetry, 181–182
 - computer imaging densitometry (CID), 181–182, 185, 202, 285
 - hyperspectral imaging, 182
 - laser ablation – ICPMS, 185–188, 285
 - PIXE, 175, 176
- antimony, 44, 67, 135
- area of window
 - effective, 14
 - physical, 3, 13
- arsenic
 - binding, 67
 - bioavailability, 217
 - DET, 176
 - high resolution, 176, 193
 - plant uptake, 250
 - sediment, 164
 - soil, 159
 - soil plant interactions, 205
 - solution, 16, 44, 135
- assembly of DGT devices, 269
 - sediments, 273
 - soils, 273
- association rate constant, 94, 229, 231
- Baltic Sea, 126, 127, 130, 138
- barium, 176
- binding layer, 67–71
 - 8-hydroxyquinoline, 67
 - activated carbon, 67
 - AG50W-X8, 67
 - Amberlite IRA910, 67
 - Amberlite IRP69, 67
- ammonium molybdophosphate, 67
- Chelex, 16, 66, 67, 135, 176
 - preparation, 194, 267
- copper ferrocyanide, 67
- Diphonix, 67
- ferrihydrite, 16, 67, 81, 135, 176
 - preparation, 194, 267
- iminodiacetate, 67, 176, 186, 194
- liquid phase, 67
- manganese oxide, 67
- methylthymol blue, 67
- Metsorb, 16, 67, 81, 176, 195
 - preparation, 268
- mixed, 67, 195, 269
- modified hydrogels, 67
- molecularly imprinted polymer, 67
- montmorillonite, 67
- multi-layer, 191
 - preparation, 194–197, 267
- silver iodide, 67, 176, 181, 195, 202
 - preparation, 268, 275
- TEVA resin, 67
- thiol-based, 67, 176, 196, 269
- titanium oxide. *See* Metsorb
- Whatman P81 membrane, 67
- XAD18, 16, 67
- zirconium oxide, 67, 176, 195, 268
- binding processes
 - approach to saturation, 107, 108
 - effect of binding strength, 109
 - effect of solution ligand, 108
 - key characteristics, 69
 - multiple layers-same material, 96, 105, 114
 - non-ideal, 70, 81, 109
- binding to hydrogels and filter membranes, 34–42
 - electrostatic, 38, 40–41
 - humic substances, 38
 - specific binding, 34, 38–40
 - trace metals, 38
 - copper, 55
- biotic ligand model, 225, 226, 248

294

biouptake
 aquatic moss, 140, 238
 fish, 238
 invertebrates, 238
 macroinvertebrates, 166
 molluscs, 238
 periphyton, 238
 plankton, 238
 bis-acrylamide, 34, 266
 blanks, 135, 273, 286
 bromide, 35
 cadmium
 binding, 67
 bioavailability, 217
 DET, 176
 free ion, 219, 220
 freshwater, 219, 220
 high resolution, 176
 kinetics, 105, 113, 114, 115
 plant uptake, 230, 245, 246, 250
 seawater, 130, 139
 sediment, 148, 166
 uptake by biota, 238
 soil, 159, 221
 2-D image, 180
 soil plant interactions, 205
 solution, 16, 35, 44, 135
 uptake by aquatic biota, 238
 calcium, 16, 35, 67, 176
 capacity
 of binding layer, 70, 81–82, 129, 198
 effective, 70, 81, 110
 intrinsic, 70, 81, 109
 of solid phase, 155
 cesium, 35, 44, 67
 charge effects
 in binding layer, 113–115
 in diffusion layer, 57–58
 Chelex resin, 3. *See* binding layer: Chelex,
 settling of beads, 116–117
 chloride, 35, 176
 chromium, 35, 44, 67, 159, 176
 bioavailability, 217
 plant uptake, 249
 uptake by biota, 238
 cobalt
 binding, 67
 bioavailability, 217
 DET, 176
 high resolution, 176
 kinetics, 105, 113, 115
 sediment, 164
 soil
 2-D image, 180
 solution, 16, 35, 44, 84
 colloids. *See* nanoparticles

Index

competition effects, 81, 84
 components
 sediment probes, 273
 soil devices, 273
 solution devices, 263
 concentration
 critical DGT for deficiency, 244, 251
 critical minimum in plant, 244
 DGT-measured
 assumptions, 10
 calculation of, 5, 272
 definition and interpretation, 11, 100
 equation, 4, 23–24, 123, 272
 effective, 161–163, 242
 equation considering DBL and effective area,
 12
 equation considering mixed ligands, 100
 free ion, 95, 146, 219, 220, 228, 236
 time weighted average, 5
 contamination, 264–265
 copper
 binding, 67
 bioavailability, 217, 218
 DET, 176
 free ion, 219, 220
 freshwater, 138, 219, 220
 high resolution, 176, 192
 kinetics, 105, 113
 plant uptake, 162, 245, 247, 249
 seawater, 126, 130, 139
 sediment, 148, 166
 uptake by biota, 238
 soil, 158, 168, 221
 2-D image, 180
 solution, 16, 35, 44, 55, 84, 135
 toxicity, 248
 uptake by aquatic biota, 235, 238
 cross-linker, 33–34
 depletion distance, 156
 deployment, 127, 275
 soil
 practicalities, 279
 preparation for, 278
 deployment time, 3, 108, 110, 111, 129–131
 in sediments, 284
 desorption, 152
 DET, 1, 34–42, 175, 189
 analytes in sediment, 164, 176, 179, 183, 184
 constrained probe, 275
 diffusion cell, 50
 diffusion coefficients, 3, 42–54, 124, 136–139
 effect of gel charge, 52–54, 57–58
 in gels and filters, 43–52
 in soils/sediments, 152
 in water, 43
 of complexes, 52

- recommended values, 51, 291
- size dependence, 43
- diffusion processes
 - in plant uptake, 228
 - in three dimensions, 154, 162
 - lateral diffusion, 14, 162
 - within DGT, 96–104
- diffusive boundary layer, 11–13, 135–136, 237, 277
 - apparent, 113
- dissociation
 - contribution to uptake flux, 228
 - Eigen mechanism, 115–116
 - ligand assisted, 115–116
 - rate constant, 94, 229, 230
- dissolved organic matter, 35, 219, 220
- distribution coefficient, 153, 166, 248
- Donnan effects, 53, 57–58, 113, 114
- dynamic techniques, 1, 234
- dynamics of resupply in soils/sediments, 163
- electrostatic effects. *See* charge effects
- elution, 3, 71–72
- flow rate of solution, 18, 22–23
- fluoride, 35
- flux
 - DGT for simple solutions, 3, 95
 - DGT with mixed ligands, 99
 - DGT measured in soils, 159
 - diffusion and internalization, 236
 - diffusion-limited, 229
 - diffusive in sediment/soil, 148
 - sustained in sediment/soil, 148
 - to plant and DGT, 243
 - uptake by biota, 224, 225, 229, 230
 - with a single ligand, 99
- fouling, 131
- free ion activity model, 225, 248
- freshwater, 42, 131, 136, 140, 219, 220
- fulvic acid, 35, 113, 138
- gold, 67
- heterogeneity of solute distributions, 6, 164
- holders
 - for plastic mouldings, 127, 128, 277
 - plastic mouldings
 - for measurements in soils, 264, 273
 - for sediment imaging. *See* sediment probes
 - for solution measurements, 3, 128, 263
- humic acid, 35, 44
- humic substances, 115
- hydrogels, 3, 32
 - preparation, 266–267
- hyperaccumulator, 159
- ionic strength effects
 - accumulation in simple solutions, 57, 133
 - accumulation of charged complexes, 113–115
 - binding performance, 82–83
- iron
 - binding, 67
 - DET, 176
 - high resolution, 176, 193
 - sediment, 148, 164, 183, 184
 - soil
 - 2-D image, 180
 - soil plant interactions, 205
 - solution, 35
- isotope composition, 141, 204, 206
 - neodymium, 140
- kinetic effects
 - binding rate, 70, 81, 110
 - in the binding layer, 105–108
 - solution complexes, 94–104
- kinetic signatures, 111–113
- lability, 6, 95–96, 147, 220, 234
 - lability degree, 95–96, 99, 101, 102, 117, 245
- lead, 67
 - bioavailability, 217
 - DET, 176
 - free ion, 219, 220
 - freshwater, 219, 220
 - high resolution, 192, 193
 - kinetics, 105, 113
 - plant uptake, 247, 249
 - sediment, 164, 166
 - uptake by biota, 238
 - soil plant interactions, 205
 - solution, 16, 35, 44, 135
 - uptake by aquatic biota, 238
- lithium, 35
- magnesium, 16, 35, 67
- manganese
 - DET, 176
 - high resolution, 176
 - kinetics, 105, 113
 - seawater, 139
 - sediment, 148, 164
 - soil
 - 2-D image, 180
 - soil plant interactions, 205
 - solution, 16, 35, 44, 84
- material diffusion layer, 3, 11
 - associated time-scale, 235
 - thickness, 19, 55, 112, 125, 235, 243
- maximum dynamic concentration, 137, 138
- membrane filters, 34, 35, 270

296

mercury, 44, 67, 166
 ethyl, 44
 high resolution, 176
 methyl, 44, 67, 72, 165, 166
 bioavailability, 217
 high resolution, 176
 plant uptake, 250
 uptake by aquatic biota, 238
 microniches, 6, 204
 mobility, 6
 modelling
 binding processes, 106–111
 DIFS, 153, 203, 281
 with multiple desorption sites, 168
 with solution speciation, 168
 of microniches, 203
 Penetration Analytical Model (PAM), 102, 105
 supply in soils and sediments, 151–157
 WHAM, 168, 219
 Modelling
 DIFS, 151
 molybdenum, 217
 binding, 67, 81
 solution, 44, 135

 nanoparticles, 58–60, 138, 217, 219, 236, 251
 nickel
 binding, 67, 109
 bioavailability, 217
 DET, 176
 free ion, 219, 220
 freshwater, 219, 220
 high resolution, 176
 kinetics, 105, 113, 114, 115
 plant uptake, 230, 246, 247, 250
 seawater, 126, 130, 139
 sediment, 148, 164
 uptake by biota, 238
 soil, 158
 2-D image, 180
 solution, 16, 35, 44
 uptake by aquatic biota, 238
 nitrate, 35, 176
 nutrients, deficiency of, 251

 particle concentration, 153
 partition coefficient. *See* distribution coefficient
 passive samplers, 2
 penetration into binding layer, 97
 performance tests, 271
 pH
 effect on binding, 82–83
 effect on measurement of metals, 134
 theoretical influence on accumulation, 103
 phosphate, 16, 35, 44, 67, 135
 DET, 176

Index

 high resolution, 176
 sediment, 184
 phosphorus
 bioavailability, 217
 freshwater, 132
 plant uptake, 251
 sediment, 164, 222
 soil, 222
 solid phase pool, 163
 planar optodes, 189, 191, 192, 193
 plant uptake, 161, 230, 242–251
 Barber–Cushman model, 232
 hydroponic, 228, 244, 246
 plutonium, 67
 polar organic compounds, 42
 antibiotics, 16, 42, 67, 161
 SMX, 42, 44
 bisphenols, 42, 67
 polyacrylamide gel, 33–34, 35, 44
 pore size, 60, 124
 preparation, 266
 potassium, 35, 67, 217, 251
 precision, 135, 180, 181, 272

 radon, 67, 176
 rare earths, 16, 44, 67
 ratio, R_c , 149, 165, 166
 reaction layer, 97, 230
 reaction processes
 within DGT, 96–104
 response time, 153, 166
 restricted gel, 34, 35, 44, 125, 266
 rhizosphere, 159, 205

 sandwich probes. *See* sediment probes: multi-layer
 seawater, 38
 sediment probes, 188–194, 274
 applied to rhizotrons, 190
 assembly and storage, 273, 274
 calculations, 285
 challenges
 ensuring contact, 197
 locating sediment water interface, 198
 material displacement on insertion, 197
 unsaturated soil, 199
 constrained, 190, 274, 275
 deployment, 281
 image generation, 199
 large format, 188
 multi-layer, 191
 selenium, 44, 67
 silver, 176, 217
 slicing of gels, 181, 284
 sorption, 152
 spatial resolution, 201
 speciation, 137, 218–223
 soils and sediments, 222, 233

- steady state, 24–27, 110
- storage and handling, 264–265
- storage of DGT devices, 269
- strontium, 35, 67, 176
- sulphate, 35, 176
- sulphide, 67, 176, 181, 184, 202, 204
- technetium, 67
- TEMED, 33, 266
- temperature effects, 27, 127–129
- thallium, 44, 176
- tin, 176
- toxicity, 223, 225, 233, 248
- transient regime, 24–27
 - effect of binding, 54–57
- tungsten, 44, 67, 176
- ultra-filtration, 138
- uncertainty
 - in DGT measurements, 135, 272, 287
 - in using DIFS, 159
- uptake by biota, 223
 - biotic ligand model, 228
 - diffusion layer, 235
 - diffusion-limited, 229
 - free ion activity model, 228
 - from sediments, 233
 - from soils, 230
 - Michaelis–Menten, 223, 229
 - related to DGT
 - sediments, 238
 - water, 238
 - uptake by DGT, 71, 72
- uranium, 16, 35, 44, 67, 176
 - bioavailability, 217
 - plant uptake, 250
- vanadium, 44, 67, 176, 200
- zinc
 - binding, 67
 - bioavailability, 217
 - DET, 176
 - free ion, 219
 - freshwater, 219
 - high resolution, 176
 - plant uptake, 230, 245, 246, 247, 249, 250
 - seawater, 130, 139
 - sediment, 148, 164
 - uptake by biota, 238
 - soil, 158, 159, 221
 - 2-D image, 180
 - soil plant interactions, 205
 - solution, 84, 135
 - toxicity, 248