

Index

- absolute convergence, 45n2, 55n7
- Addition Rule, 9–10
- almost sure convergence, 117n29
- asymptotic estimation theory, 111–116
- asymptotic mean squared error, 113
- asymptotic normality, 112, 125
 - of plug-in estimators, 120
- asymptotic relative efficiency, 113
- asymptotic standard error, 113
- asymptotic unbiasedness, 112
- atomic events, 6
- average marginal causal effect, 255
- average partial derivative, 166–168, 171, 255
- average treatment effect (ATE), 237–238
 - conditional, 249–250
 - decomposition, 238
 - plug-in estimation of, 256–257
 - sharp bounds for, 240–243
 - under random assignment, 245
 - under strong ignorability, 248–249, 253, 255
- average treatment effect among the treated (ATT), 238, 247, 273

- balance, 218, 251–252
- balance test, 252, 270n14
- bandwidth, 121
- Bayes' Rule, 11–12
 - alternative form of, 13
- Bayesian probability, 4n2
- Bernoulli distribution, 19, 46, 58, 67, 70, 135, 182
 - generalized, 180
- Bernoulli random variable, *see* Bernoulli distribution
- best linear predictor (BLP), 76–84
 - and independence, 82–84
 - and the CEF, 78–79
 - coefficients of, 87
 - estimating, 144–147
 - multivariate case, 86–87
 - properties of deviations from, 79, 87
 - under nonlinearity, 81
- bias, 103
- binary choice model, 182–185, 226, 233, 264, 279
 - CEF for, 183, 198
- bivariate normal distribution, 66n13
- bivariate relationships, 31–39
- bootstrap, 130–132, 141–142, 153–154, 203, 234
- bounds, *see* sharp bounds

- cardinality, 7, 284
- categorical distribution, 180
- Cauchy–Schwarz Inequality, 63n12
- causal inference, 235–281
 - as a missing data problem, 238–239
 - Double Robustness Theorem for, 267–269
 - doubly robust estimator for, 267, 269–270
 - propensity score for, 250–252
 - regression estimator for, 260, 261
- Central Limit Theorem (CLT)
 - for cluster samples, 139
 - for sample means, 108–110

- Chebyshev's Inequality, 55, 105
 for the sample mean, 98
- classical linear model, 180–182
 CEF for, 181, 198
 maximum likelihood estimation of, 189–192
- cluster average, 136–139
 cluster sample mean, 136–137
 sampling variance of, 137, 140
- cluster samples, 135–141
 CLT for, 139
 estimation with, 136–139
 inference with, 140–141
 WLLN for, 138
- collinearity, 132n43, 149
- complement, 6, 22, 24, 283
- complement rule, 8, 9
- conditional average treatment effect, 249–250
 under strong ignorability, 249
- conditional expectation, 67–69
 linearity of, 69, 249
 multivariate case, 85
 of a function of a random variable, 68
- conditional expectation function (CEF), 69–76
 and independence, 82–84
 and the BLP, 78–79
 for binary choice models, 183, 198
 for the classical linear model, 181, 198
 kernel plug-in estimation of, 124
 multivariate case, 85
 partial derivatives of, 165–168
 polynomial approximation of, 158, 165n11, 169
 properties of deviations from, 74–75
 under MAR, 217
- conditional independence assumption, 216, 248, 254, 255, 270–272, 274
- conditional PDF, 37–38
 multivariate case, 42
- conditional PMF, 34
 multivariate case, 42
- conditional probability, 10–12, 67, 182
 and independence, 14–15, 39
- conditional variance, 68, 74
- conditional variance function, 70, 85
- confidence interval, 124–127, 134–135, 212, 231
 asymptotic validity of, 125
 for regression, 154
 for the population mean, 127
 normal approximation-based, 125, 128, 131, 133–135, 140
 validity of, 124
- confidence level, 124
- consistency, 105
 of maximum likelihood estimation, 193
 of sampling variance and standard error estimators, 113
 of the sample mean, 105
- Continuous Mapping Theorem (CMT), 100, 107, 108, 110, 114, 116, 125–126, 140
- continuous random variable, 24
 event probabilities for, 25–27
- convergence
 absolute, 45n2, 55n7
 almost sure, 117n29
 in distribution, 108
 in probability, 99–100
- correlation, 62–63
 and independence, 64–66
 and linear dependence, 62
 properties of, 63
- covariance, 59–60
 alternative formula for, 60
 and independence, 64
 properties of, 61–62
- covariance matrix, 84
- cumulative distribution function (CDF), 21–24, 101–102
 empirical, 116–120
 estimating, 102, 117
 joint, 32–33, 40
 properties of, 22
- degenerate distribution, 47
- difference-in-means estimator, 245, 256, 257, 259
- discrete random variable, 18–21
 event probabilities for, 21
- distribution, 3–4, 20, 21
- Double Robustness Theorem
 for causal inference, 267–269
 for missing data, 229–230
- doubly robust (DR) estimator
 for causal inference, 267, 269–270
 for missing data, 228, 230
- draw, 4
- efficiency, *see* relative efficiency
- elementary event, 6
- empirical CDF, 116–117, 120
- estimation theory, 96–116
 asymptotic, 111–116
- estimator, 103–105
 bias of, 103
 consistency of, 105

- doubly robust, *see* doubly robust (DR) estimator
- inverse probability-weighted (IPW), *see* inverse probability-weighted estimators
- kernel density, 122
- kernel plug-in, 123
- lasso, 169–170, 174, 202
- MSE of, 104–105
- plug in, *see* plug-in estimator
- post-stratification, 219, 256
- regression, *see* regression
- relative efficiency of, 105
- sampling variance of, 104
- sieve, 168–169
- standard error of, 104
- unbiased, 103
- event, *see* random event
- event space, 5–8
- expectation, 45–50
 - conditional, 67–69
 - linearity of, 49–50
 - of a function of a random variable, 47
 - of a function of two random variables, 49
 - of a random vector, 48
 - of the sample mean, 97
 - properties of, 47–50
 - under MAR, 216
 - under MCAR, 213–215
- expectation operator, 45, 47
- expected value, *see* expectation
- experiment, 4
- explanatory variable, 79

- finite population, 93–94
- frequentist probability, 4
- function of a random variable, 16–17
 - conditional expectation of, 68
 - equality of, 32, 70
 - expectation of, 47, 49, 68
- functional, *see* statistical functional
- Fundamental Problem of Causal Inference, 236–239, 270

- geometric distribution, 19
- Glivenko–Cantelli Theorem, 102n14, 117n27, 120n31

- heteroskedasticity, 156
- homoskedasticity, 155, 156, 261
- hot deck imputation, 224–225, 233
- hypothesis testing, 128–130

- i.i.d., 91–92, 94–98
- identification
 - with missing data, 208–219
 - with potential outcomes, 236–256
- ignorability, *see* missing at random (MAR)
- imputation
 - hot deck, 224–226, 233
 - under MAR, 220
 - under MCAR, 214–215
 - under random assignment, 246
 - under strong ignorability, 258
- independence
 - CEFs and BLPs under, 82–84
 - conditional, 216, 248, 251, 254, 255, 270–272, 274
 - conditional probability and, 14–15, 39
 - correlation and, 64–66
 - covariance and, 59–60
 - implications of, 39, 64–65, 82
 - of events, 14
 - of random variables, 38–39
- independent and identically distributed, *see* i.i.d.
- inference, 124–135
 - with clustering, 140–141
 - with maximum likelihood estimation, 202–203
 - with regression, 151–156
- interactions, 164–165
- inverse probability-weighted (IPW) estimators, 226–228, 231, 234, 264, 266–267, 270, 275, 279
 - for causal inference, 266–267
 - for missing data, 227–228
 - stabilized, 228, 266–267

- joint CDF, 32–33
 - multivariate case, 40
- joint PDF, 36
 - multivariate case, 41
- joint PMF, 32
 - multivariate case, 40
- joint probability, 9–10
- jointly continuous random variables, 36–38, 41, 42, 49, 50, 64–65, 67–69, 72–73, 85

- kernel, 121
- kernel density estimator, 121
- kernel plug-in estimator, 123
- Kolmogorov Axioms, 6–8, 20, 183
- Kullback–Leibler Divergence, 195–196

- lasso estimator, 169–170, 174, 202
 Law of Iterated Expectations, 71–76, 155, 167, 198, 211, 212, 216, 218, 220, 227, 229, 241–243, 248, 250, 251, 258, 265, 268
 Law of Total Probability, 12–13, 15, 33
 Law of Total Variance, 73–75
 leave-one-out cross-validation, 170, 174
 likelihood function, 186, 189
 linearity of expectations, 49–50
 link function, 183n3
 log-likelihood, 187–190, 192, 194, 202
 logit model, 183–184, 233
- M-estimators, 187n7
 Manski bounds, *see* sharp bounds
 marginal effect, 165
 marginal PDF, 37
 multivariate case, 41–42
 marginal PMF, 33–34
 multivariate case, 41–42
 matching, 262–264
 maximum likelihood (ML) estimation, 185–203, 233
 consistency of, 193
 inference with, 202–203
 logic of, 185–193
 of mixture models, 198–201
 of the classical linear model, 189–192
 of the probit model, 192–193
 penalized, 201–202
 plug-in estimator, 197–198
 under misspecification, 194
 mean function, 183–184, 226, 264
 mean squared error (MSE), 56–58
 alternative formula for, 57
 asymptotic, 113
 expected value minimizes, 57–58, 151
 of an estimator, 104
 root, 57
 micronumerosity, 132–135, 149, 161, 162, 168, 274
 missing at random (MAR), 215–217
 and the propensity score, 217–219
 CEF under, 217
 expected value under, 216
 imputation under, 220–221
 reweighting under, 226–227
 missing completely at random (MCAR), 213–215
 expected value under, 213–214
 imputation under, 214–215
- missing data, 208–219
 Double Robustness Theorem for, 230, 267
 doubly robust estimator for, 228, 230
 plug-in estimator for, 220–222
 propensity score for, 217–219
 regression estimator for, 224
 with binary outcomes, 209, 212, 214, 220, 222, 239
 mixture model, 198–201
 models, *see* parametric models
 moments
 central, 51
 raw, 51
 multicollinearity, *see* collinearity
 multiplicative law for PDFs, 38
 multiplicative law for PMFs, 35
 Multiplicative Law of Probability, 10, 11, 13, 14, 35, 38
 multivariate normal distribution, 152
- negative control, *see* placebo testing
 Newton–Raphson method, 191n11
 Neyman–Rubin Causal Model (NRCM), 236, 239, 254
 normal distribution, 55–56
 bivariate, 66n13
 mean and standard deviation of, 56
 properties of, 56
 standard, 28–30
 null hypothesis, 128–130, 278
- operator, 17, 45, 47, 67, 118
 ordinary least squares (OLS) regression estimator, 145–146, *see also* regression
 outcome, 4–5, *see also* potential outcomes model
 overfitting, 161–164
 overlap
 empirical, 273–276
 population, 271–273
- p -value, 128–133, 140, 152, 154, 278, 279
 parameter space, 179, 180, 183
 parameter vector, 179, 182, 185
 parametric models, 178–203
 as approximations, 203
 binary choice models, 182–185
 classical linear model, 180–182
 maximum likelihood estimation of, *see* maximum likelihood estimation
 partial derivative at the average, 166–168, 171

Index

297

- partition, 12–13
- penalized maximum likelihood estimation, 201–202
- penalized regression, 169–170
- penalty parameter, 169, 202
- placebo testing, 270–271
- plug-in estimator, 120
 - asymptotic normality of, 120
 - for the ATE under strong ignorability, 256–257
 - for the expected value under MAR, 219–220
 - maximum likelihood, 197–198
 - plug-in sample variance as, 119
 - regression as, 143–147
 - regularity conditions, 120
 - sample mean as, 118, 210, 240
- plug-in principle, 116–124
- plug-in sample variance, 105–108
 - as plug-in estimator, 119
 - properties of, 106
- point identification, 207
- polynomial approximation of the CEF, 158, 165n11, 169
- population, 91, 95–96
 - finite, 93–94
- population mean, 93, 94, 97, 101, 103, 105, 106, 110, 115, 124, 127, 132, 133, 143, 151, 212, 214, 219, 224, 243
 - confidence interval for, 127
 - consistency of the sample mean for, 105
- population regression function, 145
- population variance, 94, 105, 116
- positivity, 271–276
- post-stratification estimator, 219, 256
- post-treatment variables, 252–254
- potential outcomes model, 236–239, 245, 248, 262, 265, 277, 278
 - generalized, 254–256
- probability density function (PDF), 24–25
 - conditional, 37–38, 42
 - joint, 36, 40
 - marginal, 37, 42
 - multiplicative law for, 38
 - properties of, 25
- probability limit, 99
- probability mass function (PMF), 18–21
 - conditional, 42
 - joint, 32, 40
 - marginal, 33–34, 41–42
 - multiplicative law for, 35
 - properties of, 20
- probability measure, 6, 7
- probability space, 6, 8
- probit model, 183
 - maximum likelihood estimation of, 192–193
- propensity score
 - for causal inference, 250–252
 - for missing data, 217–219
 - imputation with, 224–225
 - MAR and, 217–219
 - matching on, 262–264
 - maximum likelihood plug-in estimation of, 225, 264
 - strong ignorability and, 251–252
 - Winsorization, 275
- pseudopopulation, 95
- random assignment, 244–247
 - ATE under, 245
 - imputation under, 246
- random event, 4, 15
 - independence of, 14
- random generative process, 3–8, 14–16, 93
- random sampling, 92–94
 - finite population, 93–94
 - with replacement, 94
 - without replacement, 94
- random variable, 15–31
 - continuous, 24–30
 - degenerate, 47, 108, 110, 113
 - discrete, 18–21
 - equality of, 31–32
 - function of, 16
 - independence of, 38–39
 - independent and identically distributed, 91–92
 - operator on, 17, 45
- random vector, 39–40
 - covariance matrix of, 84
 - expectation of, 48
- range, 16n15, 18, 284
- realization, 4
- regression, 143–151
 - as plug-in estimator, 143–147
 - bivariate case, 144–145
 - classical standard errors for, 155–156
 - confidence intervals for, 154
 - estimator for causal inference, 258–261
 - estimator for missing data, 222–224
 - penalized, 169–170
 - properties of residuals from, 149–151
 - robust sampling variance estimator for, 153
 - robust standard errors for, 152–154
 - with matrix algebra, 147–151

- regressor matrix, 147
- relative efficiency, 105
 - asymptotic, 113
- residual, 147–149
- response propensity function, 217, 225, 230
- reweighting
 - survey, 220–222
 - under MAR, 226–227
 - under strong ignorability, 265–266
- root mean squared error, 57

- sample mean
 - as plug-in estimator, 118, 210, 240
 - Chebyshev's Inequality for, 98–99
 - consistency of, 105
 - consistency of the standard error of, 115
 - estimating the sampling variance of, 114
 - expected value of, 97
 - sampling variance of, 97–98
 - standard error of, 115
 - standardized, 109, 139
- sample points, 5
- sample space, 5
- sample statistic, 96
- sample variance
 - plug-in, 105–107
 - unbiased, 107
- sampling, *see* random sampling
- sampling variance, 104
 - estimator of, 114–115
 - of the cluster sample mean, 137, 140
 - of the sample mean, 97–98
- sandwich estimator, 153
- sharp bounds, 209–213, 231, 240–244
 - estimating, 211–212, 243–244
 - for the ATE, 240–243
 - for the expected value, 209–211
- sieve estimator, 168–169
- sigma field, 6n6
- simple event, 6
- Slutsky's Theorem, 109–110, 125, 126
- stabilized IPW estimator
 - for causal inference, 266–267
 - for missing data, 228
- stable outcomes model, 208
- stable unit treatment value assumption, 237
- standard deviation, 53
 - of the normal distribution, 56
 - properties of, 53–54
- standard error, 104
 - asymptotic, 113
 - estimator of, 113–115
 - of the regression estimator, *see* regression
 - of the sample mean, 115
- standard normal distribution, 28–30, 55
 - expected value of, 46
- standard uniform distribution, 28
- standardized sample mean, 109, 139
- statistical functional, 118
- strong ignorability, 247–258, 261, 262
 - and the propensity score, 250–252
 - ATE under, 248–249, 253, 255
 - conditional ATE under, 249
 - imputation under, 257–258
 - reweighting under, 265–266
- superpopulation, 94
- support, 30–31
- survey reweighting, 221

- t*-statistic, 129
- test statistic, 128n37
- treatment effect, *see* unit-level treatment effect
- treatment propensity function, 250, 264, 269, 273

- unbiased sample variance, 107–108
 - properties of, 108
- unbiasedness, 103–105
 - asymptotic, 112
- uniform distribution, 28
- unit-level treatment effect, 237

- variance, 51–53
 - alternative formula for, 52, 57
 - conditional, 68
 - population, 94, 105, 116
 - properties of, 52–53
 - sample, 105–108
 - sampling, 104
- variance rule, 60, 65
 - multivariate, 84, 137

- Weak Law of Large Numbers (WLLN), 99–102, 105, 107, 189, 193, 212, 243
 - for cluster samples, 138
- Weierstrass Approximation Theorem, 158
- weighting estimator, 226–228, 264–267, 273–276
- Winsorization, 275