

Cambridge University Press

978-1-107-40383-3 - Granivorous Birds in Ecosystems: Their Evolution, Populations, Energetics, Adaptations, Impact and Control

Edited by Jan Pinowski and S. Charles Kendeigh

Index

[More information](#)

Index

- Acanthus cannabina* communal roosts, 331, egg weight loss during incubation, 112, flock feeding, 331
- Acanthus flammea* basal metabolism, seasonal changes, 130, diet and feeding behaviour, 313, as a non-pest species, 335, 337, 339, as a pest species, 313, 339–40
- Acanthus* spp breeding behavior, 324, brood number, 324, cage locomotor activity, 148, diet, 324, migration, 195
- Acrodoteres cristatellus*, as a pest species, 299
- Aegithalos caudatus*, roosting behavior, 156
- aesthetics and recreation, role of birds, 206–7
- African weaver finch, *see Ploceus cucullatus*
- Agelaius phoeniceus*, 4, biomass production, 122–4, 125–6, (eggs) 122, (young) 122–4, (adults) 124, 126, breeding behavior, 86, 90, 94, 239, 321–2, 328, 334, breeding biotopes, 90, 94, 103, 207, 321, 329, diet and feeding behavior, 239–44, 293, 312, 334, distribution, 5, 6, 53, 207, energy demand, 239, flocking and roosting behavior, 86–7, 94, 102–4, 207, 210, 246–8, 267–8, 293, 312, 328–9, migration, 86–7, 267–8, as a pest species, 103–4, 207, 240, 246–7, 267–300, 335–40, population (breeding) density, 92, 102–3, 239–40, production values, 88–93, (brood number) 126, (clutch size) 88–90, (hatching success) 88–90, (fledging success) 88–90, (adult survival) 90–2, (life expectancy) 92–3, sex ratio, 88–90, size, 240–1
- Agelaius* spp breeding behavior, 334, commensalism with man, 48, diet and feeding behavior, 334
- Agelaius tricolor* breeding behavior, 317–18, 321, 324, 328–9, 334, diet and feeding behavior, 317, distribution, 316–18, flocking and roosting behavior, 317, 321, 328–9, migration, 317, as a pest species, 335–40
- agricultural communities animals, 20, crops, 15, 20, development, 15, 19–22, 50, distribution, 19–20, 23, housing, 20–1, migration, 24, settlements, 19–21
- agriculture archaeological records, 19–22, bird impact on, 209–12, development, 15, 19–22
- American goldfinch, *see Carduelis tristis*
- American Inca dove, *see Scardafella inca*
- Ammodramus deserti*, nest position and incubation of eggs, 171
- Amphispiza belli*, as a non-pest species, 335, 337
- Anas discors*, weight change, 153
- Anas platyrhynchos*, energy cost of incubating eggs, 169, swimming, 164
- Anas rubripes* energy requirement for growth, 187–8, metabolizable energy coefficient, 192
- Apodemus sylvaticus*, egg predator, 73
- Aratinga* spp, as pests, 268
- archaeological records, development of agriculture, 19–22
- Baird's sandpiper, *see Erolia bairdu*
- basal metabolism (BM), 129–34, adaptation to local climate, 133–4, daily rhythm, 129–30, seasonal changes, 130, 131, 195, sex differences, 129, and species weight and taxonomy, 130–3, of various bird species, 363–73
- Baya weaver, *see Ploceus philippinus*
- bill modifications, 303, 306–8
- biotelemetry, used to estimate DEB, 199
- BIRD simulation model, 213, 214, 220, 224, 240, 264
- black-bellied tree duck, *see Dendrocygna autumnalis*
- black-capped chickadee, *see Parus atricapillus*
- black duck, *see Anas rubripes*
- blackbirds, *see Icterids*
- blue grouse, *see Dendragapus obscurus*
- blue tit, *see Parus caeruleus*
- blue-winged teal, *see Anas discors*
- bobolink, *see Dolichonyx oryzivorus*
- bobwhite diet, 192, metabolizable energy coefficient, 192
- brambling, *see Fringilla montifringilla*
- breeding colonial, 328–30, efficiency, 321–2, initiation of, 320–1, rapid, cycles, 321–2, synchronization, 321. *See also under individual species*
- brood number, and natality rates, 323–4. *See also under individual species*
- 'brood reduction', 326
- brooding young, energy cost of, 173–4
- brown-headed cowbird, *see Molothrus ater*
- brown pelican, *see Pelecanus occidentalis*
- bullfinch, *see Pyrrhula pyrrhula*
- cactus wren, *see Campylorhynchos brunneicapillus*
- cage locomotor activity, 147
- Calamospiza melanocorys* diet, 243–4, as a non-pest species, 335, 337
- Calcarus lapponicus* energy cost of courtship, 174, as a non-pest species, 335, 337
- Calidris* spp, energy cost of incubating eggs, 171
- Campylorhynchos brunneicapillus*, energy cost of brooding young, 173–4
- Cape sparrow, *see Passer melanurus*
- Cardinalis cardinalis* 'climate space' diagram for, 154, diet, 193, metabolizable energy coefficient, 193
- Carduelis chloris* communal roosts, 331, diet and feeding behavior, 312, 331
- Carduelis tristis*, as a non-pest species, 335, 337–40
- carion crow, *see Corvus corone*
- cereal crops, early cultivation, 21
- chaffinch, *see Fringilla coelebs*
- chestnut sparrow, *see Passer eminibey*
- cinnamon sparrow, *see Passer rutilans*
- Cistothorus palustris*, calorific value, eggs, 167

Index

- climatic conditions, and the initiation of breeding activity, 320–1, 324
 clutch size; environmental influences on, 322–3; and natality rates, 322–3. *See also under individual species*
 coal tit, *see Parus ater*
Coccothraustes coccothraustes: diet and feeding behavior, 306, 331–2; flocking and roosting behavior, 332; structure of bill, 306
Coccothraustes vespertinus: basal metabolism, daily rhythm, 130; bill size and structure, 306; diet, 192, 306; metabolizable energy coefficient, 192; as a non-pest species, 337; opportunistic population movements, 263
Colinus virginianus: calorific value, eggs, 167; as a non-pest species, 335, 337
 colonies, bird: agricultural habitats, 248–9; breeding, 328–30; community attributes, 249–51, 262; energy demands, 251–3, 259, 262; feeding behavior, 295, 330–2; food consumption, 253–60, 262; migrations, 86–7; in natural ecosystems, 261–3; as pests, 267–8, 295, 327–8, 342; social organization, 327–8, 342
Columba livia: commensalism with man, 48; as a pest species, 275, 335–9
Columba palumbus, as a pest species, 268–9, 271, 274
 commensalism with man, *Passer domesticus*, 4, 9, 15–18, 22–8, 47–51, 53, 205–66 *passim*, 314–15
 common grackle, *see Quiscalus quiscula*
 common redpoll, *see Acanthis flammea*
 computer simulation models for estimating: energy requirements, 10, 11, 200–3, 213–21, 264, 343; food consumption, 10, 219–20, 264, 343; potential impact, 10, 264–6, 343–4
 cooperative breeding, 327
 coot, *see Fulica americana*
Corvus corax tibetanus, size, 146
Corvus corone, as a pest species, 335, 337–40
Corvus frugilegus, energy requirement for growth, 186–7; as a pest species, 335, 337–40
Corvus monedula, as a pest species, 335, 337–40
Corvus spp.: breeding behavior, 320–2; distribution, 313; as pests, 268, 334
Coturnix coturnix: calorific value, lean dry weight, 189; chemical control of, 274
 courtship, energy cost of, 174
 cowbird, *see Molothrus ater*
 crested mynah, *see Acridotheres cristatellus*
 critical temperature (defined): lower, 134, 136–7; upper, 134
 crop, function of, 303, 307
 crossbills, *see Loxia* spp.
 crows, *see Corvus* spp.
 daily energy budget (DEB), 10, 177–83, 195–6, 342–3; adaptation to local climate, 179–81; components, 198; estimation of, 197–203; and existence metabolism, 181; and food consumed, 191–4, 343; and growth of young, 183–91; and species weight, 182–3
Delichon urbica, nests occupied by *Passer domesticus*, 74
Dendrocygna autumnalis: calorific value, eggs, 167; energy requirement, growth of young, 183–6, 189–91; metabolizable energy coefficient, 192
Dendroica kirtlandii, population decrease caused by *Molothrus ater*, 95
Dendrocopos major, egg predator, 73
Dendropagus obscurus, energy requirement for growth, 187
 desert lark, *see Ammomanes deserti*
 desert sparrow, *see Passer simplex*
 dickcissel, *see Spiza americana*
 dietary opportunism, 308–13
 digestive tract, structural modifications, 307
 distributional opportunism, 313–19
Dolichonyx oryzivorus: brood reduction, 326; ‘distant flight’ behavior, 319; as a non-pest species, 335, 337; as a pest species, 339
 domestic fowl, *see Gallus gallus*
 domestic pigeon, *see Columba livia*
 double-crested cormorant, *see Phalacrocorax auritus*
Drosophila pseudoobscura, polymorphism, 39
Dryomys nitedula, nestling predator, 78
 dunlin, *see Erolia alpina*
 eared dove, *see Zenaidura auriculata*
 ecosystem functioning, role of birds in, 207–9
Ectopistes migratorius: colonial breeding, 328; eradication, 269
 egg laying, energy cost of, 165–6
Elomys quercinus, egg predator, 73
Emberiza citrinella: egg weight loss during incubation, 112; heat conserved by fluffing plumage, 157; as a non-pest species, 335, 337
Emberiza hortulana, heat conserved by fluffing plumage, 157
 energy-conserving conditions and activities, 153–9
 energy-demanding conditions and activities, 159–77
 energy units, conversion of, 14
Eremophila alpestris, 4, 5; basal metabolism, daily rhythm, 130; bill structure and size, 306; breeding behavior, 239, 261, 263; breeding biotopes, 263; diet and feeding behavior, 239, 242, 261, 263, 306; distribution, 7, 9; energy demand, 239, 242; migration, 242; as a non-pest species, 335, 337; population (breeding) density, 239; size, 242
Eriothacus rubecula, egg weight loss during incubation, 112
Erolia alpina: calorific value, eggs, 167; energy requirement for growth, 187; metabolizable energy coefficient, 192
Erolia bairdii: calorific value, eggs, 167; temperature regulation on hatching, 189
Euplectes orix: as a pest species, 299
Euplectes spp.: breeding behavior, 334; diet and feeding behavior, 334
 European tree sparrow, *see Passer montanus*
 evening grosbeak, *see Coccothraustes vespertinus*
 evolution, avian energetics and, 194–7
 evolutionary history, *Passer* spp., 15–19, 341

Index

- existence metabolism (EM), 140–8; at 30 and 0 °C, 142–5; adaptation to local climate, 145; cage locomotor activity, 147–8; components, 147–8; and daily energy budget, 181; heat increment on feeding (SDA), 146–8; and photoperiod, 140–1; and season, 140–1; size limits, 145–6; temperature coefficients, 141–2; of various birds species, 374–9
- fat deposits, to provide energy, 164, 195
- fecundity of birds, 102–3, 305, 319, 322–4
- feeding young, energy cost of, 174
- field sparrow, *see* *Spizella pusilla*
- fire finch, *see* *Logonostica senegalae*
- flight, energy cost of, 161–4, 196
- flightless ostrich, *see* *Struthio camelus*
- flocking and roosting behavior, 104, 295, 330–2, 342. *See also under individual species*
- fluffing of plumage, to conserve heat, 157, 196
- fly larvae, and nestling mortality, 78
- food consumption: of bird colonies, 248–63; bird impact, 209–12, 232–63; computer models, 10, 219–20, 264, 343; daily energy budget and, 191–4
- fossil record, *Passer* spp., 15, 19, 50
- Fringilla coelebs*: communal roosts, 331; diet and feeding behavior, 221, 301, 309, 331; egg weight loss during incubation, 112; energy conserved by fluffing plumage, 157; energy cost of, feeding young, 174, moulting, 176, 221; existence metabolism, components, 148
- Fringilla montifringilla*: basal metabolism, seasonal changes, 130; communal roosts, 331; flock feeding, 331
- Fulica americana*: energy requirement for growth, 187–8; metabolizable energy coefficient, 192
- Galah, *see* *Kakatoe roseicapilla*
- Gallus gallus*, calorific value, eggs, 167
- genetic variation, *Passer domesticus*, 25, 34, 39–42; evolutionary potential, 42, 44–7; geographic, gene frequencies, 41–2; relationship with phenetic differentiation, 44–5
- glossy starling, *see* *Lamprotornis* spp.
- goatsucker, energy conserved by hypothermy, 159
- goldcrest, *see* *Regulus regulus*
- golden sparrow, *see* *Passer luteus*
- grackle, common, *see* *Quiscalus quiscula*
- gray sparrow, *see* *Passer griseus*
- great tit, *see* *Parus major*
- greenfinch, *see* *Carduelis chloris*
- growth of young: energy requirement, 183–9; rapid, for breeding efficiency, 321–2; respiration-production ratios, 189–91
- hawfinch, *see* *Coccothraustes coccothraustes*
- heat increment on feeding (SDA) (defined), 146–7, 196
- heavy isotopes, used to estimate DEB, 199
- herring gull, *see* *Larus argentatus*
- Hirundo rustica*, nests occupied by *Passer domesticus*, 74
- horned lark, *see* *Eremophila alpestris*
- house sparrow, *see* *Passer domesticus*
- house wren, *see* *Troglodytes aedon*
- huddling, to conserve energy, 156–7, 195–6
- hummingbird: energy conserved by hypothermy, 158; energy cost of flight, 162–3
- hyperphagia, fat deposition due to, 164
- hyperthermy, to relieve stress at high temperatures, 158–9
- hypothermy, and energy conservation, 158–9, 195
- icterids: mating systems, 334; mortality rates, 100; as pests, 267–300 *passim*; population (breeding) densities, 101–5; social structure, 86–7, 103–4, 248. *See also individual species*
- incubating eggs, energy cost of, 168–73
- ‘information center’, roost as, 332
- insect pests, control by birds, 209, 242–4
- insolation, heat absorbed from, 153–6, 195–6
- integrative method of estimating DEB, 197–9
- jackdaw, *see* *Corvus monedula*
- Japanese quail, *see* *Coturnix coturnix*
- jay, cooperative breeding, 327
- Junco hyemalis*, diet and feeding behavior, 208
- Jynx torquilla*, and nestling mortality, 78
- Kakatoe roseicapilla*, as a pest species, 268
- Kirtland’s warbler, *see* *Dendroica kirtlandii*
- Lagopus lagopus*: daily energy budget, 182; diet, 192; metabolizable energy coefficient, 192
- Lamprotornis* spp., as pests, 268, 284
- Lanius collurio*: energy requirement for growth, 186–7, 189; metabolizable energy coefficient, 192
- lapland longspur, *see* *Calcarius lapponicus*
- lark bunting, *see* *Calamospiza melanocorys*
- Larus argentatus*: calorific value, eggs, 167; energy cost of, incubating eggs, 169–71, insulating nests, 173
- Larus atricilla*, calorific value, eggs, 167
- latent energy (defined), 179, 195
- laughing gull, *see* *Larus atricilla*
- learning, opportunism in feeding behavior affected by, 311–13
- leg structure modifications, 307–8
- lesser scaup duck, *see* *Nyroca affinis*
- linnet, *see* *Acanthis cannabina*
- ‘local enhancement’, feeding by, 330, 332
- Logonostica senegalae*, feeding habits, 294
- long-billed marsh wren, *see* *Cistothorus palustris*
- long-tailed tit, *see* *Aegithalos caudatus*
- Loxia curvirostra*, opportunistic population movements, 263
- Loxia* spp.: bill structure, 306; breeding behavior, 315, 324; brood number, 324; diet and feeding behavior, 306, 309, 313, 315, 324; distribution, 313
- mallard, *see* *Anas platyrhynchos*
- Martes martes*, egg predator, 73
- mating systems: monogamous, 328, 333; polygamous (polygynous), 328, 332–4

Index

- Melospiza melodia*, as a non-pest species, 335, 337; metabolizable energy coefficient (MEC), 191–4; migration: energy cost of, 164–5, 195–6; of pest species, 295–6. *See also under individual species*
- Molothrus ater*, 4–5; breeding behavior, 86; breeding biotopes, 86; diet and feeding behavior, 293; distribution, 6–7, 53, 207; flocking and roosting behavior, 86–7, 104, 207, 293; migration, 86–7; nest parasitism, 86, 95, 326; as a pest species, 95, 104, 207, 268, 272, 277, 293; production values, 95–9. (clutch size) 95, 97, (hatching success) 95, 97, (fledgling and juvenile survival) 95, 97, (adult survival) 97–9, (life expectancy) 93, 97; monogamous mating systems, 328, 333; morphological adaptations to seed diet, 306–8; bill size and structure, 306–7; digestive tract modifications, 307; leg structure, 307–8; sexual dimorphism, 308
- morphological relationships, *Passer* spp., 16–18; mortality and survivorship, 324–6; factors, 324–5; of young, 325. *See also under individual species*
- moult, 65; energy cost of, 175–7, 196
- Mus musculus*, polymorphism, 39
- Mycteria americana*, energy requirement for growth, 188
- natality potential, 322–4; brood number, 323; clutch size, 322–4
- nests: energy cost of building, 167; insulation, 172–3, 196; location, 171–2, 196–7; parasitism, 86, 326
- net energy cost of moult (NEM), 175–7
- Nyctea scandiaca*, heat loss increased by wind, 159
- Nyroca affinis*, energy requirement for growth, 187–8
- olive-crowned sparrow, *see Passer flaveolus*
- opportunism: dietary, 308–13; distributional, 313–19
- optimization models: dickcissel feeding patterns, 311; quelea feeding patterns, 311
- ortolan bunting, *see Emberiza hortulana*
- Ouum-Qatafa Cave (Israel), fossils, 19
- overwintering, energy stress of, 165, 195
- parakeets, *see Aratinga* spp.
- Parus ater*: diet, 193; roosting behavior, 156; metabolizable energy coefficient, 193
- Parus atricapillus*, energy conserved by hypothermy, 158
- Parus caeruleus*: diet, 193; egg weight and clutch size, 108; metabolizable energy coefficient, 193
- Parus cinctus*, energy conserved by hypothermy, 158
- Parus major*: calorific value, eggs, 167; diet and feeding behavior, 193, 312; egg weight and clutch size, 108; energy cost of incubating eggs, 170; metabolizable energy coefficient, 193
- Parus montanus*, energy conserved by hypothermy, 158
- Parus* spp., diet and feeding behavior, 312
- passenger pigeon, *see Ectopistes migratorius*
- Passer ammodendri*: morphological relationship with other *Passer* spp., 16–18; as a non-pest species, 335, 337–9
- Passer castanopterus*: commensalism with man, 48; morphological relationship with other *Passer* spp., 16–18
- Passer domesticus*, 4; ancestors, 19, 22, 24; biomass production, 107–22, 125–6, (eggs) 107–12, (young) 112–18, (adults) 118–22; breeding behavior, 53, 61–5, (egg laying and altitude 62–3, (egg laying and latitude) 62, 64, 355; breeding biotopes, 55, 60–1, 99–100; calorific value, eggs, 167; diet and feeding behavior, 53–5, 99, 210–11, 232–8, 242–6, 248, 271, 315; dispersal, 9, 25–8, 49–50, 53, 55–6; distribution, 3, 18, 53, 99–100 (in North America) 15, 24–6, 99, 314; energy demand, 221–38; evolutionary biology, 24–5; evolutionary history, 15–19, 22, 24; flocking and roosting behavior, 53–5, 210–11, 248, 338; migration, 55; morphological relationship with other *Passer* spp., 16–18; niche partitioning, variable, 49; as a pest species, 103, 210–11, 245, 268, 271, 283, 355–8; population (breeding) density, 55, 57–8, 60–1, 83–6, 100, 102–3, 105, 345–8; production values, 65–86, (brood number) 65–6, 126, 323–4, (clutch size) 67–70, 101, 323–4, (number of eggs laid) 65–70, 83, 323–4, 361, (egg loss) 70–4, 83, 100–1, 325, 358–9, 361, (nestling mortality) 74–8, 83, 85, 100–1, 325, 358–9, 361, (fledgling and juvenile survival) 78–9, 83, 85, 101, 325, 361, (adult survival) 79–82, 85, (life expectancy) 82, 84; sex ratio, 90; sexual dimorphism, 32–4, 37, 308; *see also* commensalism with man, genetic variation, phenetic variations
- Passer domesticus bactrianus*, 18; breeding behavior, 54; clutch size, 67; diet and feeding behavior, 54; egg weight, 110; flocking and roosting behavior, 54; migration, 54, 57; as a non-pest species, 335, 337–9; poisoning program against, 104; population (breeding) density, 57; weight change, 153
- Passer domesticus griseogularis*, egg weight, 110, 112
- Passer domesticus hyrcanus*, egg weight, 110, 112
- Passer domesticus indicus*, egg weight, 107, 110, 112
- Passer eminibey*, morphological relationship with other *Passer* spp., 16–18
- Passer flaveolus*, morphological relationship with other *Passer* spp., 16–18
- Passer griseus*: commensalism with man, 48; feeding behavior, 293–4; morphological relationship with other *Passer* spp., 16–18
- Passer hispaniolensis*, 4; breeding behavior, 239, 321, 328; breeding biotopes, 263; commensalism with man, 48; diet and feeding behavior, 210–11, 239, 240, 263, 267; dispersal, 9, 23; distribution, 9, 18, 23, 53; energy demand, 239, 240; evolutionary history, 16–19; flocking and roosting behavior, 60, 210–11, 267, 328, 338; migration, 9, 22; morphological relationship with other *Passer* spp., 16–18; as a pest species, 104, 210–11, 267–9, 335, 337–9; population (breeding)

Index

- density, 60, 239; production values, (brood number) 323, (egg loss) 70, 72, 325, (fledgling survival) 325; size, 48
- Passer iagoensis*: commensalism with man, 48; morphological relationship with other *Passer* spp., 16–18
- Passer luteus*: diet and feeding behavior, 294; commensalism with man, 48; morphological relationship with other *Passer* spp., 16–18; as a pest species, 287
- Passer melanurus*: commensalism with man, 48; egg loss, 70; morphological relationship with other *passer* spp., 16–18
- Passer moabiticus*, morphological relationship with other *Passer* spp., 16–18
- Passer montanus*, 4; biomass production, 107–20, 125–6, (eggs) 107–8, 111, (young) 112–18, (adults) 118–22; breeding behavior, 53–5, 61–5, 324, 328, (egg laying and altitude) 62–3, (egg laying and latitude) 62, 64, 356–7; breeding biotopes, 55, 60–1, 99–100; calorific value, eggs, 167; commensalism with man, 48; diet and feeding behavior, 53–5, 104, 210–11, 232–8, 244–5, 248; dispersal, 9, 55–6; distribution, 4, 9, 53, 99–100; energy demand, 221–38; evolutionary history, 16–18; flocking and roosting behavior, 54–5, 104, 210–11, 338; migration, 55; morphological relationship with other *Passer* spp., 16–18; as a pest species, 103, 210–11, 268–9, 283, 335–9; population (breeding) density, 55, 57, 59–61, 83–5, 99–100, 102–3, 349–54; production values, (brood number) 65–6, 126, 323–4, (clutch size) 67, 70, 323–4, (number of eggs laid) 65–7, 70, 83, 323–4, 362, (egg loss) 70–4, 100, 360, 362, (nestling mortality) 74–8, 83, 100, 325, 360, 362, (fledgling and juvenile survival) 79, 83, 85, 101, 325, 362, (adult survival) 80–3, 85, (life expectancy) 82, 84, 325; sex ratio, 90; size 48
- Passer montanus dilutus*, population (breeding) density, 60
- Passer montanus malaccensis*, egg weight, 111
- Passer montanus saturatus*: population (breeding) density, 60; egg weight, 111
- Passer montanus transcaucasicus*, population (breeding) density, 60
- Passer predomesticus*, 19
- Passer pyrrhonotus*, morphological relationship with other *Passer* spp., 16–18
- Passer rutilans*: commensalism with man, 48; morphological relationship with other *Passer* spp., 16–18
- Passer simplex*, morphological relationship with other *Passer* spp., 16–18
- Passer* spp.: commensalism with man, 16–18; diet and feeding behavior, 232–8; dispersal, 9; evolutionary history, 15–19, 341; fossil record, 19; function of crop, 307; migratory habits, 22; model simulations, population dynamics and energy demands, 221–32; morphological relationships, 16–17; as pests, 283, 334; primary zoogeographic affinity, 17; social structure, 53–5. *See also under individual species*
- Pelecanus occidentalis*, calorific value, eggs, 167
- pest management and control, 10, 102–4, 205, 267–300, 302, 340, 344; damage evasion, 279–81; population regulation, 272–6; research, 284–96, 297, 299; scaring techniques, 276–9; strategy, 281–3, 297–8; surveillance, 299–300
- pest species of birds, 10–11; characteristics, 334–40; damage, 288–9; diet and feeding behavior, 291–5, 301, 308–9, 313; distribution, 287–8; diurnal pattern of movement, 295; identification, 268–9, 271–2, 285–7; introduction of, 299–300; life history studies, 289–90; migration, 295–6. *See also under individual species*, pest management and control
- pest species of insects, control by birds, 209, 242–4
- Petronia petronia*, as a non-pest species, 335, 337–9
- Phalacrocorax auritus*: energy requirement for growth, 187; metabolizable energy coefficient, 192
- phenetic relationships, *Passer* spp. (tree diagram), 16–18
- phenetic variations, *Passer domesticus*: evolutionary potential, 42–5; interlocal, 34–9, (generalized character) 38, (plumage) 38–9, (proportions) 37–8, (size) 36–7; intralocal, 28–32; relationship with genetic differentiation, 44–5; significance to commensal granivory, 47–50
- Philetairus socius*: diet and feeding behavior, 309; distribution, 313; breeding behavior, 320–2, 324, 333, (brood reduction) 326–7, (colonial) 328, (cooperative) 327; production values, (brood number) 324, (clutch size) 322–3, 325–6, (nesting success) 325–6; roosting and huddling behavior, 156
- photoperiod: effect on existence metabolism, 140–1; and the initiation of breeding activity, 315, 320
- Ploceus cucullatus*: colonial breeding, 328; distribution, 313; energy cost of incubating eggs, 170–1; as a pest species, 268, 287, 300
- Ploceus nigerrimus*, distribution, 313
- Ploceus philippinus*, as a pest species, 268
- Ploceus* spp.: ancestral adaptations, 301; diet and feeding behavior, 301, 333–4; mating systems, 301, 333–4
- plumage: color, and absorption of solar energy, 153, 155, 196; fluffing of, 157–8, 196; geographic variation in, 38–9; moulting and renewal, energy cost of, 175–7, 195–6; winter, 157, 195
- Poephila guttata*: calorific value, eggs, 167; energy cost of, incubating eggs, 168, nest building, 167; heat absorbed by radiation, 155
- polygamous (polygynous) mating systems, 328, 332–4
- productive energy (PE), 149–53; maximum potential metabolism, 149–50; and weight change, 151–3; and working capacity, 150–1
- proportions, geographic variation in, 37–8
- Prunella modularis*, egg weight loss during incubation, 112
- Psittacula krameri*: as a pest species, 267–8, 284, 287, 299; flocking behavior, 267

Cambridge University Press

978-1-107-40383-3 - Granivorous Birds in Ecosystems: Their Evolution, Populations, Energetics, Adaptations, Impact and Control

Edited by Jan Pinowski and S. Charles Kendeigh

Index

[More information](#)

Index

- Pycnonotus jocosus*, as a pest species, 299
Pyrrhula pyrrhula: diet and feeding behavior, 301, 307, 331–2; digestive tract modification, 307; incipient pesthood, 301; roosting behavior, 331–2
- quail, see *Coturnix coturnix*
Quelea erythrops, as a pest species, 268, 284
Quelea quelea: bill size, 307, 308; breeding behavior, 7, 207, 241, 309–10, 316–17, 320–4, 328, 333–4, (brood reduction) 326; crop, 307; diet and feeding behavior, 239, 241, 294, 301, 307, 309–11, 315–16, 320, 330, 333–4; distribution, 6–7, 9, 241, 315–16; flocking and roosting behavior, 7, 104, 241, 267, 328, 330–1; migration, 7, 9, 316–17; as a pest species, 241, 267–94 *passim*, 335, 337–9; population (breeding) density, 239, 241; production values, (brood number) 324, (clutch size) 323, (fledgling survival) 325; sex ratio, 334; sexual dimorphism, 308
Quelea spp., distribution, 6–7
Quiscalus quiscula, 4–5; biomass production, (eggs) 125–6, (young) 125–6, (adults) 125–6; breeding behavior, 86, 94; breeding biotopes, 86, 94, 207; diet and feeding behavior, 211, 277; distribution, 6, 53, 207; flocking and roosting behavior, 86–7, 94, 104, 107, 293; migration, 86–7; as a pest species, 104, 268, 272, 277, 283; production values, (brood number) 125–6, (hatching success) 94–5, 125, (clutch size) 94–5, (fledgling and juvenile survival) 94–5, 125, (adult survival) 94–5, 96, (life expectancy) 93–5
- radiation, heat loss by, 153
radioisotopes, used to estimate DEB, 199
rainy season, and the initiation of breeding activity, 320–1, 324
Rattus norvegicus, nestling predator, 78
raven, see *Corvus corax tibetanus*
red-backed shrike, see *Lanius collurio*
red-billed quelea, see *Quelea quelea*
red crossbill, see *Loxia curvirostra*
red-headed quelea, see *Quelea erythrops*
red-whiskered bulbul, see *Pycnonotus jocosus*
red-winged blackbird, see *Agelaius phoeniceus*
redpolls, see *Acanthis* spp.
Regulus regulus, heat conserved by fluffing plumage, 158
renewal of plumage, energy cost of, 175–7, 195–6
ring dove, see *Streptopelia decaocto*
Riparia riparia, energy conserved by hypothermy, 159
rock ptarmigan: diet, 192; metabolizable energy coefficient, 192
‘roller feeding’, 331
rook, see *Corvus frugilegus*
roosting and flocking behavior, 104, 156, 295, 330–2, 342. See also under individual species
roosting site, energy conserved by choice of, 156, 195–6
rose-ringed parakeet, see *Psittacula krameri*
rufous sparrow, see *Passer iagoensis*
running, energy cost of, 164
- salmonellosis: and egg loss, 73; and nestling mortality, 78
sand martin, see *Riparia riparia*
Saxual sparrow, see *Passer ammodendri*
Scardafella inca, incipient pesthood, 339
Sciurus vulgaris, egg predator, 73
scrub sparrow, see *Passer moabiticus*
seeds: characteristics of, 303–6; consumption during breeding season, 303, 309; in the diet of birds, 302–6
sexual dimorphism, 32–4, 37, 308
Siberian tit, see *Parus cinctus*
Sind jungle sparrow, see *Passer pyrrhonotus*
singing, energy cost of, 174
size (weight), 10; adaptation and modification, 48–9; DEB related to, 182–3; intralocality variations, 34–7; intralocality variations, 28–32; limits, energy control of, 145–6, 197
snakes, as egg predators, 326
snowy owl, see *Nyctea scandiaca*
sociable weaver, see *Philetairus socius*
Somali sparrow, see *Passer castanopterus*
song sparrow, see *Zonotrichia melodia*
Spanish (willow) sparrow, see *Passer hispaniolensis*
specific values, nestling sparrows, 117–18, 125
Spiza americana, 4–5; breeding behavior, 239, 318, 324, 328, 334; breeding biotopes, 241, 318; diet and feeding behavior, 239, 241, 308, 311, 318, 332; ‘distant flight’ behavior, 318–19; distribution, 6, 8, 318; energy demand, 239, 311; flocking and roosting behavior, 241, 328, 332; migration, 6, 241, 311, 318; as a pest species, 268, 335–40; population (breeding) density, 239, 241, 318; production values, (brood number) 324, (nesting mortality) 326; sex ratio, 308, 334; sexual dimorphism, 308
Spizella arborea: cage locomotor activity, 148; diet, 243–4; migration, 195
Spizella pusilla, migration, 195
standard metabolism (SM), 134–9; at 0 °C, 137–8; adaptation to local climate, 138–9, 195; application, 139; lower critical temperature, 135, 136–7, 195; taxonomic differences, 139; temperature coefficients, 134–6; of various bird species, 363–73
starling, see *Sturnus vulgaris*
Streptopelia decaocto: commensalism with man, 48, 49; energy conserved by fluffing plumage, 158; energy cost of feeding young, 174
Streptopelia spp., incipient pesthood, 339
Struthio camelus, size, 146
Sturnus vulgaris: breeding biotopes, 207; diet and feeding behavior, 207, 330; distribution, 207; energy requirement for growth, 186–7; flocking and roosting behavior, 248, 293, 330; metabolizable energy coefficient, 192; pest status, 104, 268, 272, 274, 277, 283, 285, 339
sunbathing, heat absorbed by, 153–6, 195–6
survivorship and mortality, 324–6. See also under individual species
swallow, energy cost of flight, 162, 196

Cambridge University Press

978-1-107-40383-3 - Granivorous Birds in Ecosystems: Their Evolution, Populations, Energetics, Adaptations, Impact and Control

Edited by Jan Pinowski and S. Charles Kendeigh

Index

[More information](#)***Index***

- swift: energy conserved by hypothermy, 159; energy cost of flight, 162, 196
 swimming, energy cost of, 164
Sylvia communis, egg weight loss during incubation, 112
- temperature regulation, 134–9, 195; various bird species, 363–73
 territorial defense, energy cost of, 174
 time–activity–energy budgets, 200
 total metabolism, *see* daily energy budget (DEB)
 tree sparrow, *see* *Passer montanus*
 tricolored blackbird, *see* *Agelaius tricolor*
Troglodytes aedon: calorific value, eggs, 167; egg weight and order laid, 108; energy cost of incubating eggs, 169–71; insulation of nests, 173
 tuberculosis, and nestling mortality, 78
Turdus merula, nestling predator, 78
Tympanuchus cupido: as a non-pest species, 335, 337
- Uraeginthus bengalis*, feeding behavior, 293–4
- Vidua* spp., feeding behavior, 294
- white-crowned sparrow, *see* *Zonotrichia leucophrys*
- white-tailed ptarmigan: diet, 192; metabolizable energy coefficient, 192
 wild cereals, 20–2; distribution, 20
 willow ptarmigan, *see* *Lagopus lagopus*
 willow (Spanish) sparrow, *see* *Passer hispaniolensis*
 willow tit, *see* *Parus montanus*
 wood pigeon, *see* *Columba palumbus*
 wood stork, *see* *Mycteria americana*
- Xanthocephalus xanthocephalus*: mating systems, 334; recorded distress call, 277
- yellow-headed blackbird, *see* *Xanthocephalus xanthocephalus*
 yellowhammer, *see* *Emberiza citrinella*
 young (nestlings, fledglings and juveniles): energy cost of, brooding, 173–4, feeding, 174. *See also under individual species*, growth of young
- zebra finch, *see* *Poephila guttata*
- Zenaidura auriculata*, as a pest species, 268
- Zonotrichia leucophrys*, heat absorbed from insulation, 155
- Zonotrichia melodia*: diet, 193; metabolizable energy coefficient, 193