

Cambridge University Press
978-1-107-43255-0 — Noise Sensitivity of Boolean Functions and Percolation
Christophe Garban, Jeffrey E. Steif
Frontmatter
[More Information](#)

Noise Sensitivity of Boolean Functions and Percolation

This is a graduate-level introduction to the theory of Boolean functions, an exciting area lying on the border of probability theory, discrete mathematics, analysis, and theoretical computer science. Certain functions are highly sensitive to noise; this can be seen via Fourier analysis on the hypercube. The key model analyzed in depth is critical percolation on the hexagonal lattice. For this model, the critical exponents, previously determined using the now-famous Schramm–Loewner evolution, appear here in the study of sensitivity behavior. Even for this relatively simple model, beyond the Fourier-analytic setup, there are three crucially important but distinct approaches: hypercontractivity of operators, connections to randomized algorithms, and viewing the spectrum as a random Cantor set. This book assumes a basic background in probability theory and integration theory. Each chapter ends with exercises, some straightforward, some challenging.

CHRISTOPHE GARBAN is a professor of mathematics at University of Lyon 1, France.

JEFFREY E. STEIF is a professor of mathematical sciences at Chalmers University of Technology in Gothenburg, Sweden.

INSTITUTE OF MATHEMATICAL STATISTICS
TEXTBOOKS

Editorial Board

D. R. Cox (University of Oxford)
B. Hambly (University of Oxford)
S. Holmes (Stanford University)
X.-L. Meng (Harvard University)

IMS Textbooks give introductory accounts of topics of current concern suitable for advanced courses at master's level, for doctoral students, and for individual study. They are typically shorter than a fully developed textbook, often arising from material created for a topical course. Lengths of 100–290 pages are envisaged. The books typically contain exercises.

Other Books in the Series

1. *Probability on Graphs*, by Geoffrey Grimmett
2. *Stochastic Networks*, by Frank Kelly and Elena Yudovina
3. *Bayesian Filtering and Smoothing*, by Simo Särkkä
4. *The Surprising Mathematics of Longest Increasing Subsequences*, by Dan Romik

Cambridge University Press
978-1-107-43255-0 — Noise Sensitivity of Boolean Functions and Percolation
Christophe Garban , Jeffrey E. Steif
Frontmatter
[More Information](#)

Noise Sensitivity of Boolean Functions and Percolation

CHRISTOPHE GARBAN

ICJ, Université Lyon 1, Lyon

JEFFREY E. STEIF

Chalmers University of Technology, Gothenburg



CAMBRIDGE
UNIVERSITY PRESS

Cambridge University Press
 978-1-107-43255-0 — Noise Sensitivity of Boolean Functions and Percolation
 Christophe Garban, Jeffrey E. Steif
 Frontmatter
[More Information](#)

CAMBRIDGE UNIVERSITY PRESS

University Printing House, Cambridge CB2 8BS, United Kingdom
 One Liberty Plaza, 20th Floor, New York, NY 10006, USA
 477 Williamstown Road, Port Melbourne, VIC 3207, Australia
 314-321, 3rd Floor, Plot 3, Splendor Forum, Jasola District Centre, New Delhi - 110025, India
 103 Penang Road, #05-06/07, Visioncrest Commercial, Singapore 238467

Cambridge University Press is part of the University of Cambridge.

It furthers the University's mission by disseminating knowledge in the pursuit of education, learning and research at the highest international levels of excellence.

www.cambridge.org

Information on this title: www.cambridge.org/9781107432550

© Christophe Garban and Jeffrey E. Steif 2015

This publication is in copyright. Subject to statutory exception and to the provisions of relevant collective licensing agreements, no reproduction of any part may take place without the written permission of Cambridge University Press.

First published 2015

A catalogue record for this publication is available from the British Library

Library of Congress Cataloging in Publication data

Garban, Christophe, 1982–

Noise sensitivity of Boolean functions and percolation / Christophe Garban, Université Lyon 1, Lyon, Jeffrey E. Steif, Chalmers University of Technology, Gothenburg.

pages cm. – (Institute of Mathematical Statistics textbooks)

Includes bibliographical references and index.

ISBN 978-1-107-07643-3 (hardback) –

ISBN 978-1-107-43255-0 (pbk)

1. Statistical physics—Textbooks. 2. Percolation (Statistical physics)—Textbooks.
3. Algebra, Boolean—Textbooks. I. Steif, Jeffrey E. II. Title.

QC174.8.G36 2015

530.13—dc23 2014023801

ISBN 978-1-107-07643-3 Hardback

ISBN 978-1-107-43255-0 Paperback

Cambridge University Press has no responsibility for the persistence or accuracy of URLs for external or third-party internet websites referred to in this publication, and does not guarantee that any content on such websites is, or will remain, accurate or appropriate.

Cambridge University Press
978-1-107-43255-0 — Noise Sensitivity of Boolean Functions and Percolation
Christophe Garban , Jeffrey E. Steif
Frontmatter
[More Information](#)

*To Laure, Victor, Camille,
Aila, Adam, and Noah*

Contents

<i>Preface</i>	<i>page xi</i>
<i>Notations</i>	<i>xvii</i>
1 Boolean functions and key concepts	1
1.1 Boolean functions	1
1.2 Some examples	2
1.3 Pivotality and influence	3
1.4 The Kahn–Kalai–Linial theorems	4
1.5 Noise sensitivity and noise stability	6
1.6 The Benjamini–Kalai–Schramm noise sensitivity theorem	7
1.7 Percolation crossings: Our final and most important example	7
1.8 A dynamical consequence of noise sensitivity	9
1.9 Exercises	11
2 Percolation in a nutshell	14
2.1 The model	14
2.2 Russo–Seymour–Welsh	15
2.3 Phase transition	16
2.4 Conformal invariance at criticality and the Schramm–Loewner evolution (SLE)	17
2.5 Critical exponents	19
2.6 Quasi-multiplicativity	21
3 Sharp thresholds and the critical point for 2-d percolation	22
3.1 Monotone functions and the Margulis–Russo formula	22
3.2 KKL away from the uniform measure case	23
3.3 Sharp thresholds in general: The Friedgut–Kalai theorem	23
3.4 The critical point for percolation for $\mathbb{Z}^2$ and $\mathbb{T}$ is $\frac{1}{2}$	24
3.5 Further discussion	26
3.6 Exercises	26

4	Fourier analysis of Boolean functions (first facts)	29
4.1	Discrete Fourier analysis and the energy spectrum	29
4.2	Examples	31
4.3	Noise sensitivity and stability in terms of the energy spectrum	31
4.4	Link between the spectrum and influence	33
4.5	Monotone functions and their spectra	33
4.6	Exercises	35
5	Hypercontractivity and its applications	38
5.1	Heuristics of proofs	38
5.2	About hypercontractivity	39
5.3	Proof of the KKL theorems on the influences of Boolean functions	42
5.4	KKL away from the uniform measure	45
5.5	The noise sensitivity theorem	47
5.6	Proof of hypercontractivity	49
5.7	Gaussian hypercontractivity	52
5.8	Exercises	54
6	First evidence of noise sensitivity of percolation	57
6.1	Bounds on influences for crossing events in critical percolation on the triangular lattice	57
6.2	The case of $\mathbb{Z}^2$ percolation	62
6.3	Some other consequences of our study of influences	65
6.4	Quantitative noise sensitivity	67
6.5	Exercises	71
7	Anomalous fluctuations	74
7.1	The model of first-passage percolation	74
7.2	State of the art	76
7.3	The case of the torus	76
7.4	Upper bounds on fluctuations in the spirit of KKL	79
7.5	Further discussion	80
7.6	Exercises	81
8	Randomized algorithms and noise sensitivity	82
8.1	BKS and randomized algorithms	82
8.2	The revelation theorem	82
8.3	An application to noise sensitivity of percolation	86
8.4	Lower bounds on revelations	89
8.5	An application to a critical exponent	91
8.6	Does noise sensitivity imply low revelation?	92
8.7	Exercises	94

<i>Contents</i>		ix
9	The spectral sample	96
9.1	Definition of the spectral sample	96
9.2	A way to sample the spectral sample in a sub-domain	98
9.3	Nontrivial spectrum near the upper bound for percolation	100
9.4	Two more facts concerning the spectral sample in general	102
9.5	Exercises	103
10	Sharp noise sensitivity of percolation	106
10.1	State of the art and main statement	106
10.2	Overall strategy	108
10.3	Toy model: The case of fractal percolation	109
10.4	Back to the spectrum: An exposition of the proof	118
10.5	The radial case	128
10.6	Exercises	129
11	Applications to dynamical percolation	132
11.1	The model of dynamical percolation	132
11.2	What's going on in high dimensions: $\mathbb{Z}^d, d \geq 19$?	133
11.3	$d = 2$ and BKS	134
11.4	The second moment method and the spectrum	135
11.5	Proof of existence of exceptional times for the hexagonal lattice via randomized algorithms	136
11.6	Proof of existence of exceptional times via the geometric approach of the spectrum	140
11.7	Exercises	143
12	For the connoisseur	144
12.1	How noise sensitive can monotone functions be?	144
12.2	Noise sensitivity and the pivotal set	147
12.3	Influences for monotone systems and applications to phase transitions	150
12.4	The majority is stablest theorem	151
12.5	k -SAT satisfiability and sharp thresholds	153
12.6	Noise sensitivity with respect to other “noises”	156
12.7	Juntas and noise resistance	159
12.8	Social choice and Arrow's impossibility theorem	161
12.9	Non-interactive correlation distillation and Borell's reverse hypercontractivity theorem	163
12.10	Deterministic and randomized complexity for Boolean functions	165
12.11	Erdős–Rényi random graphs and strong noise sensitivity	171
12.12	Noise sensitivity and correlation with majority functions	173

Cambridge University Press
 978-1-107-43255-0 — Noise Sensitivity of Boolean Functions and Percolation
 Christophe Garban, Jeffrey E. Steif
 Frontmatter
[More Information](#)

x	<i>Contents</i>	
12.13	The BKS algorithmic noise sensitivity result	177
12.14	Black noise in the sense of Tsirelson	182
12.15	Chaos, superconcentration, and multiple valleys	186
13	Further directions and open problems	188
13.1	Randomized algorithms	188
13.2	Dynamical percolation	190
13.3	The spectral sample vs. the pivotal set	191
13.4	Noise sensitivity and exceptional times away from the independent case	191
13.5	Glauber dynamics and Ising model	193
13.6	Deterministic and randomized complexity	194
13.7	A phase transition in the k -SAT problem	194
13.8	Anomalous fluctuations	195
	<i>References</i>	197
	<i>Index</i>	203

Preface

The purpose of this book is to present a self-contained theory of Boolean functions through the prism of statistical physics. The material presented here was initially designed as a set of lecture notes for the 2010 Clay summer school, and we decided to maintain the informal style which, we hope, will make this book more reader friendly.

Before going into Chapter 1, where precise definitions and statements are given, we wish to describe in an informal manner what this book is about. Our main companion through the whole book will be what one calls a **Boolean function**. This is simply a function of the following type:¹

$$f: \{0, 1\}^n \rightarrow \{0, 1\}.$$

Traditionally, the study of Boolean functions arises more naturally in theoretical computer science and combinatorics. In fact, over the last 20 years, mainly thanks to the computer science community, a very rich structure has emerged concerning the properties of Boolean functions. The first part of this book (Chapters 1 to 5) is devoted to a description of some of the main achievements in this field. For example, a crucial result that has inspired much of the work presented here is the so-called KKL theorem (for Kahn–Kalai–Linial, 1989), which in essence says that any “reasonable” Boolean function has at least one variable that has a large influence on the outcome (namely at least $\Omega(\log n/n)$). See Theorem 1.14.

The second part of this book is devoted to the powerful use of Boolean functions in the context of statistical physics and in particular in percolation theory. It was recognized long ago that some of the striking properties that hold in great generality for Boolean functions have deep implications in statistical physics. For example, a version of the KKL theorem enables one to recover in an elegant manner the celebrated theorem of Kesten from

¹ In fact, in this book we view Boolean functions rather as functions from $\{-1, 1\}^n \rightarrow \{-1, 1\}$ because their Fourier decomposition is then simpler to write down; nevertheless this is still the same combinatorial object.

1980 that states that the critical point for percolation on $\mathbb{Z}^2$, $p_c(\mathbb{Z}^2)$, is $1/2$. More recently, Beffara and Duminil-Copin used an extension of this KKL property obtained by Graham and Grimmett to prove the conjecture that $p_c(q) = \frac{\sqrt{q}}{1+\sqrt{q}}$ for the Fortuin–Kasteleyn percolation model with parameter $q \geq 1$ in $\mathbb{Z}^2$. It is thus a remarkable fact that general principles such as the KKL property are powerful enough to capture some (not all) of the main technical difficulties that arise in understanding the phase transitions of various statistical physics models.


In the 1990s, Talagrand as well as Benjamini, Kalai, and Schramm pushed this connection between Boolean functions and statistical physics even further. In 1998, Benjamini, Kalai, and Schramm introduced the fruitful concept of **noise sensitivity** of Boolean functions. Their main motivation was to study the behavior of critical percolation, but let us briefly explain what noise sensitivity corresponds to in the more common situation of *voting schemes*. Suppose n voters have to decide between two candidates denoted by 0 and 1. They first have to agree on a voting procedure or *voting scheme*, which may be represented by a Boolean function $f: \{0, 1\}^n \rightarrow \{0, 1\}$. In France or Sweden, this Boolean function would simply be the majority function on n bits whereas in the United States, the Boolean function f would be more complicated: in fancy words, it would correspond to an iterated weighted majority function on $n \approx 10^8$ voters. The collection of all votes is a certain configuration ω in the hypercube $\{0, 1\}^n$. If the election is “close,” it is reasonable to consider $\omega = (x_1, \dots, x_n)$ as a uniform point chosen in $\{0, 1\}^n$. In other words, we assume that each voter $i \in [n]$ tosses a fair coin in $\{0, 1\}$ and votes accordingly. As such the *true* result of the election should be the output $f(\omega) = f(x_1, \dots, x_n) \in \{0, 1\}$. In reality the *actual* result of the election will rather be the output $f(\omega_\epsilon)$, where ω is an ϵ -perturbation of the configuration ω . Roughly speaking, we assume that independently for each voter $i \in [n]$, an error occurs (meaning that the value of the bit is flipped) with probability a fixed parameter $\epsilon > 0$. See Chapter 1 for precise definitions. In this language, a **noise-sensitive Boolean function** is a function for which the outputs $f(\omega)$ and $f(\omega_\epsilon)$ are almost independent of each other even with a very small level of *noise* ϵ . As an example, the *parity* function defined by $f(x_1, \dots, x_n) = 1_{\sum x_i \equiv 1 \pmod 2}$ is noise sensitive as $n \rightarrow \infty$.

As we will see, there is a very useful spectral characterization of noise sensitivity. Indeed, in the same way as a real function $g: \mathbb{R} \setminus \mathbb{Z} \rightarrow \mathbb{R}$ can be decomposed into $g(x) = \sum_{n \in \mathbb{Z}} \hat{g}(n) e^{2i\pi nx}$, one can decompose a Boolean function $f: \{0, 1\}^n \rightarrow \{0, 1\}$ into a Fourier–Walsh series $f(\omega) = \sum_{S \subseteq [n]} \hat{f}(S) \chi_S(\omega)$. (See Chapter 4 for details.) Noise-sensitive

Boolean functions are precisely the Boolean functions whose spectrum is concentrated on “high frequencies”; that is, most of their Fourier coefficients (in a certain analytic and quantitative manner) correspond to subsets $S \subseteq [n]$ with $|S| \gg 1$.

It is thus not surprising that with such a spectral characterization of noise sensitivity, a significant part of this book is devoted to various techniques that allow us to detect high-frequency behavior for Boolean functions. The techniques we introduce are essentially of three different flavors:

1. Analytical techniques based on **hypercontractive** estimates (Chapter 5)
2. A criterion based on randomized algorithms (Chapter 8)
3. A study of the “fractal” behavior of frequencies $S \subseteq [n]$ (Chapter 10)



To make the link with statistical physics, consider the following Boolean function, which is well known in computer science and in game theory because it represents the solution of the **Hex game**. In the figure above, we represent a Hex game on a 10×10 table: player A tries to go from the left boundary to the right using gray hexagon tiles, while player B tries to go from the top boundary to the bottom using black hexagon tiles. The players either take turns or, as in the **random turn hex game**, they toss a coin at each turn to decide who will move. At the end of the game, we obtain a tiling of the table as in the figure and the result of the game is then described by a certain Boolean function $f_{10}: \{A, B\}^{100} \rightarrow \{A, B\}$. Note that in the figure, player A has won. As we will see, this Boolean function (or rather the family $\{f_n\}$ defined analogously on $n \times n$ tables) is instrumental in our study of how the model of percolation responds to small random perturbations.

Boolean functions of this type are notoriously hard to study, and this book develops tools aimed at understanding such Boolean functions. In particular, we will eventually see that as $n \rightarrow \infty$, most of the Fourier transform $\hat{f}_n$ of the Hex-function f_n on an $n \times n$ table is concentrated on frequen-

cies of size $|S|$ about $n^{3/4}$. The appearance of the surprising exponent of $3/4$ corresponds to one of the **critical exponents** that are aimed at describing the fractal geometry of critical percolation. See Chapter 2.

This high-frequency behavior of the Hex-functions $\{f_n\}$ implies readily that critical planar percolation on the triangular lattice is highly sensitive to noise (in a quantitative manner given by the above $n^{3/4}$ asymptotics). This noise sensitivity of percolation has surprising consequences concerning the model of **dynamical percolation** (see Chapter 11). We will see among other things that there exist exceptional times at which an infinite *primal* cluster coexists with an infinite *dual* cluster (Theorem 11.9), which is a very counterintuitive phenomenon in percolation theory.

In Chapter 7, we give another application to statistical physics of a very different flavor: consider the random metric R on the lattice $\mathbb{Z}^d$, $d \geq 2$, where each edge is independently declared to be of length a with probability $1/2$ and of length b with probability $1/2$ (where $0 < a < b < \infty$ are fixed). Fascinating conjectures have been made about the fluctuations of the random R -ball around its deterministic convex limit. In particular, it is conjectured that these fluctuations are of magnitude $R^{1/3}$ in two dimensions and that the law describing these fluctuations is intimately related to the celebrated Tracy–Widom law that describes the fluctuations of the largest eigenvalue of large random Hermitian matrices. This book certainly does not settle this stunning conjecture but it does present the best results to date on the fluctuations of this metric using a Fourier approach.

This book is structured as follows: in Chapters 1, 3, 4, 5, 8, and 9, we introduce general tools for Boolean functions that are applicable in various settings (and are thus not restricted to the context of statistical physics). Several examples in these chapters illustrate links to other active fields of mathematics. Chapter 2 is a short introduction to the model of percolation. Chapters 6, 10, and 11 are more specifically targeted toward the analysis of the **noise sensitivity of critical percolation** as well as its consequences for dynamical percolation. Chapter 7 analyzes the fluctuations of the earlier mentioned random metrics on $\mathbb{Z}^d$, $d \geq 2$. Chapter 12 explores a large variety of interesting topics tangential to the main contents of this book. Finally Chapter 13 collects some open problems.

We assume readers have the mathematical maturity of a first-year graduate student and a reasonable background in probability theory and integration theory. Having seen some percolation would be helpful but not necessary.

Acknowledgments

This book would not be in its present shape without the help of many people whom we now wish to warmly thank. First of all, we thank Charles Newman, Vladas Sidoravicius, and Wendelin Werner for inviting us to give a course on this topic at the 2010 Clay summer school in Buzios. (This resulted in a set of lecture notes published by the Clay Institute that was the starting form of the present book.) This book would not exist without having had the opportunity to give such a course.

We also thank Diana Gillooly for her constant help at Cambridge University Press.

Many people helped us greatly with their remarks, which improved the manuscript. In this respect, we wish in particular to thank Ragnar Freij, who served as a very good teaching assistant for this course at the Clay Institute. We also thank Juhan Aru, Vincent Beffara, Itai Benjamini, Ehud Friedgut, Johan Hästad, Jeff Kahn, Gil Kalai, Jean-Christophe Mourrat, Ryan O'Donnell, Malin Palö, Yuval Peres, Gábor Pete, Jeremy Quastel, Mike Saks, Mikaël de la Salle, Yilun Shang, Zhan Shi, Arvind Singh, Vincent Tassion, and Xiaolin Zeng.

Finally, at the time of concluding the writing of this book, our thoughts go out to Oded Schramm. This book would simply not exist had it not been for Oded.

Lyon and Gothenburg, October 1, 2014

CHRISTOPHE GARBAN and JEFFREY E. STEIF

Notations

Ω_n	hypercube $\{-1, 1\}^n$
$\mathbf{I}_k(f)$	influence of the k th variable on f
$\mathbf{I}_k^p(f)$	influence of the k th variable on f at level p
$\mathbf{I}(f)$	total influence of the function f ; see Definition 1.10
$\mathbf{Inf}(f)$	influence vector of f
$\mathbf{H}(f)$	sum of the squared influences; see Definition 5.5
$\alpha_1(R)$	probability in critical percolation to have an open path from 0 to $\partial B(0, R)$
$\alpha_1(r, R)$	multiscale version of the above
$\alpha_4(R)$	probability of a <i>four-arm event</i> from 0 to $\partial B(0, R)$
$\alpha_4(r, R)$	multiscale version of the above
χ_S	character $\chi_S(x_1, \dots, x_n) := \prod_{i \in S} x_i$
$\hat{f}(S)$	Fourier coefficient $\hat{f}(S) = \langle f, \chi_S \rangle = \mathbb{E}[f \chi_S]$
$E_f(m), 1 \leq m \leq n$	<i>energy spectrum</i> of f ; see Definition 4.1
$\nabla_k f$	discrete derivative along k : $\nabla_k f(\omega) := f(\omega) - f(\sigma_k(\omega))$
$\mathcal{P} = \mathcal{P}(f)$	<i>pivotal set</i> of f ; see Definition 1.7
$\mathcal{S} = \mathcal{S}_f$	<i>spectral sample</i> of f ; see Definition 9.1
$\hat{\mathbb{Q}}_f$	<i>spectral measure</i> of f ; see Definition 9.1
$\hat{\mathbb{P}}_f$	<i>spectral probability measure</i> of f ; see Definition 9.2
$f(n) \asymp g(n)$	there exists some constant $C < \infty$ such that $C^{-1} \leq \frac{f(n)}{g(n)} \leq C, \forall n \geq 1$
$f(n) \leq O(g(n))$	there exists some constant $C < \infty$ such that $f(n) \leq Cg(n), \forall n \geq 1$
$f(n) \geq \Omega(g(n))$	there exists some constant $C > 0$ such that $f(n) \geq Cg(n), \forall n \geq 1$
$f(n) = o(g(n))$	$\lim_{n \rightarrow \infty} \frac{f(n)}{g(n)} = 0$