

Index

- accidental decannulation, 42–4
 management, 42–4
 algorithm, 43
 post-management care, 44
- ACE inhibitor-induced
 angioedema, 29, 30
- acute spontaneous urticaria, 29
- adenotonsillectomy, 184, *See also* post-tonsillectomy hemorrhage
- airway assessment, 23, *See also specific conditions*
 pregnant patient, 106
- airway exchange catheter, 129–30, 130
- airway management. *See also specific conditions*
 complications, 26
 bleeding, 26
 endobronchial intubation, 26
- failed airway, 208
 definition, 208
 management, 208
- general approach, 19–20
 getting to the event, 20–1
 obtaining patient information, 21–2
 risk assessment, 22–3
 securing the airway, 23–5
 supportive care, 26
 transportation of patient, 26–7
- airway registries, 218–19
- American Society of Anesthesiologists (ASA)
 Practice Guidelines for Management of the Difficult Airway, 10–13, 218
- difficult airway algorithm (DAA), 12
 modification for trauma setting, 14
- anesthesia. *See also* hypnotic agents; paralytic agents; sedation
 anterior mediastinal mass, 91
- cervical spine injury, 153
- foreign body aspiration, 83, 84
 pregnant patient, 107–8
 subglottic stenosis, 52
- angioedema, 28–32
 ACE inhibitor-induced, 30
 airway management, 30–2
 scenarios, 30
 stepwise approach, 31
 characteristics, 28–9
 diseases with combined urticaria and angioedema, 29–30
 examination, 31
 hereditary (HAE), 29–30
 idiopathic, 30
 mechanisms, 29
 pharmacologic treatment, 32
 post-management care, 32
- ankylosing spondylitis, 190
- anterior mediastinal mass (AMM), 89–92
 airway management strategies, 90–1
 emergency department, 90
 operating room, 91
 complications, risk assessment, 90
- antibiotic treatment, epiglottitis, 77
- apneic oxygenation, 163
- arytenoids, 6
- asthma, 94–101
 airway management, 96–100
 adjuvants, 99–100
 intubation, 98–9
 noninvasive positive pressure ventilation, 96–8
 characteristics, 94
 exacerbation severity, 95
 management algorithm, 97
 paradoxical vocal fold motion relationship, 113
 post-management care, 100–1
 risk factors, 95
 risk stratification, 94–5
- atlanto-occipital dislocation, 149
- atlas fracture, 149
- awake laryngoscopy, epiglottitis, 75–7
- axis fractures, 150
- bag-mask ventilation, 81
 congenital anomalies, 179–81
 facial trauma, 164
- Bailey maneuver, 128–9
- bleeding. *See also* post-tonsillectomy hemorrhage
 facial trauma, 158–9
 laryngoscopy complications, 26
- management algorithm, 46
- tracheostomy site, 45
 case study, 45
 management, 45–6
 post-management care, 46–7
- blunt neck injury (BNI), 135, 139–40
 airway management, 139–40
 special considerations, 141
- bougie, facial trauma management, 163
- brain herniation, 168
- brainstem compression, 166
- bronchoscopy. *See* flexible bronchoscopy; rigid bronchoscopy
- burns. *See* thermal injury
- C1-inhibitor deficiency, 29
- cavernous sinus, 3
- cervical spine anatomy, 150
- cervical spine injury, 141, 149–56
 airway management, 152–5
 anesthesia for awake intubation, 153
 asleep intubation, 154
 medications for asleep intubation, 154–5
 neck protection, 152

- rapid sequence intubation, 153
 - risks of awake intubation, 154
 - securing the airway, 152
 - diagnosis, 151
 - evaluation, 150–1
 - airway, 151
 - injury location effect on respiratory status, 151–2
 - types of, 149–50
 - cesarean section, 108
 - children
 - cricothyrotomy
 - contraindication, 51
 - emergency tracheostomy, 51
 - epiglottitis, 72
 - airway management, 74–5
 - evaluation, 73
 - foreign body aspiration, 79
 - management, 82–3
 - pediatric airway, 8
 - subglottic stenosis, 49
 - chronic obstructive pulmonary disease (COPD), 34
 - chronic spontaneous urticaria (CSU), 29
 - clinical practice guidelines, 10, 17
 - difficult airway management, 10–14
 - American Society of Anesthesiologists, 10–13, 218
 - extubation, 16–17
 - Vortex approach, 15–16
 - code bag, 20–1
 - colorectal cancer, 63
 - communication issues, 217–21
 - difficult airway letters, 220–1
 - dissemination of difficult airway information, 217–18, 220
 - confirmation of airway placement, 25
 - congenital anomalies, 172–82
 - airway management, 177–82
 - awake intubation, 181–2
 - bag-mask ventilation, 179–81
 - muscle relaxants, 182
 - supraglottic airways, 181
 - tracheal intubation, 181
 - anatomical approach, 174
 - prevalence, 174
 - congestive heart failure, 44
 - continuous positive airway pressure (CPAP)
 - morbidly obese patient, 68
 - paradoxical vocal fold motion, 116
 - conus elasticus, 49
 - crash airway algorithm, 14
 - cribriform plate, 3
 - cricoid cartilage, 5, 48
 - posterior, 49
 - cricoid pressure application, 24–5
 - cricothyroid membrane, 1, 5, 6
 - identification, 1, 5
 - cricothyroid space, 6
 - cricothyrotomy, 51, 208
 - contraindication in children, 51
 - epiglottitis, 75
 - facial trauma, 164–5
 - issues, 214–15
 - morbidly obese patient, 68
 - rapid four-step technique, 208–10
 - Seldinger technique, 213–14
 - standard technique, 210–13
 - cuff-leak test, 128
 - dashboard injury, 58
 - difficult airway letters, 220–1
 - difficult airway registries, 218–19
 - direct laryngoscopy, 20–1
 - failure of, 20
 - morbidly obese patient, 65, 67
 - dissemination of difficult airway information, 217–18, 220
 - difficult airway letters, 220–1
 - distal chip intubation scopes, 190
 - double setup, 51
 - penetrating neck trauma, 137
 - dysgnathia complex, 172
 - emergency department (ED), 21
 - endobronchial intubation, 26
 - endobronchial obstruction, 35
 - epiglottis, 6
 - epiglottitis, 71–7
 - adult airway management, 75–7
 - algorithm, 76
 - causes, 72
 - diagnosis, 73
 - fungal, 72
 - general management
 - principles, 73–4
 - manifestations, 71–3
 - pediatric, 72
 - airway management, 74–5
 - evaluation, 73
 - post-intubation
 - management, 77
 - equipment, 20–1
 - difficult airway, 191, 193
 - testing, 20
 - esophagus, 57
 - congenital anomalies, 179
 - foreign body, 49
 - ethmoid sinus, 3
 - etomidate, 21, 154
 - ex utero intrapartum treatment (EXIT), 172–4
 - examination, 23, *See also specific conditions*
 - exercise-related paradoxical vocal fold motion, 113
 - extracorporeal membrane oxygenation (ECMO)
 - asthmatic crisis, 100
 - extrathoracic tracheal stenosis, 50
 - extubation
 - asthma patients, 100–1
 - complications, 16
 - difficult airway, 126–32
 - airway exchange catheter, 129–30
 - algorithm, 132
 - Bailey maneuver, 128–9
 - guidelines, 131
 - strategies, 128–31
 - failure, 127
 - high-risk patient
 - identification, 127–8
 - risk continuum, 126
 - guidelines and algorithms, 16–17
 - morbidly obese patient, 68
 - paradoxical vocal fold
 - motion relationship, 114
 - post-tonsillectomy
 - hemorrhage, 187–8
- facial trauma, 157–65
 - airway management, 159–65
 - algorithm, 161

- facial trauma (cont.)
 awake look, 162
 bag-mask ventilation, 164
 bougie, 163
 cricothyrotomy, 164–5
 King laryngeal tube, 164
 obtaining help, 160
 positioning and opening the airway, 163
 pre-procedure checklist, 161–2
 suctioning equipment, 163
 supraglottic airway, 164
 video laryngoscopy, 164
 patterns of, 158–9
 post-management care, 165
 failed airway, 208
 definition, 208
 management approach, 208
 algorithm, 209
 fetal bradycardia, 104
 fiberoptic intubation, 190–9,
See also flexible
 bronchoscopy;
 intubation
 awake intubation, 30, 197–9
 morbidly obese patient, 68
 upper airway obstruction,
 38–9
 difficult airway equipment,
 191, 193
 handling the flexible
 intubation scope, 191
 level of consciousness, 195–6
 mechanics, 190–1
 oxygen versus suction
 application, 194
 pitfalls, 196
 route of airway access, 196
 scope diameter selection,
 192
 scope preparation, 194–5
 tracheal intubation, 196–7
 tracheal tube size selection,
 192–4
 video advantages, 191
 firearm injuries, 58
 flexible bronchoscopy, 190, *See
 also* fiberoptic
 intubation
 advantages and
 disadvantages, 82
 foreign body aspiration, 81,
 83
 airway management, 83
 children, 82–3
 pharmacologic
 management, 83
 neck injury patient, 137–8,
 140, 141
 subglottic stenosis, 52
 tracheobronchial traumatic
 injury, 60
 flexible intubation scope (FIS),
 190, *See also* fiberoptic
 intubation; flexible
 bronchoscopy
 diameter selection, 192
 handling, 191
 pitfalls, 196
 preparation, 194–5
 flumazenil, 21
 foreign body
 aspiration, 79–87
 algorithm, 86
 child, 79
 clinical manifestations,
 80–1
 complete airway
 obstruction, 85–6
 complications, 87
 epidemiology, 79–80
 management, 81–7
 partial airway obstruction,
 81–3
 pathogenesis, 80
 post-management care,
 86–7
 removal technique, 85
 esophageal, 49
 frontal sinus, 3
 glottis, 48
 stenosis, 50
 hangman's fracture, 150
 head anomalies, 175
 head injury, 166–71
 airway management, 167–71
 algorithms and pathways,
 169–71
 indications for early
 intubation, 169
 precautions, 168
 scenarios, 167–9
 pathophysiology, 166–7
 post-management care, 171
 heliox
 anterior mediastinal mass, 90
 asthmatic crisis, 50
 difficult airway extubation,
 131
 paradoxical vocal fold
 motion, 116
 subglottic stenosis, 50
 upper airway obstruction, 38
 hematoma
 postoperative, 119–24
 airway management,
 121–2
 management pathways,
 122–4
 post-management care,
 124
 subdural, 166
 hereditary angioedema (HAE)
 with C1-inhibitor deficiency,
 29
 with normal C1-inhibitor,
 29–30
 history taking, 22–3
 hypnotic agents, 25
 morbidly obese patient, 67
 side effects, 26
 hypopharynx, 4–5
 hypoxia prevention, 24, *See also*
 preoxygenation
 infectious risk, 22
 innominate artery, 57
 bleed, 45, *See also*
 tracheoinnominate
 artery fistula
 institutional algorithms, 14–15
 American Society of
 Anesthesiologists, 14
 intensive care unit (ICU), 21
 intracranial hypertension, 166,
 169–70
 signs and symptoms, 167
 intraosseous techniques, 21
 intravenous access, 21
 intubation. *See also* fiberoptic
 intubation; rapid
 sequence intubation
 (RSI)
 angioedema, 31–2
 asthmatic crisis, 98–9
 awake intubation, 11
 anterior mediastinal mass,
 91
 facial trauma, 159–60
 postoperative hematoma,
 121–2, 123–4
 risks with cervical spine
 injury, 154
 blunt neck trauma, 140
 cervical spine injury, 152–5

- anesthesia, 153
 asleep intubation, 154
 medications for asleep
 intubation, 154–5
 neck protection, 152
 risks of awake intubation,
 154
 congenital anomalies, 181–2
 Cormack and Lehane
 classification, 105, 105
 epiglottitis, 74, 77
 morbidly obese patient, 65
 paradoxical vocal fold
 motion relationship,
 113–14
 penetrating neck trauma,
 135–9
 special considerations, 139
 post-tonsillectomy
 hemorrhage, 186–7
 pregnant patient, 104–5
 failed intubation
 management, 108–10
 smoke inhalation, 145
 upper airway obstruction, 38
- Johns Hopkins difficult airway
 bracelet, 219
- ketamine
 asthmatic crisis, 98, 99–100
 cervical spine injury, 154
 paradoxical vocal fold
 motion relationship,
 114
 smoke inhalation patient,
 145
 subglottic stenosis, 52
 King laryngeal tube, 164
 Kiesselbach's plexus, 2
- laryngeal mask airway (LMA)
 cervical spine injury, 154
 facial trauma, 164
 pregnant patient, 109
 laryngopharyngeal reflux, 114
 laryngospasm, 112–17, *See also*
 paradoxical vocal fold
 motion (PVFM)
 definition, 112–13
 larynx, 5–8, 6, 7
 congenital anomalies, 178
 exposure, 24
 injury in facial trauma, 159
 innervation, 7
 musculature, 6, 7
- lidocaine, 52
 lymphoma, 92
- Mallampati class, 23
 pregnant patient, 106
 mandibular hypoplasia, 177
 manual in-line stabilization
 (MILS), 152
 effect on view during
 laryngoscopy, 152
 mask ventilation, morbidly
 obese patient, 65
 maternal deaths, 104
 maxillary sinus, 3
 MedicAlert National Difficult
 Airway/Intubation
 Registry, 218–19
 micrognathia, 172
 morbid obesity, 63–8
 airway management, 65–8
 direct laryngoscopy and
 intubation, 65
 management plans, 66–8
 mask ventilation, 65
 patient positioning, 65–6
 supraglottic airway device,
 65
 surgical airway, 68
 anatomic and physiologic
 changes, 64
 difficult airway prediction, 65
 obstructive sleep apnea and,
 64
 postoperative management,
 68
 positive airway pressure,
 68
 safe extubation, 68
 mouth, 3, 3–4
 muscle relaxants, 182
- naloxone, 21
 nasal cavity, 1
 nasal intubation, 3
 nasal malformations, 175
 nasal septum, 1
 nasopharynx, 2, 4
 neck anomalies, 177
 neck trauma, 134–41
 airway management
 algorithm, 136
 blunt neck injury, 139–41
 penetrating neck injury,
 135–9
 negative pressure pulmonary
 edema (NPPE), 40
- neostigmine, 21
 neurogenic shock, 155
 noninvasive positive pressure
 ventilation (NPPV)
 asthmatic crisis
 management, 96–8
 nose, 1–3
 blood supply, 1–2
 innervation, 2–3
- obesity. *See* morbid obesity
 obstetric patient. *See* pregnant
 patient
 obstructive sleep apnea (OSA),
 64
 omalizumab, 29
 oral cavity, 3–4
 oropharynx, 3, 4
- paradoxical vocal fold motion
 (PVFM), 112–13, 117
 anatomic findings, 113
 causes, 113–14
 asthma, 113
 exercise, 113
 extubation, 114
 intubation, 113–14
 irritants, 114
 ketamine, 114
 laryngopharyngeal reflux,
 114
 psychosocial disorders,
 114
 clinical presentation, 114–15
 evaluation and diagnosis,
 115
 treatment, 116
 paralytic agents, 24, 25
 upper airway obstruction
 and, 37
 paranasal sinuses, 3
 pediatric airway, 8, 8
 pediatric conditions. *See*
 children
 penetrating neck injury (PNI),
 135
 airway management, 135–9
 special considerations,
 139
 penetration syndrome, 80
 pharynx, 4–5
 positive airway pressure (PAP)
 morbidly obese patient, 68
 positive end-expiratory
 pressure (PEEP)
 penetrating neck trauma, 137

- post-tonsillectomy
 hemorrhage, 184–8
 airway management, 185–8
 classification, 184
 emergency treatment, 185
 frequency, 184
- pre-eclampsia, 104
- pregnant patient, 104–10
 airway assessment, 106
 airway changes, 105–6
 difficult airway management,
 107
 algorithm, 107
 failed intubation
 management, 108–10
 proposed mode of delivery,
 107–8
 cesarean section, 108
 vaginal delivery, 107
- preoxygenation, 24, 66
 blunt neck injury, 140
 facial trauma, 163
 penetrating neck wound, 137
- propofol, 21, 67
 asthmatic crisis, 98
- rapid sequence intubation
 (RSI), 24–5
 cervical spine trauma, 153
 epiglottitis, 75
 facial trauma, 159, 162
 head trauma, 169
 modified technique, 169
 morbidly obese patient, 66–7
 post-tonsillectomy
 hemorrhage, 186
- recurrent laryngeal nerve
 (RLN), 7, 57
- reintubation, 126
 failure, 127
- rigid bronchoscopy
 foreign body aspiration, 81
 airway management, 84
 children, 82, 84
 pharmacologic
 management, 84
 subglottic stenosis, 53
- risk assessment, 22–3
 infectious risk, 22
- rocuronium, 21, 25
 asthmatic crisis, 98
- sedation
 foreign body aspiration, 83
 morbidly obese patient, 67
 subglottic stenosis, 52
- upper airway obstruction
 and, 37
- Seldinger technique, 213–14
- sevoflurane, 84, 100
- smashed face, 158, 158, *See also*
 facial trauma
- smoke inhalation, 143–7
 airway management, 145–7
 algorithm, 146
 diagnosis, 144
 pathophysiology, 143
 post-intubation care, 147
 symptoms, 144
- sphenoid sinus, 3
- spinal cord injury, 149, *See also*
 cervical spine injury
- spinal shock, 155
 induction agents, 155
- stabbing injuries, 58, 134, *See*
 also penetrating neck
 injury (PNI)
- STOP Questionnaire, 64
- STOP-BANG Questionnaire,
 64
- stridor, 34–5, 112
 biphasic, 50
 causes, 36, 49
 children, 80
 general management, 50–1
 paradoxical vocal fold
 motion, 115
 potential difficult airway
 characteristics, 37
- subdural hematoma, 166
- subglottic stenosis (SGS),
 48–53
 anesthesia and sedation, 52
 causes, 49
 clinical anatomy, 48–9
 management, 50–1, 52
 rigid bronchoscopy, 53
 manifestations, 49–50
- succinylcholine, 21, 25
 asthmatic crisis, 98
 cervical spine injury, 155
 head injury, 169
 morbidly obese patient, 67
 paradoxical vocal fold
 motion, 116
 side effects, 99
- superior vena cava (SVC)
 compression, 89
- supportive care, 26
- supraglottic airway device
 morbidly obese patient, 65,
 67–8
- post-tonsillectomy
 hemorrhage, 187
 subglottic stenosis, 51
- supraglottic airway
 obstruction, 35
- supraglottitis, 71
- surgical airway, 207–15, *See*
 also cricothyrotomy;
 tracheostomy
 issues, 214–15
 open techniques, 208–13
 rapid four-step technique,
 208–10
 standard technique,
 210–13
 percutaneous techniques,
 213–14
- SVC syndrome, 90
- systemic inflammatory
 response syndrome, 63
- thermal injury, 143, *See also*
 smoke inhalation
 diagnosis, 144
- thoracic anomalies, 180
- thyrohyoid membrane, 5
- thyroid carcinoma, 119
- thyroid cartilage, 5
- thyroid gland, 57
- tongue, 28
 swelling, 28
 facial trauma, 158
- tonsillectomy, 184, *See also*
 post-tonsillectomy
 hemorrhage
- trachea, 57
 congenital anomalies, 179
 obstruction, 35
- tracheal intubation, 196–7
- tracheobronchial anatomy,
 57–8
- tracheobronchial traumatic
 injury (TTI), 56–61
 airway management, 59–61
 algorithm, 60
 diagnosis, 59
 incidence, 57
 pathophysiology, 58
 blunt trauma, 58
 iatrogenic injury, 58
 penetrating trauma, 58
 post-intubation care, 61
 tracheoinnominate artery
 fistula, 45
 management, 45–6
 post-management care, 46–7

tracheostomy. <i>See also</i> surgical airway	avoidance of sedation and paralysis, 37	conventional or proprietary stylet, 203
children, 51	clinical data significance, 36	currently available videolaryngoscopes, 201
subglottic stenosis, 51	examination, 38	facial trauma, 164
tracheostomy complications, 42–7	intubation, 38	guiding channel, 202
accidental decannulation, 42–4	post-management care, 40	integrated or detached display, 202
airway management, 42–4	surgical management, 37	morbidly obese patient, 66, 67
post-management care, 44	characteristics, 34–5	neck injury patient, 136, 141
bleeding from site, 45	level of obstruction, 35	penetrating neck trauma, 136
case study, 45	potential difficult airway characteristics, 37	outcomes, 203–4
management, 45–6	urticaria, 28–9	pole-mounted or portable, 203
post-management care, 46–7	acute spontaneous, 29	single-use vs. reusable, 202–3
transportation of patient, 26–7	chronic spontaneous (CSU), 29	use of video laryngoscope, 204
trismus, 159	diseases with combined urticaria and angioedema, 29–30	vocal cords, 6, <i>See also</i> paradoxical vocal fold motion (PVFM)
turbinates, 1	vagus nerve, 7	Vortex approach, 15–16
ultrasonography, tongue swelling evaluation, 32	video laryngoscopy, 20–1, 200–5	wheezing, differential diagnosis, 96
upper airway obstruction (UAO), 34–40	advantages, 200–1	
airway management, 36–40	blade geometry, 201	
adjuncts and rescues, 38–40	capital vs. operational budget pricing, 203	
algorithm, 39		