

INDEX OF SUBJECTS

Abbreviations, T.=tuberculosis, t.b.=tubercle bacilli, f = footnote

Abdominal tuberculosis, types of t.b. found (man) 580, table 583; abdominal origin of pulmonary tuberculosis 141; summary 176; T. of mesenteric glands in cases of phthisis 174

Abortive infections 21, 65, 69, 669 Acid-fast group of bacilli 198; acid-fast bacilli in milk 199; in tuberculous lesions in swine 404; in Johne's disease 388; acid-fast streptothrices 199 Acid-fastness, degrees of 199; may be dependent on composition of the culture medium 199

Adaptation of t.b. to conditions of artificial culture 203, 226, 286; (on broth)

209, 236; conclusion 242

Age when infection occurs (in man) 131; (in the ox) 381

Air passages, freedom of entry of air-borne bacteria into 148
Antelope, instances of T. in 482
Anthracosis, a chronic process 148; portal of entry of the carbon: natural

anthracosis in the guinea-pig 147; experimental 145, 149
Antiformin, use of in raising cultures of t.b. from sputum, etc. 218, 684
Argentine Republic, frequency of swine T. 396 f, 398
Arkansas, Oklahoma and Texas, rarity of tuberculosis in swine allowed to run

wild in the forests 396

Aspiration of t.b. into the lung, a source of error in feeding experiments 160,

162, 165, 327, 500, (in apes) 517 f
Ass, rarity of T. in 447; experimental infection 478; is it more susceptible to infection with equine bacilli? 481 f

Assortative mating 82 Attenuation of t.b.—see "Stability of type"

Attypical t.b. 214, 301 f, 354, 356, 369; from the pig 362, 406, 681; from the horse 364, 406, 470; from man, phthisis 212 f, 681; bone and joints 604, 629, 681; lupus 617, 620, 627, 635, 636, 649

Avian tubercle bacilli—see under "B. tuberculosis"

B. prodigiosus, inhalation expts. 148; feeding expts. 163
B. tuberculosis. Discovery 57; relation to other acid-fast bacteria, and to the diphtheria and streptothrix groups 198
Morphological characters 198, 224; branching 199; comparative length of human and bovine t.b. under various conditions 201; short forms sometimes in sputum 224

Cultural characters 202, 204, 209, 225; pigmentation 205, 681, 682, 685; diagnostic value of 231; relative importance of cultural characters and virulence 636, 683

Virulence for various animal species 211 et seq., table 248
Classification 196, 299, 663, 683; atypical strains—see under "Atypical";
a new sub-type 680; difference of virulence of human and bovine
types no mere difference of degree 212, 619 f, 684 Methods of staining 199; of cultivating 218, 685



688

INDEX OF SUBJECTS

Modification 286, 317, 349—see also "Stability of type"; adaptation to the conditions of artificial cultivation 226; of bovine bacilli to glycerin broth 209, 222, 236

Numbers. Comparative numbers of t.b. in the lesions of various species of animals 464, 535; relation of numbers to susceptibility of the species 464 f; presence in normal tissues 72, 597, 605; visible but not viable in old lesions 71, 660; dissemination after subcutaneous

injection 253, 672 The Human Type. hipection 253, 672

he Human Type. Virulence for calf 253; results of exceptional severity 254, 258; chronic lesions in eyes and joints 259; minute non-progressive lesions common 253; death after intravenous injection 257; influence of dose in intravenous injections, table 265; effect of enormous doses injected subcutaneously 250, 256; virulence for the rabbit 276; death after intravenous injection 276; instances of progressive tuberculosis 272, 280; exceptional results 250; virulence

of progressive tuberculosis 277, 280; exceptional results 279; virulence for the pig 403, 408, table 405

The Bovine Type. Virulence of strains from the ox itself compared with that of those from—man 353, 358; pig 362; horse 364; proportion of deaths caused by bovine t.b. in various kinds of human T. 658; are bovine bacilli less virulent for man than human t.b.? 659; lesions produced in calf 252, in rabbit 273; doses employed for differ-

entiation of type 269, 283

The Avian Type. Cultural characters 209; variability 210; approxi-The Avian Type. Cultural characters 200; variability 210; approximation to those of human type 210, 205 f; pellicle formation on broth 211; ease of emulsification 230. Found in: various birds 537; localized lesions of the pig 403, man 646, 665, 666; ox 388; mice 451, 459; rats 443, 449; rabbits 427, 428; no relation to Johne's disease 388; or to lupus 646. Duration of life in culture 221 and in generative susceptibility to T. of different kinds of birds 526; T. rare in geese and ducks 521; and in wild birds 527; instances recorded in lapwing 527, sparrow, kestrel, partridge 528; unknown in grouse 528; why common among domesticated birds 529

Lesions of tuberculosis. General character and distribution 520: lesions

Lesions of tuberculosis. General character and distribution 529; lesions

Lesions of tuberculosis. General character and distribution 529; lesions in—lungs and kidneys 529, 531; intestines 532; bones and joints 534; skin 532, 535, 656; certain glands of the neck 534

Lymphatic glands. Do birds possess any? 533; lymphoid tissue in the intestine 532, 533; and mesentery 533

Comparative susceptibility of birds to infection with different types of t.b.

Types of t.b. found in birds 536; in parrots 536, 538; in birds of prey 537; experimental inoculation—of fowls and pigeons 541; of parrots 538, 540; of canaries 539; of various other birds 340

Portals of entry 542; transmission through the egg 120; t.b. in the droppings 543 droppings 543
Transmission of infection by mice and rats 544

Bone and joint tuberculosis (human) 601, 609, tables 679, 680; conclusions 609, 681; in the rabbit 387 f

Brandon, flint-knapper's phthisis at 98

Bronchial glands, more frequently tuberculous than mesenteric glands 170; tuberculosis of (human) 584; table and conclusions 586

Calf. Frequency of T. 379; age at infection 381; as a means of differentiating the types of mammalian t.b. 249 et seq.—see also "Ox"

Camel, instances of T. in 482

Caseous lesions, emulsions of injected into guinea-pigs without effect 70,

660

Cat. Frequency of T. 503; distribution of the lesions, and portals of entry 504; T. in cats of consumptives 503; instance of infection with sputum,

(Filleau and Petit's case) 503 f

Comparative susceptibility to infection with the three types of t.b. 505; expts. of R.C.T. table 506, 507; effect of avian t.b. on the cat 509; why the cat has nine lives 530 f



INDEX OF SUBJECTS

689

Cattle, comparative susceptibility of different breeds 376 f, 250 Cervical glands, tuberculosis of (human) 589; types of t.b. found 590, table 599; relative frequency of bovine infection in different countries 595, table Cattle, comparative susceptibility of different breeds 600 Frequency of infection in the pig 402

Chimpanzee, comp. suscept. to infect. with the three types of t.b. 514; expts.

with avian t.b. 519; passage expts. with bovine bacilli 350 Chopin, harsh treatment experienced on account of phthisis 48

Chyle, examination for t.b. after experimental feeding in dogs 159, 161 Civilization, its relation to susceptibility to T. 528 Coal-dust and smoke, said to be good for consumptives 97 f Coal-miners, their low mortality from phthisis 88 f Cohnheim's law 168

Congenital tuberculosis, in birds 120; ox 125; man 128, 132; conclusion 134

Contagion, why not more evidence of in phthisis 67; extreme views once held 47, 49 f; views of Young and Rush 49; James Clark 50; Laennec 50 f; Bennet 54; summary 107

Cow, frequency of infection 373, 375, 381; in different breeds 376 f; in cows of Jersey 375 f; excessive lactation a predisposing cause of T. 383

Cultivation of t.b. 215; the antiformin method 218, 685

Culture media, inconstancy of 227; egg medium 217

Curability of tuberculous lesions in—lymphatic glands 597, 661; spinal column 604 f; hones and joints 670; peritoneum 660

column 604 f; bones and joints 679; peritoneum 660

Decline of tuberculosis 14; accelerating 28; diagram 16; cause of the Decime of tuperculosis 14; accelerating 28; diagram 16; cause of the decline 17; comp. decline in males and females respectively in different decades 31; diagrams 32, 38

Denmark, swine tuberculosis in 396, 397; tuberculin tests in cattle 379

Differentiation, of avian and mammalian t.b. 185, 244, 245; of human and bovine t.b. 187, 247

Diphtheria bacilli, related to B. tub. and the acid-fast group 198

Dissemination of t.b. after subcutaneous injection 373, 673

Dissemination of t.b. after subcutaneous injection 253, 672

Doctors, frequency of phthisis among 75, 77 f; in continental hospitals 78

Dog. Reputed immunity, therapeutic use of its serum 486; instances of T. 486; frequency of canine T. in different countries 487, 489

Sources of infection. Sputum of consumptive owners 487, 491; instances of T. in butcher's dogs 491, 492; portals of invasion 492; difficulty of infecting by feeding 493; conclusion 494

Lesions, nature and distribution of 480. T. in the heart 400, 401 f

Lesions, nature and distribution of 489; T. in the heart 490, 491 f

Comp. suscept. to infect. with the three types of t.b. 496; risk of aspiration into the lungs in feeding experiments 160; passage experiments 351

Dose of t.b., influence of on the severity of the infection 60; effect of small and large doses on the calf 61, 261; on the pig 408; why small doses may fail to infect 63; selection of doses to be used for the differentiation of the types of t.b.—in the ox 266, 269; rabbit 276, 283; methods of measuring 251, 262 f; relation of numbers of t.b. to the volume of the dose of culture 262 f

Ducks and geese rarely suffer from T. 527

Ducks and geese rarely suffer from T. 527 Dust, as a cause of pulmonary disease 97; mechanical theory of dust phthisis 98; chemical theory 100

Dysgonic, meaning of 204

Dysgonic bacilli of low virulence for ox and rabbit 406, 604, 649, 681

Edinburgh and Scotland, large proportion of infection with bovine t.b. 595, 598, 608, 610, 681

Egg, transmission of infection through 120; fate of t.b. injected into eggs

Egg medium for the cultivation of t.b. 217 Elephant, instance of T. in 483 Elimination of susceptible stocks from a population 19

44

690

INDEX OF SUBJECTS

Eradication of tuberculosis 30, 109
Etiology of tuberculosis, the old view 50; seed v. soil 59, 109; dose and susceptibility—see under each heading
Eugonic, meaning of 204
Eugonic bacilli of high virulence for ox and rabbit 330, 636

Faeces, t.b. in the droppings—of cattle 400; of birds 543 Feeding experiments, danger of aspiration into lungs—see "Aspiration" Fish-tubercle bacilli 197f; stability of type of 302; presence in normal frogs 302

Flint-knapper's phthisis 101
Functional activity and local predisposition to tuberculosis 491 f; of uterus 127; mammary gland 384, 385 f, 491 f; skeletal muscle 510;

General tuberculosis 577; types of t.b. found, table 579; instance of general tuberculosis in the horse (Koch) 463 f

Glucose, action of B. tub. on, a substitute for glycerin in culture media 233 f Glycerin in culture media as a differentiator of mammalian t.b. 206, 225, 331 f; does its continued use tend to blunt the distinction between the types? 217; does it attenuate? 296; separation of human bacilli from

types? 217; does it attenuate? 296; separation of human bacilli from mixtures by growth on glycerin media 331; chemical changes in glycerin in tubercle cultures 238, (Harden) 239

Goat. Frequency of T. 412, 413; early expts. 412; outbreak of T. in a herd 413; relative susceptibility to infection with the three types of t.b. 416; susceptibility to infection by feeding 417; liability of stomach and gastric glands to be involved 418; influence of pregnancy and parturition 417 f; susceptibility to infection with avian t.b. 419; instances of intestinal disease caused by avian bacilli 419 f, 420; tuberculin test 270

Grinshill, inequality of incidence of phthisis on males and females at 100 Grouse, permeability of the intestinal mucosa 152; T. unknown 528

Guinea-pig. Spontaneous T. rare 431; relative susceptibility to infection with human and bovine t.b. 435; peculiar distribution and character of the lesions 431; extensive necrotic areas in liver and spleen 432; pleural effusion 434; site of earliest lesions in intraperitoneal injection 434;

effusion 434; site of earliest lesions in intraperitoneal injection 434; susceptibility to infection with avian t.b. 245, 437; absence of tubercles 437; intraperitoneal injection may cause fatal toxæmia 440; summary 443; liability to die of pleural effusion 387 f
Pseudo-tuberculosis 434

Heredity 83, 467 f, 543
Horse. Frequency of T., early-recorded instances 461. Peculiarity of the lesions of T. 462; enlargement of spleen mistaken for sarcoma or lymphadenoma 462; rarity of caseation and calcification 464; large numbers of bacilli 464

Sources of infection 465; infection from the cow 466; a possible instance of infection spreading from one horse to another 467 f; congenital

of infection spreading from one horse to another 467 f; congenital infection 467 f

Comp. suscept. to infect. with the three types of t.b. 468; types of t.b. found in the horse 467; compared with those from the ox 364; an avian strain (Nocard) 468; a human strain 471, 665; atypical strains of low virulence 470; experimental infection 473; is there a special equine type? 477, 481; capacity for resisting experimental infection 475; experiments with avian bacilli, table 474, 475; etiology of equine T., summary 475

Heart, is it specially liable to infection in this species? 491 f

Hungary, rarity of T. in forest swine 396

Immunity, of domesticated mammals (the dog excepted) to infection by the human tubercle bacillus 668; of certain organs—in the ox 386, in various animal species 387 f



INDEX OF SUBJECTS

691

Immunization, of cattle 22, 669; dangers of 674; Weber and Titze's case for the Cambridge case 677; of human races by elimination of the more susceptible stocks 19; by abortive infections caused by (a) small doses 21, 23, 669; (b) bovine bacilli 24 f; (c) dead and dying bacilli 669 f; influence of urbanization on the frequency of immunizing infections 22 Industrial phthisis 96

Industries, influence of the growth of on the sex distribution of T. 41; com-

parative death-rates from phthisis in various 88 f, 97 f
Infection, frequency of in man 59; by dried sputum 66; by droplets
ejected by consumptive patients 66; age at which infection occurs (a) in
man 129; (b) in ox 381
Infectiousness of phthisis—see "Contagion"

Inhalation, experiments with B. prodigiosus 148; with tubercle bacilli 150,

390; expts. in the dog 500

Inhalation and swallowing of t.b. which is the surest means of infection? 150
Injection of t.b., methods 245; estimation of dose 251; intravenous injection
in birds 246; intraperitoneal injection of the rabbit, etc. 285 f; intramammary injection of cow 390; with human t.b. 673; intravenous
injection, its limits as a test for pathogenicity (a) in birds 244; (b) in calf
391, 257, 264; summary 391; (c) in rabbit 276; (d) in goat 418
Insufflation—see "Aspiration"
Intestinal tuberculosis infraquency in Germany, 103; infection with bovine

Intestinal tuberculosis, infrequency in Germany 192; infection with bovine bacilli more liable to cause ulceration than human 517

Jersey, T. rare in the cattle of the island of; milk of Jersey cow too rich and plentiful for its offspring 383 f; susceptibility of the Jersey breed 250 Johne's disease 388; cultivation of bacillus 216, 388; alleged to be caused by avian t.b. 388

Kestrel, instance of T. in 528

Lapwing, instance of T. in 527 Latency of T. 68; latent T. lighted up by violence 68, 72, 80; in congenital T. 132, 134

Lemurs, natural T. 511; experimental 514 Lesions of T., healed and unsuspected during life 59, 69, incapable of causing T. on injection 660, 679; distribution different in different animals 387 f; distribution unusual in the guinea-pig 431; peculiarities of the lesions of T. in the horse 462; in birds 529; in the rat 444; mouse 459; in man and ox compared 546

Leysin, treatment of T. by sunlight at 655

Lime, alleged protective influence of inhaled lime 97 f

Limestone dust innocuous to masons 98; to cement makers 97 London, poverty and overcrowding in relation to T. 25

Lungs, constant presence of bacteria in normal lungs 148; portal of entry in pulmonary anthracosis 145-149. (See also "Pulmonary tuberculosis") Lupus. Annual mortality attributed to lupus 615 f; frequency of infection of lymphatic glands 615, and other parts 654 f; is lupus due to direct inoculation of skin? 616, 654 f; attempts to produce lupus experimentally

R.C.T. 617; Griffith's investigation 615; first case investigated by R.C.T. 617; Griffith's investigation for the R.C.T. 619; Griffith's second investigation 635; foreign researches 648; evidence of attenuation in many of the strains of t.b. from lupus 636; relation of virulence to (a) duration of disease 626, 639; (b) situation of the lesion 631, (tables) 633, 642; (c) treatment with X-rays, etc. 628, 629; are the bacilli which cause lupus attenuated, or do the bacilli become attenuated during their residence in skin? 627, 645, 647; has progressive attenuation been observed? 627; possible sources of attenuated bacilli 645, horse 645, birds 646; passage of attenuated strains 623 uated strains 623

44-2

692

INDEX OF SUBJECTS

Frequency of infection with bovine and human bacilli respectively in males and females 634 f Summary and conclusions 651

Lymphoid tissue in the mucous membranes, its function 152; lymphatic glands, their bacteria-arresting function 153; extent to which the glands are permeable by bacteria 154; experimental investigation of their permeability 158; conclusion 162–167; lesions of lymphatic glands as indications of the portal of entry of t.b. 168

Relative frequency of infection of bronchial and mesenteric glands

in man 169

Types of t.b. in T. of various groups of glands in man, tables—abdominal 583; cervical 599; bronchial 586; two or more sets 589 Lymphoid tissue and lymphatic glands in birds 533

Mammary gland, tuberculosis of in various animals—ox 384, 386; pig 385 f, 401; goat 385 f; rabbit 385 f, 429; man 385 f; severity of result of intramammary injection of bovine bacilli into the functional mammary gland of the cow 390; result of injecting human bacilli 392f; persistence of human t.b. after such injection 673

Excessive demand made on the mammary gland of the cow 383; and other animals 385 f; relation between susceptibility to infection and excessive functional activity 384 f

Man. Tuberculous lesions of man contrasted with those of the ox 546; his susceptibility to infection with T. compared with that of various animals 61 Comp. suscept. to infection with the three types of t.b. 193, 549, 654; distribution of the various types of t.b. among the various forms of human tuberculosis, summary 666; estimate of the deaths caused

by bovine bacilli 659, table 658 Manners, improvement in, one of the causes of the decline of T. 26

Marital infection 79

Mason's phthisis 98 Meat inspection 116

Meningitis (human), relation to general tuberculosis 569; types of t.b. found

in, table 575; conclusion 572 Mesenteric glands, types of t.b. found in human glands—see "Abdominal tuber-culosis" 580, table 583; t.b. in mes. glands of animals fed with t.b. 153; culosis" 580, table 583; t.b. in mes. glands of animals fed with t.b. 153; bacteria in normal mesenteric glands 157; relation to peritoneal cavity

436
Methylene blue as a stain for t.b. 199
Milk, control of the supply 117, 671; use of unsterilized milk in Edinburgh 596; t.b. in milk coming from normal udders 386, 676; escape of t.b. into the mammary gland after subcutaneous injection 674

Miller's phthisis 98 f

Mixed cultures, experiments with 326, 331 Mixed infections 301; mixtures of human and bovine t.b. found (a) in sputum

327, 563, (table) 564; (b) in viruses undergoing modification 334, 335, 337, 342; mixtures of avian and bovine t.b. 31

Monkeys. Susceptibility to T. 509, greater than that of man 61; character of the lesions 510; susceptibility of the skeletal muscles 387 f, 510; types of t.b. in spontaneous tuberculosis 510, 511; experimental infection 512; comparative virulence of human and bovine bacilli—M. rhesus 513; susceptibility to infection with avian chimpanzee 515; gibbon 515; susceptibility to infection with avian bacilli 517; exp. cutaneous T. 512, 518

Portals of infection. Griffith's feeding experiments 165; insufflation or aspiration into the lungs 166; habits of the monkey in feeding

166; passage experiments 350

Reaction to tuberculin 166

Mortality from Tuberculosis. Reliability of statistics 9-13; the real mortality in children 10; in old age 2; annual mortality in England and Wales 1, digram with proprious countries with Iroland and Francisco diagram 4; in various countries 1; in Ireland and France 26 f; in Prussia (from phthisis) 2 f; ratios of deaths from pulmonary and other forms of



INDEX OF SUBJECTS

693

T. 140 f; 550; actual number of deaths from T. in E. and W. in 1911 3 f;

Age and Sex distribution 3, 7; in the early years of life 5; changes in the liability of the two sexes to die of T. 41; period of maximum mortality later in males 8; retardation of the maxima in the two sexes 8; ratio of female to male mortality from T. in various places 86

Decline of the mortality from T. 8, diagram 16; in the two sexes in successive decades 31; increase in some countries 19; periodic waves in the curve 15; during the siege of Paris 15f; summary 37 Mortality in various occupations (male) 96f, 88f; (female) 43 Mouse. Reaction to infection with t.b. compared to that of the rat 449;

general resistance high 449; susceptibility nearly equal to infection with all three types of t.b. 449

Comparative susceptibility of different species of mouse 450;

omparative susceptibility of different species of mouse 450; greater susceptibility of field mouse as compared with white mouse (Koch)

450, 451 Natural infection with avian t.b. 451

Lesions, poverty of 450, 454; nodule formation after intraperitoneal injection of bovine t.b., lesions caused by avian t.b. 457; lesions caused by avian and mammalian bacilli contrasted 455

Comp. suscept. to infect. with the three types of t.b. Natural infection

451; experimental infection 453

The mouse as a carrier of avian infection to fowls 453

Muscles, their immunity from T. in most species of animal 386; t.b. in the flesh of animals slaughtered for food 386 f, 505; instances of tuberculosis in the muscles of the monkey 387 f, 510; and in those of the pig 400

Nose, efficiency as an air filter 148 Numbers of t.b. in the tuberculous lesions of various animals—man, monkey 464 f; ox, pig 465 f; sheep 421; horse 464; rat 445; mouse 457; birds 535; relation of susceptibility of a species and the numbers of t.b. in its lesions 464 f

Influence of numbers of t.b. on (a) successful artificial cultivation 208 f, 219; (b) the severity of the result of injection of animals—see "Dose, influence of"

Occupation; mortality from T. in various occupations (males) 96 f, 88 f; (females) 43

Odour of cultures of t.b. 209

Esophageal tube, use of in experimental T. by Calmette 143 f; criticism 144 f

Œsophagus, ligature of in anthracosis experiments 146, 149

Ovary, tuberculosis of 124

Overcrowding in relation to T. in London 25
Ox. As a means of differentiation of type of t.b. 249; frequency of T. in 373; in different countries 375 et seq.; rarity of T. in cattle of Java, Japan, and Jersey 375; age distribution of T. 379; age at infection 381; discrepancy between slaughter-house returns and tuberculin tests 374;

discrepancy between slaughter-house returns and tuberculin tests 374; complete post-mortem long and tedious 374 f; tuberculosis compatible with healthy appearance 377 f

Predisposing causes of infection 382; excessive lactation 383; milk of Jersey cow too rich for the calf 383 f

Portals of infection 385; character of the lesions 548; immunity of certain organs 386; immunization dangerous 387 f

Types of t.b. found in the ox 388; avian bacilli said to have been found 388; Johne's disease not due to avian bacilli 388

Experimental infection 249 et seq.; summary (with bovine bacilli) 389; (with human bacilli) 391; intravenous infection with human t.b. followed by fatal results 257, 391; exceptionally severe result of subcutaneous injection of human t.b. 254; inoculation of functional

© in this web service Cambridge University Press

www.cambridge.org

694

INDEX OF SUBJECTS

mammary gland (a) with bovine bacilli 390; (b) with human bacilli 392, 673, 674 f

Reaction to tuberculin 249, 329, 338, 673

Relative susceptibility of different breeds 376 f, 250

Paris, mortality from T. during the siege 15 f

Parrot 535; lesions 535; in the skin 543; infection with human t.b. 538; susceptibility to infection with all three types of t.b. 538; infection with

avian t.b. in zoölogical gardens 538

Partridge, instance of T. in 528

Passage experiments—see "Stability of type"; expts. with "attenuated" bacilli from (a) lupus 623; (b) horse 473; (c) pig 406

Permeability by bacteria of intestinal mucosa 153, 168; affected by nematoid worms and coccidia 152, 411; of lymphatic glands 153

Phthisis, relative frequency as compared with other forms of tuberculosis 140 f. 550; associated with disease of cervical and mesenteric glands 174.

140 f, 550; associated with disease of cervical and mesenteric glands 174; portals of entry 140; summary 176; nature of certain forms of occupational phthisis 96; infectivity of, Chapters IV and V; instances related by—Hermann Weber 52; Dr Sprigge 53; Bennet 55; Frequency of swine T. in various countries 396 et seq., in U.S.A. and

Argentina 396 f, in London 398, in Brighton 399; is swine T. increasing?

Argentina 390 i, in London 357,
397 f

Sources of infection and portals of entry 396, 399, 401; relation of swine T. to the dairy industry 396, 399 f

Types of t.b. found in natural T. 403, 405; avian strains 404, 405; human strains 403, 404; atypical bovine strains 403, 404; comp. virulence of bacilli of bovine type found in pig and ox 362

Succeptibility to experimental infection with the three types of t.b. 403;

virulence of bacilli of bovine type round in page and Susceptibility to experimental infection with the three types of t.b. 403; Susceptibility to experimental infection with the three types of t.b. 403; Sustain 409; comwith avian t.b. 411, 412; with human t.b. 408; sputum 409; comparative susceptibility of young and old pigs 408; susceptibility of pig compared with that of the calf 408

Pigmentation of cultures of t.b. human type 209, 681, 682

Population, growth of 8 f

Portal lymph gland, relation to drainage of peritoneal cavity in guinea-pig 434 Portals of entry of infection, Chapters viii and ix; in phthisis, summary 176; in general tuberculosis and meningitis 178; in T. of bones and joints 180, 601; cervical gland T. 590; lupus 616

Post-mortem examination of ox 374

Potter's phthisis 106
Poverty in relation to T. in London 25
Pseudo-tuberculosis in the guinea-pig 434
Pulmonary tuberculosis, death-rates in various countries 2; in Ireland and
France 26 f; in Prussia 8 f

Relative number of deaths from pulmonary and other forms of T. 140 f, 550 f

Types of t.b. found in pulmonary tuberculosis: strains of t.b. from tuberculous organs 551, table 557; from sputum 558, table 564; instances of infection with bovine bacilli—alleged instances by Creighton 547; Arloing's case 559; the de Jong-Stuurmann case 565; Mohler and Washburn's case 564; Griffith's three instances 565, 567; summary 568

Quiescent tuberculosis, reactivation by violence 68, 80

Rabbit. Rarity of T. in the wild state 426; the cause of this 429; spontaneous T. in laboratory animals 426; an outbreak of T. in a stock of

tame rabbits (Rothe) 427

Comp. suscept. to infect. with the three types of t.b., summary 428

Experim. infect. with human t.b. 272, 276, 428; exceptional results 279, 281; are these caused by certain strains? 282 f; instances of progressive infection 277, 280; fatal pulmonary T. after intravenous



INDEX OF SUBJECTS

695

injection 276; lesion caused by the human type of t.b. 276, 429; liability of lungs and kidneys 278; T. of the functional mammary gland 385 f; alleged escape of lymph glands 283 Experim. infect. with bovine t.b. Extreme susceptibility, the effect of

minute doses 274; lesions caused by bovine t.b. 275, 429; their distribution 275; T. of functional mammary gland 385 f; susceptibility to infection by feeding 275, 428, 430

As a means of differentiating the types of t.b. 185, 188, 271, 273; reaction to tuberculin 372 f; methods of infection 275, 284, 482; doses used

283, 285

Permeability of intestinal mucosa affected by coccidiosis 153; weak

spots in the rabbit's defences 387 f

t. Instances of spontaneous T. 443; peculiarity of its lesions 445, cause of this 445; enormous numbers of t.b. 445 f

Comp. suscept. to infect. with the three types of t.b. 446; lesions

caused by avian and mammalian bacilli compared 446; difficulty of infecting by feeding 448
Reaction curve (Smith) 231 et seq.; conclusion 241

Sanatoria, their influence on the eradication of T. 114

Sandstone and limestone, dust from the former alone dangerous to masons 98 Seed and soil, relative importance of 59; which is easier to control? 109
Segregation 27, 45, 111; value of partial segregation 112; influence of in
the eradication of leprosy 112

Selective breeding, its influence on resistance to T. in cattle 383; in birds 529
Serum from various animals as culture media for t.b. 216
Sheep, recorded cases of T. 420; lesions caused by Strongylus rubescens liable to be mistaken for tubercle 420; susceptibility to experimental infection with various types of t.b. 422; lesions 421; scarcity of t.b. in the tubercles 421; suggested explanation of the comparative immunity 421 Sheffield, disparity of the mortality from T. of men and women in 89, 90;

compulsory notification 2 f; mortality among grinders 31, 90, 96

Silicosis 97, 100, 102
Skin, cutaneous T. in parrots 543; in the fowl 532; in birds 656 note
Slaughter-house statistics, unreliability of 374, 378; complete post-mortem examination long and tedious 374 f
Sparrow, instance of T. in 528

Sputum, precautions necessary when investigating types of t.b. contained in 161; types of t.b. found in 558, 564; bovine t.b. found in 559, 565; dried and pulverized sputum as a cause of T. 65, 66, 140, 671; indiscri-

minate spitting 23

minate spitting 23
ability of type. Attenuation, evidence of 114, 471, 620, 635; attenuation under artificial culture 286, 288, 290, 292, 295; stability of virulence on change from animal to culture tube 287; an instance of attenuation in the records of the R.C.T. 294 f; attenuated strains from man (lupus), horse, pig 214, 355, 635; summary 298; influence of glycerin 296; alleged attenuation in cold-blooded animals 302

On "passage" through various animals, Chapters xv, xvi, xvii; difficulties of investigation 300, 326; alleged conversion of mammalian into avian bacilli 304; attempts to convert avian into mammalian t.b. 310; alleged conversion of human into bovine bacilli in goats and calves 317; experiments of—Gesundheitsamt 319, 325; Stability of type.

mammalian t.b. 310; alleged conversion of human into bovine bacilli in goats and calves 317; experiments of—Gesundheitsamt 319, 325; Dammann and Müssemeier 320; Eber 322; Royal Commission (1st series) 329, (2nd series) 345; passage through the rabbit 347; attempts to modify bovine bacilli 349; in the monkey and chimpanzee 350; dog 351; do bacilli of bovine type from man differ at all from those from the ox? 358; stability of bovine type in man (a) cultural characters 357 f; (b) virulence 360

Virulence of strains of bacilli of bovine type from various animals compared with those from the ox itself—man 358, see also 354, 356; pig 362; horse 364

pig 362; horse 364

696

INDEX OF SUBJECTS

Summary of "passage" experiments 366
Submaxillary gland, the common seat of localized T. in the pig 402
Susceptibility to T., of man compared with that of other animals 61; uncivilized races 19,49 f; susceptibility and civilization 528; susceptibility of a race diminished by repeated elimination of its more susceptible indi-

viduals 19; susceptibility of domesticated and wild animals compared 668 f; T. in wild animals in captivity 483, 527, 528

Susceptibility influenced by various causes. Behring's view 69; influence

of parturition in the goat 417 f; of artificial selection and excessive

Want of correspondence between the susceptibility of a species as measured by experiment and by its tendency to suffer naturally from the disease—in the rabbit 429; guinea-pig 431; goat 415;

sheep 421; rat 443; monkey 511

Susceptibility of certain hard-worked organs. Mammary gland of cow 384 and footnote; pig and rabbit 385 f; heart of dog 491 f; climbing muscles of monkey 387 f, 510; susceptibility of virgin and pregnant uterus compared 127; comparative susceptibility of various organs 386; may differ in different species, to each its weak spots 387 f; the susceptible organs in rabbits and guinea-pigs contrasted 429

Susceptibility of earlier-born members of a family alleged to be greater than that of the later-born 84 f

greater than that of the later-born 84 f Difference of susceptibility of different individuals of the same species-

calves 266, 269 f; rabbits 282; influence of age in the ox 379, 382; pig 408

Texas, rarity of T. among swine allowed to run wild 396

Therapeutic action of sunlight 655 note

Therapeutic action of various substances—smoke 98; sulphur 98; lime 97;

serum of ass 461, 478; of dog 486

Thoracic duct, T. of 179

Tissue emulsions, their virulence compared with that of pure cultures 287; method of estimating numbers of t.b. in 252 f, 262 f

Tonsils, T. in the, (man) 73, 175, 590; (pig) 401

Toxæmia, evidence of after injection of functional mammary gland of cow 390

Tubercel bacillus—see "B. tuberculosis"

Tuberculin tests, reliability of in various animals, ox 240, (Fber) 200, 208;

Tuberculin tests, reliability of in various animals—ox 249, (Eber) 329, 338; cows injected with human t.b. 673; goat 270; rabbit 272 f; monkey 166; discrepancy between tuberculin tests and slaughter-house returns 374 Types of t.b. 196, 684; stability of—see under "Stability of type"; are all strains included in one and the same type identical (a) in cultural characters 357, 683 (b) in virulence 282 f, 361 f, 353, 358, 362, 364—see also Chapter xxv; a fourth type 197 f; a new sub-type 681

Underground Railway as a sanatorium for consumptives 98 f United States, frequency of T. in cattle 377, 378; in swine 397, 398; frequency of infection with bovine bacilli in human cervical gland tuberculosis 600 Uterus, lesions commonly present in when infection of the cow is transmitted to its calf 126; comparative susceptibility of pregnant and virgin uterus 127; infection by human t.b. spreading from peritoneum to uterus in the rabbit 279

Violet coloration of cultures of fish-tubercle bacilli 209 Virulence, difficulty of accurate estimation owing to variable resistance of different individual test animals 267; v. of bacilli in pure culture and tissue emulsions compared 287; stability of—see "Stability of type" Difference of v. of bovine and human types not merely one of degree

Weak spots in the defensive mechanisms of various animals 387 f



INDEX OF AUTHORS

Abel 503

Achard 541 Adami. On the lymphatic defence of the animal body 153; invasion by intestinal bacteria 155

Addison 50 Albrecht. Frequency of infection at various periods of early life 130 Allbutt, Sir Clifford. Views on contagion 52, 77; frequency of unsuspected lesions 70
Andrews. Post-mortem appearance of the lungs of two ganister miners 104
Archibald. T. in the camel 482

Arlidge. Potter's phthisis 103; pulmonary diseases induced by dust 106 Arlidge. Frequency of T. in cattle in France 378; feeding experiments in the dog 492; case of pulm. T. in man yielding bovine bacilli 559 Arloing and Courmont. An attenuated strain of t.b. 288 Arloing and Tripier 525

Arnold. Origin of pulmonary anthracosis 146 Auché and Chambrelent. Congenital tuberculosis 132

Aufrecht 176

Baermann and Halberstaedter. Experimental tuberculosis of skin 650 Baginsky. Intestinal origin of phthisis 141; rarity of intestinal T. in

Germany 192
Bang, B. T. in cattle in Denmark (congenital) 125; (proportion reacting to tuberculin at different ages) 377, 379, (calves) 381; T. in the dog 489 Bang, O. The "reaction curve" 233; Johne's bacillus 385; conversion of mammalian into avian t.b. 307; growth of avian t.b. on broth 211 Bartel and Spieler 176
Bataillon, Dubard and Terre. The fish-tubercle bacillus 302

Batier 176

Beck. Glycerin in culture media a cause of attenuation 296

Beckmann 176
Beevor, Sir Hugh. On contagion in phthisis 57; why girls suffer from T. more than boys 39; causes of the decline of T. 25
Behring. Intestinal origin of phthisis 68, 69, 141, 142, 176
Behring, Römer and Ruppel. Conversion of human into bovine t.b. in goat

317

Beitzke. Œsophageal tube in feeding experiments 144; channels of infection 176; repetition of Grober's experiment 176; types of t.b. found in the

lymphatic glands of children 554
Bennet, Dr J. Henry. Etiology of phthisis 54
Bisanti and Panisset. Absorption of t.b. from the dog's intestine 161
Bland-Sutton and Gibbes. T. in birds of Zoölogical Gardens 526



698

INDEX OF AUTHORS

Blaxall. Frequency of T. in calves purchased by the L.G.B. for the preparation of vaccine lymph 381 Bollinger 188

Bowditch. Subsoil drainage

Bowditch. Subsoil drainage 24
Bright. On the tubercular diathesis 50
Broadbent. Incidence of phthisis on rich and poor 27; latency (case of quiescent T. reactivated through an accident) 68; segregation 110
Brusasco. T. in dog 487
Buchanan. Subsoil drainage 24; statistics of swine tuberculosis 397
Bulloch. On contagion in phthisis 58; types of t.b. in sputum 563
Bulloch and Greenwood. Quotation from Thomas Young 49 f
Bulstrode. On contagion in phthisis 58; criticism of Cornet 70; prevention

Bulstrode. On contagion in phthisis 58; criticism of Cornet 79; prevention

of T. III

Burckhardt. Types of t.b. found in man (in cervical glands) 591, (bones and joints) 603; the rabbit as a means of differentiating types of t.b. 271 irnet. Attenuated t.b. 650; immunization by abortive infection 669 f Burnet. Attenuated t.b. 650; in Burns, Sir John. Segregation 27

Cadiot. T. in the dog 490, 491
Cadiot, Gilbert and Roger. Infection of guinea-pig with avian t.b. 438;
T. in the goat 412; in the parrot 535; types of t.b. in the parrot 536, 538
Calmette. Hereditary transmission of infection from male parent 124; intestinal origin of phthisis 143, 151
Calmette and Guérin. Experimental production of pulmonary T. 143; use of œsophageal tube 143; experimental anthracosis 145, 147
Calmette, Guérin and Deléarde. T.b. in normal glands 73
Carozzi Immunity of sulphur workers in Sicily 08 f

Carozzi. Immunity of sulphur workers in Sicily 98 f
Chauseé. Feeding experiments in dogs 493
Chauveau. Early experimental tuberculosis 51; the alimentary canal as portal of entry of t.b. 141; experimental T. in the ass 478; also 188
Clark, Sir James. Pathology of tuberculosis in 1855 50; on contagiousness of phthisis 51

of phthiss 51
Coates. Frequency of T. in children 10, 12 f
Cobbett. Portals of infection—congenital tuberculosis, rarity in guinea-pig
and rabbit 125; three cases in the ox 126. Accessibility of lungs to
air-borne bacteria 148; inhalation experiments with B. prodigiosus 148, with
B. tub. 150. Penetrability of mucous membranes and lymphatic glands—
feeding experiments with B. prodigiosus 163; B. tub. 150, 161; indian
ink 146; fate of carbon mixed with cream in the intestine 148; accidental
aspiration into lungs in feeding experiments 162, 163; influence of
intestinal worms on the absorption of bacteria in groups rabbit shoop. aspiration into langs in feeding experiments 102, 1031, instituted of intestinal worms on the absorption of bacteria in grouse, rabbit, sheep 153

Bacterial content of normal blood and organs 148, 156

Rabbit—extreme susceptibility of to small doses of bovine t.b. 274; intraperitoneal injection, method employed 285; spontaneous avian

tuberculosis 427

Dog-exceptional result of a feeding experiment 493 f

Tubercle bacilli-method of broth culture 220; material from caseous

Tubercle bacilli—method of broth culture 220; material from caseous lesions on injection may be incapable of infecting 76. Stability of type—influence of glycerin on virulence 296; experiments with mixed cultures of tubercle bacilli 331. Passage—experiments in the ox 345, 348; rabbit 347

Dissemination of bacilli after subcutaneous injection 673

Cohnheim. Early experimental tuberculosis 51; Cohnheim's law 168

Collis. Certain forms of industrial phthisis due to dust—in millers 98 f, masons 98, flint-knappers 101; distinction between dangerous and harmless stone dusts 98; chemical theory of dust phthisis 99; reason of non-infectiveness of stone-worker's phthisis 102; on pulmonary cirrhosis unaccompanied by tubercle 103

Cornet. Infection by dried and pulverized sputum 66, 140; incidence of T. on the members of the staffs of foreign hospitals 78; channels of infection in pulmonary tuberculosis 176

in pulmonary tuberculosis 176



INDEX OF AUTHORS

699

Courmont and Dor. Passage experiments with avian bacilli in the rabbit 313; intraperitoneal contrasted with subcutaneous injection of avian t.b. into the rabbit 438

Creighton. Supposed characters of the lesions of bovine tuberculosis in

man 547

Opposition to Koch at the London Congress 191 Crookshank. Types of t.b. in sputum 563 Cruickshank.

Dammann and Müssemeier. Stability of type of t.b. 320; susceptibility

Dammann and Mussemeier. Stability of type of t.b. 320; susceptibility of the sheep to infection with the two mammalian types of t.b. 423

Dammann and Stedefeder. T. in the elephant 483

Deaver. T. of human mammary gland 384 f

Deetz. Statistics of swine tuberculosis in Berlin 397 f

Delbanco. T. in the parrot 538

Delépine. Frequency of T. among cows 376; age distribution of T. in cattle 380; instability of type of t.b. 307 f; relation of bovine to human tuberculosis 101 culosis 191

Delille, Dr. A. Therapeutic influence of sunlight 655

Derry. Tuberculosis in early Egypt 46
Dieterlen. Types of t.b. in sputum 562, 563
Dinwiddie. Relation of animal to human tuberculosis 189 f; comparative susceptibility of the pig to infection with human and bovine bacilli 409 f Dorset, M. Egg medium for the cultivation of t.b. 217 Von Dungern and Smidt. Exp. T. in the gibbon 515

stwood. Cultural characters of human and bovine t.b. 204; method of raising broth cultures 220; estimation of dose of bacilli by weight and volume compared 262 f; cultural characters of t.b. of bovine type from Eastwood.

man and the ox compared 357
Eastwood and F. Griffith. Injection of caseous material from tuberculous glands may fail to infect 71, 661; t.b. in apparently normal glands 73.

Frequency of tuberculous infection in children 11 f; relative frequency of infection of various groups of lymphatic glands in children 172; portal of entry in general tuberculosis 179, 180.

Post-mortem examination of a case of lupus 139; types of t.b. in human home and injust tuberculosis 679.

bone and joint tuberculosis 679

Swine tuberculosis 399; types of t.b. found in 404; feedin ments with (a) bovine bacilli 408, (b) human bacilli 410. feeding experi-

Eber. Stability of type of t.b.—modification experiments 322, 328; of t.b. present in pulmonary tuberculosis 556; T. in the dog 487 Eichhorn. Outbreak of T. in a herd of goats 413 Elderton. Assortative mating 82 types

Ellenberger and Baum. Lymph system in birds 533

Instances of transmission of infection in human tuberculosis Emmanuel.

Fibiger and Jensen. The distinction between the three types of t.b. 197 f; reaction curve 234, 237

Ficker 161

Filleau and Petit. A remarkable case of T. in a cat infected with sputum 503 f

Findel. Infection by feeding and inhalation compared 151

Finlay.

Experimental anthracosis 149
Infection by droplets of sputum 66, 140; infection by feeding Flügge. and inhalation compared 150

Ford. Bacterial content of normal tissues 155

Fowler, Dr Kingston. On latency in tuberculosis 68
Fox, Dr Wilson. Case of congenital tuberculosis in a guinea-pig 125 f
Fraenkel. Intestinal origin of phthisis 141
Fraser. Types of t.b. found in human bone and joint tuberculosis 606, 608, 609

www.cambridge.org



700

INDEX OF AUTHORS

Friedberger and Fröhner. Canine tuberculosis mainly a pulmonary disease 489; frequency of T. in the cat 503; source of infection in feline tuberculosis 504
Fröhner and Eberlein. T. in the parrot 535
Frosch and Hertha. Frequency of T. in the goat 412

Frothingham 189 f

Gadow, H.

H. On the lymphatic systems of birds 533 Proportion of deaths of children caused by tuberculosis T. of lymphatic glands of children 73; relative frequency in different groups 170; t.b. in mesenteric glands of a newly born infant 129
Gaffky and Koch. Experiments on mice 451
Galen. Infectiousness of phthisis 47, 49

Ganghofner. Age at infection 129

Ganghofner. Age at infection 129
Gärtner. Inoculation of canaries' eggs 123, 539; insusceptibility of fowl to mammalian T. 541; fate of t.b. in the fowl 542
Gaultier. Experimental T. in the ass 479
Gebhardt. Infection by feeding and inhalation compared 150
Gibbes, Heneage. Virulence of tuberculous material from man and the ox compared 188; see also Bland-Sutton
Giordano, Dr Alfonso. Immunity from phthisis of sulphur workers in Sicily 98 f
Goodale. Tuberculosis of the tonsil 73
Gosio. Type of t.b. in lupus 648

Goodale. Tuberculosis of the tonsil 73 Gosio. Type of t.b. in lupus 648 Grancher and Ledoux-Lebard. Injection of dead avian t.b. in the guineapig 440 Graves. On "the scrofulous habit"

Greenwood. Child mortality in Blackburn 95
Griffith, A. S. Frequency of infection at different age periods of early life 131; caseous or calcareous glands may contain no living t.b. 70, 660.

Penetrability of intestinal mucosa and lymph glands, expts. in (a) dogs 153, 159, 164; (b) pigs, cat, monkey 160; fallacies of feeding experiments 165, 500; conclusions 165, 167, 173

Comparative frequency of infection of various groups of lymph glands in

children 172, 179

Tubercle bacilli. Cultivation. Egg-medium 217; use of antiformin

children 172, 179
Tubercle bacilli. Cultivation. Egg-medium 217; use of antiformin 218, 685; relative importance of cultural characters and virulence 636; the reaction curve 232, 237, 238; growth of bovine t.b. in broth 227, 237; new sub-type of t.b. 680
Classification of animal species according to their reaction to various types of tubercle bacilli 247. Rabbit. Extreme susceptibility to minute doses of bovine t.b. 274; intravenous injection with human t.b. 276 f; subcutaneous inoculation with human t.b. 280; method adopted 283; frequency of infection of mammary gland 385 f; value for differentiating types of t.b. 271. Pig. Susceptibility of young and old compared 408; susceptibility of pig to infection with human t.b. compared 408; susceptibility of pig to infection with human t.b. compared with that of the calf 409 f; Goat. Spontaneous tuberculosis 413, 414. Dog. Feeding expts. 493. Cat 508. Birds 541; T. of certain glands in the neck 534; parrot 538 f. Mice 456. Monkey. Comparative virulence of human and bovine t.b. 513. Stability of type, of mammalian t.b. in the fowl 308; passage experiments in the calf 346; rabbit and rat 347. Man. Types of t.b. found in (1) sputum 503, 682; (2) in the lymphatic glands of children suffering from (a) pulmonary tuberculosis 556; (b) meningitis 574, 576; (c) general tuberculosis 588; cervical glands 597; bones and joints 601, 605, 679, 680. Relative frequency of bovine infections in England and Scotland 598, 605, 683. Lupus, atypical strains of t.b. 614; first investigation (for R.C.T.) 618; tabular summaries of results 626, 632; second 598, 605, 683. Lupus, atypical strains of t.b. 614; first investigation (for R.C.T.) 618; tabular summaries of results 626, 633; second



INDEX OF AUTHORS

701

investigation (at Cambridge) 635; tabular summaries of results 639, 642; action of artificial light on t.b. 634

Griffith, A.S. and F. Stability of type of tubercle bacilli—modification experiments with bovine t.b. in (1) the monkey and chimpanzee 350; (2) dog 351

Swine tuberculosis. Types of tubercle bacilli found in the pig 403; localized tuberculosis 402, 399; T. in the mammary gland of the sow 385 f; t.b. in the joints 400; intestinal origin of T. 402.

Dissemination of t.b. after subcutaneous injection 673; do t.b. in the blood get into the milk without lesion of mammary glands? 677; T in a sparrow 528 blood get into the milk without lesion of mammary glands? 677; T. in a sparrow 528

Griffith, F. Duration of life of cultures of tubercle bacilli 221

Stability of type of t.b. Modification experiments with (a) avian t.b. in the (1) calf 310; (2) various animals 311, 312; (b) mammalian t.b. in the fowl 399; bovine t.b. in the dog 352

Virulence of avian t.b. for goat 419, 420; pig 412; horse 474, 475, 476; guinea-pigs 440; rabbit 428; monkeys 518; chimpanzee 519; dog 501; cat 509; rat 449; mouse 456

Horse. Types of t.b. found in the horse—atypical strains 469 et seq.; experimental infection 475 experimental infection 475
Chimpanzee and other monkeys. Types of t.b. found in spontaneous tuberculosis 511 Rabbit. Spontaneous infection with avian bacilli 427 Wild animals kept in captivity, instances of T. in 482 Parrot 538 Grober. Portals of entry in phthisis—indian-ink experiment 175 Grund. Growth of bovine t.b. on broth 227; the reaction curve 234, 236 De Haan. Immunity of wild monkey from T. 510; rarity of T. in cattle of Java 375
Haldane, Martin and Thomas. T. in tin-miners 97f; low infectiveness of miners' phthisis in Cornwall 100
The horse Frequency of infection at various periods of early life 130 Hamel. Relative number of deaths from pulmonary and other kinds of T. in various countries 550 f Hamilton. Nature of phthisis caused by stone dust 104 Hammond Smith. T. in a kestrel 528 Harbitz 176
Harden. Bio-chemistry of B. tub. 239
Hay, Matthew. Deaths from phthisis attributed to bronchitis 2 f, 13 f; decline of T. 17 f; greater liability of girls than boys to phthisis 39 Heberden. Infectiousness of phthisis 49 Héricourt and Richet. Effect of avian t.b. on monkey 517; therapeutic use of dog's blood in tuberculosis 486, 461 Heymans. Reactivation of lesions due to human t.b. in the rabbit 277 Hippocrates 47 Hobday and Belcher Holth. Johne's bacillus 389 Humphry, Sir George. Collective investigation of infectiousness of phthisis 53, 80 Hutchens, Prof. H. Comparative length of human and bovine t.b. 200 Hutchinson, Sir Jonathan 68 Hutyra and Marek. T. in the parrot 538 Jensen. T. in the dog 489; in the cat 503
Johne. Expt. T. in the ass 478
Johne and Frothingham 388
De Jong. Spontaneous T. in the mouse 451
De Jong and Stuurmann. Case of pulmonary tuberculosis alleged to be caused by bovine bacilli 559; summary of the case 565

Kelly 24



702

INDEX OF AUTHORS

Kersten and Ungermann. Swine tuberculosis 406; portals of entry in 401 Kitasato. Rarity of T. in cattle of Japan 375; types of t.b. in sputum 563 Early experimental T. 51 Klebs. Klein. Distinction between human and bovine tuberculosis foreshadowed 187 Kleine. Atypical t.b. from T. of skin 354, 648 f, 683 f Klenke Klimmer. Alleged modification of fish-tubercle bacilli 303
Koch, R. Decline in the death-rate from phthisis in Prussia 8 f; opinion in Germany as to contagiousness of phthisis 55; infection by dried pulverized sputum 66, 140; institutional segregation 28, 111; partial segregation and the stamping out of leprosy 112
Relation of animal and human tuberculosis—first opinion opinion expressed in London in 1901 190, 191 f; cause of Koch's early failure to recognize differences of virulence in human and early failure to recognize differences of virulence in human and bovine t.b. for the cat 505; rabbit 510

Swine tuberculosis—portals of entry 401

Susceptibility of field mice to experimental tuberculosis 450; case of gen. tub. in a horse 463; experimental tuberculosis in dog 487; cat 505 f; rarity of spontaneous T. in laboratory rabbits 426; case of T. in wild rabbit kept in captivity 430 f; lesions of monkey and man compared 510; lupus 617; instance of gen. tub. in the horse 463 f; criticism of sputum experiments of the Roy. Comm. 559 and Schütz Distinction between human and bovine tuberculosis 190 f; Koch and Schütz. Distinction between human and bovine tuberculosis 190 f; experiments on the sheep 423
Koch, Schütz, Neufeld and Miessner. Attenuation of t.b. 288
Koch, Schütz, Neufeld and Miessner. Attenuation of t.b. 288
Koch, M. and Rabinowitsch. T. in Birds of the Zoölogical Gardens, Berlin 526; Parrot 535; types of t.b. found in the parrot 537, 538; in birds of prey 537, 540; experiments in canaries 539; T. rare in young birds 121; liability of bird's lungs to infection 531

Hereditary transmission 124; fate of t.b. introduced into eggs 122; inoculation of eggs 123; t.b. seen in bird's ovary 124

Avian tubercle bacilli. Cultural characters approximate to those of human type 205 f. pellicle formation on broth cultures 211; effect human type 205 f; pellicle formation on broth cultures 211; effect of avian t.b. on the guinea-pig 441

Spontaneous Tuberculosis in wild rat 443; mouse 542; expts. in mice 453; transmission of infection to fowls by mice 544. Passage experiments with avian bacilli in guinea-pigs and rabbits 313 Johne's disease 388 Kortum 51 Kossel. Types of t.b. from sputum 562; mixtures of human and bovine t.b. found in sputum 327
Kossel, Weber and Heuss. Infection by feeding and inhalation compared
150. Human and bovine tubercle bacilli—cultural characters compared
205. Stability of virulence of bovine bacillus in artificial culture 289; passage experiment with human t.b. in the goat 319; criticism of Behring's passage experiments 318; differences in individual susceptibility in calves 270 f Swine tuberculosis 406, 409 f

Human tuberculosis—types of t.b. found in meningitis 572 Kraus and Grosz. Experimental cutaneous tubercle in the monkey 512, 650 *Ibid*. 650 Kraus and Kren.

Views as to contagiousness of phthisis 50; cause of the prevalence of phthisis among nursing sisters 78 Lannelongue 541 Lartigan 189
Latham, A. A case of cutaneous tuberculosis 138 f
Legg, Dr T. H. Ganister miner's phthisis 105
Lehnert. T. in a colt kept with cows 466

Avian t.b. from the lung of an ox 388

Léniez 525



INDEX OF AUTHORS

703

Leudet and Petit. Failure to infect dogs by inhalation 492 Lewis. The reaction curve 233; types of t.b. in cervical gland tuberculosis 591 Recovery of an old bovine strain from a mixture of t.b. 222; Lindemann. difficulty of preparing culture media of constant value 228; attenuation of t.b. 288; types of t.b. found in sputum 562, 563; a mixture of human and bovine t.b. found in sputum 328 ngard and Leese. T. in the camel 482 Lingard and Leese. Link 501 Lipschütz. Avian t.b. in man 656
Lishman. T. in vertebrae of horse 463 Lister. On Koch's announcement re relation of human and animal tuberculosis 191
Litterer. Types of t.b. in cervical gland tuberculosis 592
Longstaff. Criticism of evidence of contagion in marital infection 81
Löwenstein. An atypical strain of t.b. (?avian) from sputum 212 f; other instances of avian t.b. in man 646 f Lubenau. Egg medium 217 Lyon. Deaths from T. in old age 3

McFadyean. On Calmette's theory of intestinal origin of phthisis 145; McFadyean. On Calmette's theory of intestinal origin of phthisis 145; summary of the results of the feeding experiments of the R.C.T. 164; frequency of T. in the Ox 375; in various parts of England 375; infrequency in young cattle 380; T. compatible with appearance of health 377 f; T. in the Horse 462; recognition of enlargement of spleen as a manifestation of T. 462; Equine T. mainly an abdominal disease 463; portals of entry 465; T. in the bones 463 f; atypical strains of t.b. in the horse 471; experiment on an ass 481; on the existence of an equine type of t.b. 481 f
McFadyean, Sheather and Edwards. Johne's disease 389
Macfadyen, Alan. Comp. virulence of human and boving the for the monkey

ther and Edwards. Johne's disease 389 Comp. virulence of human and bovine t.b. for the monkey Macfadyen, Alan. 512, 517 f

Macfadyen and MacConkey. T.b. in normal lymph glands 72; in mesenteric

glands of newly born infant 129
Maffucci. Possibility of infection through the egg 122; inoculation of eggs 123; insusceptibility of fowl to infection with mammalian t.b. 540; early distinction of avian from mammalian t.b. 187

early distinction of avian from mammalian t.b. 187
Mainwaring. Infectiousness of phthisis 47
Malin. Early-recorded case of T. in the dog 486
Malm. T. in cattle of Norway 377
Markus. Johne's disease 388; T. in the horse 465
Marsden, Dr S. Tuberculosis death rates 96
Martin, Dr Sidney. Virulence of human and bovine tubercle for the ox 188
Maundry. T. of human mammary gland 385 f
Melvin. T. in the ox, frequency of in U.S.A. 375, 378; unfrequent in calf 381; frequency of T. in the goat 413
Metchnikoff. Racial susceptibility 20 Racial susceptibility 20 Metchnikoff.

Meyer 389 Milroy, Garvin. Subsoil drainage 24

Mitchell. Types of t.b. in human tuberculosis—meningitis 574; general T. 579; abdominal T. 583; cervical glands 595

579; abdominal T. 583; cervical glands 595
Möller. Lupus 648
Möllers. Types of t.b. in sputum 562, 563; T. of bones and joints 604
Mohler and Washburn. The reaction curve 233; swine tuberculosis 396; epidemic of swine T. in Oregon 411; T. in mammary gland of the pig; t.b. in the droppings of cattle 400 f; of birds 543; an atypical strain of t.b. from sputum 564
Moore, Prof. B. Infectiousness of phthisis 77
Moore, V. Tuberculous nodules in the mesentery of birds 533; insusceptibility of the fowl to mammalian T. 541
Morgagni 48

Morgagni 48



704

INDEX OF AUTHORS

Therapeutic influence of sunlight 655 Morton, R. 47
Much. Immunizing influence of bovine infections 24 f Murphy, Sir Shirley. Overcrowdin lost some of its virulence? 18 Mursajeff. T. in the parrot 538

Naegeli. Tuberculous lesions unrecognized during life 21, 69; T. in newly born infants 128; age when infection most frequently occurs 129; decline of ratio of deaths to infections as age advances 21

Are bacteria present in normal blood and tissues? 154 Dold and Lindemann. Repetition of Eber's "passage" experiments Neisser. Are bacteria present in Neufeld, Dold and Lindemann.

Newsholme. Reliability of statistics of T. 10; alleged diminution of virulence of t.b. 19; on the communicability of phthisis 58; sub-soil drainage 24; influence of urbanization on the prevalence of T. 27; control of tuberculosis 110; segregation 111; seed and soil, which is easier to control? 109 economic application of available resources 109, 110; success of partial segregation in the control of leprosy in Norway 113; influence of segregation on the reduction of the mortality in the two sexes 44; criticism

of the evidence of the non-infectiousness of phthisis 77 Nicolas and Descos. T.b. in the chyle of dogs fed with infected soup 161 Niven. On the unequal incidence of the mortality from T. on the two sexes 43 Nocard. Bacterial content of normal blood and tissues 154; early recognito the cultural peculiarities of avian t.b. 187; relation of bovine to human T. 191; alleged conversion of mammalian into avian t.b. in the fowl 304. Ox. Frequency of T. in cattle in France 375; unfrequency of lesions in the muscles 386 f. Horse. Peculiarity of the lesions 462; primary pulmonary T. 463; portals of entry 465; sources of infection 466, 469; reason why T. does not spread from horse to horse 460. T. in the Dog 487; Cat 504; Birds 541

Oberwarth 161 Oehlecker. The rabbit as a means of differentiating human and bovine bacilli 271; alleged escape of lymphatic glands of rabbits injected with human t.b. 283; best dose and method of injecting rabbits 284; human tuberculosis—cervical glands 590; bones and joints 603
Oliver, Sir T. Nature of the "phthisis" caused by stone dust 104; intestinal

origin of phthisis 142 th. T.b. in normal tissues 72

Orth and Rabinowitsch. Absorption of t.b. from alimentary canal 162, 164
Osler, Sir W. An early Greek reference to tuberculosis 46
Ostertag. Frequency of T. in cattle in Berlin 378; unfrequency of lesions in the muscles of the ox 386 f

Palamidessi 540

Pansini. Avian t.b. in the ox 388; in man 646
Park and Krumwiede. Cultural characters of human and bovine t.b. 205;

rk and Krumwiede. Cultural characters of human and bovine t.b. 205; choice of media for differentiation 229; inconstancy of culture media 228; reaction curve of t.b. 236; adaptation 226 f; at what age are cultural differences most marked? 228; attenuation 288; types of t.b. concerned in human tuberculosis 549; in pulmonary T. (sputum investigations) 563; meningitis 571; general T. 578; abdominal T. 581; cervical glands 592; bones and joints 603. Inoculation of the rabbit 284; its value in determining types of t.b. 272

Parsons 58
Pasteur. Infection of sheep with anthrax spores facilitated by a diet of thistles 99 Pearson, Karl.

The tubercular diathesis 83, 84; marital infection 81; communicability of phthisis 58; heredity 58



INDEX OF AUTHORS

705

Petit. T. in the dog 487; in the cat 503 Petit and Basset. T. in the cat 504 Pfander 121 Pfeiffer and Friedbergher. Infection by feeding and inhalation compared 151 Philip, Dr R. W. 528
Plate. Absorption of t.b. from the intestine 161
Pomay. Epidemic of T. among fowls and turkeys which spared the geese and ducks 527 Pope. Marital infection 81 Porcher and Desoubrey. Portal. On latency 68 Bacterial content of dog's blood 154 Preyss. Infection by feeding and inhalation compared 150 Pybus. T. of the tonsil 590

Querli. T. among American Indians 77

Rabinowitsch. Attitude towards the question of types of t.b. 511; t.b. in apparently normal glands 73; types of t.b. in the monkey 511; virulence of avian t.b. for the monkey 517; attempts to cause lupus in the monkey with avian t.b. 651; passage of human t.b. through the rabbit 346; see also M. Koch and R.

Ransome Reliability of statistics of T. 10 ft periodic waves in the death.

nsome. Reliability of statistics of T. 10 f; periodic waves in the death-rate from phthisis 15 f; prevention of T. 112 Ransome.

Ravenel. Feeding experiments on dogs (penetrability of intestinal mucosa) 153, 159; bovine bacilli in human T. 189; exp. T. in the sheep 422;

in the horse 473
Raw, Dr Nathan. Immunizing influence of bovine infections 24 f Raw, Dr Nathan. Immunizing influence of bovine infections 24 f
Reichenbach. Infection by inhalation and feeding compared 151
Ritchie. Nature of miner's phthisis 103; frequency of t.b. in 102
Riviere, Dr Clive. Immunizing influence of bovine infections 24 f
Rivolta. Cultural peculiarities of avian t.b. (early recognition of) 187;
insusceptibility of the fowl to infection with human t.b. 540
Römer. Lesions caused by avian t.b. in the guinea-pig 438; passage experiments with human t.b. in the goat 317; experimental T. in mice 452
Roller. Therapeutic influence of sunlight 655
Roller 525

Roloff 525

Rosenberger. T.b. in normal glands 73
Rosenberger. T.b. in normal glands 73
Rothe. Proportion of deaths from T. among children IIf; outbreak of T. in a stock of tame rabbits 427; T. in man—relative frequency of infection in various lymphatic glands 170; pulmonary T. 554; meningitis 570; general T. 578; abdominal T. 582; T. of bronchial glands 584; T. in two or more groups of glands 588
Rothe and Bierotte. Types of t.b. found in lupus 648
Rush Dr. B. Contagiousness of phthisis 40, 50

Rush, Dr B. Contagiousness of phthisis 49, 50

Sand, Georges. Experience of the consumptive Chopin in Spain and Portugal

Sandilands, Dr J. E. Female occupation and child mortality in Kensington 95

San Felice. Avian t.b. in the ox 388
Schlossmann and Engel. Penetrability of mesenteric glands by t.b. 158
Schmid. Relative numbers of deaths from pulmonary and other kinds of T. in various countries 550 f

Schmorl and Geipel 134
Schroeder and Cotton. Healthy appearance of tuberculous cattle 377 f
Schroeder and Mietsch. Atypical t.b. 301 f; persistence of human t.b. in the udder of a cow 674
Schrum. Experimental T. in the dog

De Schweinitz. Attenuation of t.b. 288 Scott. Human manuscript.

Scott. Human mammary tuberculosis 384 f Scurfield. Explanation of the high F. to M. death-rate from phthisis in Scotland 87; on the ratio of F. to M. death-rate from T. in Sheffield 89;

© in this web service Cambridge University Press

www.cambridge.org



706

INDEX OF AUTHORS

mortality from T. in various towns 91; influence of flats on outdoor

exercise 95 Sehlbach. Frequency of infection in the first year of life 130

Shattock. Human mammary tuberculosis 384 f
Shattock, Seligmann, Dudgeon and Panton. Stability of type of mammalian
t.b. in the fowl 307; T. in the birds of the Zoölogical Gardens, London
526; no T. in the wild pigeon 528; instance of T. in a wild bird 527;
T. in the parrot 535; insusceptibility of fowls and pigeons to mammalian

T. 541, 542
Sibley. T. in the birds of the Zoölogical Gardens, London 526; no T. in

the wild pigeon 528; peculiarity of the lesions of T. in birds 529 f Siedamgrotsky. T. in cattle of Saxony 378 Silvius, Franciscus. First description of tubercles 47 Smith, G. Elliot and Ruffer. Tuberculosis in an Egyptian mummy 46 Smith, Theobald. Differentiation of human and bovine t.b. 189; reaction

curve 231, 234, 236, 238 Smith, Th. and Brown. Types of t.b. in cervical gland tuberculosis 591 Sprigge. Spread of infection from a case of phthisis in a dress-maker to her assistants 53 Steinmeyer. Relation of meningitis to general tuberculosis 570 f

Steinmeyer. Relation of meningitis to general tuberculosis 570 f Sticker. Experimental T. in the dog 501
Stockman. Experimental T. in the ass 478; T. in the dog 487; Johne's disease in the sheep 388
Straus. Pulmonary T. in the pig 401; expt. T. in the mouse 452; T. in the dog 487; canine T. mainly a pulmonary disease 489; a forecast of the type of t.b. found in lupus 615; portal of entry in lupus 615 f Straus and Gamaléia. Differentiation of avian and mammalian t.b. by means of the dog 246, 501; fate of avian t.b. in the guinea-pig 313; effect of avian t.b. injected into guinea-pigs 438; insusceptibility of the fowl to infection with mammalian t.b. 540
Stuurmann. Johne's disease believed to be caused by the avian bacillus 388—see also de Jong 559, 565
Summons. Miner's phthisis and pulmonary tuberculosis in S. Africa 106
Symmers. On the intestinal origin of phthisis 142; experimental anthracosis 146

Symmers. On the intestinal origin of phthisis 142; experimental anthracosis 146

Feeding expts. on dogs 492 Poliability of statistics 10 f; Tappeiner.

Tatham. Reliability of statistics 10 f; occupational mortality females 43 f; potter's phthisis 106

Templeman. Female occupation and child mortality in Dundee 95
Thomas, Dr Lynn. Therapeutic influence of sunlight 655
Thomassen. T. in the dog of a butcher 49
Thompson. Frequency of tuberculosis in the goat 414
Thorne Thorne. Subsoil drainage 24
Thompson. Female occupations of plague 47 occupational mortality among

Thucydides. Belief in the contagiousness of plague 47
Titze. Feeding expts. with avian t.b. on pigs 411; on a foal 475
Titze and Weidanz. Expts. on the dog 494, 499; exceptional result on a

feeding experiment 500 Todd. Views on development of the fœtus 50
Tonking. T.b. in the sputum of phthisical Cornish miners 102
Transvaal Medical Society. Fibroid phthisis and pulmonary tuberculosis

106; T.b. in the sputum of gold-miners 102 Treves. On the escape of the lymph glands in lupus 615 f

Trousseau 19

Trudeau. Attenuation of t.b. 288
Twort. Dead bacilli or their extracts as constituent part of culture media

Twort and Ingram. Cultivation of Johne's bacillus 216, 389

Ungermann. Proportion of deaths of children caused by T. 11f; frequency of infection in early life 130; relative frequency of infection of various groups of glands 170; exam. of the glands of a newly born infant 128;



INDEX OF AUTHORS

707

types of t.b. in the lymphatic glands of children in—pulmonary tuberculosis 555; meningitis 573, 576; general T. 587; abdominal T. 582; bronchial glands 585; two or more groups of glands 588; cervical glands 593; difficulty of preparing constant culture media 228; a strain of t.b. of human type which grew badly on broth 230

Vagedes 189 Vallée. Attenuation of a virulent strain of t.b. from the horse 288

Van Steenberghe and Grysez. Experimental anthracosis 146
Villemin 51, 52; first recognition of the difference between human and bovine tuberculosis 185; experimental T. in the dog 487; cat 505; doubt as to the existence of T. among birds 525
Viquerat. Exp. T. in the ass 478

Viseur. Exp. infection of the cat 505

Walsham. Tuberculous glands in phthisis 174; T. in the tonsil 175
Wartin and Cowie. Congenital tuberculosis 133
Watson, Sir Thomas 19, 50
Weber. Duration of life of cultures 221; passage of human t.b. through calf and pig 319; summary of cases of T. investigated up to 1907 at the Gesundheitsamt 553; abdominal T. 581; cervical gland T. 590; caseous material from glands may be not infective 71; value of rabbit for determination of type of t.b. 272
Weber and Baginsky. T.b. in normal glands 73
Weber and Bofinger. Transmission of infection through the egg 121; fate of t.b. injected into eggs 122; hereditary infection in avian T. 124 ft.

The Avian Tubercle Bacillus. Effect on the guinea-pig 439, 440; modification experiments in mammals 314; strain obtained from a pig 406; transmission of avian T. by mice 544. Mouse. Spontaneous T. 451; experimental T. 452, 457. Birds. Liability of lungs to attack 531
Weber and Dieterlen. Types of t.b. in sputum 562
Weber and Steffenhagen. An attenuated strain of t.b. from a human metacarpal bone 356, 629

carpal bone 356, 629
Weber and Taute. Fish-tubercle bacilli in frogs 302, 303; Types of t.b. in cervical gland tuberculosis 594; a strain of human t.b. from lupus 648
Weber and Titze. Infection by experimental feeding and inhalation compared 151; severe results following injection of human t.b. into calves 259; type of bacillus unchanged in these cases 320; milk infective after

immunization of heifer 675
Weber, Titze and Weidanz. T. in the parrot 538; in canaries 539; in

Weber, Titze and Weidanz. T. in the parrot 538; in canaries 539; in various small birds 540
Weber, Dr Hermann. Transmission of infection from husband to wife 52, 79; instance of quiescent T. reactivated by violence 80
Weichselbaum and Bartel. T.b. in apparently normal glands 73
Wheatley. Stone-worker's phthisis 103; not infective 100, 101
Whitla, Sir William. Intestinal origin of phthisis 142
Whitla and Symmers. Experimental anthracosis 146
Williams, Maxwell. Bovine T. in Edinburgh 94
Williams, Theodore. Latency 68; immunity of doctors and nurses in hospitals for consumption 75; intestinal origin of phthisis 142
Wilson, E. A. No T. in the wild grouse 528
Woodhead. The part played by the alimentary canal as portal of entry in phthisis 141; pulmonary T. secondary to glandular disease 175; spread of T. from gland to gland 175

Young, Thomas. On the contagiousness of phthisis 49 Young and Walker. Tuberculosis in cattle 376

T.b. from the pig 406; from the horse 471 Zundel 525

© in this web service Cambridge University Press

www.cambridge.org