

Cambridge University Press

978-1-107-45660-0- Rabies: Its Place Amongst Germ-Diseases, and its

Origin in the Animal Kingdom

David Sime

Index

[More information](#)

INDEX

- Absorbent corpuscles of circulation, 98, 119, 122, 125, 133, 134
- Acute or wholly paralytic rabies, 51—53, 208
- Aërophobia, 41
- Afferent or sensory system of nerves, fundamental seat of disturbance in rabies, 36—45
- Ague, 147—152; a recurrent, centric germ-disease, non-prophylactic, 148; morbid results of its virus, irrespective of fever, 116; significance of its characteristic periodicity, 149
- Ague of negro, 176, 177; as a final form, 181, 234, 235
- Alcohol, a typical virus, 106; morbid results of, when protecting from intoxication, 114, 116; graduated accumulation of, *sine quâ non* of alcoholic protection, 109, 110
- Alcoholic immunity, amount of, test of drinking habit, 107; its simultaneous morbid effects, 116
- Alcoholic protection, evanescence of, 111
- Alcoholism, 106—110; its alliance to rabies, 107—109; acute form, important half of germ-disease, 106; chronic or long-standing, protecting from intoxication, 114
- Amœboid cells, behaviour of, to toxic material, 163; vegetable, 162, 163
- Amœboid order of cell, 158, 162, 163; of connective-tissue, absorbing the micro-organisms of centric preventive germ-disease, neither as phagocytes nor as germinators, 121; its relation to pathogenic microbes, of a threefold character, 134; its phagocytic function, 158, 162; as cause of the external or exclusively peripheral germ-disease, 90, 100; of the prophylaxis of the prophylactic order of germ-disease, 139, 157, 168; of the ultimate attenuation of the excessively recurrent preventive germ-disease, 180
- Analgesia, with extreme prostration, fundamental feature of paralytic rabies, 36, 38, 39, 40, 41
- Anfractuous circulation of invertebrate, retained in vertebrate, kingdom, 145
- Annales de l'Institut Pasteur*; Metschnikoff's theory of the vaccinal element, 169; Pasteur's, 170
- Anthrax; Pasteur's research on, 101, 102; a germ-disease, prophylactic, 101, 139; its modified micro-organism, as eliminated from a gelatinous medium, 172; a triumph over the prophylactic order of germ-disease, 101; a vaccination, 101, 102, 172; in frog, cold-blooded and, artificially, warm-blooded, 134
- Anti-toxin or serum treatment, 159; its value in preliminary and intermittent stages of germ-disease, 131, 159; its failure, when used late, 160, 168
- Areolar spaces of lymphatic system, subcutaneous and subvenous, its limits, 145
- Ascending neuritis, rabies as an, 92, 94
- Ascending or intermittent paralytic rabies, 51
- Attenuating power in rabbits, possible consequences of its absence, 199
- Attenuating power of dog, 216, 218; of the monkey, 217; proof of intrinsic soundness and strength of central nervous substance, 240, 255
- Attenuation of rabies, 4, 84—87, 215
- Attenuators or attenuating division of

Cambridge University Press

978-1-107-45660-0- Rabies: Its Place Amongst Germ-Diseases, and its

Origin in the Animal Kingdom

David Sime

Index

[More information](#)

- animal kingdom, 5, 204—221 ; characterised by exceptional strength and integrity of nervous system, 239
- Attenuators' rabies, convulsive and convulso-paralytic, 208, 209 ; secondary and derived, rather than primary, 220, 235
- Attraction of phagocytes or amoeboid order of cell, succeeding repulsion, vital and permanent, 163, 164, 165
- Bacillary products as the cause of germ-disease, 106, 107
- Bacillus anthracis*, as a prophylactic micro-organism, modifiable, 101, 102
- Bacterial agency, in living tissues of plants, 191, 195 ; indispensable characteristic of vegetable and animal kingdoms, 181 ; in relation to allotropic forms and conditions, 185 ; to catalysis, 192 ; to the mineral as well as the vegetable kingdom, 185 ; to the removal of dead animal and vegetable matter in soil, 184
- Bacterial life, cause, direct or indirect, of the pre-digested pabulum of plants, and of their toxic principles, 185 ; in relation to the evolution of secreting organs, 179—203 ; more extrinsic or peripheral in plants than in animals, 181—191
- Bacterial zone of the soil, 186
- Bactericidal qualities of the serum of a protected animal, 159 ; may be occasioned by infinitesimal quantum of bactericidal material, 173 ; peculiar to the preventive order of germ-disease, 159 ; cf. also Germicidal
- Bark of furiously mad dog, diagnostic, 39
- Belladonna, 186
- Bell-wether ; his *raison d'être*, 242
- Berlin muzzling of dogs, its results, 262, 263
- Bert (Paul), on rabies-virus of salivary gland, 7 ; of the bronchial mucus, 9
- Bignami, on causation of ague, 148
- Bite of rabid dog not necessarily rabic, 2, 3, 11, 209
- Blood-stream and lymph-stream, the routes of conveyance of the rabies-microbe, 96
- Blood-stream of rabidised animal rarely rabic, 96 ; only in respect of an intensifier, 145
- Blood-system, two currents, its liquor sanguinis and its wandering amoeboid cells, 96 ; corpuscular system of, germinating-soil of prophylactic order of micro-organism, 96 ; phagocytic in respect of preventive order, 135, 162
- Bouchard, on rabic condition of lymphatic glands, 9 ; and lymphatic fluid, 145
- Boussingault, on incapacity of Papilionaceous plants to assimilate free nitrogen of air, 192
- Bradford case of hydrophobia, its extremely prolonged incubation period, 212
- Brown, Prof., on cessation of rabies in herbivora with death of victim, 224
- Brown-Sequard on rabies as "an ascending neuritis," 34, 45, 92
- Bulbar nerves supplying glosso-pharyngeal region, 45
- Bulbar nerve-tissue, theoretic and practical importance of, as a primary site, 28
- Bulbar rabies, 28, 29
- Bulbar substance, the most constantly implicated by rabies-microbe, at the end of incubation, 27, during the disease, 25, in death, 23, 24 ; containing the microbe in every stage of growth to decay, 124
- Bull-dog's tenacity of "pluck," 239
- Burdoni-Uffreduzzi of Turin, on rabic condition of pancreatic gland, spleen, and liver, 9
- Butterwort order of plants, its peptic power, 193
- Cancer, a peripheral and centric germ-disease of a connective-tissue nidus, 144
- Canine character and source of rabies, 204, 206
- Canine-rabies, 24, 223 ; convulsive in character and diagnostic of attenuators, 23, 24, 141, 142 ; extreme rarity of a wholly paralytic element in, and in that of attenuators, 2, 16, 18, 210, 211 ; extrinsic in source rather than intrinsic, 220 ; fading, final form, 181, 234 ; initial, originating form of rabies examined, 204—210 ; initial form of, of an outbreak paralytic and the most transmissible ; why, 238 ; not necessarily canine, 221 ; of country districts rather than of the towns, 262 ; canine or convulsive rabies more dangerous to the intensifying division than profoundest paralytic rabies, 223, 224 ; more phenomenal in rabbit than the completely paralytic rabies in dog, 233 ; aptitude for, contradistinct from that of rabbit or intensifiers,

Cambridge University Press

978-1-107-45660-0- Rabies: Its Place Amongst Germ-Diseases, and its Origin in the Animal Kingdom

David Sime

Index

[More information](#)*Index*

277

- 214, 215; serial transmission of, through rabbits, 198; signs and symptoms of, however prevalent, no test of its typical stability, 207; statistics of infective, of Battersea Dogs' Home, and of the London police authorities, 255
- Cat, the, as a hunter, 238
- Centricgerm-diseases, 137—141; preventive and prophylactic, 138, 139, 140; former, germinating exclusively in centric connective-tissue, 138, 141, 142; latter, exclusively in wandering, corpuscular system of circulation, 139, 140
- Centric nervous substance, specific seat of germination of rabies-microbe, 240
- Cerebro-spinal infection in completely convulsive, in convulso-paralytic, in completely paralytic rabies, 23, 25, 28, 53; cerebro-spinal system, as offshoot (differentiated) of sympathetic, 61
- Cervical site, the, seized by the hunting class, the most infective, 247
- Chemiotaxis, characteristic of amœboids of animal and vegetable kingdoms, 163
- Circulation, anfractuous, of invertebrate, retained in vertebrate, kingdom, 145; cf. also Absorbent corpuscles, Blood-stream, Blood-system
- Comité d'Hygiène, on variability of rabies-infection, 11, 33
- Completely convulsive rabies, bulbar in site, 50, 128, 209; non-infective by salivary secretion, 53, 210; viewed with tetanus, the sensory lesions of former, the motor of latter, 37
- Comte's metaphysical and theological modes of explanation, 205
- Connective-corpuscles of peripheral connective-tissue, absorbents of preventive micro-organisms, 121, 141, 144; media of conveyance of rabies-microbe, 96, 98, 125; not of germination, 133, 135, 137; the more they abound, the more favourable the site for infection, 144, 145
- Connective-tissue, the lower, or simpler in structure, the more favourable to the germination of the preventive order of micro-organisms, 153
- Connective-tissue, peripheral and centric, seat of germination of the entire preventive and recurrent order of germ-disease, 136, 141, 146, 152, 153; gelatinous composition of, 154
- Connective-tissue, composition of, of interstitial framework of centric organs, and of synovial membranes and sacs, 142
- Connective-tissue of spleen, as site of germination of malarial fever, 151; of cerebro-spinal substance, as that of rabies-microbe, 154
- Connective-tissue, the most lowly organised in animal frame, 153; the least injured by a virus, and the most fitting germinating soil, 153; the seat of germination of the preventive order of germ-disease, 137, 138, 144—147
- Conveyance of preventive micro-organisms, mode of, favourable to their deposit in the connective-tissue of centric structures, 144
- Conveyance of rabies-microbe, mode and route of, to cerebro-spinal substance from periphery, 90—98
- Convulso-paralytic canine-rabies, 208; its infective rate, 210
- Cope, on cessation of rabies amongst deer with the death of victim, 224; on deer-rabies at outset of Richmond Park epidemic, 225
- Croton-oil, as a product of plant-life of bacterial agency, 189; primarily, as a pathological, ultimately, as a physiological product, 190
- Darwin, on specialised sensitiveness of sundew, 202; on exceptional strength and stability of monkey's nervous organisation, 239
- Dead micro-organisms, after a germ-disease, or after preventive inoculations, as "the vaccinal matter," 169
- Dearth of germ-disease in animal kingdom, and of toxic principles in vegetable kingdom, where dearth of micro-organisms, 184
- Death inevitable in even slightest convulsive rabies; why, 23
- Death, sudden, of reinoculated dogs, but not from rabies, 114, 115
- Deer-rabies, 225, 226
- Deptford case of convulso-paralytic canine-rabies, 17, 261
- Descending neuritis, the character of rabies, 45, 94; its virus-germ being centrifugal, not centripetal, 94
- Desiccation of rabic nerve-substance kills, progressively, its rabies-germ, 124, 170

Cambridge University Press

978-1-107-45660-0- Rabies: Its Place Amongst Germ-Diseases, and its

Origin in the Animal Kingdom

David Sime

Index

[More information](#)

- Differentiation of specific plant-tissues for toxic principles with synchronous concentration of latter, 190
- Diminution in amount and potency of rabies-microbe during pre-incubation period, 95, 123, 124
- Diphtheria, a preventive germ-disease, 90, 91; of peripheral connective-tissue cultivation, 93, 135
- Diphtheritic membrane, 136
- Dispersal of rabies, its character, through canine-race, 222; through rabbit or intensifiers, 223
- Dog, as an attenuator of rabies, 208—228; contracts rabies, irrespective of race or breed, and of any intrinsic aptitude for the disease, 214; liability of, to rabies diametrically opposite to that of intensifiers, 214; its significance on origin of rabies in the animal kingdom, 215
- Dog, folly of summarily destroying a suspected, 2, 3, 4
- Dog, the, of the East, 255
- Dog-race, intrinsic liability to rabies of, examined, 215; cf. Mongrel-dogs
- Dog-race, bearing of, to rabies-microbe, contrasted with that of rabbit, 216, 218, 239
- Dowdeswell's investigation of canine-rabies, 16; on refractoriness of dog to rabies, 214, 219
- Dread after dog-bite, predisposing cause of intensification of hydrophobia, 3—5; the persistent feature of hydrophobia, 44
- Drosera, evolution of the secreting organ of, 193—195
- Drunkenness, its "mad" and "dead" varieties, 107
- Duboué, on rabies-virus in salivary glands, 7; on rabies as "an ascending neuritis," 92
- Dumb or paralytic canine-rabies, 39, 107, 208
- Duration of protection from rabies-virus, or from inoculative treatment, its character and mode of protection, 105, 111, 112
- Efferent or motor nerves, at their roots in spinal substance, special seat of disturbance in tetanus and in strychnine-poisoning, 40
- Elephantiasis and its connective-tissue element, 147
- Elimination of the virus of a pathogenic micro-organism, an irregular process, but final, 128; more or less sudden and simultaneous, 131; occurring neither during pre-incubation nor incubation periods, 125, 126, 127
- Elimination of bactericidal material in blood-stream, or establishment of a bactericidal condition of specific tissues, pathological, 174
- Elimination, ever-renewable, special character of a secreting organ, 191
- Environment, immediate, of plant-roots, natural habitat of micro-organisms, 183
- Erichsen, on fear of swallowing in hydrophobia, 44
- Erysipelas, a preventive germ-disease, 90, 93; recurrent, 117; of peripheral connective-tissue cultivation, 134, 135
- Evolution, exclusively on intrinsic modifying variability of living tissues, 195; of a pathogenic to a physiological micro-organism, that of rabies, 198, 200; to higher and complex organisation, from attainment of secreting organ, 202, 203
- Extension or migration of rabies-microbe through nerves, 45, 92; when it occurs, 94
- Face and neck, why exceptionally dangerous sites of rabies-infection, 34
- Farm-yards, badly drained stables, gardens, natural "home" of tetanus-bacillus, 91
- Fermentation, bacterial, 181
- Fibrocytes or connective-corpuses derived serially from vegetable cells, 158; less phagocytic with respect to preventive micro-organisms than leucocytes or lymphocytes, 136, 137, 144
- "Fixed," so-called, character of ordinary canine-rabies, 215, 218, 233, 234
- Fleming, Prof., on canine character and origin of rabies, 206; his statistical investigation of canine-rabies, 251, 252, 253
- Flügge, on tetanus-infection, 91
- Fol's discovery of the rabies-microbe, 86
- Foreign material in stomach at autopsy, 209; its meaning, 210
- Fox and the wolf frequently contract rabies, why, 263; and an "immune" animal resists the disease, 97, 157
- Fractional virus-germ of rabies, 92, 83; restored to its completed virus-germ on a second transmission through an intensifier, 83

Cambridge University Press

978-1-107-45660-0- Rabies: Its Place Amongst Germ-Diseases, and its

Origin in the Animal Kingdom

David Sime

Index

[More information](#)*Index*

279

- Frazer's, Surgeon-major, case of hydrophobia, prolonged incubation period, 211
- French Commission on Pasteur's experiments before them; the positive and the negative results disclosed, 12, 13
- Frost-bite, as a condition of tetanus, 90, 269, 270; as a possible condition of rabies, 272, 273
- Furious, or wholly convulsive, canine-rabies, 38, 209; abnormal to rabbit, 258
- Galtier, on rabies-virus in bucco-pharyngeal mucous membrane and in lingual glands, 9; in parotid and salivary glands, 7; on rabbit-rabies, 40; on its copious salivary secretion, 5, 41
- Galton, Sir Douglas, on "light from the sky" as bactericidal, 183
- Gamaleia, on rabies, as an ascending neuritis, 92
- Gardner's connecting-cells of a plant, 182, 186; as a specialised train, for conveyance of nutritious material and of toxic products, 187; and, very rarely, even of special micro-organisms, 194
- Gelatinous composition of connective-tissue, peripheral and centric; its significance in the germination of micro-organisms, 154
- Gelatinous substance, film of, around connective-corpusele, 144; possibly around red blood-corpusele, 144; its significance in absorption of the preventive micro-organisms, 144
- Germ-disease, of exclusively centric connective-tissue cultivation, alone possessing a pre-incubation period, 122; dearth of, in animal kingdom, 184
- Germ-diseases, either preventive or prophylactic in type, 99—118; of the Middle Ages mainly prophylactic; and their attenuation on transmission, prophylactic, 103; order of, to which rabies belongs, 99—117
- Germicidal principle in serum; as a product of the pathogenic micro-organism? 170, 171
- Germicidal property of plasma sanguinis and of specific tissues, 159, 160, 172; unproduced in dead tissue; inference, 172, 173; why germicidal, 168, 169, 173; cf. also Bactericidal
- Germinating-tissue of stubbornly recurrent preventive germ-disease, ultimately specialised into a nidus for the pathogenic micro-organism, 199, 200
- Germination, exclusively in fixed connective-tissue, characteristic of the preventive order, peripheral and centric, 139; exclusively in interior of the wandering absorbent corpuscles, characteristic of the prophylactic order, 121, 122, 139, 140; of rabies-microbe, exclusively in nervous substance, 19—32, 127; in the connective-tissue of nervous substance, 154; only after pre-incubation period, 127
- Gilbert, on incapacity of Papilionaceous plants to assimilate the free nitrogen of the air, 192
- Glandular system rabidised from centric nerve-source, 19; after, not before, incubation, 93
- Golgi, on the causation of malarial fever, 148
- Goulstonian Lectures* on "the chemical pathology of diphtheria," by Sidney Martin, 171
- Hellriegel micro-organisms in root-nodules of Papilionaceous plants, 192; their influence and significance, 192, 194
- Hemp, Indian, 182
- Hens and birds, their characteristic rabies, 16; of lumbar site and of intermittent paralytic character, 26, 130
- Herbivora, the intensifying division of the animal kingdom, 5, 198, 224, 226; their feebleness determining the intensifying property, 227, 228, 229, 230
- "Histology of the cell-wall," Gardner's research, 186
- Horsley's, Sir Victor, delineation of deer-rabies at end of epidemic, 225; his experiments with dog-virus on rabbits, 14; testing the incubation period of canine-rabies, 212; the special rabies of the Richmond Park deer outbreak, 223, 224; of the attendant Goffi who died of paralytic hydrophobia in spite of inoculative treatment, 245
- Humoralists, on germ-disease, 160 f.
- Hunted animal, why its nervous organisation so fitted for germinating the rabies-microbe, 240—245; and that of a hunting animal so ill fitted; yet rabies so prevails in the dog, 236—250; horror of, for the hunting, 242, 243, 244

Cambridge University Press

978-1-107-45660-0- Rabies: Its Place Amongst Germ-Diseases, and its

Origin in the Animal Kingdom

David Sime

Index

[More information](#)

- Hunted class of animal, comprising the entire herbivora, 236, 237; relationship of, to the hunting class of animal, 245—250; liability of, to rabies, without conspicuous outbreaks of the disease, 238
- Hunter, John, on refractoriness to rabies of dog, and of man, 10, 11; his estimate of the mortality of hydrophobia examined, 11, 12
- Hunting class of animals, attenuators of rabies, 237
- Hunting and non-hunting dog, 256, 257
- Huxley, on red blood-corpuscle, 149
- Hydrophobia, but a canine-rabies in man, 8, 207; a convulsive swallowing, 44; its fundamental lesion, a specific poisoning of the sensory side of the cerebro-spinal axis, 41, 42; of a strong-nerved man, a very different disease from the paralytic hydrophobia of a neurotic, 240; unamenable to inoculative treatment twelve days before the advent of the disease, 27, 250; dread after dog-bite, 3—5, 44; terror predominant feature of, 42; thirst in, 52; cf. also Canine-rabies
- Hyperæsthesia with intense excitability the fundamental feature of convulsive rabies, 36—39, 41, 43
- Hypertrophied connective-tissue of spleen in malarial fever, 151; of testicle in leprosy or tubercle, 147, 151, 153
- Hypertrophy of connective-tissue, increasing its adaptability for germinating the preventive order of micro-organisms, 151
- Identity, with specific character, of micro-organisms of the prophylactic and of the preventive order, 180
- Idiopathic hydrophobia, 269; tetanus, 91, 269
- Immune animal, or natural protection, 113
- Immunity, general, of the preventive germ-disease, contrasted with prophylaxis, 129—169; how induced, 130, 134, 168—175; contrasted with specific prophylaxis, 157
- Incubation of prophylactic and of peripheral preventive micro-organisms, immediate and regular, 121
- Incubation period and symptoms of convulsive and paralytic rabies, fundamental difference between, 38—43, 223; between tetanus or strychnine-poisoning and rabies, 133
- Incubation period of rabies, genuine, opposed to its pre-incubation period, 122; as normal and unvarying as that of tetanus, 122
- Incubation period of rabies and of centric preventive micro-organisms delayed and irregular, 127—131; extremely prolonged, 212; latency of, 127, 128; of ordinary canine-rabies, 74, 206, 214, 233; its irregularity in canine-rabies, 97, 211, 212, 213; of ordinary rabbit-rabies, 74, 223, 233; of hydrophobia, 211; inversely as the potency of the disease, 95; a law of germ-disease, 211; the Pasteurian period of germ-disease, iii
- Infection of cerebro-spinal axis during pre-incubation, and incubation periods, 20—35; of the bulb, or the superior or anterior tracts of cerebro-spinal axis, 23—28; of the lumbar swelling, or the inferior or posterior tracts, 26, 28, 32, 71; of the dog or the hunting animal in the later stages of incubation, 249, 250; risk of, from profoundest paralytic rabies, during latest stages of incubation, 28; sites of, 26, 34
- Infective rates of completely convulsive, of convulso-paralytic, and of completely paralytic rabies, 208, 209, 210; determined by amount of paralysis in case, 19, 210, 211
- Influenza maligna, as a recurrent germ-disease of the centric preventive order, 141
- Infusorial monads of sponge, 162
- Inoculation, intravenous, M. Pasteur's experiments, 96; in relation to the lumbar-swelling and to the development of intermittent paralytic rabies, 71; subdural, as test of incubation period, 212
- Inoculations, intracranial and intravenous, 212; the former, feeble in its results than intravenous or subcutaneous, why, 69—72
- Insectivorous plants, 193
- Insects, their venomous secretions, why, 195—197
- Instability of canine-rabies, 207—219
- Intensification of rabies on transmission, 4, 84—87, 222—233; through rabbits, methodic, progressive, and endless, 228,

Cambridge University Press

978-1-107-45660-0- Rabies: Its Place Amongst Germ-Diseases, and its

Origin in the Animal Kingdom

David Sime

Index

[More information](#)*Index*

281

- 231, 232, 233; table of, by its transmission through the rabbit, 233; laws of, 232; illustrations from M. Pasteur's investigation, 232, 233
- Intensifiers, the, or the intensifying division of the animal kingdom, 5, 222—235; their special rabies, paralytic, and of the cerebro-spinal and sympathetic systems, 28, 52, 53, 55
- Intensifying property, an infallible indication of intrinsic nerve-weakness, 240
- Intermittent character of rabies-saliva, 12, 14; of the rabies of hens and of birds, why, 26, 27, 51; similar to the intermittent character of ague, pyæmia and of many preventive germ-diseases, 26, 130
- Intermittent rabies, an ascending paralytic rabies of lumbar-swelling infection, 72
- Intoxication; cf. Drunkenness, Mad
- Jenner's vaccine, 118; his vaccination the *beau-ideal* of prophylaxis, 104
- Juniper "essence," 182
- Kelvin, Lord, on origin of living forms on earth, 183
- Kerslake, on the regions where rabies does not occur, 265
- Klein, on the occasional phagocytic power of the tubercle-bacillus, respecting its absorbent corpuscle, 152
- Koch, on causation of malarial fever, 148; on the spores of the paludal micro-organism in the intimate tissue of the spleen, 150; Koch's tuberculin, or the virus-elimination of the *Bacillus tuberculosis*, 67, 91, 106, 131; his tuberculin-test, as one of the most infallible in research, 107, 132, 147, 173; in blood-stream, as a cause, even in infinitesimal amount, of the elimination of the virus of germinating *Bacilli tuberculosis*, 131, its significance, 173; as an analogue of what occurs in the final outburst of every germ-disease, 131; in very graduated course, likewise protecting against tubercle, 110
- Kölliker, on the relation of the sensory or posterior nerves to the sympathetic ganglia, 61
- Lankester, Prof. Ray, on modifying influence of vaccine on small-pox, 105
- Latency of pre-incubation and incubation periods, why, 127, 128
- Laveran, on causation of ague, 148
- Lawes, on the incapacity of the Papilionaceous plant to assimilate free nitrogen of the air, 192
- Leopard, lion, tiger, or the giant *Felidae*, not genuine hunters, 238
- Leprosy, a germ-disease of a connective-tissue nidus, 847
- Leucocytes and wandering corpuscular cells of the highest animals, peculiar to the lowest planes of animal organisation, 163, 167; lineal descendants of vegetable cells, 158; seat of germination of the prophylactic micro-organism, 139, 162; phagocytic, even in respect of the prophylactic order of micro-organisms, after germination in their interior, 162; normally so in respect of the micro-organisms, which germinate exclusively in connective-tissue, 135, 162
- Leucocytic system, limitation of, as a cause of prophylaxis, 165, 166
- Liability of dog to rabies diametrically opposite to that of intensifiers, 214; its significance on origin of rabies in the animal kingdom, 215; of the hunted classes to rabies, without conspicuous outbreaks of the disease, 238
- Liability to a preventive germ disease increased after protection, 117
- Life, organic incorporation of primitive forms of, into one complex compound life within an organism, 199
- Ligament, tendon, bone, connective-tissue structures, 143
- Limitation of leucocytic system as a cause of prophylaxis, 165, 166
- Limited range of the prophylactic order of germ-disease over animal kingdom, its extreme significance, 103, 104; a clue to prophylactic investigation, 102, 119, 162
- Liquor sanguinis, a uniformly-flowing stream, 97; its influence in conveying the virus and the rabies-microbe to the cerebro-spinal axis, 96, 97; its bactericidal condition, after protection, 168—175
- Lister, Lord, on Gardner's researches in respect of "The histology of the cell-wall," 186; M. Pasteur's experimentation regarding the superabundance of micro-

Cambridge University Press

978-1-107-45660-0- Rabies: Its Place Amongst Germ-Diseases, and its

Origin in the Animal Kingdom

David Sime

Index

[More information](#)

- organisms in overcrowded life, vegetable and animal, 184; the invisible minuteness of many specific micro-organisms, 201; his antiseptic treatment, 118; in the suppression of pyæmia and germ-disease, particularly preventive, 146
- Living, not dead, tissues, constituting a germicidal condition, and eliminating a germicidal principle into the blood-stream, 173
- Local Government Board, report to, its definition of the protective and the preventive properties of the rabies-virus, 105; on the alleged averting of small-pox by vaccination, 106; on the prophylactic character of Pasteur's treatment, 104; on Pasteur's statistics of canine-rabies, 253
- Long's, Mr, influence in suppressing the canine-rabies of England, 220, 253
- Lords, report to House of, respecting the surgical treatment of rabic wounds, 254
- Lower or posterior extremities, a less important site of infection than upper or anterior, 26
- Lumbar-swelling, as a primary centre of the rabies-microbe, 24—28
- Lupus, 181, 137
- Lymphatic stream, an important relic in the vertebrates of the invertebrate circulation, 145; a medium of conveyance of rabies-microbe and of the preventive micro-organisms, 98; through its wandering connective-corpuseles, the most direct route of extended infection in preventive germ-disease, 137
- Lymphatic system; its fluid, even of an attenuator, never failing to rabidise, 145; observations of Galtier and of Bouchard, 9; cf. also Areolar spaces
- Mad drunkenness, 107
- Madness, canine, "furious" and "dumb," 107
- Magendie's transmission experiments of canine-rabies through dogs, 15, 216, 218
- Malarial fever, as a germ-disease of the spleen, 150; of the negro or of tropical natives, as a fading, final form, 181, 198
- Malarial micro-organism in blood-cells, Manson's chart, 148
- Mannaberg, Manson, and Marchiafava, on the causation of malarial fever, 148
- Manson, on the degenerating and degenerated forms of the paludal micro-organism in blood-cells, 149
- Marrow of bones, lymphatic glands and spleen, as the sources of the phagocytic leucocytes, 140
- Marrow of bones as osseous lymphatic glands, 166
- Martin, Sidney, on the chemical products of pathogenic micro-organisms, 171
- Measles, why prophylactic, 179
- Medulla oblongata* or "bulb," as the prime seat of germination, 24, 25
- Melun series of rabies - transmissions through rabbits, 229, 230
- Memory of pains soon forgotten, of pleasures lingers and expands, 241
- Metschnikoff's observations on *Daphne*, 134
- Microcosm of rabies as it exists in nature in the drying marrows of rabidised rabbits, 228
- Micro-organism, of every preventive germ-disease, unmodifiable, 101—106; law of the virulent results of a pathogenic, proportioned to its incubation, 211; law of the protection afforded by a virus, 109
- Micro-organisms; their agency in plants and animals, its enormous importance, 192, 203; in plants, extrinsic, 191; very rarely centric, or directly in plant-tissues, 191; pathogenic and physiological, 202; latter far more numerous than the former, 202; located in secreting organs, and at the source of their secretion, 201, 202, 203; associated with rabies-microbe in rabic saliva, 8; transmuting agency of, in mineral or inorganic as well as in organic substances, in gases and elements of the atmosphere, 184
- Micro-organisms of rabies, centrifugal in growth rather than centripetal, 94; equal, as monads, in germinating capacity, 81, 86; individually, unmodifiable, 86, 88, 101—106; pathogenic, but in the most persistent intensifiers ultimately may become physiological, through specialisation of germinating tissues and of micro-organism, 199
- Minuteness of the rabies-microbe, extreme, 87, 202

Cambridge University Press

978-1-107-45660-0- Rabies: Its Place Amongst Germ-Diseases, and its
Origin in the Animal Kingdom

David Sime

Index

[More information](#)*Index*

283

- Modified form of a prophylactic micro-organism, 167, 168; retaining the prophylactic property of the virus-germ, of which it is a modification, 84, 156
- Modifying character of the individual micro-organism, peculiar to that of the prophylactic order of germ-disease, 87, 88, 89
- Mongrel-dogs, as the source of rabies in the dog-race, 254
- Monkey of tropical Africa, a most potent attenuator of the rabies-microbe, 181, 217
- Monkey-rabies, a convulsive rabies, and as a final form of the disease, 235
- Morphia-solution, absorbed by root-tissues, conveyed to remotest structures, 183
- Mortification or necrosis, as a predisposing cause of tetanus, 90
- Mosquitoes, the mode of conveyance of the paludal micro-organism, or the means of infection of malarial fever, 148
- Mumps, an exceptionally prophylactic germ-disease, 166; why, 179
- Muzzle; in suppressing the canine disease; significance of the results, 220, in Sweden and in Germany, 262
- Natural preventive conditions, extrinsic and intrinsic, 10
- Nerve; fragments of, peripheral terminations, occasionally found rabid, 93; function of sensory, blended with that of sympathetic, 62
- Nerve-instability of the hunted class, 241; essential to the intensification of rabies, 245
- Nerves, cf. Afferent, Bulbar, Centric, Cervical site, Efferent, Peripheral, Pneumogastric, Sciatic, Spinal, Terminal
- Nervous substance, in the motor, the sensory, and the sympathetic systems, not alike sensitive to the virus of rabies, 54, 55
- Nervous system; its vital influence in specific centres on the leucocytic system of animals, 164; in the establishment of specific phagocytosis and in the production of prophylaxis, 164, 165, 167; condition of, determining the character of tetanus and rabies, 269, 270
- Neuroglia, a most primitive connective-tissue structure, 142, 143, 144; as specific site of germination of the rabies-microbe, 154
- Nocard's dialysis of rabid salivary secretion, 7, 98
- Normal, convulsive canine-rabies, 21, 214; as sensibly one in its virulence and germinating capacity, considered, 81, 82
- Normal hydrophobia, the convulsive rabies of the dog, 41
- Normal rabbit-rabies, 21, 40, 213, 231; the paralytic rabies of intensifiers, 27, 28, 40, 41, 233
- Normal tracts of migration of the rabies-microbe after infection, and after incubation, 93—95, 119—135
- Nux vomica, its strychnine, 187; of bacterial agency amidst root-tissues, 192
- Onsets, sudden, of virus, a series of, necessary for the full induction of rabies, particularly in its intermittent forms, 128; likewise characteristic of many preventive germ-diseases, 131, 159, 160
- Opium in the poppy as a product of bacterial agency, 190
- Organ in plant or animal, storing a virulent principle, specialised for storing, not elaborating it, 188
- Organism, plant or animal, perfected through suffering, 197
- Origin of consciousness and life, increasing mystery of, 183, 205; of elementary nervous system from attainment of secreting system, 202; of rabies in intensifiers, 251—268; of secreting-organ in plant and animal, 200, 202; of serpent-venom, through bacterial disease, 196
- Origin, spontaneous, of canine-rabies, considered, 204, 205
- Outbreak, extensive, of canine madness originating from a paralytic, not a convulsive, rabies, 218, 219
- Paludal micro-organisms in blood-cells, degenerated and degenerating, 149
- Paralytic element, importance of, over the convulsive element, even in canine-rabies, 19, 219; in rabies, surety of a rabid salivary secretion, 56
- Paralytic hydrophobia, 244
- Paralytic rabies, involves the cerebro-spinal and the sympathetic nervous systems, 50, 51, 128, 208; extremely rare in the

Cambridge University Press

978-1-107-45660-0- Rabies: Its Place Amongst Germ-Diseases, and its

Origin in the Animal Kingdom

David Sime

Index

[More information](#)

284

Index

- dog, 2; the most stable and the most capable of transmission through the dog of any form, 219; the initiating canine-rabies, 218, 219, 234, 237
- Paralytic, convulso-, canine-rabies of high infective rate, at Deptford and Pantin, 14, 17, 208
- Paralytic, profound, rabbit-rabies, 40; the most persistent and specific variety, 77, 230—233; its extreme severity masking it as a rabies, 260; its rarity in nature, 268, 273
- Parotid glands,—rabie, 7, 93
- Pasca's seasonal statistics of canine-rabies, 264
- Pasteur, on canine character and source of rabies, 201, 206; cerebro-spinal infection of canine-rabies, 23; constantly rabic condition of the bulb as compared with the salivary secretion, 20—35; why selected for experimental purposes, 35; on dog-rabies, as "the only rabies," 206; desiccation of rabic marrows, 170; fermentation, and spontaneous generation of rabies, 202, 204; intensifying property of the rabbit, 229—232; intermittent rabies of the hen, 26; intracranial, intravenous and subcutaneous inoculations, their difference in imparting the disease, 70; lumbar-swelling, as a primary seat of the intermittent paralytic rabies, 71; massive infection in blood-stream, with occasional protection, 129; monkey, indicating the animal's superlative attenuating power, 217; peripheral sites of infection, as determining factors of the sites of cerebro-spinal infection, 29, 30, 31; quantitative character of protection from rabies-virus inoculations, 109, 111; quantitative character of the rabies-virus-germ, 68; of intravenous and subcutaneous inoculations, as compared with the intracranial, 72; rabic condition of the blood, 9; its extreme rarity as compared with that of nervous substance or of salivary secretion, 9; rabbit, indicating the animal's superlative intensifying property, 228, 229, 230; refractoriness of the dog and the possibility of escape from infection, 14, 15; subdural inoculation as testing the incubation period of ordinary canine-rabies of the streets, 213, 214; variations in the length of incubation from diminution in an initial virus-germ, 78; various forms of rabies, induced by various amounts and sites of infection at the periphery, 29—33; various sites of the cerebro-spinal substance as seats of germination, 22—29
- Pasteur's discovery of the rabies-microbe, 87; his research, i—iv; initial experiments from salivary gland of a girl who died of paralytic hydrophobia, 7, 8; his treatment of anthrax that of a prophylactic germ-disease, 101; and of the entire realm of germ-disease that, either of a preventive or of a prophylactic order, 118; of rabies, as a germ-disease, of essentially prophylactic type, 101; being in reality that of a genuinely preventive germ-disease, 110, 118; his views of the germ-diseases "common to man and animals," 102; of the nature of the "vaccinal matter" in drying rabic marrows, 169
- Perfume of flowers and leaves, 185
- Peripheral infection, most regular, 30; nearness of, to cerebro-spinal axis the more effective the rabies induced, 69
- Peripheral terminations of nerves, as the starting-points of propagation of the rabies-microbe, 93, 96
- Persistency of recurrence, for many ages, of a preventive germ-disease, in the attainment of prophylaxis, 176, 180; or, failing prophylaxis, in the establishment of a secreting-organ, 199, 200, 201
- Persistency of virulence, test of a typical rabies, 76
- Pfeiffer's microbe of influenza, 87, 201
- Phagocytes, lineal descendants of amœboid order of cell in plants and in lowest planes of animal organisation, 158; attraction of, succeeding repulsion, vital and permanent, 163, 164, 165
- Phagocytic function common to the amœboid order of cell, 158, 162, 163; very special, 165—168; oldest function of animal organism, 158, 167; as a cause of prophylaxis, 166; of its persistent endurance, 166; of prevention itself, after prolonged recurrence of a preventive germ-disease, 175—177, 198—200
- Phagocytic prophylaxis as a probable ending of the malarial fever of the negro, 147;

Cambridge University Press

978-1-107-45660-0- Rabies: Its Place Amongst Germ-Diseases, and its

Origin in the Animal Kingdom

David Sime

Index

[More information](#)*Index*

285

- and a possible ending of rabbit-rabies, 198
- Phagocytosis or microbe-devouring, 157—159, 165, 167; characteristic of prophylactic order of germ-disease, 167; physiological, not pathological, 165; vital and permanent, not mechanical nor chemical, 163, 165
- Physiological accumulations of toxic products in a plant, fixed in amount and unaltering in character, 190
- Physiological micro-organisms, as ultimate evolution of the pathogenic, 201, 202
- Pitcher-plant, 193
- Plants, active principles of, significance of their elimination to deciduous structures, 182; insectivorous, 193
- Pneumogastric nerve, rabie, 93
- Pneumonia, a pulmonary erysipelas, 135
- Poppy, European and Indian, 186
- Power, on wounds favourable to tetanus, 90
- Predigestion by micro-organisms of animal and vegetable matter in soil, 184
- Pre-incubation period, importance of, for inoculative treatment, 97; and for research, iii
- Pre-incubation period of rabies, 119—127, 212; a process of conveyance by connective-corpuscles, 120; its varying duration, and why, 120—122; through the pre-incubation period no germination, 123, 124; no virus eliminated, 125; latency of, 127, 128
- Preventive conditions, intrinsic, of the animal kingdom, 11, 34; extrinsic, 10, 34
- Preventive germ-diseases, 99, 100; exclusively cultivated in fixed connective-tissue, 131—142; of external or peripheral connective-tissue cultivation, 134, 135, 136, 139; of universal (peripheral and centric) cultivation in connective-tissue, 137, 142, 162; of exclusively centric or internal connective-tissue cultivation, 141, 142; their primary habitat in badly-drained soils, 271; liability to, increased after protection, 117
- Preventive inoculations as potent after as before incubation, 105; their protection quantitative, 105, 110, 156, 157; unenduring, and essentially pathological, 174, 175; the germ-disease recurrent, 175, 177
- Preventive micro-organism, law regulating the germinating site of, and determining the special tissues which its virus deteriorates, 152
- Preventive order of germ-disease of widest range over the animal kingdom, 183; recurrent, 176; cf. Onsets
- Prophylactic order of germ-diseases, 100—116, 138; centric, never peripheral, 138, 139; limited range of, over animal kingdom, its extreme significance, 103, 104; a clue to prophylactic investigation, 102, 119, 162; deterioration of, through transmission for many ages, 179; of likewise the excessively recurrent preventive germ-disease, 180, 181; phagocytosis in both cases determining the attenuation, 181
- Prophylactic, as compared with the preventive order of civilised life, exceptionally disastrous amongst savages, why, 234
- Prophylactic property, a highly specialised function in leucocytes, 103, 163, 165, 167; permanent and vital, 165, 167
- Prophylactic treatment, clue to possibility of a, 102, 103
- Prophylactic virus-germ, simple, not quantitative, 88; individually, most modifiable, 89; may be modified into a vaccination, 89; the individual prophylactic micro-organism, compared with rabies-microbe, 88, and with micro-organisms of the preventive order, 102; prophylactic micro-organism, amount or site of, does not determine the character of the disease, 87, 112
- Prophylaxis, how occasioned, 140, 159, 162, 163, 167; characteristic of prophylactic order of germ-disease, 167; may be also induced by the continuous, recurrent action of a preventive virus, 176
- Protection from a prophylactic virus-germ, enduring character of, 112; the vital nature of its prophylaxis, 175
- Protection from rabies-virus, its evanescence, 111
- Protection of preventive class of germ-diseases, evanescence of, 111, 112
- Pugh on incapacity of Papilionaceous plant to assimilate the free nitrogen of the air, 192
- Pyæmia, a preventive germ-disease of uni-

Cambridge University Press

978-1-107-45660-0- Rabies: Its Place Amongst Germ-Diseases, and its Origin in the Animal Kingdom

David Sime

Index

[More information](#)

- versal connective-tissue cultivation, 137, 138, 146; site of infection in no way determines character of the disease, 137
- Quain's Anatomy*, on the bulbar nerves, and their origin in the bulb, 45; disparity of size between anterior and posterior roots of spinal cord, 48, 49; the motor roots and their freedom from ganglia, 59; the sensory roots and their incorporation with the ganglionic system, 59; the special connective-tissue structure of the interstitial framework of centric organs, of synovial membranes and sacs, and of valves, 60
- Quantitative character of rabies-virus, in the production of the disease, and of prevention; Pasteur's experiments, 64—80, 109—111
- Quantitative virus, not virus-germ, and of a preventive germ-disease, alone producing prevention, 156
- Quantity of the rabies-microbes of which it is composed constituting a virus-germ unit, 88, 89; this, in specific types of the disease, specialised and organised to its full possibility, 82, 84, 89
- Quantity of tetanus-bacillus in peripheral connective-tissue sufficient to tetanise, infinitesimal, 125; of rabies-microbe varying and various, 62, 64
- Rabbit, the best ground for the study of intensification, 229; the most potent intensifier, 222, 227, 228; possible consequences of absence of attenuating power in, 199
- Rabbit, the, in Australia, 265, 266, 267
- Rabbit-rabies, rarity of a wholly convulsive element in, or in that of intensifiers, 233; cf. Normal rabbit-rabies
- Rabbits, absence of rabies in countries without, 265
- Rabic condition of saliva, an indication of implication of the sympathetic system, 210; of lymphatic fluid in later stages of incubation, 145; intermittent character, 12, 14
- Rabic nerve-structures, comparative investigation of, 21—29
- Rabies, of the cat and of the fox, 244; a germ-disease of the country rather than of cities, 262; absence of, in countries without rabbits, 265; centric, not peripheral, in cultivation, 133; in the connective-tissue of the cerebro-spinal axis, or the neuroglia, 154, 155; malady of the hunted rather than of the hunting class of animals, 236, 237; of intensifiers rather than attenuators, 222—235
- Rabies, of the dog, convulsive, 205, 233; of the deer, convulso-paralytic, 224, 225, 226; the hen, intermittent and subacutely paralytic, 26; the rabbit, profoundly paralytic, 198, 222—235; the sympathetic system, always paralytic and infective, 50; characteristic of intensifiers, 28, 50—63
- Rabies in cerebro-spinal axis, anatomical classification of, 28; disclosures of bulbar centres of germination, 46; rabies from various amounts and sites of infection, comparative investigation of the forms of, 29—35; prevalence of a special form of, in attenuators, no test of its stability, 207; summary ending of outbreak of, in attenuators and intensifiers, 227, but for opposite reasons, 227; of the Middle Ages, theory of, 205
- Rabies or hydrophobia, analgesic or hyper-æsthetic, its lesion of sensory system underlying all lesions, 38—49; a descending neuritis from cerebro-spinal centres, 45, 46; an intoxication of nervous system like acutest alcoholism, 107; considered, as a preventive germ-disease, 100—112, 177; as a prophylactic germ-disease, 100—106; whether convulsive or paralytic, not necessarily canine, 219, 220, 257; order of germ-diseases to which it belongs, 99—117; cf. also Acute, Ascending, Attenuation, Attenuators', Bulbar, Canine, Completely convulsive, Convulso-paralytic, Dispersal, Dumb, "Fixed," Furious, Idiopathic, Instability, Intensification, Intermittent, Paralytic, Spinal
- Rabies-infection, external or peripheral sites of, the more important the nearer to bulkier sections of cord, 32—38; face and neck exceptionally dangerous sites of, 34
- Rabies-micro-organism, a microbe of living not of dead tissue, 124, 169; of nerve-structure, exclusively, 20, 92; sensory rather than motor side of the cerebro-

Cambridge University Press

978-1-107-45660-0- Rabies: Its Place Amongst Germ-Diseases, and its

Origin in the Animal Kingdom

David Sime

Index

[More information](#)*Index*

287

- spinal axis, 38, 47; its deterioration in pre-incubation period, 95, 123, 124; its elaboration of virus, when and where, 125, 126; its extreme minuteness, 87, 202; its moribund or degenerating condition in rabic saliva, 8, 124; in bulb, 124; its multiform structure as a virus-germ, 86, 98; speciality of the multiform virus-germ, the cause of the speciality of the typical varieties of the disease, 86—98; extension or migration through nerves, 45, 92; when it occurs, 94; medium by which borne to central nervous axis, 93, 94, 96, 98; by what nerve-tracts, from cerebro-spinal to sympathetic system, 59; amount of, in central nervous substance in convulsive and in paralytic rabies, 21, 25, 28; amount of rabies-virus due to amount of rabies-microbe, 70, 82; amount of rabies-virus, an animal capable of tolerating, and how far it can tolerate it, 113, 114, 115, 116; constancy of, in bulb, as compared with salivary gland, 23, 24; difference in composition, form, and force of, 83; rabies-microbe, or that of any centric preventive germ-disease, why not destroyed by the phagocytes of the circulation, 120; cf. also Conveyance, Virus-germ, Normal tracts
- Rabies-protection, graduated accumulation of rabies-virus, *sine quâ non* of, 109
- Rabies-virus, or product of the specific micro-organism, deadliest of ferments, 110; causing the disease, and all morbid changes, 106, 125; occasioning protection or prevention, 90—96, 108—112; the protection or prevention, like that from the virus of every preventive germ-disease, being pathological, 175; unending, 112, 174; with the disease recurrent, 112; morbid results of, which prevents or protects, 114, 115; cf. also Duration
- Rayleigh's, Lord, with Prof. Ramsay's, elimination of argon, 192
- Recurrence, persistent, of malarial fever, 175; of the preventive order of germ-disease, 117; why, 141, 142, 151, 175
- Red blood-cells, their possible origin, 149
- Reinoculation, periodic, absolutely necessary for universal prevention, 112, 113, 114; why impracticable, 115, 116; its ultimate effect, 116—119
- Repulsion from, with subsequent attraction towards irritants, of the amoeboid order of cell, 163; of highest animals, 164; vital, not chemical or mechanical, 165
- Restoration of an initial rabies, from a second transmission through rabbits of its fractional virus, 79, 84
- Rheumatism, acute, as a centric preventive germ-disease, 141; its recurrence, 146
- Richmond's, Duke of, Governor of Canada, hydrophobia from rabid fox-bite, 42; his dread of water, 44
- Richmond Park epidemic of deer-rabies, 224, 225
- Root-cells of plants, their absorptive power, 183
- Roots of bulbar and lumbar sensory nerves the nidus of the rabies-microbe, 49
- Ruffer on diphtheritic membrane, 136
- Saliva of paralytic rabies, liquid condition of, 10
- Salivary gland, considered as the specific seat of the rabies-microbe, 1—19
- Salivary secretion, absorbable in paralytic rabies, copious in amount, and rabic, 5, 17, 56; churned in convulsive rabies, diminished in amount, non-absorbable, and frequently non-rabic, 5, 56; earliest involved of any secretion, and the most seriously, why, 57, 58, 62; main channel for elimination of the rabies-microbe, 19; its rabies-micro-organisms in every stage of degeneration, 8, 124; its effectiveness due to amount of paralysis in the case, 17, 219; its non-effectiveness to the sparse amount of rabies-microbe, and exclusively located in the bulbar sites, 15, 210, 219
- Sanderson, Burdon, on fungoid spores and the leucocytes of *Daphne*, 134
- Sciatic nerve, rabic, 93
- Seat of germination, experimental determination of, in bulbar and lumbar substance, 24—29, in cerebro-spinal substance, 23—25, in sympathetic substance, 50—56
- Secreting-organs implicated in paralytic, seldom or never in completely convulsive rabies, any more than in tetanus or strychnine-poisoning, 52, 53
- Secreting-organs peculiarly characteristic of the animal organism, 195; rare in the extreme in the vegetable kingdom,

Cambridge University Press

978-1-107-45660-0- Rabies: Its Place Amongst Germ-Diseases, and its

Origin in the Animal Kingdom

David Sime

Index

[More information](#)

- 193, 195; attainment of, initial factor of evolution, 202
- Secreting-structures in plants and animals, in relation to bacterial agency, 195
- Secreting-tissues, fermenting foci; their secretion, a fermentation, 200, 201; their specialised tissue habitually inhabited by a specific micro-organism, constituting it a specific secreting-organ, 201
- Secretion of a secreting-organ, why unlimited in amount, and specific in character, 19, 60—64
- Secretions, the so-called, of the plant organism, 188
- Sensibility, disturbance of, fundamental feature of rabies, 37, 43
- Sensory, junction of, with the ganglionic sympathetic system; its importance in the production of acute, paralytic rabies, and in the implication of the glandular system, with infective forms of the disease, 52, 53, 55, 57, 62
- Sensory centres of bulb primary and very special seats of germination, 43, 44
- Sensory nerves, incorporation of, with sympathetic ganglia, 55—60
- Serpent-venom, how it originated, and how it is maintained, 195, 196, 197
- Serum, after a preventive germ-disease, bactericidal, 159; why, 168—177
- Serum treatment; cf. Anti-toxin
- Sharpey, on serous spaces or the “anfractuous cavities” of the lymphatic system, 145; on structure, connective-tissue, of serous sacs, membranes and valves, 143
- Sheep, clearing the gunwales of a steam-boat, why, 243
- Signs and symptoms of canine-rabies, however prevalent, no test of its typical stability, 207
- Soil, the natural habitat of micro-organisms, 183; bacterial zone of, 186
- Soil-life of the rabies-microbe, 271
- Solidists, the, on germ-disease, 160
- Sormani's discovery of the rabies-microbe, 89
- Specialisation of germ-disease to milder forms, through transmission, 179, 180; of particular tissues for storing toxic products, characteristic of vegetable kingdom, 189
- Spinal nerves rabic, 93
- Spinal rabies, 29
- Spleen, as a centre of the leucocytic system, 140; as the main seat of malarial fever, 150; softened and atrophied lymphoid element of, in malarial fever, 151, also of specific organs, infected with cancer, leprosy, tubercle, etc., 151
- Spontaneous generation, no explanation, 204, 219, 267
- Spontaneous protection, 130
- Spores of “the paludal parasites,” how they are so constantly conveyed to the intimate structure of the spleen, 120, 146, 150; of exclusively peripheral micro-organisms, occasionally conveyed to centric connective-tissue, 137, 146, 150
- Spores in the ultimate tissue of the spleen found at the earliest stages of malarial fever, 150
- Statistics of infective canine-rabies of Battersea Dogs' Home, and of the London police authorities, 255
- Stelling, on the anatomical origin of the bulbar nerves, 46, 47
- Sting of nettle; its relation to habitual bacterial agency, 187, 190
- Strain to the nervous system of the persistently hunted existence, 241, 244
- Strychnine of nux vomica, as a product of bacterial agency, 190
- Superinduced liability after any virus-protection, 116, 117
- Suzor, on convulsive, 38, and paralytic canine-rabies, 40; on Hunter's historic “instance” of hydrophobia, 11, 12; on statistics of the Comité d'Hygiène, 33
- Swallowing, act of, greatest agony of rabies or hydrophobia; why, 44, 45
- Sympathetic system, an important relic of the nervous system of the invertebrate kingdom, 60; less highly specialised than any part of the cerebro-spinal substance, 55, its relation to secretion and secreting-organs, 19, 55, 203
- Taste and the other special senses exaggerated in convulsive, perverted in paralytic rabies, 38, 39
- Terminal nerves, their rabic condition as compared with that of spinal substance, 94
- Terrier's indomitable courage, 239

Cambridge University Press

978-1-107-45660-0- Rabies: Its Place Amongst Germ-Diseases, and its

Origin in the Animal Kingdom

David Sime

Index

[More information](#)*Index*

289

- Terror, predominant feature of hydrophobia, 42
- Tetanus and strychnine-poisoning, compared with convulsive and paralytic rabies, 37
- Tetanus bacillary product and strychnine, 188
- Tetanus, a germ-disease of the country rather than of cities, 91, 270; extremely rife in villages of India, 188; its bacillus of peripheral connective-tissue cultivation, 90, 91, 93, 135; its site of infection in no way determining the character of the disease, 31
- Thirst in hydrophobia and convulsive rabies, excessive; why, 52
- Tobacco, habitual use of, its results, 164, 166; its specific protection, 167
- Toxic principles of a plant, fixed amount and unvarying strength, the special character of, 188, 191, 195; dearth of, in vegetable kingdom, where dearth of micro-organisms, 184
- Toxic plant-products and the viruses of pathogenic micro-organisms, specially neurotic poisons, 182; how conveyed to the remotest structures, 183; primarily pathogenic, ultimately physiological, even in plants, 189, 190, 191; why incapable of being stored in animals, 182
- Toxicity-tones, 161—168; from which prevention or protection, absolute, 109, 110, 174; chemical and pathological, not physiological, 168, 175; transient and evanescent, 175; with likewise, a condition of subsequent unprotection, also equally transient, 175
- Transmission, serial, through the dog, of ordinary convulsive rabies, even of profound paralytic rabies, progressively attenuated to extinction, 218; of canine-rabies through rabbits, 198
- Tubercle-bacillus, a micro-organism of peripheral and centric connective-tissue cultivation, 182; its latent stage, 131, 144
- Umbelliferous plants, why some deadly, some harmless and nutritious, 185
- Uniformity of symptoms, lesions, incubation and infective rate, characteristic only of the profoundest paralytic rabies, 219, 226, 230
- Unit-microbe of preventive germ-disease, unmodifiable; of prophylactic germ-disease pre-eminently modifiable, and capable of being transmuted into a vaccination, 84
- Vaccinal protection or prophylaxis due to phagocytosis, 156—162
- Vaccine, the modified micro-organism of small-pox, 101, 105; the prophylactic property of the original micro-organism fully retained, 106; its virulent property reduced to a minimum, 88, 89, 105; “vaccinal matter,” the, in rabies-inoculations, not the dead micro-organisms, 169; not the virus of the pathogenic micro-organism, 168, 169; what it is, 156, 160
- Variability of rabies in attenuators; of incubation and pre-incubation periods, 211, 212, 213, 214; of infective rate, 210, 211; of lesions, 209, 210; of ordinary canine-rabies from even intracranial inoculation, 212—214
- “Varieties” or modifications of ordinary canine-rabies of the streets, considerable, 214; from amount of infection rather than from innate difference of breed or race, 215
- Venom of serpent, physiological, that of rabid animal, pathological, but both of bacterial agency, 197
- Virus, elaborated at later stages of incubation-period proper, 127; outflow of, from incubating micro-organisms, a final series of phenomena, 128; eliminated, simultaneously and repeatedly, in sudden outbursts from the ripest germinating-sites, 130, 131; never in peripheral tissues, 67, 68; nor in itself bactericidal, 169; obtained, before Pasteur’s research, exclusively in salivary gland, 7; action of, continuous or excessive, on specific tissues, as cause of the germicidal condition of serum or tissues, 174; effect of the secretion of, on the process of germination, 127; graduated course of, protects, 107, 108, of rabies-virus, the *sine quâ non* of prevention and protection, 108, 109, 110, 126; cf. Onsets
- Virus-germ of rabies, also quantitative in infection, both in intensifiers and attenuators, 84, 85; various and varying in

Cambridge University Press
978-1-107-45660-0- Rabies: Its Place Amongst Germ-Diseases, and its
Origin in the Animal Kingdom
David Sime
Index
[More information](#)

- structure and composition, 85, 86 ; amor-
phous, 86 ; collective and multiform, 86 ;
fragmentary and completed, 83, 86 ; frac-
tional, 82, 83 ; restored to its completed
virus-germ on a second transmission
through an intensifier, 83 ; individually
unvarying, collectively most varying,
82—85 ; compared with virus-germ of
prophylactic and other germ-diseases, 85 ;
units of special, how constituted, 80—87 ;
cf. also Rabies-micro-organism
- Whooping-cough, 166, 179
- Wilfarth, on the micro-organisms of the
root-nodules of Papilionaceous plants,
192
- Yeast fermentation, analogous to virus-
elaboration in germ-disease, 106, 107 ;
and to the bacterial fermentation, oc-
curring incessantly in the soil amidst
plant-roots, 181—187
- Youatt on the symptoms of canine-rabies, 60