

Index

- absolute risk reduction (ARR), 57
- acceptability assessment, 118
- access to medicines, 292–3
- accountability, 25
- action efficacy, 136
- activities of daily living, 248
- adults, 227
 - determinants of health, 228
 - health promotion, 239–41
 - mortality and morbidity causes, 228–39
 - cancer, 229–30
 - cardiovascular diseases, 230–2
 - diabetes, 155
 - mental health disorders, 236–9
 - obesity, 234–6
- advocacy, 142, 143
- age
 - as an epidemiological variable, 48
 - at menopause, 72
- ageing
 - demography of old age, 245–6
 - end-of-life care, 254
 - health and, 246–9
 - health and social care, 252–3
 - prevention of age-related conditions, 249–52
 - dementia, 252
 - falls, 251–2
 - primary prevention, 249–50
 - secondary prevention, 250
 - tertiary prevention, 251
- ageing population, 35, 245–6
 - policy responses, 254–5
- AIDS, 291
- amniocentesis, 150
- analytical studies, 62, 64
 - interventional studies, 66–9
 - observational studies, 64–6
- aortic aneurysm, 151
 - screening, 155–6
- attitudes to work, 22
- attributable risk (AR), 58–9
 - smoking study, 59–60
- audit, 192–3
 - significant event audit, 196
- avoidable mortality, 41
- Beauchamp and Childress principles, 125
- behaviour change
 - in practice, 137–9
 - in theory, 135–7
 - nudge approach, 139
 - stages in, 136–7
- bias
 - in screening programmes, 159–60
 - information bias, 71–2
 - lead time bias, 160
 - length time bias, 160
 - selection bias, 71, 159–60
- body mass index (BMI), 234
- breast cancer, 148
 - screening, 151–2, 157
- burden of disease, 38–9, *See also* morbidity; mortality
 - cancer, 229
 - cardiovascular diseases, 230–1
 - diabetes, 232
 - global burden, 285–7
 - mental health disorders, 236–7
 - obesity, 234
- cancer, 229–30
 - registration, 84
- cardiovascular diseases, 134–5, 230–2
 - burden of disease, 230–1
 - policy, 232
 - prevention, 231–2
 - risk factors, 231
- Care Quality Commission (CQC), UK, 279
- case fatality, 53–4
- case reports, 63
- case series, 63
- case-control studies, 64–5
- causal factors, 74–5
- censuses, 39–40, 85
- cervical cancer, 148
 - screening, 150, 155–6
- chance, 70
- change
 - behaviour change in practice, 137–9
 - behaviour change in theory, 135–7
 - in public health practice, 25–7

- evaluation of effects of change, 100
 - implementation, 99–100
- innovation adoption, 19–20
- psychology of, 19–20
- theories of, 18
- child mortality, 209, 212, 214
 - global issues, 287–8, 289
- child population, 211–12
- child public health, 209
 - challenges, 213
 - determinants of child health, 213–19
 - behaviour and lifestyle, 218–19
 - child care, 216–17
 - ethnicity, 215
 - family structure, 216
 - individual and congenital factors, 215
 - media, 219
 - parenting, 215–16
 - peer relationships, 217
 - physical environment, 218
 - poverty and deprivation, 217–18
 - public services, 219
 - social capital, 218
 - global issues, 287–9
 - health policy, 224
 - health promotion, 219–21
 - obesity case study, 220–3
 - mortality and morbidity causes, 212, 214
- Chlamydia trachomatis*, 166
- chorionic villus sampling (CVS), 150
- citizens' juries, 110
- climate change, 304, *See also* sustainable development
 - health co-benefits of action, 305
- clinical governance, 194–5
- clinical leadership, 24–5
- clinical trials, 67–8
 - randomised controlled trials (RCTs), 66
- Cochrane Database, 95
- cohort studies, 65–6
- communicable disease control, 165–72
 - global issues, 290–1, 292
 - outbreak investigation, 174
 - principles, 164–5
 - surveillance, 172–4
 - transmission control, 166–9
 - vaccination, 169–72, 173
- community development, 141
- community trials, 68–9
- confidence interval (CI), 70
- confounding, 72–4
- confounding variables, 64
- Consumer Assessment of Healthcare Providers and Systems programme, USA, 193
- cost effectiveness assessment, 119–26
- cost utility analysis, 121
- cross functional teams, 22
- cross-sectional studies, 63
- data, 79
- death rate, 79–81, *See also* mortality
 - directly standardised death rate (DSR), 80–1
 - standardised mortality ratio (SMR), 80–1
 - under-5 mortality rate, 82
- decision making, 116, *See* funding decisions
- dementia, prevention, 252
- Demographic and Health Surveys (DHS) Project, 40
- demographics
 - disease burden, 38–9
 - fertility, 31–2
 - health transitions, 36–8
 - life expectancy, 34–5
 - methodological issues, 39–41
 - migration, 33–4
 - mortality, 32–3
 - old age, 245–6
- demography, 29
 - population structure, 30
 - population trends, 31
- deprivation, children, 217–18
- descriptive studies, 62, 63
 - individual studies, 63
 - population studies, 63
- desired health outcomes, 184–5
- diabetes, 155
 - burden of disease, 232
 - prevention, 233
 - risk factors, 233
- Diabetes Control and Complications Trial (DCCT), 68
- diabetes screening, 155–7
- diabetic retinopathy, 149
- direct costs, 119
- directly standardised death rate (DSR), 80–1
- Directors of Public Health, 6
- disability, dimensions of, 247
- disability-adjusted life years (DALYs), 38–9, 287
- disaster planning and response, 179–81
- disease burden., *See* burden of disease
- disease prevention., *See* prevention
- disease-specific approaches, 296
- Down's syndrome screening, 150
- Duncan, William, 1–2
- ecological fallacy, 63
- economic evaluation, 119–23
- education, 192–3
- effectiveness assessment, 117–18, *See also* cost effectiveness assessment
 - effectiveness assessment
- emergency planning and response, 179–81
- emotional intelligence, 15
- end-of-life care, 254
- environmental public health, 177–9
- epidemiology, 43–4
 - comparing risks, 54–7
 - concepts, 61
 - disease frequency measures, 49–53

- epidemiology, (cont.)
 - incidence, 50–3
 - prevalence, 50
 - rates, 49
 - history, 44–5
 - interpretation of study results, 69–75
 - chance, 70
 - error, 70–4
 - validity, 74–5
 - number needed to treat (NNT), 57–8
 - population concept, 46–7
 - population impact measures, 58–9
 - smoking study, 59–60
 - types of studies, 62–9
 - analytical studies, 64
 - descriptive studies, 63
 - study designs, 62
 - variables, 47–9
- error, 70–4
 - random error, 71
 - systematic error, 71–4
- ethnicity, child health and, 215
- evidence-based health-care (EBHC), 90–1
 - appraising the evidence, 96–9
 - asking a focussed question, 91–2
 - assessing evidence of effectiveness, 117–18
 - changing practice, 99–100
 - evaluating the effects of change, 100
 - limitations, 100–1
 - tracking down evidence, 92–6
- evidence-based medicine (EBM), 91
- external validity, 74
- falls, prevention in old age, 251–2
- family structure, child health and, 216
- Farr, William, 44
- fertility, 31–2
 - estimation, 40
- Finagle's law, 78
- financial incentives, 100
- financing health systems, 294–6
- focus groups, 110
- followers, 18
- functional teams, 21
- funding decisions, 115–16
 - acceptability assessment, 118
 - assessing evidence of effectiveness, 117–18
 - assessing need, 116–17
 - cost-effectiveness assessment, 119–26
 - economic evaluation, 119–23
 - ethical approaches, 124–6
 - identifying the proposed service, 116
 - value-based decisions, 123–6
- gender inequalities, 297
- General Household Survey, UK, 85
- germ theory, 2
- global issues
 - access to medicines, 292–3
 - barriers to health-care access, 297–8
 - burden of disease, 285–7
 - communicable diseases, 290–2
 - health improvement, 294–7
 - delivery, 293–4
 - organising and financing health systems, 294–6
 - people-centred or disease-specific services, 296–7
 - maternal health, 288, 290
 - measuring progress, 298–300
 - non-communicable diseases, 291–2
 - poverty, 286
 - poverty reduction priorities, 300
 - preventable illness and death among children, 287–9
 - reproductive health, 288–90
- governance, 294
- Graunt, John, 44
- handicap, 247
- health, 139–40
 - ageing and, 246–9
 - causes, 308–11
 - knowledge use by health professionals, 311–12
 - poverty relationship, 285
- health-care
 - in old age, 252–3
 - problems with, 188–9
- health inequalities, 257–8, 280–2
 - approaches, 268–9
 - causes, 259–64
 - access to services, 264
 - material circumstances, 260–2
 - social status, 77–8
 - gender inequalities, 297
 - goals of public health, 265
 - implications for public health practice, 258, 269–70
 - life course perspective, 265–6
 - lifestyle and behaviour, 263–4
 - policy approaches, 264–5
 - responsibility for action, 266–8
- health needs assessment, 108–14
 - agreeing local priorities, 110–11
 - comparative needs assessment, 109–10
 - epidemiological needs assessment, 108–14
 - objectives, 109
 - planning and strategy development, 111–13
 - theoretical perspectives, 104–7
 - health or health-care, 105
 - individual or population, 105
 - need, supply and demand, 105–6
- health policy., *See* policy
- health profile, 78
- health promotion, 139–44
 - adults, 239–41
 - child health promotion, 219–20, 221
 - obesity case study, 220–3
 - community development, 141

- health education, 140–1
- healthy public policy, 141–2
- legislative action, 141
- mental health, 238–9
- responsibility for, 142–4
- tobacco control programme, 140
- workplace case study, 239–41
- health protection, 163–4, 281–2
 - communicable disease control, 165–72
 - principles, 164–5
 - emergency planning and response, 179–81
 - environmental public health, 177–9
 - outbreak investigation, 174
 - surveillance, 172–4
- Health Survey for England, 85
- health transitions, 36–8
- health-related social marketing, 138
- hip fracture, 251–2
- HIV, 291
- horizontal teams, 22
- hospital episode statistics (HES), 83–4
- human resources, 297–8
- Illich, Ivan, 2
- immigrant screening, 152
- immunisation, 2
- impairment, 247
- incidence, 50–3
 - prevalence relationship, 53
- incidence rate, 50–1
- incremental costs, 120
- Index of Multiple Deprivation, 86
- indirect costs, 119
- individual studies, 63
- inequality measures, 86–7, *See also* health inequalities
- infant mortality, 81
- infectious diseases, 166, *See also*
 - communicable disease control
 - drug resistance, 168
 - morbidity statistics, 83
 - transmission control, 166–9
 - vaccination, 169–72
- influenza pandemic preparedness, 181
- information bias, 71–2
- innovation adoption, 19–20
- instrumental activities of daily living, 248
- internal validity, 74
- International Classification of Diseases and Related Health Problems (ICD), 44
- interventional studies, 62, 64, 66–9
 - clinical trials, 67–8
 - community trials, 68–9
- interviews, 110
- job satisfaction, 22
- Joint Strategic Needs Assessments, 87, 117
- knowledge, 3
 - use by health professionals, 311–12
- lead time bias, 160
- leadership, 15–18
 - clinical leadership, 24–5
 - in public health practice, 25–7
 - styles, 16–17
- lean technique, 195
- legislation, health promotion, 141
- length time bias, 160
- life expectancy, 34–5, 82
 - income relationship, 35
 - medical care contribution, 3–4
- life tables, 34–5, 41
- lifestyle factors
 - children, 218–19
 - health inequalities, 263–4
- likelihood ratios, 157
- Lind, James, 44–5
- literature searches, 94–6
- local health surveys, 86
- lung cancer, 71–2
- management, 13–14
 - clinicians' role, 24–5
 - in public health practice, 25–7
 - tasks, 14
- marginal costs, 120
- market incentives, 277–8
- Marmot Review, 266, 268, 280
- maternal health, 288
- maternal mortality, 33, 81, 288, 290
 - Pakistan, 69
- McKeown, Thomas, 2–3
- media influence on child health, 219
- medical care contribution, 3–4
- Medline, 95
- menopause, age at, 72
- mental health, 236–9
 - burden of disease, 236–7
 - policy, 239
 - prevention of disorders, 238–9
 - risk factors, 237–8
- meta-analysis, 98
- migration, 33–4
 - measurement, 41
- mild cognitive impairment (MCI), 252
- millennium development goals (MDGs), 298–300
- Monitor, 279
- morbidity, 83–4, 231, *See also* burden of disease
 - adult morbidity causes, 228–39
 - child morbidity, 212
 - global issues, 287–8
 - statistics, 83–4
- mortality, 32–3, *See also* burden of disease; death rate
 - adult mortality causes, 228–39
 - avoidable mortality, 41
 - case fatality, 53–4

- mortality, (cont.)
 - child mortality, 209–12, 214
 - global issues, 287–9
 - infant mortality, 81
 - maternal mortality, 33, 81, 288, 290
 - Pakistan, 69
 - measurement, 40–1, 78–82
 - neonatal mortality, 81
 - perinatal mortality, 81–2
 - Pakistan, 69
 - post-neonatal mortality, 81
 - proportional mortality, 81
 - smoking relationships, 59–60
 - standardised mortality ratio (SMR), 80–1
- motivational needs, 24
- National Child Measurement Programme (NCMP), UK, 220
- National Institute of Health and Clinical Excellence (NICE), UK, 279
- National Patient Safety Agency (NPSA), UK, 193
- natural history of disease, 129–30
- needs, 104–5
- needs assessment, 116–17, *See also* health needs assessment
- neonatal mortality, 81
- new public health movement, 3
- NHS, 6–7
 - structural reform, 278–9
- nicotine gum, 55–6
- notifiable diseases, 174
- number needed to treat (NNT), 57–8
- obesity, 234–6
 - burden of disease, 234
 - childhood, health promotion case study, 220–2, 223
 - effects on health, 235
 - policy, 233–6
 - prevention, 235–6
 - risk factors, 234–5
- observational studies, 64–6
 - case-control studies, 64–5
 - cohort studies, 65–6
- Odds Ratio, 54–5
- old age., *See* ageing
- opportunity costs, 119
- oral contraceptives, pulmonary embolism
 - relationship, 65
- organisational behaviour, 20–4
- organisational culture, 21
- organisational types, 21
- outbreak investigation, 174
- P* value, 70
- Pakistan, maternal and perinatal mortality, 69
- panels, 110
- parenting, child health and, 215–16
- payment for performance, 194
- peer relationships, children, 217
- perinatal mortality, 81–2
 - Pakistan, 69
- period prevalence, 50
- PESTLE analysis, 18
- PICO approach, 92
- Plan-Do-Study-Act (PDSA) cycle, 192
- planning, health needs assessment and, 111–13
- point prevalence, 50
- policy, 273
 - ageing population and, 254–5
 - cancer, 229–30
 - cardiovascular diseases, 232
 - child public health, 224
 - context for policy making, 274–5
 - England, 276–9
 - developments, 279–82
 - government priorities, 280
 - market incentives, 277–8
 - NHS structural reform, 278–9
 - objectives, 281
 - health inequalities, 264–5
 - health promotion, 141–2
 - implementation, 275–6
 - meaning of, 273–4
 - mental health, 239
 - obesity, 233–6
- population, 46–7
 - child population, 211–12
 - health status, 77–8
 - inequality measures, 86–7
 - local health surveys, 86
 - morbidity measurement, 83–4
 - mortality measurement, 78–82
 - population surveys, 85
- population attributable risk, 59
- population censuses, 39–40, 85
- population growth, 287
- population structure, 30
- population studies, 63
- population surveys, 40, 85
- population trends, 31
 - fertility, 31–2
 - life expectancy, 34–5
 - migration, 33–4
 - mortality, 32–3
- post-neonatal mortality, 81
- poverty, 286
 - child health and, 217–18
 - health inequalities and, 260–2
 - health relationship, 285
 - poverty-reduction priorities, 300
- power, 71
 - sources of, 16
- predictive values, 156–7
- prevalence, 50
 - incidence and disease duration relationship, 53
- prevention, 128–9
 - age-related conditions, 249–52

- dementia, 252
- falls, 251–2
- primary prevention, 249–50
- secondary prevention, 250
- tertiary prevention, 251
- behaviour change in practice, 137–9
- behaviour change in theory, 135–7
- cancer, 229
- cardiovascular diseases, 231–2
- diabetes, 233
- in clinical practice, 133–5
- levels of, 130–1
- mental health disorders, 238–9
- obesity, 235–6
- strategies, 131–3
- prevention paradox, 132
- primary prevention, 130
- professionalism, 24–5
- programme budgets, 120
- proportional mortality, 81
- proportionate universalism, 268
- prostate cancer, 152
- public health, 6
 - child public health, 209
 - competence development, 312–13
 - core functions, 295
 - domains of, 4–6
 - environmental public health, 177–9
 - historical background, 1–3
 - new public health movement, 3
- Public Health England, 6, 282
- pulmonary embolism, oral contraceptives relationship, 65
- qualitative methods, 93–4
- quality-adjusted life years (QALYs), 82, 121
- disadvantages, 122
- Quality and Outcomes Framework (QOF)
 - payment system, 100
- quality of care, 183–4
 - definition, 184
 - dimensions of, 185
 - evaluation, 185–8
 - improvement, 190–6
 - clinical governance, 194–5
 - education and audit, 192–3
 - lean technique, 195
 - market-based approaches, 193–4
 - regulation and standards, 190–1
 - root-cause analysis, 196
 - significant-event audit, 194–6
 - six sigma, 194–5
 - measurement, 184–5
 - problems with, 188–9
- question, 91–2
- questionnaire surveys, 110
- random error, 70, 71
- randomised controlled trials (RCTs), 66, 93
 - appraisal, 96
 - rapid appraisal, 110
- registration of births and deaths, 40
- regulation, 190–1
- relative risk, 54–7
 - smoking study, 59–60
- reproductive health, 288–90
- resource distribution, 305–6
- risk, 50
 - attributable risk (AR), 58–9
 - case fatality, 53–4
 - comparing risks, 54–7
 - population attributable risk, 59
- risk factor thresholds, 133–5
- root-cause analysis, 196
- Rose, Geoffrey, 131
- sampling error, 71
- screening, 132–3, 146–7
 - criteria for programme establishment, 147–52
 - the condition, 148–9
 - the programme, 151–2
 - the test, 149–50
 - the treatment, 150–1
 - harms and benefits, 147
 - monitoring programmes, 159–60
 - sources of bias, 159–60
 - other forms of screening, 152
 - screening test performance, 154–8
 - likelihood ratios, 157
 - predictive values, 156–7
 - sensitivity, 154–6
 - specificity, 154–6
 - test outcomes, 154
 - successful UK programmes, 152–3
- scurvy, 45
- secondary prevention, 131
- selection bias, 71
 - in screening programmes, 159–60
- self-directed teams, 22
- self efficacy, 136
- sensitivity, 154
 - calculation, 154–6
- significant event audit, 194–6
- six sigma, 194–5
- smoking
 - cessation, 55–6
 - mortality relationships, 59–60
 - tobacco control programme, 140
- Social Action Research Project (SARP), 269
- social capital, 218
- social care in old age, 252–3
- social exclusion, 297
- social marketing, 138
- specificity, 154
 - calculation, 154–6
- standardised mortality ratio (SMR), 40, 80–1

standards of care, 190–1	teams, 21–2
stillbirth rate, 81	tertiary prevention, 131
strokes, 132	therapeutic era, 2
surveillance, 172–4	thrombolysis, 98–9
survival rates, 53–4	tipping point, 20
sustainable development, 304–8	transactional leadership, 15
food, 305	transformational leadership, 15
framework for action, 306–8	
health co-benefits of action, 305	under-5 mortality rate, 82
public health role, 308–13	user consultation panels, 110
causes of health, 308–11	
competence development, 312–13	vaccination, 169–73
knowledge use by health professionals, 311–12	validity, 74–5
redistributing resources, 305–6	value-based decisions, 123–6
travel, 305	vertical teams, 21
systematic error, 70–4	
confounding, 72–4	Whitehall study, 66, 264
information bias, 71–2	whole-systems approach, 280
selection bias, 71	workplace health promotion, 239–41
systematic reviews, 96–9	
	years of life lost (YLL), 82