

CONTENTS

CHAPTER XII

COMPOUND MATRICES

§§		PAGES
98–99.	Compound matrices; constituent matrices. Special notations for scalar, quasi-scalar, zero and one-rowed matrices; matrix one of whose orders is zero. Multiplication of compound matrices	1–6
100	Compartite matrices: parts; compartite matrix in standard form; successive parts; the rank of a compartite matrix is the sum of the ranks of its parts; special notations for a compartite matrix whose parts are quasi-scalar matrices	6–8
101.	The conjugate reciprocals and inverses of certain matrices. Some special square matrices	8–13
102.	The primaries of a minor determinant; horizontal and vertical primaries; number of primaries; notation for primaries. Primary subdeterminants and primary superdeterminants	13–18
103.	Elimination of a variable from a system of inequalities	18–19
104–107.	Possible ranks of a matrix containing a given minor matrix; of a minor of a matrix whose rank is given; rank of a matrix which contains an undegenerate simple minor formed by the addition of 0's to an undegenerate square matrix. Diagonal minors. Possible ranks of a symmetric matrix containing a given diagonal minor; of a diagonal minor of a symmetric matrix whose rank is given	19–36

CHAPTER XIII

RELATIONS BETWEEN THE ELEMENTS AND MINOR DETERMINANTS OF A MATRIX

108.	Connections between the short rows of an undegenerate matrix: identities which serve to express every element of the matrix as a rational function of degree 1 of any one regular simple minor determinant Δ and the elements and primaries of Δ ; the identities for a matrix whose orders differ by 1. Connections between the rows of a degenerate matrix	37–46
------	--	-------

CONTENTS		xiii
§§		PAGES
109.	Relations between simple minor determinants: identities which serve to express every simple minor determinant D of an undegenerate matrix as a rational function of degree 1 of any one regular simple minor determinant Δ and the primaries of Δ ; the standard identities; ascription of arbitrary values to the elements and primaries of any regular simple minor determinant; utility of these identities; determinants of the primaries of a simple minor determinant. Corresponding identities for superior simple minor determinoids	46-66
110.	<i>Sylvester's</i> identities: identities which serve to express every superdeterminant D of a regular minor determinant Δ as a rational function of degree 1 of Δ and the primary superdeterminants of Δ ; determinants of primary superdeterminants	66-70
111.	Identities which serve to express every subdeterminant D of a regular minor determinant Δ as a rational function of degree 1 of Δ and the primary subdeterminants of Δ ; determinants of primary subdeterminants	70-73
112.	Relations between simple minor determinants: identities which give the sums of terms formed from a product ΔD of two simple minor determinants by replacing s rows of Δ by rows of D , and the s rows taken from D by s rows of Δ	73-75
113.	Equivalence of two similar undegenerate matrices: two such matrices are mutually equivalent when and only when their correspondingly formed simple minor determinants are proportional; sign of equivalence	75-78
	Spacelets of homogeneous space represented by undegenerate matrices	78-80
114.	Criteria for the equivalence of two systems of linear algebraic equations. Mutually orthogonal and mutually normal undegenerate matrices and spacelets.	80-85
115.	Relations between the elements of a matrix of rank r : equations which serve to express every element as a rational function of degree 1 of any one regular minor determinant Δ of order r and the elements and primaries of Δ	85-89
116.	Identical relations between the elements of any matrix: the fundamental identity which serves (<i>see</i> Appendix A) to express every element of the matrix as a rational function of any one regular minor determinant Δ and the elements, primaries and primary superdeterminants of Δ ; deduction of all previously obtained relations from the fundamental identity; factorisation of a matrix. Necessary and sufficient conditions that a matrix may have rank r . Regular subdeterminants and superdeterminants of a regular minor determinant; standard arrangement of the rows of a matrix. Rank and other properties of a matrix of primary superdeterminants	90-98
117.	Relations between the minor determinants of order r of a matrix whose rank is r : equations which serve to express every minor determinant of order r as a rational function of degree 1 of any one regular minor determinant Δ of order r and the primaries of Δ	98-106

§§		PAGES
117 <i>a.</i>	APPENDIX A. Utility of the relations obtained in Chapter XIII : ascription of arbitrary values to the elements, primaries and primary superdeterminants of any regular minor determinant Δ ; if the order of Δ is r , and if the matrix of its primary superdeterminants has rank x , then the matrix has rank $r+x$. Summary of the relations of Chapter XIII	515–520
 CHAPTER XIV SOME PROPERTIES OF SQUARE MATRICES 		
118.	Properties of a product of square matrices: recapitulation	107–108
119.	Co-joint complete matrices of the minor determinants of a square matrix: definition; fundamental property; a square matrix and its reciprocal are two co-joint matrices. Co-joint matrix of a complete matrix of the minor determinants of a rectangular matrix	108–112
120.	Determinant of a complete matrix of the minor determinants of a square matrix. Reciprocals of two co-joint matrices	112–114
121–122.	Relations between any two anti-correspondent minor determinants, any two anti-correspondent matrices of the minor determinants, and any two corresponding complete matrices of the minor determinants of two co-joint complete matrices of the minor determinants of a square matrix. Rank of a complete matrix of the minor determinants of a rectangular matrix	114–123
123.	Expansions of certain bordered determinants in terms of the simple minor determinants of the bordering rows	123–127
124.	Properties of the reciprocal of a square matrix: determinant and reciprocal of the reciprocal matrix; rank of the reciprocal matrix; relations between any two anti-correspondent minor determinants, any two anti-correspondent matrices of the minor determinants, and any two corresponding complete matrices of the minor determinants of a square matrix and its reciprocal; reciprocal of any derangement of a square matrix	127–133
125.	Properties of a symmetric matrix of rank 1: the diagonal elements do not all vanish; if the matrix is real, the non-vanishing diagonal elements all have the same sign; expression of the matrix as a product of two mutually conjugate one-rowed matrices	133–135
126.	Some general properties of symmetric matrices: properties of diagonal minor determinants; rank determined by diagonal minor determinants; if the matrix has rank r , it has a regular diagonal minor determinant of order r ; standard arrangement of the rows	135–141
127.	Some general properties of skew-symmetric matrices: diagonal elements are all 0's; properties of diagonal minor determinants; rank determined by diagonal minor determinants, is always even; if the matrix has rank r , it has a regular diagonal minor determinant of order r ; standard arrangement of the rows	141–145

CONTENTS

xv

§§	PAGES
128	The symmetric matrix of order 3: notation for minor determinants 145–146
129.	The symmetric matrix of order 4: notation for minor determinants; reciprocals of square minors of order 3 of the matrix and its reciprocal 147–156
130.	Identical relations between the elements of a square matrix: the fundamental identity; properties of a matrix of the super-determinants of a minor determinant; applications of the fundamental identity to symmetric matrices of orders $m, 3$ and 4 157–164
130 <i>a.</i>	APPENDIX B. The Pfaffian of a skew-symmetric matrix of even order: definition; Pfaffian co-factors; expansions of a Pfaffian; a skew-symmetric matrix of even order is the square of its Pfaffian; bordered skew-symmetric determinants expressed as products of two Pfaffians; symmetrically bordered skew-symmetric determinants; reciprocals of skew-symmetric matrices . 521–530

CHAPTER XV

RANKS OF MATRIX PRODUCTS AND MATRIX FACTORS

131.	Rank of a matrix product in which one of the extreme factor matrices has rank equal to its passivity; applicability of the results to a product of functional matrices 165–167
132.	Possible ranks of the solutions of a given matrix equation of the first degree: the equations $AX=C$, $XB=C$, $AXB=C$; the symmetric equation $A'XA=C$. Formulae for the general solutions of these equations 168–177
133.	Restrictions on the rank of any matrix product: the rank of the product matrix cannot exceed the rank of any factor matrix; the sum of the rank and the passivities of the product cannot be less than the sum of the ranks of the factor matrices 177–179
134–135.	Possible ranks of the product matrix and the factor matrices in a matrix product: products of two matrices; products of three matrices; any matrix product 179–194
136–137.	Possible ranks of the product matrix and the factor matrices in a symmetric matrix product; symmetric products of two matrices; symmetric products of three matrices; any symmetric product 194–205
138.	Equivalences of two matrices, and equivalent systems of linear algebraic equations: conditions for the horizontal or vertical equivalence of two matrices; equivalences of two similar square matrices; sign of equivalence 205–208
	Spacelets represented by matrices which are not necessarily undergenerate 208–209
139.	Joins and intersections of spacelets in homogeneous space of $n-1$ dimensions or rank n : join and complete intersection of two spacelets; formulae for two spacelets, their join and their intersection; relation between the ranks of their join and intersection; possible ranks of the join and intersection of two spacelets of given ranks which are otherwise arbitrary, which both lie in or both contain a given spacelet, which both lie in one given spacelet and both contain another given spacelet;

xvi	CONTENTS	PAGES
§§	spacelets having a given complete intersection with a given spacelet ; mutually complementary spacelets ; mutually incident spacelets	209–219
	Join and complete intersection of any number of spacelets ; spacelets which do not intersect either of two given spacelets ; spacelets restricted to be real	219–222
	Joins and intersections of matrices	222–223
140.	Connections between matrices ; connections between spacelets ; unconnected spacelets, no one of them intersects the join of the others ; spacelets which do not intersect one given spacelet and have a given complete intersection with another	223–227

CHAPTER XVI

EQUIGRADIENT TRANSFORMATIONS OF A MATRIX WHOSE ELEMENTS ARE CONSTANTS

EQUIGRADIENT TRANSFORMATIONS OF ANY MATRIX.

141.	Definitions of an equigradent transformation and its inverse ; equigradent matrices, have the same rank ; symmetric equigradent transformations ; symmetrically equigradent matrices ; equigradent transformations between two similar matrices ; composition of equigradent transformations, resultant equigradent transformation ; elementary equigradent transformations ; derangements, unitary transformations, quasi-scalar and scalar transformations ; equigradent transformations of a compartite matrix and its parts ; correspondences between equigradent transformations of a matrix with constant elements and linear transformations of bilinear and quadratic algebraic forms ; completion of any two mutually inverse matrices ; conversion of any equigradent transformation and its inverse into two mutually inverse equigradent transformations of similar matrices ; some general properties of equigradent transformations	228–241
	SOME SPECIAL EQUIGRADIENT TRANSFORMATIONS IN Ω OF ANY MATRIX IN Ω WHOSE ELEMENTS ARE CONSTANTS.	
142.	Unitary transformations converting the matrix into a bipartite matrix one of whose parts is a given non-zero element or a given undegenerate square minor ; special cases	241–252
143.	A more general unitary transformation converting the matrix into a compartite matrix of standard form whose non-zero parts are all undegenerate square matrices	252–260
144.	A corresponding non-unitary equigradent transformation	260–264
145.	The corresponding transformations of any symmetric or skew-symmetric matrix in Ω whose elements are constants: conversion into a bipartite or compartite matrix by symmetric unitary (or corresponding non-unitary) equigradent transformations in Ω ; special cases	264–271

§§	CONTENTS	xvii PAGES
	REDUCTION OF ANY MATRIX WHOSE ELEMENTS ARE CONSTANTS TO STANDARD FORMS BY EQUIGRADENT TRANSFORMATIONS.	
146.	Reduction of any matrix in Ω by transformations in Ω ; reduction to a standard form by unrestricted equigradent transformations in Ω ; reduction to a standard form by derangements and unitary (or corresponding non-unitary) equigradent transformations in Ω ; two matrices are equigradent when and only when they have the same rank	272–278
	Equigradent transformations between two similar undegenerate quasi-scalar or square matrices whose determinants are equal in value; reduction of a square matrix whose determinant is ± 1	278–280
147.	Reduction of any symmetric matrix in Ω by symmetric transformations: reduction to a standard form by derangements and unitary (or corresponding non-unitary) transformations in Ω ; reduction to a standard form by unrestricted symmetric equigradent transformations (not in Ω); two symmetric matrices are symmetrically equigradent when and only when they have the same rank	280–295
148.	The signants and signature of a real symmetric matrix: defined; the positive and negative signants remain invariant in all real symmetric equigradent transformations of the matrix; reduction of a real symmetric matrix to a standard form by such transformations; there exists a real symmetric equigradent transformation between two real symmetric matrices when and only when they are equisignant; the positive and negative signants of a real symmetric partite matrix are respectively the sums of the positive and negative signants of its parts	295–300
149.	Definite and indefinite real symmetric matrices: defined; general formula for all definite matrices; all non-vanishing diagonal minor determinants of order s of a definite matrix have the same sign; every symmetrically formed complete matrix of the minor determinants of a definite matrix is definite; roots of a real and definite quadratic form; essentially positive and essentially negative real quadratic forms; semi-definite matrices	300–303
150.	Reduction of any skew-symmetric matrix in Ω by symmetric transformations in Ω : reduction to a standard form by derangements and unitary (or corresponding non-unitary) equigradent transformations in Ω ; reduction to a standard form by unrestricted symmetric equigradent transformations in Ω ; two skew-symmetric matrices are symmetrically equigradent when and only when they have the same rank	303–308

CHAPTER XVII

SOME MATRIX EQUATIONS OF THE SECOND DEGREE

§§		PAGES
	GENERAL EQUATIONS.	
151.	Matrix equations of the second degree : definitions and descriptions	309–310
152–153.	$XY=AB$: Determination of all solutions of the equations $\overline{x}_m [y]_n = \overline{a}_m [b]_n, \quad [x]_m^r [y]_r^n = [\alpha]_m^r [b]_r^n, \quad [x]_m^s [y]_s^n = [\alpha]_m^r [b]_r^n,$ the given factors on the right in the last two equations having rank r ; general solutions of the first two equations	310–315
154.	$XY=C$: Determination of all solutions of the equations $\overline{x}_m [y]_n = [c]_m^n, \quad [x]_m^r [y]_r^n = [c]_m^n, \quad [x]_m^s [y]_s^n = [c]_m^n,$ the given matrix $[c]_m^n$ having rank 0 or 1 in the first equation, and rank r in the second and third equations; general solutions of the first two equations	316–320
	SYMMETRIC EQUATIONS.	
155.	$X'X=I$, where I is a unit matrix : Semi-unit matrices ; determination of all semi-unit matrices, of all real semi-unit matrices ; enlargement of a given semi-unit matrix (or real semi-unit matrix) by the addition of long rows (or real long rows) ; number of arbitrary parameters in a general semi-unit matrix ; every derangement and every long minor of a semi-unit matrix is a semi-unit matrix. Square semi-unit matrices ; derangements of a unit matrix ; compound square semi-unit matrix with four real and purely imaginary constituents, all constituents are square ; most general square semi-unit matrices of orders 2 and 3	320–330
	Rotations of a rigid body represented by square semi-unit matrices of order 3 ; theorems on rotations about a fixed point	330–332
156–157.	$X'X=A'A$: Determination of all solutions of the symmetric equations $\overline{x}_m [x]_m = \overline{a}_m [a]_m, \quad \overline{x}_m^r [x]_r^m = \overline{a}_m^r [a]_r^m, \quad \overline{x}_m^s [x]_s^m = \overline{a}_m^r [a]_r^m,$ the given factors on the right in the last two equations having rank r ; general solutions of the first two equations ; formulae giving all solutions of the third equation of any possible rank ρ ; formulae giving all solutions of rank ρ of the equation $\overline{x}_m^s [x]_s^m = \begin{bmatrix} 1, & 0 \\ 0, & 0 \end{bmatrix}_{r, m-r}^{r, m-r}$	333–344
158–159.	$X'X=0$: General formulae for all solutions of rank ρ of the equation $\overline{x}_m^s [x]_s^m = 0$; standard formula for all undegenerate solutions of rank r of the equation $[x]_r^n \overline{x}_n^r = 0$	344–352

CONTENTS

§§		PAGES
160.	$X'X=C$: Determination of all solutions of the symmetric equations $\begin{bmatrix} x \\ x \end{bmatrix}_m = [c]_m^m, \quad \begin{bmatrix} x \\ x \end{bmatrix}_m^r = [c]_m^m, \quad \begin{bmatrix} x \\ x \end{bmatrix}_m^s = [c]_m^m,$ the given symmetric matrix $[c]_m^m$ having rank 0 or 1 in the first equation, and rank r in the second and third equations; general solution of the first equation; particular solutions and general solution of the second equation; semi-real solutions of the second equation when $[c]_m^m$ is real. Special symmetric equations of the forms $X'X=A'BA$, $X'BX=A'A$, $X'BX=A'CA$. . .	352-363
161.	$X'AX=C$: Determination of all solutions of any symmetric equation of this form	363-365
	SPECIAL EQUATIONS	
162.	Expressions for a symmetric matrix of order 2 as a product of two square factors	365-369
163.	Some special equations of the form $[x]_m^2 [y]_2^m = [a]_m^2 [b]_2^m$. . .	369-373
164.	Expressions for a symmetric matrix whose rank does not exceed 2 as a product of two 2-rowed factors; applications to symmetric matrices of orders 3 and 4	373-377

CHAPTER XVIII

THE EXTRAVAGANCES OF MATRICES AND OF SPACELETS
 IN HOMOGENEOUS SPACE

165.	The degeneracy of any matrix: defined	378
	The extravagance of an undegenerate matrix: possible values; general formulae for an undegenerate matrix whose orders and extravagance are given; condition that two similar undegenerate matrices may have the same extravagance; non-extravagant, completely extravagant, and plenerily extravagant undegenerate matrices; general formulae for these; possible ranks and extravagances of undegenerate matrices connected with a given undegenerate matrix	378-385
	The horizontal and vertical extravagances of any matrix: of a matrix of rank r expressed as a product of passivity r ; possible values; horizontally (or vertically) equivalent matrices have the same horizontal (or vertical) extravagance; general formulae for a matrix of which the rank and one or both of the extravagances are given; condition that two matrices may have the same rank and the same horizontal (or vertical) extravagance; matrices whose extravagances are both zero	385-393
	The extravagance of a symmetric matrix: possible values; general formulae for a symmetric matrix whose rank and extravagance are given; ranks of the powers of a non-extravagant symmetric matrix	393-395

§§		PAGES
166.	Minor determinants of the matrices $[1, a]_m^{m, n}$ and $[a, 1]_m^{n, m}$: simple minor determinants; minor determinants of order s	395–397
167.	Properties of two mutually normal undegenerate matrices: they have the same extravagance; the corranged simple minor determinants of either one are proportional to the anti-correspondent affected simple minor determinants of the other; general formulae for two mutually normal undegenerate matrices; possible values of their ranks and common extravagance; one of them is completely extravagant when and only when the other is plenary extravagant. Possible ranks and extravagances of two mutually orthogonal undegenerate matrices. Definitions of horizontally (or vertically) orthogonal and normal matrices; all matrices horizontally (or vertically) normal to a given matrix are horizontally (or vertically) equivalent; possible ranks and extravagances of any two mutually orthogonal matrices	398–411
168.	Reduction of an undegenerate matrix (of extravagance ρ) to an equivalent similar undegenerate matrix whose long rows are mutually orthogonal; number of extravagant long rows in the equivalent matrix (is equal to ρ); non-extravagant rows at unit intensity; reduction of a non-extravagant (or real) matrix to an equivalent semi-unit (or real semi-unit) matrix	412–415
169.	The cores of an undegenerate matrix (of extravagance ρ): defined; are mutually equivalent completely extravagant matrices (of rank ρ); two mutually normal undegenerate matrices have the same cores, these being their complete intersections; determination of the cores	415–422
170.	The extravagance, core and plenum of a spacelet in homogeneous space of $n - 1$ dimensions or rank n . Preliminary remarks: Spacelets, their joins, intersections and incidences; mutually complementary spacelets; unconnected spacelets; orthogonal and normal spacelets; general formula for a spacelet which lies in the join of two given spacelets and has a given complete intersection with one of them	422–425
	The extravagance of a spacelet: defined, is the degree of its orthogonality with itself; possible values; non-extravagant, completely extravagant and plenary extravagant spacelets; mutually orthogonal non-extravagant spacelets are unconnected; two spacelets have the same rank and the same extravagance when and only when each is convertible into the other by a semi-unit transformation	425–427
	The core of a spacelet (of extravagance ρ): is a completely extravagant spacelet (of rank ρ) which is the locus of all points which lie in the spacelet and are orthogonal with it; general formulae for a spacelet and its core; the core of the join of unconnected mutually orthogonal spacelets is the join of their cores; possible ranks and extravagances of a spacelet which lies in a given spacelet and has a given complete intersection with its core	427–430
	The plenum of a spacelet: is the plenary extravagant spacelet which is the normal to its core; is the smallest spacelet which contains the given spacelet and all points orthogonal with it	431–432

CONTENTS

§§		PAGES
	Properties of normal spacelets: two mutually normal spacelets have the same extravagance, the same core and the same plenum, their common core being their complete intersection, and their common plenum being their join; general formulae for two mutually normal spacelets, their common core and their common plenum; the join (or complete intersection) of the normals to any number of given spacelets is the normal to the complete intersection (or join) of the given spacelets; ranks of the intersection and join of the normals to two given spacelets; two spacelets are non-intersecting when and only when their normals are mutually complementary. Complementary theorems . . .	432–434
	Interpretations of the terms ‘extravagant’, ‘core’, ‘orthogonal’, ‘normal’ as denoting relations to the absolute quadric . . .	434–435
171.	Standard representations of completely and plenarily extravagant spacelets; a plenarily extravagant spacelet is the join of its core and a real spacelet; anti-cores of a spacelet . . .	435–440
172.	Unconnected mutually orthogonal solutions of any system of homogeneous linear algebraic equations: methods of determining a complete set of such solutions; number of extravagant solutions in each set . . .	440–446
173.	Possible extravagances of two non-intersecting spacelets: their extravagances are independent, i.e. each spacelet can have independently of the other spacelet any extravagance consistent with its rank; the extravagances of unconnected spacelets are independent . . .	446–450
174.	Possible values of the rank and extravagance of a spacelet which lies in a given spacelet of homogeneous space ω_n . A spacelet which contains a plenarily extravagant spacelet must itself be plenarily extravagant . . .	450–456
175.	Possible values of the rank and extravagance of a spacelet of homogeneous space ω_n which contains a given spacelet. A spacelet which lies in a completely extravagant spacelet must itself be completely extravagant . . .	456–462
175a.	APPENDIX C. Equigradent transformations in which one of the transforming factors is a semi-unit matrix: unilaterally semi-unit equigradent transformations of a matrix; semi-unit transformations of a matrix; equigradent transformations of a spacelet; semi-unit transformations of a spacelet . . .	531–534

CHAPTER XIX

THE PARATOMY AND ORTHOTOMY OF TWO MATRICES AND OF TWO SPACELETS OF HOMOGENEOUS SPACE

176. The paratomy, orthotomy and cross rank of any two spacelets in homogeneous space of $n-1$ dimensions or rank n .
 The paratomy, orthotomy and cross rank of two matrices which represent spacelets: are those of the two spacelets which they represent. The mutual paratomy (or degree of intersection) of two spacelets: is the rank of their complete intersection; possible values for two arbitrary spacelets of given ranks. The

xxii	CONTENTS	PAGES
§§	mutual orthotomy (or degree of orthogonality) and the cross rank of two spacelets : defined ; relations between them ; interpretations ; the extravagance of a spacelet is its orthotomy with itself. The two cases in which each of two spacelets is incident with the normal to the other.	
	Possible values of the cross rank and mutual orthotomy of two arbitrary spacelets of given ranks ; greatest and least values of the orthotomy	463–468
177.	Possible values of the cross rank and mutual orthotomy of two spacelets which both lie in a given spacelet ; which both contain a given spacelet.	
	Spacelets which lie in or contain a given non-extravagant spacelet, or which lie in one given non-extravagant spacelet and contain another : correspondences between such spacelets and spacelets of a complete homogeneous space ; correspondences of ranks and cross ranks ; of extravagances and orthotomies	468–474
178.	Possible values of the cross rank and mutual orthotomy of two non-intersecting spacelets of given ranks ; greatest and least values of the orthotomy ; possible values for two non-intersecting spacelets of given ranks which both lie in a given non-extravagant spacelet. Corresponding results when one of the two spacelets is given	474–478
179.	Possible values of the cross rank and mutual orthotomy of two mutually complementary spacelets of given ranks ; greatest and least values of the orthotomy ; possible values for two spacelets of given ranks which both lie in a given non-extravagant spacelet and are mutually complementary in it	478–480
180.	Properties of two spacelets of given ranks which have a given complete intersection ω_p , and which both lie in the plenum of their intersection ; representation of each spacelet as the join of two mutually orthogonal non-intersecting spacelets, one of which is ω_p ; cross rank and mutual orthotomy of these two joins ; extravagance and core of each join.	
	Possible values of the cross rank and mutual orthotomy of the two spacelets ; greatest and least values of the orthotomy ; possible extravagances of the two spacelets, the extravagance of each is independent of the other.	
	Possible values of the mutual orthotomy of two spacelets whose complete intersection is a given non-extravagant spacelet ; of two real spacelets whose complete intersection is given	480–489
181.	Properties of any two spacelets of given ranks which have a given complete intersection ω_p : representation of each spacelet as the join of three unconnected spacelets of which the first is ω_p , the second is orthogonal with ω_p , and the third lies outside the plenum of ω_p ; cross rank and mutual orthotomy of these two joins ; extravagance and core of each join.	
	Possible values of the cross rank and mutual orthotomy of the two spacelets ; greatest and least values of the orthotomy ; possible extravagances of the two spacelets, the extravagance of each is independent of the other.	
	Orthotomy and extravagances of two spacelets having a given join	490–506

	CONTENTS	xxiii
§§		PAGES
182.	Possible simultaneous values of the paratomy, cross rank and orthotomy of two entirely arbitrary spacelets; of two entirely arbitrary real spacelets; possible simultaneous values of any two of these quantities	506–509
183.	Properties of mutually orthogonal spacelets: they have the same complete intersection as their cores; the core of their join is the join of their cores; their cores are their intersections with the core of their join	509–514
117 <i>a.</i>	APPENDIX A. Utility of the relations obtained in Chapter XIII .	515–520
130 <i>a.</i>	APPENDIX B. The Pfaffian of a skew-symmetric matrix of even order	521–530
175 <i>a.</i>	APPENDIX C. Equigradent transformations in which one of the transforming factors is a semi-unit matrix	531–534
	INDEX	535–555