

## INDEX

- Accommodation coefficient,  
 calculated values, less than measured values, 2  
 definition, 1, 2  
 effect of surface impurities, 4, 60, 94, 95-7, 99, 102-3  
 mass of gas and solid atoms, dependence on, 2  
 measurement, experimental arrangement, 10  
 of helium, measured values, 4  
 temperature variation of, 4-5  
 of neon, measured values, 6, 60-1, 95  
 sensitive to adsorbed films, 6-7, 11, 60, 95-9, 102-3  
 temperature variation of, 61  
 of rotational energy, 7  
 relation to kinetic theory, 7-9  
 relation to potential energy of gas atom and solid, 6  
 rough surface, 6 f.n.  
 theory of measurement, 7-9  
 variation with heat treatment of wire, 12, 59-60, 85
- Adsorption, heat of, *see* Heat isotherm of, *see* Isotherm
- Ammonia,  
 adsorption on barium fluoride, 123-4  
 heat of adsorption of, 118  
 molecular constants of, 115
- Argon,  
 adsorption on caesium iodide, 122
- Atomic hydrogen, *see* Hydrogen
- Bare surfaces,  
 importance of, 3, 10, 12, 13
- Caesium,  
 adsorbed on tungsten, effect on electron emission, 77  
 each atom held on metallic surface, equivalent to electric dipole, 105  
 electric moment per atom adsorbed on metal, 105  
 stability of adsorbed film of, on tungsten, effect of oxygen, 77
- Carbon dioxide,  
 heat of adsorption of, on barium fluoride, 124
- Collision of gas atom and solid,  
 classical theory of, 2  
 quantum theory of, 5
- Contact potential,  
 difference between bare and hydrogen covered tungsten, 30, 58  
 formation of hydrogen films on tungsten, 57  
 in study of monolayers, experimental procedure, 57  
 sodium films on tungsten, 57  
 to measure evaporation of oxygen from tungsten, 77  
 variation with fraction of surface covered, 58
- Crystal planes,  
 aged tungsten, important plane for, 16  
 geometrical arrangement of atoms in, for tungsten, 17, 18, 87  
 determines potential field for adsorbed atom, 20  
 number of atoms in, for tungsten, 16, 17
- Detailed balance, principle of,  
 atomic hydrogen, production of, processes concerned, 66-7, 70-4  
 deduction of Langmuir adsorption isotherm, 21
- Dipoles,  
 adsorbed on solid, statistical analysis for, 109-14  
 electrostatic forces between, 107-9  
 heat of adsorption, *see* Heat induced by ionic crystal, 104  
 lattice of, interaction energy, 107  
 mutual depolarizing action of 105, 114  
 effect of, on heat of adsorption, 114, 118-19
- Emission of electrons, *see* Thermionic emission
- Energy, interaction,  
 continuous variation over surface, for adsorbed particle, 36, 38-40  
 fixed, statistical theory for, 24-33, 38  
 for immobile film, 32-3  
 for mobile film, 28

- Energy, interaction (*cont.*)  
 of hydrogen on tungsten, only short-range forces important, 29–30  
 of lattice of dipoles, 107  
 of linear chain, 43
- Energy, interchange,  
 between helium and tungsten, 4  
 between solid and gas, 1, 3  
 between solid and gas, depends on law of force, 5  
 between solid and gas, depends on masses, 2, 5  
 value of accommodation coefficient, relation to, 2
- Energy, potential,  
 between gas atom and solid surface, represented by Morse function, 5  
 continuous variation over surface, 38–40  
 determined by structure of surface, 20  
 electrostatic, 107–9
- Energy, states of minimum,  
 difference between mobile and immobile films, 22–3  
 effect of thermal motion, 23, 54  
 for linear chain, 49, 51–4  
 for square array, 50  
 when occupation of a site precludes occupation of its neighbours, 87
- Evaporation,  
 rate of, if interactions negligible, 21  
 to determine apparent heat, 77–81  
 when occupation of a site precludes occupation of its neighbours, 91  
 theory of, of adsorbed particles, 67–70  
 of atoms, 67, 71  
 of molecules by combination of adsorbed atoms, 67, 72  
 of molecules from atomic film, 67, 73
- Film, adsorbed,  
 accommodation coefficient, effect of, on, 4, 6, 12, 94–9, 102–3  
 forces between molecules, relation to variation of heat of adsorption, 19  
 structure of, on known substrate, 3, 59, 95–8
- Film, adsorbed (*cont.*)  
*see also* Immobile film, Mobile film, Monomolecular layer
- Heat of adsorption,  
 allowing for continuous variation of energy terms, 44–8  
 experimental results, application to, 48–51  
 deduced from rate of evaporation, 77, 91  
 different from apparent heat of evaporation, 81, 89, 91  
 effect of dipole forces on, 104, 110–24  
 experimental procedure, 14–16  
 for simple adsorption, into mobile film, 27–8  
 indication of repulsive forces between adsorbed particles, 19  
 measurement of, experimental arrangement, 13–14  
 of hydrogen on tungsten, 18, 19  
 of mobile and immobile films, on localized sites, difference between, 21–3, 35, 37  
 reduction of, due to electrostatic forces, 115, 120, 122  
 theory of, allowing for continuous variation of potential energy, 44–8, 54  
 for constant dipole moment, 106, 111–12  
 for immobile film, 23, 32–3, 45–8, 88  
 for mobile film, 22, 27–9, 44–7, 87–9  
 use of wire as calorimeter, 13  
 when occupation of site precludes occupation of its neighbours, 87–9
- Heat of evaporation,  
 deduced from rate of evaporation, 77, 91  
 different from true heat of adsorption, 81, 89, 91
- Helium,  
 accommodation coefficient of, 4  
 temperature variation of, 4–5  
 interchange of energy with solid, adsorption effects absent except at low temperatures, 4
- Hydrogen,  
 accommodation coefficient, on tungsten, measured, 12, 85

- Hydrogen (*cont.*)
- accommodation coefficients, variation with heat treatment of surface, 12, 85
  - adsorption of, rapid on tungsten, 11, 12
    - exothermic process, 75
    - raises temperature of filament, 15
  - amount adsorbed, on surface planes of tungsten, 16–18
  - film adsorbed on tungsten, only nearest neighbours contribute to interaction energy, 29–30
  - heat of adsorption, measured variation, 18–19, 75
    - interpretation, 34–5, 48–51
  - molecular adsorption, 86
  - monolayer on tungsten, immobile, 35, 51, 55
  - ortho-para* conversion of, on tungsten, 86
- Hydrogen, atomic,
- produced by hot tungsten filament, 65
  - rate of production, 65
    - effect of pressure on, 66
    - effect of temperature on, 66, 75
    - theory and experiment compared, 76, 85
    - theory of, 67, 70–5
  - trapped by film of molybdenum oxide, 65
- Immobile film,
- contrasted with mobile film, 21 f.n., 22–3, 35, 37, 59, 92–4
  - definition of, 22
  - equivalent to random distribution of occupied sites, 30
  - gaps in final film, 18, 33–4, 55, 92–4
    - as centres of catalytic activity, 55 f.n.
    - determined empirically, 56, 92
    - determined statistically, 55–6, 92
  - heat of adsorption, compared with experiment, 35
    - contrasted with that of mobile film, 23, 47, 89
  - for adsorption with dissociation, 30, 33–4, 93
  - for continuous variation of energy terms, 45–8
  - for simple adsorption, 23, 30
  - occupation of site precludes occupation of its neighbours, 91, 93
- Immobile film (*cont.*)
- rate of formation, occupation of site precludes occupation of its neighbours, 90, 93
    - of film formed by dissociation, 57, 59, 93
  - statistical treatment, 30–3, 55–6, 92, 101
- Indicator of adsorbed films, measured accommodation coefficient as, 7, 60, 95–9, 102–3
- Interaction energy, *see* Energy
- Isotherm, of adsorption,
- for simple adsorption into mobile film, 27
  - from kinetic considerations, 74
  - long-distance order, effect on, 84
  - neglecting interactions, Langmuir, 21, 82
  - occupation of a site precludes occupation of its neighbours, 98
  - statistical derivation, 27, 82, 98
- Kinetics of formation of monolayer, immobile film, formed with dissociation, 56–7
- mobile film, formed with dissociation, 61–4
  - when occupation of site precludes occupation of its neighbours, 89–91
- of adsorption of hydrogen, 57–9
- of adsorption of oxygen, 100–3
- Kinetics, chemical,
- condition of quasi-chemical equilibrium, 35
  - use of, to examine nature of approximation of local configurations, 25
- Kinetic theory,
- and accommodation coefficient, 7–9
  - collision of gas atom and solid surface, 2
  - exact calculations, conditions under which possible, 1, 9
  - superseded by quantum theory of collisions, 3
- Local configurations,
- method of, introduced in theory of alloys, 25
  - used to determine equilibrium distribution, 24–5, 31, 109–11
  - used to determine heat of adsorption, 28, 32, 111–14, 116–19

- Long-range forces,  
 effect of, on equilibrium properties  
 of monolayer, 104  
 statistical analysis of, 124  
*see also* Dipoles
- Mobile film,  
 approximation of local configura-  
 tions, use of, 25  
 contrasted with immobile film,  
 21 f.n., 22-3, 35-7, 59, 92-4  
 definition of, 21  
 equilibrium distribution of particles  
 in, 24-7  
 heat of adsorption, for simple ad-  
 sorption, 22, 27-8  
 for adsorption with dissociation,  
 28-30, 92  
 for continuous variation of energy  
 terms, 44-7  
 when occupation of site precludes  
 occupation of its neighbours, 92  
 interpretation of experimental heats  
 of adsorption, 35, 48-51, 78-81
- Monomolecular layer,  
 adsorption isotherm, Langmuir, 21  
 of oxygen, 98  
 statistical deduction of, 82  
 changes in, effect of, on accommo-  
 dation coefficient, 12, 60-1, 94,  
 95-7, 99  
 contact potential, effect on, 57-9  
 different types, on array of localized  
 sites, 21-3, 35, 37  
 effect on surface reactions, 1  
 evaporation and condensation pro-  
 cesses, 21, 57, 62-3, 66-7,  
 70-4  
 formation, kinetics of, 55-7, 61-4  
 formation, rate of, 21, 57, 63  
 formed at very low pressure, 11, 13,  
 96  
 formed on clean surface, 1, 12-13  
 interchange of energy with gas  
 atoms, effect on, 1  
 measured accommodation coeffi-  
 cient, effect on, 2, 4, 6, 11, 12,  
 60-1, 94, 95-7, 99  
 molecular or atomic film, depends  
 on temperature, 86  
 occupation of site precludes occu-  
 pation of its neighbours, 86,  
 89-92  
 of hydrogen on tungsten, 11, 16-19,  
 35, 58
- Monomolecular layer (*cont.*)  
 of oxygen on tungsten, 59-61, 94-9  
*see also* Heat of adsorption, Immo-  
 bile film, Mobile film
- Morse function,  
 constants of, physical significance, 6  
 determination from temperature  
 variation of accommodation  
 coefficient, 6  
 definition, 6-7  
 potential energy of gas atom and  
 solid, represented by, 6, 68
- Neon,  
 accommodation coefficient, measur-  
 ed, 6, 12, 95, 96 f.n.  
 effect of adsorbed impurities on,  
 6-7, 12, 60, 95-7  
 as indicator of adsorbed films, 7, 60,  
 96-7
- Nitrous oxide,  
 heat of adsorption, on barium  
 fluoride, 124
- Oxygen,  
 effect of trace of, on accommodation  
 coefficient of neon, 60, 94, 96  
 isotherm, of adsorption, 98  
 molecular adsorption, 86  
 monolayer of, on tungsten, rate of  
 formation, 59  
 gaps in final film, 59, 96  
 heat of adsorption, 59-60  
 partial pressure of, relation to  
 accommodation coefficient of  
 neon, 96  
 structure of adsorbed film, 95-8
- Partition function,  
 immobile film, 31  
 mobile film, 25-7  
 systems with long-range forces,  
 109-11  
 when occupation of site precludes  
 occupation of its neighbours,  
 92, 98  
*see also* Vibrational partition func-  
 tion
- Potential energy, *see* Energy
- Roughness factor, 6
- Solutions, theory of,  
 contrasted with that of adsorbed  
 monolayers, 35-7

## INDEX

133

- Solutions, theory of (*cont.*)  
 random mixture an approximation  
 to mixture with non-zero heat  
 of mixing, 35–6
- Sulphur dioxide,  
 adsorption, on charcoal, 122  
 on barium fluoride, 123  
 heat of adsorption of, 123, 124
- Surface,  
 bare, importance of, 3, 10, 12, 13  
 rapid adsorption of hydrogen on,  
 12  
 behaviour simplest in helium atmo-  
 sphere, 4  
 cleaned by flashing, 14, 17  
 impurities, effect of, 2, 4, 12  
 necessary to keep free of, 3  
 interchange of energy with gas  
 atoms, 1, 3  
*see* Immobile film, Mobile film,  
 Monomolecular layer
- Thermionic emission,  
 effect of adsorbed caesium on, from  
 tungsten, 77  
 used to measure amount of adsorbed  
 oxygen, 77
- Tungsten,  
 adsorption on, of helium, 4  
 of neon, 6  
 contact potential of, with adsorbed  
 hydrogen film, 58  
 dipoles formed, with adsorbed  
 caesium, 105  
 removal of impurities from, 11, 14,  
 17  
 surface plane, in aged tungsten, 16  
 thermionic emission from, effect of  
 adsorbed caesium, 77  
 use of, as calorimeter, 13
- Tungsten powder,  
 interpretation of experimental heats  
 of adsorption, 36  
 surface contaminated by impurities,  
 12  
 use of, unsatisfactory, in adsorption  
 measurements, 12
- Vibrational partition function,  
 effect on variation of heat of  
 adsorption, 51 f.n.  
 independent of state of occupation  
 of neighbouring sites, 24,  
 51 f.n.