

Cambridge University Press

978-1-107-63012-3 - Theory of Differential Equations: Part II: Ordinary Equations,
not Linear: Vol. III

Andrew Russell Forsyth

Frontmatter

[More information](#)

THEORY
OF
DIFFERENTIAL EQUATIONS.

Cambridge University Press

978-1-107-63012-3 - Theory of Differential Equations: Part II: Ordinary Equations,
not Linear: Vol. III

Andrew Russell Forsyth

Frontmatter

[More information](#)

Cambridge University Press
978-1-107-63012-3 - Theory of Differential Equations: Part II: Ordinary Equations,
not Linear: Vol. III
Andrew Russell Forsyth
Frontmatter
[More information](#)

THEORY
OF
DIFFERENTIAL EQUATIONS.

PART II.
ORDINARY EQUATIONS, NOT LINEAR.

BY
ANDREW RUSSELL FORSYTH,
Sc.D. (CAMB.), HON. Sc.D. (DUBL.), F.R.S.,
SADLERIAN PROFESSOR OF PURE MATHEMATICS,
FELLOW OF TRINITY COLLEGE, CAMBRIDGE.

VOL. III.

CAMBRIDGE:
AT THE UNIVERSITY PRESS.

1900

All rights reserved.

Cambridge University Press
978-1-107-63012-3 - Theory of Differential Equations: Part II: Ordinary Equations,
not Linear: Vol. III
Andrew Russell Forsyth
Frontmatter
[More information](#)

CAMBRIDGE UNIVERSITY PRESS
Cambridge, New York, Melbourne, Madrid, Cape Town,
Singapore, São Paulo, Delhi, Tokyo, Mexico City

Cambridge University Press
The Edinburgh Building, Cambridge CB2 8RU, UK

Published in the United States of America by Cambridge University Press, New York

www.cambridge.org
Information on this title: www.cambridge.org/9781107630123

© Cambridge University Press 1900

This publication is in copyright. Subject to statutory exception
and to the provisions of relevant collective licensing agreements,
no reproduction of any part may take place without the written
permission of Cambridge University Press.

First published 1890
First paperback edition 2011

A catalogue record for this publication is available from the British Library

ISBN 978-1-107-63012-3 Paperback

Cambridge University Press has no responsibility for the persistence or
accuracy of URLs for external or third-party internet websites referred to in
this publication, and does not guarantee that any content on such websites is,
or will remain, accurate or appropriate.

Cambridge University Press

978-1-107-63012-3 - Theory of Differential Equations: Part II: Ordinary Equations,
not Linear: Vol. III

Andrew Russell Forsyth

Frontmatter

[More information](#)

CONTENTS.

CHAPTER XI.

REDUCED FORMS OF SYSTEMS OF EQUATIONS OF THE FIRST ORDER IN
THE VICINITY OF SINGULARITIES OF THE DERIVATIVES.

ART.		PAGE
145.	System of equations in vicinity of ordinary position for the derivatives	1
146—148.	Reduction to a canonical form of the general system $\frac{dx_1}{X_1} = \frac{dx_2}{X_2} = \frac{dx_3}{X_3} = \dots$	2
149.	Various cases of exceptional combinations arising out of singularities of a system of two equations of first order	9
150.	Case I: when the set of values is ordinary for one equation, and accidental of first kind for the other .	11
151.	Case II: ordinary for one, accidental of second kind for other	13
152.	Case III: accidental of first kind for both	15
153.	Case IV: accidental of first kind for one and of second kind for other	15
154.	Case V: accidental of second kind for both	17
155, 156.	Equations determining the leading terms in the integrals of the reduced form in Case V	19
157.	Space-diagrams, as extension of Puiseux diagrams	23
158—160.	Reduced forms of systems in Case V	26
161.	Results for system of n equations	31
162.	Reduction of remaining forms for a system of two equations	32
163.	Examples	35

Cambridge University Press

978-1-107-63012-3 - Theory of Differential Equations: Part II: Ordinary Equations,
not Linear: Vol. III

Andrew Russell Forsyth

Frontmatter

[More information](#)

vi

CONTENTS

CHAPTER XII.

THE INTEGRALS OF THE REDUCED FORMS OF A SYSTEM OF EQUATIONS,
CHIEFLY OF TWO DEPENDENT VARIABLES.

ART.	PAGE
164, 165. Complete integral of a special system of two equations in a canonical reduced form: with inferences suggested as to the complete integral of a general system of two equations	40
166. Two alternative canonical forms of a system of two equations: with transformation	47
167—169. Regular integrals of the system when the critical quadratic has unequal roots; neither, either, or both, of the roots being a positive integer	49
170, 171. Regular integrals of the system when the critical quadratic has a repeated root, whether a positive integer or not	62
172. On the possibility of non-regular integrals	69
173—176. Non-regular integrals of the system when the critical quadratic has unequal roots, neither being a positive integer	73
177, 178. Likewise when one (but not the other) of the unequal roots is a positive integer	87
179. Likewise when both of the unequal roots are positive integers	95
180—182. Non-regular integrals of the system when the critical quadratic has a repeated root, which is not a positive integer	96
183—185. Likewise when the repeated root is a positive integer	105
186. Case remaining over from § 173	116
187. Enunciation of the corresponding theorems for a system of any number of equations	116
188. Remaining typical forms of Chap. XI: with examples	118

CHAPTER XIII.

SYSTEMS OF EQUATIONS WITH MULTIFORM VALUES OF THE
DERIVATIVES; SINGULAR SOLUTIONS.

189. Systems of equations of the first order and any degree	121
190. Form of the equations in the vicinity of branch-points	122
191. The integrals when a certain Jacobian does not vanish	124
192. Likewise when it does vanish	127
193, 194. Summary of results: with geometrical interpretation	133

Cambridge University Press

978-1-107-63012-3 - Theory of Differential Equations: Part II: Ordinary Equations,
not Linear: Vol. III

Andrew Russell Forsyth

Frontmatter

[More information](#)

CONTENTS

vii

ART.		PAGE
195.	Method of proceeding when the earlier method does not lead to determinate results	138
196.	Examples	140
197.	When the Jacobian of a system of two equations vanishes for an integral, another equation of the first order also is satisfied	148
198.	When the four equations of § 197 are satisfied, then a solution is provided	150
199, 200.	Singular solutions; general theorem as to conditions	153
201.	Examples	159
202.	Corresponding theorem for a system of n equations: various classes of singular solutions, with examples	162
203—206.	Analytical relation of the functions (whether singular integrals or not), defined by the four equations of § 197, to the complete integral	166
207.	Integrals of equation of the first order deduced from primitive	176
208.	Derivation of singular integrals from complete integral of a system of two equations	178

CHAPTER XIV.

EQUATIONS OF THE SECOND ORDER AND THE FIRST DEGREE.

209.	Regular integrals of an equation $w' = f(w', w, z)$, as determined by initial conditions	184
210.	Integrals (regular and non-regular) of an equation $z^2 w'' = f(zw', w, z)$, where f is a regular function: the critical quadratic	187
211.	Likewise for an equation $z^m w^{(m)} = f(z^{m-1} w^{(m-1)}, \dots, zw', w, z)$	192
212.	Likewise for an equation $zw'' = f(w', w, z)$	193
213, 214.	Non-regular integrals of $w' = f(w', w, z)$: the simplest cases	196
215.	Accidental singularity of the first kind of f , determined by the assigned initial values	198
216.	When this arises independently of an initial value of w'	200
217.	Examples	201
218.	Accidental singularity of the second kind of f , determined by the assigned initial values	208
219—221.	Reduced typical forms of the equation	210
222.	The integral of the reduced form $t^2 v'' = F(tv', v, t)$, with examples	215
223, 224.	The remaining reduced forms of § 221	220
225.	Discussion of the indeterminate case of § 218	222

CHAPTER XV.

EQUATIONS OF THE SECOND ORDER AND ANY DEGREE.

ART.		PAGE
226.	Initial conditions satisfying $F(c, a, \beta, \gamma) = 0$ for an equation $F(z, w, w', w'') = 0$ of any degree	227
227.	An integral, when γ is a simple root and finite	229
228.	A set of integrals, when γ is a multiple root and finite	230
229, 230.	Puiseux diagram applied when F is evanescent: with examples, and some omitted cases	232
231.	An integral when γ is infinite	242
232, 233.	Summary of results, and geometrical interpretation	244
234—236.	The w'' -discriminant of $F = 0$, and associated equations, in relation to possible singular integrals of $F = 0$	247
237.	A general class of equations possessing singular integrals	252
238, 239.	An equation of the second order may have two distinct kinds of singular integrals: their geometrical relations to one another when the variables are real: examples	253
240.	Corresponding theorems for equations of any order	260
241—243.	Analytical relation of singular integrals to a primitive for equation of second order, with examples	262
244, 245.	Classification of integrals derivable from primitive of an equation of the second order; with examples	269

CHAPTER XVI.

EQUATIONS OF THE SECOND ORDER WITH SUB-UNIFORM INTEGRALS:
WITH SOME GENERAL CONSIDERATIONS.

246.	Introductory remarks upon equations of second order the integrals of which have descriptive properties: <i>sub-</i> <i>uniform</i> functions	276
247.	Picard's equations $w'' = R(w, w')$ with sub-uniform integrals: first set of conditions, that the equations must have the form $w'' = A_0 + A_1 w' + A_2 w'^2$, where A_0, A_1, A_2 depend on w only	279
248.	Limitations upon A_0, A_1, A_2 : all the finite poles must be simple	282
249—253.	Further conditions, when A_0, A_1, A_2 have simple finite poles	285
254—256.	Further conditions, when A_0, A_1, A_2 have no finite poles	294
257.	Discussion of the equation $w'' = bw^2 + cw + d + hw'$	298

Cambridge University Press

978-1-107-63012-3 - Theory of Differential Equations: Part II: Ordinary Equations,
not Linear: Vol. III

Andrew Russell Forsyth

Frontmatter

[More information](#)

CONTENTS

ix

ART.		PAGE
258.	Picard's method of integration when the integral is uniform, applied by Mittag-Leffler to the equation of § 257: with other examples	301
259.	General considerations on integrals compatible with an equation $F(w'', w', w, z) = 0$; with the partial differential equation characteristic of compatible integrals	306
260, 261.	Inferences deduced from the characteristic	307

CHAPTER XVII.

GENERAL THEOREMS ON ALGEBRAIC INTEGRALS: BRUNS'S THEOREM.

262.	Statement of the theorem to be proved	311
263.	Outline of the argument	313
264.	The differential equations of the problem of n bodies, and their classical integrals	314
265.	The integrals for two bodies, being an exceptional case	316
266.	Definitions of <i>integral</i> , <i>solution</i> , <i>sub-integral</i> , <i>composite solution</i> , <i>particular solution</i> , <i>function satisfying the differential equations</i>	318
267.	Provisional introduction of equations of more general form: the irrational variable introduced	322
268.	Reduction of algebraic integrals to functions, which satisfy the differential equations and are rational in the variables and the arbitrary constants	323
269—271.	A function, containing arbitrary constants, can be algebraically composed from functions, containing no arbitrary constants	326
272.	Definition of <i>homogeneity</i> : a function, containing no arbitrary constants, can be composed from homogeneous functions	332
273.	Rejection of particular solutions: property of solutions retained; they must involve one special set of the variables	335
274—276.	Every solution can be algebraically composed of solutions, which are polynomials in that set of variables, are of only odd or only even order, and are free from imaginaries	338
277.	If a solution does not determine an integral, the sub-integral which it contains can be changed into an integral on multiplication by an appropriate factor	344
278.	Proof of the statement in § 277, when the coefficients of the highest terms in the polynomial do not involve the irrational variable	346

Cambridge University Press

978-1-107-63012-3 - Theory of Differential Equations: Part II: Ordinary Equations,
not Linear: Vol. III

Andrew Russell Forsyth

Frontmatter

[More information](#)

x

CONTENTS

ART.		PAGE
279.	The statement of § 277 not necessarily true, if the irrational variable occurs: example	351
280—282.	Establishment of Poincaré's rule for determining cases such as the example in § 279	352
283, 284.	All the cases, which occur under Poincaré's rule, are imaginary for the problem of three bodies, and for the problem of n bodies	358
285.	Summary of results obtained for the equations of § 267.	360
286.	Reintroduction of the dynamical equations: preliminary limitation on the form of ϕ_0 , taken as the leading aggregate of an integral ϕ , which is a polynomial in the velocities	361
287, 288.	The partial differential equations satisfied by ϕ_0 , to avoid logarithmic terms in ϕ : the complete solution of this system	364
289, 290.	Further transformation of ϕ_0 , and partial expression by classical integrals	370
291, 292.	Further partial differential equations satisfied by ϕ_0 , to secure that the rest of ϕ is polynomial in the velocities: the complete solution of this system	374
293.	Final transformation of ϕ_0 ; the establishment of Bruns's theorem	380
294.	Concluding remarks	383
	INDEX TO PART II.	385