

Cambridge University Press

978-1-107-63012-3 - Theory of Differential Equations: Part II: Ordinary Equations,
not Linear: Vol. III

Andrew Russell Forsyth

Index

[More information](#)

INDEX TO PART II.

(Part II occupies volumes II and III of the present work. The figures refer to the pages in these volumes.

The Table of Contents at the beginning of volume II and of volume III may also be consulted.)

- Accidental singularity of first kind, of function of two or more variables, II, 49; form of differential equation in vicinity of, II, 52; can form a continuous aggregate, when determined by the equation, II, 56; it is an algebraical critical point of the integral, save for three general exceptions, II, 62—68; when it is a point of indeterminateness, with tests, and methods of settlement, II, 71, 72, 77; general summary of effect of, upon integral of the equation, II, 79, 80;
for a system of two equations, III, 11—16;
for an equation of second order and first degree, III, 199—207.
- Accidental singularity of second kind, of function of two or more variables, II, 49; form of differential equation in the vicinity of, II, 53; are isolated, when determined by the equation, II, 56; final form of equation in vicinity of, II, 88; leading term of integral in vicinity of, II, 89; reduced forms of equation, (*see* reduced forms);
effect of double, on forms of system of two equations, III, 17 et seq.;
for equation of second order and first degree, III, 208; reduced forms in vicinity of, III, 213, 214; integral in vicinity of, III, 215.
- Aggregate of leading terms in homogeneous integral, (*see* leading aggregate).
- Algebraic integrals, equations of first order with, II, 338; determinable by means of Abelian integrals of the first kind, that belong to an associated Riemann surface, II, 341;
of systems of equations, general theorems on, Chapter XVII; reduction of, to integrals, involving no arbitrary parameter and polynomial in variables, III, 323 et seq.
- Analytical relation between singular solutions and primitive for equation of first order, II, 249; for equation of second order, III, 262.
- Apparence uniforme*, defined, III, 279.
- Appell, III, 260.
- Approximation, Picard's method of, II, 41.
- Autonne, II, 333.
- Bendixson, II, 126, 187, III, 8.
- Biermann, II, 14.
- Binomial equations, whose integrals have their critical points fixed, II, 296—308; whose integrals are uniform, II, 325—333.
- Birational transformation of Riemann surface, II, 292.
- Boole, II, 190.
- Bouquet, (*see* Briot and Bouquet).
- Branches of a function near algebraical branch-points, II, 58; will be supposed limited in number, II, 58; consequent form of a differential equation, II, 59.

- Branching of a function at a point of indeterminateness may be definite, or indefinite, II, 70.
- Branch-points, of equation of first order, II, 224; special sets of, II, 234; integrals in the vicinity of, Chapter VIII; when fixed, determine classes of equations, Chapter IX; equations devoid of parametric, (*see* parametric branch-points);
- of system of equations of first order, and integrals in the vicinity of, Chapter XIII;
 - of equation of second order and any degree, III, 230 *et seq.*
- Briot and Bouquet, II, 40, 41, 60, 62, 83, 140, 149, 156, 158, 176, 178, 182, 188, 224, 309, 311, 325, 332, 333.
- Briot and Bouquet's conditions that integrals of equation of the first order should be uniform, II, 323; applied to binomial equations, II, 325.
- Bruns, III, 311, 312, 313, 314, 316, 318, 380.
- Bruns's theorem in the problem of n bodies, III, 383.
- Canonical equivalent of system of equations, II, 14—17, III, 5; in vicinity of double accidental singularity of second kind for a system of two equations, III, 21; reduced forms of last, III, 26.
- Cauchy, II, 40; *passim*, for general existence-theorem.
- Cauchy's theorem on the existence of regular integrals, II, 26; does not exclude the possibility of integrals that are not regular, II, 44, (*see also* regular integrals).
- Cayley, II, 8, 264, 288, 318, 325; III, 134.
- Chrystal, II, 62, 189.
- Classes of singular solutions, (*see also* singular solutions); first, III, 153; second, III, 157; of system of any number of equations, III, 162.
- Classical integrals in the problem of n bodies, III, 312, 314, 370, 380.
- Classification of integrals, for a single equation, III, 176; for a system of two equations, III, 178; for an equation of second order, III, 269—273.
- Clebsch, II, 288, 338.
- Compatible with a given equation of first order, equations that are, II, 333; determinable by a partial differential equation, II, 335; can be expressed by means of any two, II, 336; effect of, when integrals are algebraic, II, 340;
- with a given equation of second order, equations that are, III, 306.
- Complete integrals and singular integrals of a system of two equations, how related, III, 166, 183.
- Composite solution, III, 320.
- Composition of algebraic integrals, III, 323 *et seq.*
- Constants may, in a particular case, be the only integrals of a system satisfying conditions, II, 31.
- Continuation of regular integral, II, 42.
- Critical curves, in geometrical interpretation of equations of the second order, III, 254.
- Critical equation for integrals of equation of any order and first degree, III, 193.
- Critical lines, in geometrical interpretation of equations of the second order, III, 254.
- Critical points of a differential equation, defined, II, 43, (*see also* exceptional points); algebraical, of an integral, II, 63.
- Critical quadratic, for system of two reduced forms, III, 45—47; as affecting integrals of, Chapter XII; for special forms of equation of second order and first degree, III, 188, 193.
- Critical reduced forms of equation in vicinity of accidental singularity of second kind, II, 96; obtained by means of leading term of integral (q.v.), II, 97; various types for the simplest case, II, 98—104; and for other cases, II, 111—114, 115—120, 121; general summary of results, II, 122—124; examples, II, 124—139.
- Curves defined by a differential equation, (*see* geometrical applications).
- Darboux, II, 264, 333, III, 137.
- Definite branching, (*see* branching of a function).
- Diagrams of Puiseux, (*see* Puiseux diagram).
- Discriminant of an equation of the first order, as affecting the integral determined by initial values, II, 241; its relation in general to the equation, II, 245; it does not in general provide a solution, II, 246; but only for particular sets of values, II, 249; if conditions are satisfied, it can provide a solution, II, 249; Hamburger's investigation of relation between integral of the equation and root of the discriminant, II, 249; can provide singular solutions and particular solutions, II, 257; how such solutions are determined, II, 261;
- of an equation of second order and any degree, connected with singular solutions, III, 247.

- Dixon, III, 121, 157, 161, 183, 253, 255, 260, 261, 262, 274.
- Dominant equations; system of, in proof of Cauchy's theorem, II, 28; in obtaining the integral of reduced forms of a single equation, II, 144, 152, 163; of a system of equations, Chapter XII.
- Dominant function defined, II, 27.
- Equation of first order, (*see* volume II); of second order, and of higher order, (*see* volume III).
- Equations of first order, having no parametric branch-points, Chapter IX; if of first degree, have Riccati's form, II, 271; if of higher degree, satisfy Fuchs's conditions, II, 284; when the genus is zero, the equations are rationally transformable to Riccati's equation, II, 287; when genus is unity, it can be integrated by transformations and quadrature, II, 291; when genus is greater than unity, integral is algebraic, II, 292; binomial, II, 296;
 - with uniform integrals, II, 310; Briot and Bouquet's conditions for, II, 323; how integrated, II, 317; binomial, II, 325.
- Equations of second order with sub-uniform integrals, III, 278—285; tests are not complete, III, 286; general conclusions, III, 292, 297, 298.
- Equations compatible with equation of first order, II, 333—344; with equation of second order, III, 306—310.
- Essential singularity of a function of two or more variables, II, 49; for an equation of first order, are isolated when determined by the equation, II, 55, and is an essential singularity of its integral, II, 210.
- Essential singularity of integrals of a system of equations may be parametric, II, 48, 216; of a single equation must be isolated fixed points, II, 211—216, 266—269; but for a system may be a continuous aggregate, II, 222; for system of two equations, III, 11.
- Evanescence of equations for initial values, how to determine a solution, III, 138.
- Evanescent form of equation of second order and any degree; how to determine the form of the integral, III, 232; with examples, III, 234—242.
- Exceptional points of a differential equation, defined, II, 43; their source, in the form of a single equation of the first order, II, 47; parametric exceptional points may occur in the integral, II, 48.
- Existence of regular integrals, Cauchy's theorem as to the, II, 26, (*see* regular integrals, non-regular integrals).
- Existence-theorems: what is required in general for their establishment, II, 3.
- Faà de Bruno, II, 8.
- Fine, II, 224, III, 121.
- First class of singular solutions, III, 153, 163.
- First kind of accidental singularity, (*see* accidental singularity).
- First order, single equation of, and first degree, Chapters III—VII;
 - single equation of, and degree higher than the first, Chapters VIII—X;
 - two equations of, and first degree, Chapters XI, XII;
 - system of equations of, and degree higher than first, Chapter XIII.
- Fixed critical (or branch-) points, (*see* parametric branch-points).
- Focal points, III, 137.
- Focal surface, in geometrical illustration of two simultaneous equations, III, 134, 137.
- Fransén, III, 276.
- Fuchs, II, 40, 45, 60, 69, 73, 81, 82, 249, 270, 285, 325, III, 9.
- Fuchs's conditions that an equation of first order and any degree shall be devoid of parametric branch-points, II, 285—296; association of Riemann surface with, II, 286.
- Genus of equations, which satisfy Fuchs's conditions as to parametric branch-points, II, 286; when zero, II, 287; when unity, II, 288; when greater than unity, II, 291.
- Geometrical applications, II, 126, III, 39, 133, 140, 227, 228, 245, 254, 384.
- Goursat, III, 23, 40, 46, 121, 137, 157, 159, 161, 183, 259, 261.
- Grünfeld, III, 8.
- Hamburger, II, 224, 249, 262, III, 3, 170, 177, 183; on relation between integral of an equation of the first order and root of the discriminant, II, 249—261.
- Hermite, II, 318, 325.
- Hill, M. J. M., III, 177.
- Homogeneous integrals in problem of n bodies, III, 332; can be taken as polynomial functions, III, 340; general properties of, III, 361.
- Horn, II, 83, 187, III, 8, 9, 47.
- Indefinite branching, (*see* branching of a function).

- Indeterminateness, point of, II, 60; as associated with functions, II, 69; branching of functions near such a point, II, 70; effect upon expressibility of the function, II, 71; when an accidental singularity of the first kind is, II, 72, 77; examples of, II, 73, 79; effect of, upon uniqueness of regular integrals, II, 81; for integral of the third typical form, II, 187; for integrals of remaining forms in general, II, 201, 204; with example of exception, III, 202.
- Integral, defined for problem of n bodies, III, 319, 335, 341.
- Integral in vicinity of accidental singularity of first kind, (*see* accidental singularity of first kind): in vicinity of accidental singularity of second kind, (*see* accidental singularity of second kind).
- Integral of a single equation cannot have parametric essential singularities, II, 211; when algebraic, (*see* algebraic integrals); when uniform, II, 310; Briot and Bouquet's conditions for, II, 323; belong to one of three classes, II, 318.
- Integral, singular, (*see* singular solution).
- Integrals, algebraic, of a system of $2n$ equations, III, 312, Chapter XVII; Bruns's theorem on, III, 383.
- Integrals, classes of, for a single equation, III, 176; for two simultaneous equations, III, 178; regular, (*see* regular integrals).
- Integrals of equation of first order in the vicinity of a branch-point, Chapter VIII; summary of results, II, 239; affected by the discriminant, II, 241; relation between, and root of the discriminant, II, 249; general classes of, II, 261.
- Integrals of reduced forms of system of two equations, various cases, Chapter XII; regular, III, 49–69; non-regular, III, 69–116; for system of any number, III, 116–118.
- Integrals of reduced forms of equation of the second order, III, 215–222.
- Integrals of equation of second order, classes of, III, 269–273.
- Integrals, equation of second order with sub-uniform, III, 278; tests for, III, 279–285; further conditions necessary, III, 286; with summary of results, III, 286–303.
- Integrals of equation of any order and any degree, III, 260; classes of, III, 261.
- Integralgleichungen*, III, 314.
- Invariantive equation in linear transformations, III, 3.
- Irrational sub-integrals in the problem of n bodies are complex, III, 351 *et seq.*
- Jacobi, II, 19.
- Jacobian, of system of two equations, effect of upon integrals, when it vanishes for initial values, III, 127–133; when it vanishes in general, III, 148; special note upon, III, 158; of system of any number of equations, III, 163; limiting form of, for equation of second order and any degree, III, 228.
- Jordan, II, 1, 19, 35, 41, 140, 163, 310, III, 9, 47, 109.
- Kinds of singularities of a function of two or more variables, II, 49; general characteristics of, for a single equation, II, 55; effect of, in considering the variations of the independent variable, II, 57, (*see also* accidental singularity, essential singularity).
- Königsberger, II, 1, 14, 41, III, 8, 9, 17, 23, 40, 47, 87, 195.
- Korkine, II, 333.
- Kowalevsky, II, 41.
- Lagrange, III, 259, 312, 320.
- Leading aggregate of homogeneous integral, III, 362–382; determined by two sets of characteristic equations, III, 366, 378.
- Leading term of integral in vicinity of accidental singularity of second kind, II, 89, 107; obtained by means of Puiseux diagram, II, 91, 108; applied to obtain typical reduced forms of equations for the simplest case, II, 97–104; in other cases, II, 107–114, 115–120, 121; examples, II, 124–139.
- Lie, III, 384.
- Lipschitz, II, 40.
- Majorante* (dominant), II, 27.
- Mayer, III, 121, 157, 159, 160, 165, 183, 261.
- Méray, II, 40.
- Mittag-Leffler, III, 276, 301, 302, 304.
- Moigno, II, 40.
- Non-regular integrals, existence of, not excluded by Cauchy's theorem, II, 44; example of, satisfying same conditions as regular integrals, II, 45; possessed by equations of first order in first typical reduced form, II, 149, 158; by the second typical reduced form, II, 181; of the third one, in general, the only integrals, II, 187–201; special case of the third, II, 198.

- Non-regular integrals of reduced forms for system of two equations, *iii*, 69–116; for system of any number, *iii*, 116–118.
- Non-regular integrals of equation of second order and first degree, *iii*, 196.
- Normal form, for a system of equations, Weierstrass's, *ii*, 14.
- Normal forms of systems of equations, *ii*, 6, 14, 17.
- Osculants in theory of plane curves, associated with equations of the second order, *iii*, 227, 254–256, 269–273.
- Painlevé, *ii*, 47, 209, 211, 212, 217, 224, 266, 270, 333, 338, 344, *iii*, 184, 276, 312, 383.
- Painlevé's theorem on fixity of essential singularities of a single equation of first order, *ii*, 211, 266; examples, *ii*, 218; not necessarily valid for a system of equations, *ii*, 48, 219, 220.
- Parametric branch-points of an integral, *ii*, 48.
- Parametric branch-points, equations having integrals that are devoid of, Chapter *ix*; Fuchs's conditions, *ii*, 285, (*see* equations of first order, having no parametric branch-points).
- Parametric essential singularity of integrals of a system, *ii*, 48.
- Partial differential equations, sets of, characteristic of leading aggregate in homogeneous integral of problem of n bodies, *iii*, 366, 378.
- Particular solutions of equation of first order, *ii*, 257; tests for, *ii*, 261; examples, *ii*, 265.
- Particular solution of equation of second order, criterion for, *iii*, 267.
- Particular solutions in problem of n bodies, *iii*, 320, 321, 337.
- Phragmén, *ii*, 310.
- Picard, *ii*, 19, 40, 41, 45, 47, 60, 126, 140, 212, 270, 292, 310, 338, *iii*, 8, 9, 46, 276, 278, 301, 302, 304, 383, 384.
- Picard's equations of second order with sub-uniform integrals, *iii*, 279–285; further conditions, with results, *iii*, 292–303.
- Poincaré, *ii*, 27, 41, 126, 140, 149, 158, 270, 292, 296, 333, *iii*, 8, 311, 312, 343, 351, 384.
- Poincaré's theorem on integrals of equations that satisfy Fuchs's conditions as to fixity of branch-points, *ii*, 295; on Bruns's theorem as to problem of n bodies, *iii*, 311, 312, 343, 351, 384.
- Problem of n bodies, *iii*, 312; classical integrals of, *iii*, 312, 314; algebraic integrals of, Chapter *xvii*; Bruns's theorem on, *iii*, 383.
- Problem of three bodies, *iii*, 351, 358, 372; Bruns's theorem in, *iii*, 383.
- Problem of two bodies, *iii*, 316; Bruns's theorem does not apply, *iii*, 383.
- Puiseux diagram, generalised into space diagrams, *iii*, 23.
- Puiseux diagram used to determine leading term of integral for equation of first order, *ii*, 91, 108, 116–118, 121, 232, 236; examples of, *ii*, 124–139; generalised, *iii*, 23, 138.
- Puiseux diagram used for equation of the second order, *iii*, 209, 222, 232, 234, 243.
- Quadratic for system of two reduced forms, critical, (*see* critical quadratic).
- Raffy, *ii*, 325.
- Rational sub-integrals can be changed to integrals in problem of n bodies, *iii*, 344.
- Real integrals alone retained in problem of n bodies, *iii*, 343.
- Reduced forms of equation of first order in vicinity of accidental singularity of second kind for the simplest case, *ii*, 98–104; and for other cases, *ii*, 111–114, 115–120, 121; general summary of results, *ii*, 122–124; examples, *ii*, 124–139.
- Reduced forms of equation of second degree in vicinity of accidental singularity of second kind, *iii*, 214; integrals of, *iii*, 215–222.
- Reduced forms of system of two equations in vicinity of double accidental singularity of the second kind, with examples, *iii*, 26–39; integrals of, Chapter *xii*.
- Reduction of algebraic integrals to integrals, which involve no arbitrary constants and are polynomial in the variables, *iii*, 323 *et seq.*
- Regular, as applied to functions: the term defined, *ii*, 3;
as applied to the solution of linear differential equations, *ii*, 73, 193, *iii*, 9.
- Regular function of several variables near a zero value, Weierstrass's theorem on form of, *ii*, 19; special case of two variables, *ii*, 24, 50.
- Regular integrals, Cauchy's theorem as to the existence of, *ii*, 26; can become mere constants in a particular case, *ii*, 31; they are unique as regular integrals, *ii*, 33; continuation of, *ii*, 42; Fuchs's theorem to determine

- whether they are the only integrals satisfying initial conditions, II, 81, and an example, II, 82.
- Regular integral possessed by equation in first typical reduced form, II, 146, 156, 171; with conditions when one particular coefficient is a whole number, II, 156, 170; by the second typical form, II, 181; by the third typical form, II, 187; and by special case of the third, II, 198.
- Regular integrals of reduced forms for system of two equations, III, 49–69; for system of any number, III, 116.
- Regular integrals of equation of second order and first degree, III, 184–196.
- Relation between singular integrals and complete integrals, for a single equation, II, 249; for a system of two equations, III, 166, 183.
- Riccati's equation, all the branch-points are fixed, not parametric, II, 271–276, 307.
- Riemann surface, associated with Fuchs's conditions that equation shall be devoid of parametric branch-points, II, 286, 296; birational transformation of, II, 292; for binomial equations, II, 297; with Briot and Bouquet's conditions for uniform integrals, II, 317; used first by Clebsch, II, 338; applied to determine algebraic integrals of an equation of first order, II, 341.
- Riquier, II, 40.
- Salmon, II, 8, 287.
- Satisfy differential equations in the problem of n bodies, when a function is said to, III, 321.
- Sauvage, III, 8.
- Schwarz, II, 333.
- Second class of singular solutions, III, 157, 163.
- Second kind of accidental singularity, (*see* accidental singularity).
- Second order, equation of, and first degree, Chapter xiv; existence-theorem for regular integrals of, III, 186; existence-theorems for regular integrals of particular cases of, III, 187, 193; non-regular integrals of, III, 196; in vicinity of accidental singularity of first kind, III, 199–207; in vicinity of accidental singularity of second kind, III, 208 et seq.; indeterminate case, III, 222.
- Second order, equation of, and degree higher than the first, Chapter xv; regular integrals of, III, 229; multiform integrals of, III, 230, 242; interpreted geometrically, III, 245; singular solution of, III, 247.
- Second order with sub-uniform integrals, equations of, III, 279–285; further conditions and results, III, 286–303.
- Second order, equations compatible with equation of, III, 306–310.
- Serret, III, 159, 182, 183, 259.
- Simart, II, 19, III, 276.
- Single equation of first order, (*see* first order).
- Singular solutions of equation of first order, II, 224, 243, 257; tests for, II, 261; examples, II, 264.
- Singular solutions of equation of second order and any degree, III, 247; condition for, III, 248; class of equations which necessarily possess, III, 253; can be of two kinds, III, 253; how to construct equations which possess, III, 255; represent osculants, III, 254–256; how analytically connected with primitive, III, 262.
- Singular solutions of system of equations, III, 162; of equations of any order and any degree, III, 261.
- Singular solutions of two simultaneous equations, III, 23, 148; conditions necessary and sufficient for, III, 153; are of two classes, with the further conditions for the second class, III, 156; how related analytically to the primitive, III, 166, 183.
- Singularities of a single equation of first order, II, 43; of a function of more than one variable, II, 49.
- Singularities of systems of equations, III, 1; possibilities among the various combinations for a system of two equations, III, 10; integrals in vicinity of simplest among the combinations of, III, 11–16; form of, in vicinity of double accidental singularity of second kind, III, 17–21.
- Skew curves in space, used to illustrate two simultaneous differential equations, III, 133–137.
- Solution defined for problem of n bodies, III, 319, 335; composite solution, III, 320; particular solution, III, 320, 321.
- Solutions, singular, (*see* singular solutions).
- Space diagrams, as generalisation of Puiseux diagrams, used for system of two equations, III, 23, 138.
- Sub-integrals, III, 314, 319, 336, 341 et seq.; when rational, can be changed to integrals, III, 344 et seq.; when irrational, are complex, III, 351 et seq.
- Sub-regular integral, in ordinary linear differential equations, II, 73, 193.
- Sub-uniform integrals, equations of second order with, III, 278–285; further conditions, with results, III, 286–303.

Cambridge University Press

978-1-107-63012-3 - Theory of Differential Equations: Part II: Ordinary Equations, not Linear: Vol. III

Andrew Russell Forsyth

Index

[More information](#)

INDEX TO PART II

391

- System of equations, not of first degree, integrals of, Chapter XIII; singular solutions of, III, 148–164; singularities of, (*see* accidental singularity, essential singularity).
- Thomé, II, 73.
- Three bodies, problem of, (*see* problem of three bodies).
- Two bodies, problem of, (*see* problem of two bodies).
- Typical form for a system of equations, where the functions are regular, II, 17.
- Typical forms of system of equations near accidental singularity, (*see* reduced forms).
- Typical reduced forms of equations, in vicinity of accidental singularity of second kind, for the simplest case, II, 98–104; for other cases, II, 111–114, 115–120, 121; general summary of results, II, 122–124; examples, II, 124–139.
- Typical reduced forms of single equation in the vicinity of an accidental singularity of second kind; integrals of the first, in various cases, II, 141–178; regular integral of the second, II, 178; non-regular integrals of the second, II, 181; integrals (if any) of the third are in general non-regular, II, 187–201; integrals of the remaining typical forms are in general non-regular, II, 201; Summary of results for integrals of, in all cases, II, 207.
- Uniform integrals, equations of the first order with, Chapter X; Briot and Bouquet's conditions for, II, 323; how obtainable, II, 317; belong to one of three classes, II, 318.
- Unique determination of regular integrals, II, 33; Jordan's proof of, for a system, II, 35; fundamental assumption used in proof, II, 45; uniqueness of such integrals as affected by point of indeterminateness, II, 81.
- Wallenberg, II, 270, 296, 307, 308, 309, III, 276, 304, 305, 306.
- Weierstrass, II, 3, 14, 15, 19, 41, 42, 49, 83, III, 9; theorem on form of a regular function near a zero value, II, 19, with special case of two variables, II, 24, 42; applied to transform differential equations, II, 50.