

Index

- Abortion Act 1967, 9, 13
 fetal anomaly, 153
 abortion counselling, *see* support and counselling
 abortion incidence complications method (AICM), 25
 Abortion Law Reform Act (2008) (Vic), 240
 abortion methods, 89
 grief resolution and, 89
 historical approaches, *see* historical perspectives
 medical abortion, *see* medical abortion
 mortality and, 123–4
 surgical abortion, *see* surgical abortion
 acceptability
 TOPFA (termination of pregnancy for fetal anomaly), 157–8
 vacuum aspiration abortion (VA), 72
 access to abortion, *see* human rights
 acute intermittent porphyria, 137
 Advancing New Standards in Reproductive Health (ANSIRH), 251
 Africa, *Protocol to the African Charter on Human and Peoples' Rights on the Rights of Women in Africa*, 247
 age profiles, 29–30
 second trimester abortions, 204–7
 allergic reactions, local anaesthesia, 99
 American Congress of Obstetricians and Gynecologists (ACOG), 251
 professional responsibilities, 188–9
 anaemia, 136
 post medical abortion, 128–9
 anaesthesia
 complications related to, 126
 general, *see* general anaesthesia
 international approaches, 102
 local, *see* local anaesthesia
 for medical abortion (TOPFA), 156
 medical co-morbidity and, 134
 indications, 134
 obesity and, 135
 pre-operation, 134
 types of anaesthesia, 134
 analgesia
 dilatation and evacuation (D&E), 90, 91
 early medical abortion (EMA), 67, 103
 medical abortion (higher gestation), 85
 non-steroidal anti-inflammatory drugs (NSAIDs), 85
 vacuum aspiration abortion (VA), 73
see also pain management
 antibiotics
 complications and, 123
 dilatation and evacuation (D&E), 95
 prophylactic, 127
 early medical abortion (EMA), 66
 vacuum aspiration abortion, 73
 World Health Organization (WHO) guidance, 129
 sexually transmitted infections (STIs), 66
 anxiety, pain and, 102–3
 association and causality, 163–5
 analogous associations, 165
 biological mechanism, 164
 consistency, 164
 definition, 163
 dose-response effect, 164
 epidemiology and, 164
 research evidence, 164
 specificity, 164
 strong association, 164
 temporal factors, 164
 Association of Reproductive Health Professionals (ARHP), 251
 asthma, 136
 attitudes to abortion, 10–14
 Catholic, 11
 feticide, 146–8
 health professionals, 12–14, 212
 judgementalism in abortion care, *see* health professionals
 legal, 12
 philosophical, 10–11
 public, 12
 repeat abortions, 193, 216–17
 stigma, *see* stigma
 Australia, 236–43
 Australian Capital Territory (ACT), 240
 historical approaches, 237–9
 legal context, 236–7
 mifepristone, 63
 Offences Against the Person Act (OAP) (1861), 237
 Tasmania, 241–2
 conscientious objection, 242
 Reproductive Health (Access to Terminations) Act 2013 (Tas), 241
 Victoria, 240–1
 Abortion Law Reform Act (2008) (Vic), 240
 conscientious objection, 241
 termination of pregnancy for fetal anomaly (TOPFA), 240
 Western Australia, 239
 autonomy (of woman), 16
 biparietal diameter, 56
 bleeding
 anaemia, 128–9
 dilatation and evacuation (D&E), 95
 disseminated intravascular coagulopathy (DIC), 125
 haemorrhage, *see* haemorrhage
 breast cancer, 168, 169, 177–8

- caesarean section, uterine scars, 139
- Canada, telemedicine (Women on Web), 228–9, 231–2
- cancer, breast, 168, 169, 177–8
- cardiovascular disease, 138
- Catholicism, 11
- Center for Reproductive Rights, 251
- cervical laceration, surgical abortion, 124
 dilatation and evacuation (D&E), 94
- cervical preparation
 dilatation and evacuation, *see* dilatation and evacuation (D&E)
 medical co-morbidity, 134
 obesity, 135
 surgical problems and, 139
 vacuum aspiration abortion, *see* vacuum aspiration abortion (VA)
- Chiba needle, 148, 149, 150
- Chlamydia trachomatis*, 66, 127
- 'choice' argument
 autonomy (of woman), 16
 women's rights and, 16
- coercion, 44, 48
- Committee on the Elimination of Discrimination against Women (CEDAW), 189
- complete abortion, rates and complications, 65
- compliance, 107
 follow-up clinic visit, 107
- complications of abortion
 continuing pregnancy, *see* follow-up
 of dilatation and evacuation, *see* dilatation and evacuation (D&E)
 factors associated, 31
 abortion method, 123–4
 abortion provider's skill, 123
 antibiotics and cervical priming, 123
 gestational age, 123
 in safe abortions, 123–4
 future pregnancy
 complications, 95
- incomplete abortion,
see retained products of conception (RPOC)
- medical abortion, 128–30
 continuing pregnancy, 130
 early, 128
 heavy or prolonged bleeding, 128–9
 higher gestation, 84–5, 130
 incomplete abortion, 129–30
 infection, *see* infection postabortion (endometritis)
 uterine rupture, *see* uterus
- second trimester abortion, 157, 202
- surgical abortion, 124–8
 anaesthesia-related complications, 126
 cervical laceration, 124
 continuing pregnancy, 127–8
 dilatation and evacuation (D&E), *see* dilatation and evacuation (D&E)
 disseminated intravascular coagulopathy (DIC), 125
 haematometra, 126
 haemorrhage, 125
 incomplete abortion, 126
 infection, *see* infection postabortion (endometritis)
 mid-level provision, 221–2
 uterine perforation, *see* uterus
- vacuum aspiration abortion (VA), *see* vacuum aspiration abortion (VA)
 vasovagal reaction, 124
see also morbidity (abortion-related)
- Concept Foundation, 251
- conscience/conscientious acts, 187–90
- conscientious
 commitment, 187–8
- conscientious objection, 187, 204, 212–14
- Australia (Victoria), 241
 General Medical Council (GMC) (UK), 213
 Nursing and Midwifery Council (NMC) (UK), 213
 Royal College of Obstetricians and Gynaecologists (RCOG), 213
 Tasmania, 242
 USA, 213–14
- definition, 188
 guidelines, American Congress of Obstetricians and Gynecologists (ACOG), 188–9
- refusal of care, 188–90
 Committee on the Elimination of Discrimination against Women (CEDAW), 189
 health system obligations, 189–90
 professional responsibilities, 188–9
- right of, 188
 right to, 187
 stigma and, 187
- consensual issues
 principle of retraction, 21
 telemedicine, 233
see also consent
- consent 66, 72, 90, 227, 232, 233
 dilatation and evacuation (D&E), 90
 vacuum aspiration abortion (VA), 72
see also consensual issues
- continuing pregnancy
 medical abortion, 130
 surgical abortion, 127–8
see also follow-up
- contraception
 advice at time of abortion, 181–2
 anti-progesterone drugs as, 62
 efficacy, 179–80
 emergency contraception, 179
 medical co-morbidity, 140
 partner violence and, 199
 postabortion, 66, 68, 72, 110–11, 114–21
 after medical abortion, 111, 118
 after surgical abortion, 96, 111, 117–18
 commencement, 110–11, 117–19
 cost-effectiveness, 119
 current research, 119–120
 hormonal methods, 117
 intrauterine device (IUS/IUD), 111, 116–7, 118
 long-acting reversible contraceptive (LARC), 115, 119
 mid-level providers, 223
 ovulation, return of, 115
 repeat abortions and, 198–9
 risk of subsequent abortion and, 115–7
 repeat abortions and, 193, 198, 199
 unsafe abortions and, 114

- counselling, *see* support and counselling
- cremation and burial
 fetal remains, 216
 Institute of Cemetery and Crematorium Management (ICCM), 216
 TOPFA (termination of pregnancy for fetal anomaly), 159
- criminal abortion (historical context), 8
- culture, sex-selective abortion and, 20
- Czech Republic, 171
- decision-making process
 coercion, 44, 48
 contradictory feelings and complexity, 37–8
 informed decision, 43
 in legally restricted jurisdictions, 50
 male progenitors and, 35–6
 method of abortion, 84
 grief resolution, 89
 vacuum aspiration abortion (VA), 72
 planning and pregnancy, 34
 setting for EMA, 65
 support and counselling, 43–4, 178, 179
 additional support requirements, 45–6
 case example, 47–8
 duration, 46
 function of, 46–7
 key aspects, 44
 with significant others, 48
 timing, 37
 TOPFA, *see* TOPFA (termination of pregnancy for fetal anomaly)
- deep sedation, 101, 102
- demographics (abortion), *see* epidemiology
- denied abortion, 170–1, 248
 effects on the child born, 171
 effects on the woman, 170–1
 outcomes and, 42–3
 second trimester, 202
- diabetes mellitus, 135–6
- digoxin, feticide, 94, 145
 administration procedure, 94, 150, 150–1
 failed feticide, 151
- Dilapan-S[®], 91
- dilatation and evacuation (D&E), 88–97
 background, 88–9
 cervical preparation, 90–2
 Dilapan-S[®], 91
 laminaria, 91
 mifepristone, 92
 misoprostol, 92
 osmotic dilators' placement, 91–2
 pain management, 91
 complications, 81, 94, 157
 bleeding, 95
 cervical laceration, 94
 incomplete abortion, 95
 infection, 95
 long-term sequelae, 95
 maternal mortality, 94
 uterine perforation, 94
 uterine scar rupture and, 86
- dilatation and extraction (D&X), 88, 93
- feticide, 94
- indications, 88
- operative procedure, 92–3
 forceps, 93
 instruments, 92
 vacuum aspiration, 93
- placenta accreta, 90
- postabortion care, 95–6
- pre-procedure assessment, 90
 consent, 90
 pain management, 90
 ultrasound scanning in, 90
- prevalence, 88, 89–90
- safety and effectiveness, 89
 maternal mortality, 89
 setting, 88
- dilatation and extraction (D&X), 88, 93
- disseminated intravascular coagulopathy (DIC), 125
- dizygotic twins, 56–7
- doctors, *see* health professionals
- double decidual sac sign (DDSS), 53–4, 55
- Down syndrome, Australia (Vic), 240
- early medical abortion (EMA), 62–70
 analgesia, 67, 103
 antibiotics, prophylactic, 66
 contraindications, 66
 drugs
 internet, from the, 69
 methotrexate, *see* methotrexate
 mifepristone, *see* mifepristone
 misoprostol, *see* misoprostol
 'off-label' drugs, 63
 follow-up, 68
 by telephone, 68
 gestation sac in, 66
 infection, 68
 management, 67–8
 mortality and, 123
 pain during, 103
 process, 64
 protocols, 66–7
 setting, decision-making process, 65
- ectopic pregnancy, 166
- Education for Choice, 251
- electric pump (EVA), 76
- email, 229
- embryo
 cardiac activity, 54
 definition, 54
 moral worth, 19
 ultrasound, 54–5, 56
- emergency contraception, 179
- emotional wellbeing
 contradictory feelings, 37–8
 postabortion, 39, 111
 preabortion, 37–8
 ultrasound image viewing, 58–9
see also support and counselling
- enacted stigma, 185
- endometritis, *see* infection
 postabortion (endometritis)
- epidemiology, 24–33
 abortion ratio, 28–9
 association and causality, *see* association and causality
 demographic factors, 29–30, 35, 114
 age profiles, *see* age profiles
 marital status, 29
 social determinants, 30
 historical perspectives, *see* historical perspectives
 legal abortions, 10
 legal considerations and, 29
 measurement, 24–5
 abortion incidence
 complications method (AICM), 25
 regression equation approach, 25

- epidemiology (*cont.*)
 prevalence, *see* prevalence
 repeat abortions, *see* 'more than one abortion'
 unsafe abortion, 24
- epilepsy, 138
- ethics, 16–23
 conscience, *see* conscience/
 conscientious acts
 in counselling, 50
 personhood, criteria for, 17–18
 moral worth, 19
 potentiality, *see* potentiality
 policies, *see* policies
 professional
 responsibilities, 188–9
- rights
 individual's, *see* rights
 ownership rights, 17, 21
 sex selection, 206–7
 viability, *see* viability
- Exhale, 251
- failed pregnancies, ultrasound
 scanning, 57–8
- felt stigma, 185
- fertility, 165, 176–7
- fetal anomaly, 143–4, 153–62
 counselling following
 diagnosis, 155–6
 medical conditions, 154, 155
 patterns and trends, 153–5
 prevalence of abortion, 154
 prevalence of anomalies, 153–4
 termination of pregnancy for,
see TOPFA (termination
 of pregnancy for fetal
 anomaly)
- fetal reduction, *see* feticide
- fetal remains, 216
- feticide, 142–52
 definition, 142
 dilatation and evacuation
 (D&E), 94
 failed feticide, 151
 fetal anomaly
 termination for, *see* TOPFA
 (termination of
 pregnancy for fetal
 anomaly); *see also* fetus
- fetal reduction, 56, 143
 risks, 143
- higher gestation abortion and,
 143–4, 151
 indications, 142–4
 medical abortion (higher
 gestation), 85–6
- methods, 85, 144–5
 digoxin, *see* digoxin
 mechanical, 144
 pharmacological, 144, 145–6
 potassium chloride,
see potassium chloride
 (KCl) (intracardiac)
 risks, 145–6
- opinions/attitudes, of health
 professionals, 146–8
- research evidence, 144
- Royal College of Obstetricians
 and Gynaecologists
 (RCOG)
 recommendations, 85–6,
 143–4, 145, 146, 159
- selective feticide, 143
- settings, 146–8
- Society of Family Planning (SFP)
 (USA), 144
- unintended live births, 143, 159
 RCOG guidelines, 143–4
- fetus
 definition, 54
 moral worth, 19
 pain, 178
 ultrasound (first trimester), 55
 viability, *see* viability of fetus
- FIAPAC, 252
- fibroids, uterine, 139
- FIGO (International Federation
 of Gynecology and
 Obstetrics), 252
- Finland, 204
- first trimester
 abortion during, *see* early
 medical abortion (EMA)
 ultrasound scanning,
see ultrasound scanning
- follow-up, 106–13
 complications, detection of,
 109–10
 confirmation of a successful
 abortion, 106–7
 exclusion of a continuing
 pregnancy, 107–9, 110,
 127–8, 130
 self-assessment, 107, 109
 urinary tests, *see* pregnancy
 guidelines (RCOG, WHO), 106
- indications, 106
- patient compliance, 107
- procedure types
 early medical abortion
 (EMA), *see* early medical
 abortion (EMA)
 medical abortion, 106
- outpatient abortion, 68
 termination of pregnancy for
 fetal anomaly (TOPFA),
 159–60
 vacuum aspiration abortion
 (VA), 78, 106
- telemedicine, 233
- ultrasound scanning, 107
- forceps, 93
 dilatation and evacuation
 (D&E), 93
 McClintock forceps, 93
 ring forceps, 93
- Frankenhauser plexus, 98
- funding
 historical, 8
 NHS funded abortions, 133
 second trimester abortions,
 204
- gemeprost, 82
- general anaesthesia, 102
 characteristics, 102
 levels of sedation, 100
 deep sedation, 101, 102
 medical co-morbidity and, 134
 postoperative pain,
 decreasing, 102
 regimens, 102
- General Medical Council (GMC)
 (UK), conscientious
 objection, 213
- general practitioners (GPs),
 judgementalism,
 215–16
- gestational age
 early abortion, *see* early medical
 abortion (EMA)
 higher gestation, *see* higher
 gestation abortion
 late/second trimester abortions,
see second trimester
 abortion
 terminology, 201
- gestational limit, 181, 201, 202
- gestation sac, 53, 54
 double decidual sac sign
 (DDSS), 53–4, 55
 in early medical abortion, 66
 mean sac diameter (MSD), 53
 pseudosac, 55
 yolk sac, 54
- grief resolution, 89
- Guttmacher Institute, 252
- Gynuity Health Projects, 252
- haematometra, 126

- haemorrhage
 disseminated intravascular
 coagulopathy (DIC), 125
 misoprostol and, 128
 surgical abortion, 125
- hCG measurement, postabortion,
 107, 115
- healthcare providers, *see* health
 professionals
- health professionals, 211–18
 attitudes to abortion,
see attitudes to abortion
 categories of healthcare
 providers, 217
 mid-level providers, *see* mid-
 level providers
 midwives, 221, 222–3
- conscience and, *see* conscience/
 conscientious acts
- conscientious objection,
see conscience/
 conscientious acts
- information provision, 214–15
 judgementalism, 215–16
 general practitioners (GPs),
 215–16
 non-judgemental care, 215
- medical abortion and, 216
 mortality and, 123
- personal challenges, 211–12
 ‘inoculation theory’, 212
 length of service, 212
 therapeutic bond, 211
- professional
 responsibilities, 188–9
- role of, 211
- shortage of, 219, 224
- specific challenges, 216–17
 advanced gestation, 216
 fetal remains, 216
 more than one abortion, 216–17
- support and counselling, role
 in, 217
- unpopular work, 214–15
 ‘stigma by association’, 214
- Hegar dilator, 74
- hepatic impairment, 136
- higher gestation abortion
 definition, 81
 medical abortion, *see* medical
 abortion
- termination of pregnancy for
 fetal anomaly (TOPFA),
 154–5, 160–1
- see also* second trimester
 abortion; third trimester
 abortion
- high sensitivity urine pregnancy
 test (HSUP), 107, 109
- historical perspectives, 6–15, 175
 attitudes, *see* attitudes to
 abortion
 in Australia, 237–9
 frequency of abortion, 8–10
 funding, 8
 maternal mortality, 9
 methods of abortion, 6–8
 criminal abortion, 8
 infanticide, 7
 medical, 6–7, 62, 80–1
 self-abortion, 7
 surgical, 7
 procurement of abortion, 12
 stigma, 9
- home, abortion at, *see* outpatient
 abortion
- Hong Kong, 170–1
- human chorionic gonadotrophin
 (hCG), 107, 115
- human immunodeficiency virus
 (HIV) infection, 138
- human rights, 244–50
 access to abortion, 248–9
 denied abortion, 248
 social inequality, 248
*International Conference
 on Population and
 Development (ICPD),
 see International
 Conference on Population
 and Development (ICPD)*
- law reform, 246–8
- legal rights, 245–6
 governance, 245
*Protocol to the African Charter
 on Human and Peoples’
 Rights on the Rights of
 Women in Africa*, 247
- public health
 limitations, 248
 political consensus
 and, 244–5
 social inequality, 248
 unsafe abortion, 244–6
- reproductive rights, 244
 right to conscience, 187
- hypertonic saline, 80–1
- Ibis Reproductive Health, 252
- illegal abortions, *see* unsafe
 abortion
- incomplete abortion, *see* retained
 products of conception
 (RPOC)
- Individual Level Abortion Stigma
 (ILAS) scale, 186
- infanticide, 7, 19
- infection postabortion
 (endometritis), 127
 antibiotics, *see* antibiotics
 medical abortion, 129
 early medical abortion, 68
 mortality and, 129
 retained products of conception
 (RPOC) and, 127
- information, patient, 214–15
 telemedicine, 233
- informed decision, 43
- ‘inoculation theory’, 212
- Institute of Cemetery and
 Crematorium
 Management
 (ICCM), 216
- internalized stigma, 185
- international approaches
 Africa, 247
 anaesthesia, 65, 72, 73, 102
 Australia, *see* Australia
 Canada, 228–9, 231–2
 Czech Republic, 171
 Finland, 204
 Hong Kong, 170–1
 Nepal, 224
 prevalence (safe/unsafe), 26
 Romania, 31
 Scandinavia, 36
 Sweden, *see* Sweden
 Tasmania, *see* Australia
 UK, *see* UK
 USA, *see* USA
- International Conference on
 Population and
 Development (ICPD)*
 reproductive rights, 244
 unsafe abortion, 244–5
- International Consortium for
 Medical Abortion
 (ICMA), 252
- International Federation of
 Gynecology and
 Obstetrics (FIGO), 252
- International Planned Parenthood
 Federation (IPPF), 252
- internet
 medical abortion and, 229
 early medical abortion
 (EMA), 69
 website only (Women on Web),
 229–30, 231–2
- intrauterine device (IUD), 111
- Ipas, 253

- Karman cannula (VA), 74, 75
- laminaria, 84
 dilatation and evacuation (D&E), 91
- late abortions, *see* second trimester abortion
- legal considerations
 attitudes to abortion
 Australia, *see* Australia
 availability of care and, 219–20
 banning abortion, 181
 decision-making process and, 50
 epidemiology and, 24–5, 29
 fetal anomaly (TOPFA), 153
 gestational age, 201
 gestational limit, 181, 201, 202
 human rights and, *see* human rights
 international, 203
 mortality rate and, 123
 repeat abortions and, 196
 safe/unsafe abortions and, 31
 second trimester abortions, 203, 209
 sex selection, 19
 task shifting/sharing, *see* mid-level providers
 telemedicine abortion service, 232–3
- legislation, *see* Abortion Act 1967; policies
- lidocaine, 99
- local anaesthesia (surgical abortions), 99–101
 additives for PCB (paracervical block), 100–1
 allergic reactions, 99
 amides, 99
 compared to placebo or no intervention, 100
 injection
 site, 99, 101
 technique, 101
- long-acting reversible contraceptive (LARC), 115, 119
- long-term outcomes, *see* outcomes
- low sensitivity urine pregnancy test (LSUP), 107, 108–9
- male progenitors
 decision-making and, 35–6
 rights, 20–1
- manual pump (MVA), 76, 77
- Marie Stopes International (MSI), 253
- marital status, 29
- McClintock forceps, 93
- mean sac diameter (MSD), 53
- medical abortion
 complications, *see* complications
 of abortion
 early, *see* early medical abortion (EMA)
 for fetal anomaly, 156
 follow-up, 106
 health professionals and, 216
 higher gestation abortion (after 9 weeks), 80–7, 89
 9 to 13 weeks' gestation, 83
 13 to 24 weeks' gestation, 83–4
 definition, 80
 feticide before abortion, 85–6
 laminaria, 84
 methods, 81
 osmotic dilators, 84
 pain management, *see* pain management
 preabortion, 84
 assessment, 84
 prevalence, 80
 prostaglandins, *see* prostaglandins
 safety, 80
 history of, 6–7, 62, 80–1
 mid-level providers, role of, *see* mid-level providers
 pain management, *see* pain management
 postabortion care, contraception, 111, 118
 rates of complete and complications, 65
 refusal to provide abortion care, 188
see also conscience/conscientious acts
- settings, 64, 220
 clinic, 64
 contraindications, 65
 home, *see* outpatient abortion
 medical co-morbidity, 135
 telemedicine, *see* telemedicine
- vs surgical abortion, 64–5
 unsafe abortion, 85
 vacuum aspiration, *see* vacuum aspiration
- medical co-morbidity, 133–41
 anticipated surgical problems, 139–40
- physical disabilities, 140
 trophoblastic disease, 140
 twin pregnancy, 140
 uterine conditions, 139
- conditions, 133
 contraception, 140
 general principles of
 management, 134–5
 anaesthesia, *see* anaesthesia
 cervical preparation, 134
 environment, 135
 ultrasound, 134
- specific medical conditions, 135–8
 acute intermittent porphyria, 137
 anaemia, 136
 asthma and respiratory diseases, 136
 cardiovascular disease, 138
 diabetes, 135–6
 epilepsy, 138
 hepatic impairment, 136
 human immunodeficiency virus (HIV) infection, 138
 obesity, *see* obesity
 platelet disorders, 137
 psychiatric conditions and substance misuse, 138
 renal disorders, 136
 sickle cell disease, 137
 thrombophilia, *see* thrombophilia
 thyroid disease, 136
see also morbidity
- mental health, 138, 168–70
 abortion and, 20
 myths, 177
 'postabortion syndrome', 168–9
 risk factors, 177
- methotrexate, 68–9
 administration, 69
 teratogenic potential, 69
- mid-level providers, 219–26
 definition, 219
 legal restrictions to task shifting/sharing, 220
 medical abortion
 experiences of mid-level providers, 222–3
 role in, 220
 settings, 220
 training, 221
- midwives, 221, 222–3
 Nepal, 224

- postabortion care, 223
 contraception, 223
- surgical abortion
 experiences of mid-level
 providers, 221–2
 morbidity, 221–2
 Sweden, 222–3
 training, 221
- task sharing, 220, 224–5
 task shifting, 219–20
 training of, 220–1
- midwives, mid-level provision,
 221, 222–3
- mifepristone, 82
 administration (EMA), 67
 in Australia, 63
 history of, 62–3
 indications, 63
 9 to 13 weeks' gestation,
 abortion at, 83
 13 to 24 weeks' gestation,
 abortion at, 83–4
 dilatation and evacuation
 (D&E), 92
 teratogenic effect, 130
- miscarriage, 64, 165–6
- misinformation, *see* myths and
 misinformation
- misoprostol, 82
 gestation age
 9 to 13 weeks' gestation,
 abortion at, 83
 13 to 24 weeks' gestation,
 abortion at, 83–4
- haemorrhages and, 128
- history of, 63
- procedures
 dilatation and evacuation
 (D&E), 92
 early medical abortion,
 67, 103
 vacuum aspiration abortion
 (VA), 73
- retained products of conception
 (RPOC), 126, 129
- as a single agent
 9 to 13 weeks' gestation, 83
 13 to 24 weeks' gestation, 83
 early medical abortion, 69
- teratogenic effect, 130
- unsafe abortions, 69
- uterine scar rupture, 86
- Misoprostol in Obstetrics and
 Gynecology, 253
- monozygotic twins, 56–7
- moral issues, *see* ethics
- morbidity (abortion-related), 30–1,
 65, 122–32
 antibiotics and, 123
 digoxin, feticide with, 94
 follow-up, detection at, 109–10
 incomplete abortion,
see retained products of
 conception (RPOC)
- mortality, *see* mortality
- myths, 176
- prevalence, 122–3
- rates, 65
 in safe abortions, 123–4
- second trimester abortions, 202
- stigma and, 190
see also complications of
 abortion; medical co-
 morbidity
- 'more than one abortion', 193–200
- attitude of healthcare providers,
 193, 216–17
- comparative studies, 197–8
- contraception and, 193, 198–9
- definition, 194
- demographics, 195, 196–7, 204
 liberalization of law and, 196
 steady state, 196–7
- intervals, 197
- prevalence, 194, 195, 197
 worldwide statistics, 194–6
- risk factors, 115–7, 198
 postabortion contraception,
 115–7
 socio-economic factors, 115
- terminology, 193–4
- mortality, 10, 11, 30–1, 219
 abortion methods and, 123–4
 dilatation and evacuation
 (D&E), 89, 94
 early medical abortion
 (EMA), 123
 misoprostol, 69
 causes, 122, 123
see also morbidity
- historical, 9
- infanticide, *see* infanticide
- infection, 129
 early medical abortion, 68
- legal considerations
 legality and, 30
 liberalization of abortion law
 and, 123
- myths, 176
- Romania, 31
- safe procedures, 122
- stigma and, 190
- unsafe abortions, 30, 114, 122
- uterine scar rupture, 86
- multiple-order gestations
 fetal reduction, *see* feticide
 ultrasound scanning, 56
- myths and misinformation,
 175–83
 contraception
 advice, 181–2
 efficacy, 179–80
 emergency, 179
 fertility, 176–7
 fetal pain, 178
 harm from abortion, 176
 higher gestation abortion,
 180–1
 legal considerations, 181
 mental health issues, 177
 mortality and morbidity, 176
 prevalence of abortions, 180
 reliable sources of
 information, 176
 support and counselling
 (preabortion), 178–9
- National Abortion Federation
 (NAF), 253
- National Health Service (NHS),
 funded abortions, 133
- Nepal, mid-level providers,
 224
- non-judgemental care, 215
see also health professionals
- non-steroidal anti-inflammatory
 drugs (NSAIDs), 85
- dilatation and evacuation
 (D&E), 91
- Nursing and Midwifery
 Council (NMC)
 (UK), conscientious
 objection, 213
- obesity, 135
 anaesthesia and, 135
 cervical preparation, 135
 higher gestation abortion,
 135
- Offences Against the Person
 Act (OAP) (1861,
 Australia), 237
- 'off-label' drugs, 63
- organizations, 251–4
 ACOG, *see* American
 Congress of
 Obstetricians and
 Gynecologists (ACOG)

- organizations (*cont.*)
 Advancing New Standards in Reproductive Health (ANSIRH), 251
 Association of Reproductive Health Professionals (ARHP), 251
 Center for Reproductive Rights, 251
 Concept Foundation, 251
 Education for Choice, 251
 Exhale, 251
 FIAPAC, 252
 FIGO (International Federation of Gynecology and Obstetrics), 252
 Guttmacher Institute, 252
 Gynuity Health Projects, 252
 Ibis Reproductive Health, 252
 International Consortium for Medical Abortion (ICMA), 252
 International Federation of Gynecology and Obstetrics (FIGO), 252
 International Planned Parenthood Federation (IPPF), 252
 Ipas, 253
 Marie Stopes International (MSI), 253
 Misoprostol in Obstetrics and Gynecology, 253
 National Abortion Federation (NAF), 253
 Nursing and Midwifery Council (NMC) (UK), 213
 Planned Parenthood Federation of America, 253
 Population Council, 253
 Postabortion Care (PAC) Consortium, 253
 RCOG, *see* Royal College of Obstetricians and Gynaecologists (RCOG)
 United Nations, 254
 WHO, *see* World Health Organization (WHO)
- osmotic dilators, 84
 Dilapan-S®, 91
 dilatation and evacuation (D&E), 91
 laminaria, *see* laminaria placement (D&E), 91–2
 outcomes, longer-term, 163–74
 adverse, risk factors, 45
 association and causality, *see* association and causality
 breast cancer, 168, 169, 177–8
 complications/morbidity, *see* complications of abortion; morbidity
 denied abortion, *see* denied abortion
 ectopic pregnancy, 166
 fertility, 165, 176–7
 miscarriage, 165–6
 physical health, 165–70
 placenta praevia, 166
 placental abruption, 166
 pre-eclampsia, 167
 preterm delivery, 167–8
 psychological, 39, 165–70
 emotional issues, *see* emotional wellbeing
 identification of, 40
 mental health, *see* mental health
 support and counselling, *see* support and counselling
 in USA, 171
 outpatient abortion
 home, 36, 64, 67
 follow-up, 68
 support and counselling, 65
 ovulation, after abortion, 110–11, 115
 ownership rights, 17, 21
 pain
 anxiety and, 102–3
 early medical abortion (EMA), 103
 factors affecting, 98
 Frankenhauser plexus, 98
 mechanism, 98
 postoperative, 102
 pain management, 98–105
 medical abortion
 first trimester, 103
 higher gestation, 85, 103
 non-steroidal anti-inflammatory drugs (NSAIDs), 85
 surgical abortion, 99–102
 adjunct medications, 101
 anaesthesia, *see* anaesthesia
 non-pharmacological pain management, 102–3
 procedural sedation (moderate IV sedation), 101
 types, 98, 104
 anaesthesia, *see* anaesthesia
 analgesia, *see* analgesia
 deep sedation, 98
 placebo, 100
 paracervical block (PCB)
 additives for, 100–1
 sites, 99
 partner(s)
 support, 205
 violence, 199
 patient compliance, 107
 patient information, *see* information, patient
 philosophical approaches, attitudes to abortion, 10–11
 physical disabilities, 140
 placental conditions
 placenta accreta
 abortion and, 86
 dilatation and evacuation (D&E), 90
 placenta praevia, 166
 placental abruption, 166
 planned parenthood, abortion requests and, 36
 Planned Parenthood clinic (Iowa, USA), 227–8, 230–1
 Planned Parenthood Federation of America, 253
 platelet disorders, 137
 policies, 20
 counselling and, 42–3
 Population Council, 253
 postabortion care
 complications, *see* complications of abortion
 contraception, *see* contraception
 dilatation and evacuation (D&E), 95–6
 hCG measurement, 107, 115
 mid-level providers, *see* mid-level providers
 patients, information for, 112
 Postabortion Care (PAC) Consortium, 253
 ultrasound, role of, 59–60
 RCOG guidelines, 59
 ‘postabortion syndrome’, 168–9
 post-mortem examination, TOPFA (termination of pregnancy for fetal anomaly), 158
 postnatal abortion, *see* infanticide

- potassium chloride (KCl)
 (intracardiac)
 feticide, 85, 94, 145
 procedure, 148–50
 Chiba needle, 148, 149, 150
- potentiality, 18–19
 moral worth, 19
 ‘pro-life’ approach and, 18
- preabortion care
 decision-making, *see* decision-making process
 support and counselling,
see support and counselling
 ultrasound scanning, 58–9
- pre-eclampsia, 167
- pregnancy
 continuing (postabortion),
see follow-up
 contradictory feelings, 37–8
 decision-making, 34
 ectopic, 166
 miscarriage, 64, 165–6
 preterm delivery, 167–8
 progesterone, 62
 twin pregnancy, 56–7, 140
 ultrasound scanning,
see ultrasound scanning
 urinary pregnancy tests, 107
 high sensitivity (HSUP),
 107, 109
 low sensitivity (LSUP), 108–9
 ‘qualitative’, 107
 semi-quantitative, 109
- preterm delivery, 167–8
- prevalence, 25
 abortion for fetal anomaly
 (TOPFA), 154
 by legality, 28
 ‘more than one abortion’
 population, *see* ‘more
 than one abortion’
 myths, 180
 NHS funded abortions, 133
 of TOPFA (termination of
 pregnancy for fetal
 anomaly), 154
 unsafe abortions, 26, 28
- procedural sedation, *see* pain
 management
- procurement of abortion, 12
- progesterone, 62
 receptor modulators,
see mifepristone
- ‘pro-life’ approach, potentiality
 and, 18
- propofol, 102
- prostaglandins, 81
 administration, 81
 analogues, 81–2
 gemeprost, 82
 misoprostol, *see* misoprostol
 sulprostone, 82
- Protocol to the African Charter
 on Human and Peoples’
 Rights on the Rights of
 Women in Africa*, 247
- providers, *see* health professionals
- Providers Share Workshop, 186–7
- pseudosac, 55
- psychological impact
 of delayed abortions, 202–3
 of denied abortion,
see denied abortion
 mental health, *see* mental health
 termination of pregnancy of fetal
 anomaly (TOPFA), 160
- psychosocial factors
 abortion requests, 34–41
 demographics,
see epidemiology
 male partners, 35–6
 planned parenthood, 36
 Scandinavia, 36
 social perspectives, 38–9
- public attitudes, *see* stigma
- public health issues, *see* human
 rights
- ratio, abortion, 28–9
- regression equation approach, 25
- renal disorders, 136
- repeat abortions, *see* ‘more than
 one abortion’
- Reproductive Health (Access to
 Terminations) Act 2013
 (Tas), 241
- reproductive rights, 244
- research on abortion,
 limitations, 37
- retained products of conception
 (RPOC)
 after medical abortions, 129–30
 management, 129
 after surgical abortions, 126
 dilatation and evacuation
 (D&E), 95
 vacuum aspiration abortion
 (VA), 77
 endometritis and, 127
 fetal anomaly, abortions for, 157
 haemorrhage and, 125
 management, 126
 risk of failed abortion, 65
 symptoms, 126
 ultrasound, role of, 59, 126
- right of conscience, 188
- rights
 human, *see* human rights
 male progenitors, 20–1
 ownership, 17, 21
 pregnant women’s, 16–17
- ring forceps, 93
- risk of failure (abortion), 65
see also retained products of
 conception (RPOC)
- Romania, mortality, 31
- Royal College of Obstetricians
 and Gynaecologists
 (RCOG), 253
 conscientious objection, 213
 feticide, recommendations
 pertaining to, 85–6,
 143–4, 145, 146, 159
 follow-up, 106
 periabortion antibiotic
 prophylaxis, 127
 ultrasound scanning in abortion
 care, 59
- RU-486, *see* mifepristone
- safety
 dilatation and evacuation
 (D&E), *see* dilatation and
 evacuation (D&E)
 safe procedures, 80, 122, 123–4
 vacuum aspiration abortion
 (VA), 71–2
see also complications of
 abortion
- Scandinavia, psychosocial factors
 in abortion, 36
- second trimester abortion, 201–10
 antecedent factors, 204–8
 age, 204
 non-western countries, 206
 sex selection, 206–7
 UK, 206
 complications and adverse
 effects, 157, 202
 contextual issues, 203–4
 denied abortion, 202
 feticide and, 143–4, 151
 funding, 204
 gestational limit, 181, 201, 202
 indications, 180
 legal considerations, 203, 209
 methods, 88

- second trimester abortion (*cont.*)
 medical abortion, *see* medical abortion
 pain management, 103
 settings, 146, 147
 surgical abortion, *see* surgical abortion
- morbidity, 130
 myths, 180–1
 obesity and, 135
 partner support, 205
 regulations, 203
 stigma and, 201–2
 termination of pregnancy for fetal anomaly (TOPFA), 154–5
 terminology, 201
 USA, 204–5
see also higher gestation abortion
- sedation
 levels, 100
 conscious, *see* pain management
 deep, 101
 general anaesthetic, *see* general anaesthesia
 procedural, *see* pain management
- selective feticide, 143
 self-abortion, 7
 semi-quantitative urinary pregnancy test, 109
- settings for abortion
 feticide, *see* feticide
 higher gestation abortion, 146, 147
 medical abortion, *see* medical abortion
 statistics, 146
 TOPFA (termination of pregnancy for fetal anomaly), 159
 vacuum aspiration abortion (VA), 73
- sex-selective abortion, 19–20, 206–7
 culture and, 20
 legal considerations, 19
- sexually transmitted infections (STIs)
Chlamydia trachomatis, 66, 127
 screening, 66
- sickle cell disease, 137
 social inequality, 248
 Society of Family Planning (SFP) (USA), feticide, 144
- socio-economic factors, 30, 35
 of male partner, 35
 risk factors for subsequent abortion, 115, 116
- staff, *see* health professionals
- stigma, 12, 39, 184–92
 conceptualization, 184–5
 definition, 184
 domains, 185
 enacted stigma, 185
 felt stigma, 185
 internalized stigma, 185
 conscience and, 187
 experience of, 39–40
 historical perspectives, 9
 late abortion and, 201–2
 measurement, 185–6
 Individual Level Abortion Stigma (ILAS) scale, 186
 methodologies, 185–6
 qualitative tools, 186
 quantitative tools, 186
 Stigmatizing Attitudes Beliefs and Actions Scale (SABAS), 186
 mortality/morbidity and, 190
 reduction interventions, 186–7, 190
 culture of support, 187
 Providers Share Workshop, 186–7
 ‘stigma by association’, 214
 Stigmatizing Attitudes Beliefs and Actions Scale (SABAS), 186
- substance use, 138
- suction evacuation/curettage, *see* vacuum aspiration abortion (VA)
- sulprostone, 82
- support and counselling
 culture of support (stigma reduction intervention), 187
 ethics in, 50
 fetal anomaly, 155–6
 health professionals, role of, 217
 outpatient abortion, 65
 partner support, 205
 postabortion, 48
 function and timing, 49
 preabortion, 42–51
 compulsory counselling, 42
 counselling techniques, 47–8
 functions of, 46
 identification of needs, 44, 45
- intense level, 44–6
 myth, 178–9
 policy context, 42–3
 resistance to, 46
- professional care, 49–50
 boundaries and personal feelings, 50
 core principles, 49
 qualifications, 49
 telemedicine, 233
 TOPFA (termination of pregnancy for fetal anomaly), 160
see also emotional wellbeing
- surgical abortion, 7
 complications, *see* complications of abortion
 for fetal anomaly, 156, 157
 higher gestation, 89
 vs medical abortion, 64–5
 methods
 dilatation and evacuation (D&E), *see* dilatation and evacuation (D&E)
 vacuum aspiration, *see* vacuum aspiration abortion (VA)
- mid-level providers, role of, *see* mid-level providers
- pain management, *see* pain management
- postabortion care, 48
 contraception, 96, 111, 117–18
 pre-procedure medication, 101
 rates of complete and complications, 65
 ultrasound, role of, 59
- Sweden, 171
 mid-level provision, 222–3
- taboo, *see* stigma
- task sharing, *see* mid-level providers
- task shifting, 219–20
 Tasmania, *see* Australia
- technology, fetus viability and, 21
 telemedicine, 227–35
 applications, 227
 definition, 227
 development guidelines, 232–3
 consent/information/support, 233
 follow-up, 233
 gestational dating and investigations, 233

- hardware/software, 233
 regulatory and licensing
 issues, 232–3
 evaluation strategies, 230–2
 medical abortion, 227–30, 232
 Canada (Women on Web),
 228–9, 231–2, 254
 clinic to clinic, 227–8
 clinic to woman at
 home, 228–9
 internet and email, 229
 internet website only, 229–30
 Planned Parenthood clinic
 (Iowa, USA), 227–
 8, 230–1
 telepharmacy, 228
 telephone, follow-up (EMA), 68
- teratogens
 methotrexate, 69
 mifepristone, 130
 misoprostol, 130
- termination of pregnancy for fetal
 anomaly, *see* TOPFA
 (termination of
 pregnancy for fetal
 anomaly)
- therapeutic bond, 211
- third trimester abortion, 160–1
 pain management, 103
 termination of pregnancy for
 fetal anomaly (TOPFA),
 154–5, 160–1
- thrombophilias, 137
 venous thromboembolism
 (VTE), 137
- thyroid disease, 136
- TOPFA (termination of pregnancy
 for fetal anomaly)
 Australia (Victoria), 240
 cremation and burial, 159
 decision-making factors, 155–8
 availability, 158
 complications and adverse
 effects, 157
 post-mortem
 examination, 158
 preference and
 acceptability, 157–8
 gestational age, 154–5, 160–1
 hospital follow-up visit, 159–60
 legal criteria, 153
 medical conditions, *see* fetus
 methods, 155, 156
 anaesthesia, 156
 feticide, 158–9
 medical abortion, 156
 setting, 159
 surgical, 156, 157
 prevalence of, 154
 psychological impact, 160
 counselling, 160
 retained products of
 conception, 157
 in third trimester, 160–1
- training
 of mid-level providers, *see* mid-
 level providers
 second trimester abortion, 89
 ultrasound targeted training, 58
- transabdominal (TA) transducers, 52, 56, 57
- transvaginal (TV) transducers, 52, 54, 55
- twin pregnancy, 56–7, 140
- UK
 gestational limit, 181
 repeat abortion statistics, 196
 second trimester abortions, 206
- ultrasound (first trimester), 57
- ultrasound scanning, 52–61
 abortion care and, 58–60
 image viewing, patients,
 58–9
 intraoperative assessment, 59
 postabortion, 59–60
 preabortion assessment, 58–9
 RCOG guidelines, 59
 targeted training, 58
 applications, 53, 59–60
 in failed pregnancies, 57–8
 first trimester, 53–5
 embryo and fetus, 54–5,
 56, 57
 gestation sac, *see* gestation sac
 medical co-morbidity, 134
 multiple gestations, 56–7
 objectives, 52
 postabortion, 107
 procedures
 dilatation and evacuation
 (D&E), 90
 vacuum aspiration abortion,
 use in, 72
 of retained products of
 conception (RPOC),
 59, 126
 sonoembryological
 development, 55–6
 biparietal diameter, 56
 techniques, 52
 transabdominal (TA)
 transducers, 52, 56, 57
 transvaginal (TV)
 transducers, 52, 54, 55
 unintended live births, *see* feticide
- United Nations, 254
- unsafe abortion, 24, 175
 contraception,
 postabortion, 114
 definition, 122
 human rights and, 245–6
*International Conference
 on Population
 and Development*
 (ICPD), 244–5
 interventions, 246
 legal abortion and, 31
 medical abortion, 85
 misoprostol, 69
 mortality and, 30, 114, 122
 prevalence, 26, 28
 as a public health issue, 244
- urinary pregnancy tests,
see pregnancy
- USA
 conscientious objection, 213–14
 outcomes, 171
 Planned Parenthood clinic
 (Iowa), 227–8
 second trimester abortions,
 204–5
- uterus
 fibroids, 139
 perforation
 dilatation and evacuation
 (D&E), 94
 management, 125
 risk factors, 124
 surgical abortion, 124–5
 rupture, medical abortion
 (higher gestation), 84–5,
 86, 130
 scar rupture, 139
 dilatation and evacuation
 (D&E) and, 86
 maternal mortality, 86
 misoprostol, 86
 placenta accreta, 86
 uterine sounding (VA), 75
- vacuum aspiration
 abortion, *see* vacuum aspiration
 abortion (VA)
 dilatation and evacuation
 (D&E), 93
 retained products of conception
 (RPOC), 126

- vacuum aspiration abortion
 (VA), 71–9
 acceptability, 72
 complications, 76–8, 123–4
 incomplete evacuation, 77
 definition, 71
 efficacy, 71
 history of, 71
 indications, 71
 post-procedure management, 78
 follow-up, 78, 106
 pre-procedure, 73
 antibiotics, 73
 cervical dilation, 75
 cervical preparation, 73
 cervical traction, 74–5
 informed consent, 72
 initial assessment, 72–3
- misoprostol, 73
 pain management, 73
 procedure, 74–6
 aspiration, 75–6
 bimanual examination, 74
 cervical dilation, 75
 cervical traction, 74–5
 equipment/instruments, 74
 Hegar dilator, 74
 Karman cannula, 74, 75
 uterine sounding, 75
 safety, 71–2
 settings, 73
 ultrasound scanning, 72
 vacuum source, 71
 electric pump (EVA), 76
 manual pump (MVA), 76,
 77
- vasovagal reaction, surgical
 abortion, 124
 venous thromboembolism
 (VTE), 137
 viability of fetus, 21–2
 definition, 22
 technology dependent, 21
 Victoria, *see* Australia
 Western Australia, 239
 Women on Web, 228–9,
 231–2, 254
 World Health Organization
 (WHO), 254
 follow-up, 106
 safe abortion, 129
 yolk sac, 54