

Index

- acetylcholine, age-related decrease, 67
- activities of daily living (ADL), 86, 89, 90
- aerobic exercise, 114–115
- age-related macular degeneration (AMD), 87
- age-related sleep patterns, 74
- Alzheimer's disease (AD)
 - aerobic exercise benefits, 114–115
 - dental caries, 100
 - dysphagia, 46
 - energy requirement increase, 45
 - epilepsy as risk factor, 32
 - falls association, 5, 7
 - frailty, 112, 113, 114
 - gait impairment, 5
 - inflammation role in development, 101, 113–114
 - late-onset, delirium in, 16
 - prediction, from frailty index, 114
 - seizures, 30–31, 32, 34
 - pathogenesis, 33
 - sleep disturbances, 74, 75, 76
 - urinary incontinence, 62, 63
 - visual dysfunction, 86
 - pathophysiology, 86, 87–88
 - symptoms/types, 90, 91
 - visual hallucinations, 91
 - visual variant (VVAD), 88
 - visuospatial dysfunction, 90
 - weight loss and malnutrition, 40, 42
 - causes, 43–45
 - nutritional supplementation, 47–48
- amyloid plaques, in retina, 88
- amyloid precursor protein, 33
- anterior cingulate cortex (ACC), 45
- anti-epileptic therapy, 34–35, 102
- antimuscarinic drugs, 66–67
- antipsychotics, 22, 33, 102
- apathy, 43, 44, 101
- ApoE4 allele, 43, 76
- arthritis, 47
- attentional skills, delirium management, 22
- axonal sprouting, 33
- behavioural disturbances
 - in delirium, 19
 - falls associated, 7
 - impact on oral health, 101
 - weight loss and malnutrition, 44–45
- benzodiazepines, 6, 22
- bladder
 - normal control/physiology, 61–62
 - overactive, 61, 66–67
- body mass index (BMI), 47, 113
- brain structural changes, weight loss and, 43, 45
- CADASIL, 88
- carer stress/mental health problems, 80
- carers, burden of
 - oral health problems, 101
 - sleep disturbances, 80
 - urinary incontinence, 67–68
 - weight loss, 43
- case studies
 - delirium, 23
 - epileptic seizures, 35–36
 - fecal incontinence, 70
 - falls, 10–11
 - frailty, 115–116
 - oral disease, 106–107
 - sleep disturbances, 81–82
 - urinary incontinence, 69–70

- visual dysfunction, 93–94
- weight loss and malnutrition, 52
- cholinergic deficiency, 17
- cholinesterase inhibitors, 33
 - falls associated, 7
 - sleep disturbances and, 76
 - urinary incontinence and, 66
 - weight loss and malnutrition, 46
- circadian rhythm changes, 74, 76, 77
- cognitive impairment
 - exercise program benefits, 114–115
 - falls associated, 3, 4
 - frailty association, 111, 112
 - mild *see* mild cognitive impairment
 - visual acuity problems, 89
- colour vision impairment, 91
- community
 - activity programs, 79
 - fall reduction/prevention, 9
 - prompted voiding program, 66
- comorbidity, 1
 - in dementia, literature search, 1–2
- Confusion Assessment Method (CAM), 20, 22
- constipation, 61, 68
- continence, 60–61, 63–64
 - see also* urinary incontinence (UI)
- continence aids, 67
- contrast sensitivity, 88–89
- Creutzfeldt–Jakob disease (CJD), 32, 88, 90
- dawn dusk simulation (DDS) light
 - therapy, 77
- daytime sleepiness, 75–76
- delirium, in dementia, 15–24
 - assessment, 20–21
 - case studies, 23
 - clinical features, 15, 18, 19–20
 - definition, 15
 - dementia cause/effect relationship, 16–17
 - diagnostic difficulties, 18, 20
 - epidemiology, 15–16
 - etiology, 16–17
 - impact on dementia, 15, 18–19
 - management, 21–22
 - mortality, 18
 - pathogenesis, 17
 - persistent, 15, 18, 20
 - prevention, in hospital, 21
 - recommendations, 22–23
 - risk and risk factors, 15, 17–18, 19
- Delirium Room, 21
- dental caries, 99, 102
 - epidemiology, 100, 101
 - medications impact on, 102
- dental hygiene/care *see* oral health; oral hygiene
- dental plaque, 99
- dentistry, 105
 - minimal intervention, 105
- dentures, 99, 100, 106
- detrusor hyperactivity, 61, 66–67
- diarrhea, 61
- dietary intake, alterations, 43–44
 - see also* malnutrition
- diffuse Lewy body disease (DLBD)
 - delirium *vs* dementia, 19
 - gait impairment and falls, 5, 7
 - neurovascular instability, 6
 - seizures, 31–32
 - urinary incontinence, 63
- donepezil, 66
- Down syndrome, seizures, 32
- drugs *see* medications
- dual-tasking, 6
- dysphagia (swallowing difficulties), 43, 46
 - management/interventions, 50
- Eating Behaviour Scale, 47
- eating environment, 48–49
- eating patterns, disordered, 43–44, 45
- education of staff, 48
 - delirium management, 22
 - dental hygiene, 105
 - falls prevention, 8–9
 - nutritional supplementation, 47, 48
- energy
 - increased requirements, 45
 - intake, timing, 49
- enteral feeding, 50–51
 - hand feeding *vs*, 51
- environmental factors, nutritional intake
 - improvement, 48–49
- epileptic seizures (and epilepsy), 30–36
 - in AD *see* Alzheimer's disease
 - case studies, 35–36
 - definition, 30
 - diagnosis, 34
 - epidemiology and prevalence, 30–32
 - etiology, 33
 - falls associated, 35
 - features, 34
 - management, 34–35
 - recommendations, 35
 - risk and risk factors, 31, 34
 - as risk factor for dementia, 32–33
 - seizure types, 31, 34
 - in specific dementias/conditions, 31–32
 - treatment (drugs), 34–35

122 Index

- exercise programs
 - delirium incidence reduction, 21
 - fall prevention, 9
 - frailty and cognitive decline reduction, 114–115
 - prompted voiding program with, 65–66
- falls, in dementia, 3–11
 - assessment, 8
 - behavioural disturbances, 7
 - case studies, 10–11
 - definition, 3
 - dementia pathology and, 5, 6
 - epidemiology, 3–4
 - etiology, 4–5
 - gait impairment, 5
 - management, 8–9
 - medication effects, 6–7, 9
 - preventative interventions, 8–9, 10
 - recommendations, 10
 - risk and risk factors, 3, 7, 8
 - seizures associated, 35
 - sequelae/effects, 3, 4
- fecal impaction with overflow, 61, 68
- fecal incontinence, 61, 68
 - case study, 70
 - epidemiology, 62
 - management, 68
 - prevalence, and risk factors, 68
- feeding, enteral *see* enteral feeding
- feeding behaviour changes, 44
- fibre (dietary), intake, 68
- finger foods, 49
- food(s)
 - familiar, and familiar environment for, 48–49
 - finger, 49
 - refusal, 44
 - types, to improve eating/nutrition, 49
- frailty, 111–116
 - case studies, 115–116
 - causes, 111
 - definition/criteria, 111, 112
 - epidemiology, 111–113
 - etiology, 113–114
 - management, 114–115
 - as noncognitive feature of AD, 114
 - recommendations, 115
- frontotemporal dementia (FTD)
 - dysphagia, 46
 - eating pattern changes, 43–44
 - urinary incontinence, 63–64
 - weight loss and malnutrition, 40
- functional incontinence, 61
- gait
 - impairment, 5
 - normal, 5
 - see also* walking
- gingival hyperplasia, 102
- glucose metabolism, 45
- haloperidol, 21, 22
- handgrip strength, decreased, 113
- hemispatial neglect, 91
- hip fractures, 4, 19, 21
- hippocampal lesions, falls and, 5
- hospitalization
 - after falls, 4
 - delirium precipitation, 16–17, 18
 - delirium prevention, 21
 - protein/energy intake reduced, interventions, 50
- Huntington's disease, seizures, 32
- hyperbolic state, 45
- hypotension, orthostatic (postural), 6
- incontinence, 60–70
 - see also* fecal incontinence; urinary incontinence (UI)
- incontinence pads/aids, 67
- individualized social activity interventions (ISAI), 79
- indwelling catheter (IDC), 67
- inflammation, 101, 113–114
- lean body mass, reduced, 113
- levetiracetam, 35
- light therapy, 77
 - melatonin with, 78
- macular retinal nerve fibre layer, 87
- macular volume, reduction, 87
- malnutrition, 40
 - assessment, 47, 51
 - behavioral symptoms associated, 44–45
 - case studies, 52
 - causes/pathogenesis, 41, 42–46
 - dementia impact, 40–42
 - disordered eating patterns, 43–44
 - dysphagia, 46
 - intervention and management, 46, 47–51
 - in nursing home residents, 41
 - recommendations for management, 51–52
 - see also* weight loss
- meals, 49–50
- medial temporal lobe, abnormalities, 45

- medications
 - falls associated, 6–7, 9
 - impact on oral health, 102–103
 - seizure threshold reduced, 33
 - for sleep disturbances, 77–78
 - for urinary incontinence, 66–67
 - weight loss/malnutrition association, 46
- melatonin
 - delirium management, 22
 - sleep disturbance intervention, 78
- mild cognitive impairment (MCI)
 - frailty and reduced motor function, 112
 - retinal nerve fibre layer, 87
 - sleep pattern changes, 75, 81–82
- Mini Nutritional Assessment (MNA), 47
- mouthwash, 104
- multitasking (dual-tasking), 6
- muscle strength decrease, 111, 113
- nasogastric tubes, 50
- National Institute of Clinical Excellence,
 - delirium diagnosis, 20
- neurofibrillary tangles, 75
- neuroleptics, falls associated, 6
- neurovascular instability, 6
- night monitoring system (NMS), 80
- nursing homes and residential care
 - malnutrition in, 41
 - oral hygiene and dental care, 105
 - urinary incontinence prevalence, 62
 - visual impairment, 92
- nutritional disorders *see* malnutrition; weight loss
- nutritional state, dementia impact, 40–42
- nutritional supplementation, 47–48, 50
- optic flow, impairment, 91
- optic nerve, changes, 87
- oral disease, 99–107
 - assessment, 104, 105
 - case studies, 106–107
 - epidemiology, 100
 - management, 103
 - symptoms and signs, 103
 - treatment, 104–105
- oral health
 - behavioural changes impact, 101
 - benefits, 99
 - goals, 103–104
 - impact of dementia on, 100–102
 - medication impact on, 102–103
 - recommendations, 106
- oral hygiene
 - daily care recommendations, 104, 106
 - poor state, 99, 100
 - staff education on, 105
- oral ulceration, 102
- orthostatic hypotension, 6
- oxybutynin, 66–67
- Parkinson's disease
 - dysphagia, 46
 - gait impairment and falls, 5, 7
 - neurovascular instability, 6
 - urinary incontinence, 63
- pelvic floor exercises, 66
- perceptual disturbances, 19
- percutaneous endoscopic gastrostomy (PEG)
 - tubes, 50–51
- periodontal disease, 102
- physical activity
 - increased nutritional intake, 50
 - sleep disturbance management, 79
 - see also* exercise programs
- physical restraints, fall prevention, 9
- polysomnography, 76
- postural hypotension, 6
- prompted voiding (PV) programs, 65–66
 - design and goals, 65
 - outcome and resource requirements, 65–66, 68
- REM sleep, decrease, 75, 76
- residential care *see* nursing homes and residential care
- respite care, 80
- restlessness, night time, walking impact, 79
- retinal nerve fibre layer (RNFL), thickness, 87
- risperidone, 78
- saliva, reduced, 102
- saliva stimulators, 106
- sedation, in dentistry, 105
- sedatives, 77
- seizures *see* epileptic seizures (and epilepsy)
- sialorrhea, 102
- sleep
 - age-related changes, 74
 - deprivation in carers, 80
- sleep disturbances in dementia, 74–82
 - carer burden, 80
 - case studies, 81–82
 - epidemiology, 75–77
 - intervention and treatment, 77–79
 - pathology, 74–75

124 Index

- sleep disturbances in dementia (*cont.*)
 - recommendations, 81
 - types, 76
- stress incontinence, 61
- sucking reflexes, 102
- sundowning, management, 22
- swallowing problems *see* dysphagia
- synaptic hyperactivity, 33
- 'talking while walking' test, 6
- teeth
 - low number, dementia association, 101
 - preservation, benefits, 105
- urge incontinence, 61
- urinary incontinence (UI), 8, 60–70
 - assessment, 64
 - caregiver burden, 67–68
 - case studies, 69–70
 - causes, 60, 63, 64, 66
 - definition, 60
 - dementia types and, 62, 63–64
 - epidemiology, 62
 - impact of dementia on, 63–64
 - intervention and management, 64–68
 - continence aids, 67
 - medications, 66–67
 - PV program *see* prompted voiding (PV) programs
 - prevalence and increase in, 60, 62
 - recommendations, 69
 - stress incontinence, 61
 - types, 61
- urinary obstruction with overflow, 61
- vascular dementia
 - delirium in, 16
 - gait impairment and falls, 5
 - urinary incontinence, 62
 - weight loss and malnutrition, 40
- visual acuity impairment, 89, 91
 - recommendations, 92
- visual dysfunction, 86–94
 - assessment, 87, 92
 - case studies, 93–94
 - definition, 86
 - features/types, 88–91
 - impact in dementia, 88
 - interventions, 93
 - pathology, 86–88
 - recommendations, 92, 93
 - in residential care, 92
 - types of visual problem, 86, 88
- visual field defects, 86, 89–90
- visual hallucinations, 88, 91–92
- visual motion, optic flow impairment and, 91
- visuospatial dysfunction, 90
- walking
 - impact on nighttime restlessness, 79
 - speed decrease, frailty, 111, 112
 - see also* gait
- wandering, falls and, 7
- weight
 - dementia impact on, 40–42
 - gain, 42, 46
- weight loss, 40–53
 - assessment, 47, 51
 - behavioural symptoms and, 44–45
 - case studies, 52
 - causes/pathogenesis, 41, 42–46
 - cholinesterase inhibitors, 46
 - dietary intake alterations, 43–44
 - disordered eating patterns, 43–44
 - epidemiology and prevalence, 42
 - frailty and, 113
 - increased energy needs, 45
 - intervention and management, 46, 47–51
 - education of staff, 48
 - enteral feeding, 50–51
 - environmental factors, 48–49
 - multifactorial strategy, 49–50
 - nutritional supplementation, 47–48
 - targeted approaches, 49–50
 - preceding dementia, 41–42
 - recommendations for management, 51–52
- xerostomia (dry mouth), 99, 100, 101
 - drugs causing, 102