

Cambridge University Press

978-1-107-65024-4 - Theory of Differential Equations: Part I: Exact Equations and Pfaff's Problem

Andrew Russell Forsyth

Index

[More information](#)

## INDEX.

(The figures refer to pages.)

## AUTHORS QUOTED :

- Bach, 33.  
 Bäcklund, 230.  
 Bertrand, 18, 33, 109, 227.  
 Biermann, 299, 312, 316, 321.  
 Binet, 148.  
 Boole, 67, 299, 305, 307.  
 Bour, 1.  
 Brioschi, 211.  
 Cauchy, 86, 118, 183, 211, 294.  
 Cayley, 81, 99, 115.  
 Christoffel, 274.  
 Clebsch, 72, 76, 78, 82—86, 117, 168, 173, 175, 179, 181, 194 et seq., 233, 244, 245, 252, 257, 266, 298, 321.  
 Collet, 25, 34.  
 Condorcet, 33.  
 Darboux, 88, 181, 227, 238, 294 et seq.  
 Deahna, 51.  
 De Morgan, 20.  
 Dirksen, 88.  
 du Bois-Reymond, 81, 82.  
 Engel, 280, 299, 805, 806.  
 Euler, 18, 80, 88, 55, 56, 67, 80, 109.  
 Fais, 35.  
 Frisiani, 82.  
 Frobenius, 51, 54, 87, 88, 181, 272 et seq., 294, 299.  
 Gauss, 81, 112, 251.  
 Grassmann, 82, 83, 86, 121 et seq., 251, 329.  
 Hamburger, 85, 331.  
 Hamilton, 86, 118, 183.  
 Hesse, 35.  
 Imschenetsky, 33, 299, 307.  
 Jacobi, 1, 40, 41, 76, 78, 79, 81—83, 86, 115, 117, 118, 120, 138, 153, 173, 181, 183, 209.  
 Joachimstahl, 33.  
 Jordan, 239, 298.  
 Kowalewski, 331.  
 Kronecker, 35.  
 Lagrange, 33, 242.  
 Laurent, 35.  
 Legendre, 242.  
 Le Pont, 59.  
 Lexell, 33.  
 Lie, 78, 82, 84, 86—88, 168, 181, 183, 230 et seq., 248 et seq., 278, 294.  
 Lipschitz, 274.  
 Mansion, 183, 230, 306.  
 Maximowitch, 58.  
 Mayer, 1, 59, 62, 67, 72, 76 et seq., 81, 86, 200, 215, 230, 233 et seq., 257, 258, 294, 331.  
 Monge, 80, 102, 307.  
 Natani, 27, 30, 32, 55, 56, 59, 78, 82, 83, 85, 86, 143 et seq., 180, 187, 190, 191, 200, 208, 278, 312.  
 Pfaff, 81—83, 89 et seq., 180.  
 Pittarelli, 85.  
 Pujet, 33.  
 Raabe, 88, 193.  
 Sarrus, 83.  
 Scott, 81, 95, 106, 275.  
 Stodockiewicz, 58.  
 Stoffel, 33.  
 Tanner, 117, 142, 299.  
 Vályi, 321.  
 Voss, 33, 34, 299.  
 Weiler, 32.  
 Winckler, 33.  
 Algebraical substitutions, 274;  
   changed into differential substitutions, 281.  
 Arbitrary element in integral equivalent of single equation, 146;  
   of non-integrable system, 316;  
   used to modify single equation, 157;  
   used to modify non-integrable system, 319.  
 Arithmetic mean of two invariant integrals in transformation of simultaneous forms is a sufficient invariant, 278.  
 Bertrand's integration of an exact equation, 319.

- tion in three variables, 18, 109, 226, 227;  
 generalised to exact equation in  $n$  variables, 20—24.
- Biermann's** determination of minimum number of equations in equivalent of non-integrable system, 312, 316.
- Bilinear** differential covariant, 273, 294;  
 changed to algebraical covariant, 274.
- Binet's** simultaneous variations, 148 (note).
- Canonical form**, of system of exact equations, 43;  
 reduction of two inexact equations in four variables to, 325—327.
- Cauchy** substitutions, 59, 86;  
 used by Lie to transform conditioned expression, 258, 269;  
 do not affect character of normal form, 258, 269.
- Cayley's** symbolical solution of equations subsidiary to first reduction of expression, 100.
- Character** of a normal form, persistent, 249;  
 conditions which determine the, 253—255, and inferences from conditions, 255—257;  
 unaffected by Cauchy substitution, 258, 269;  
 inferred from results of Natani and Frobenius, 278 (note).
- Class** of an expression, 290;  
 examples relating to, 293.
- Clebsch's** methods, note on complete systems of partial differential equations in, 76, 78;  
 number of integrations required in, 79;  
 equations inferred from Lie's method, 266 (note);  
 of treating Pfaff's problem, Chap. viii, see table of contents;  
 abstract and general character of, 85, 86, 181.
- Collet's** determination of integrating factor, 25.
- Complete** reduction of expression, after Gauss, to contain minimum number of terms, 112—114.
- Complete** systems of linear and homogeneous partial differential equations, in generalisation of Bertrand's method, 21—25;  
 in Frobenius' conditions for exact equations, 52;  
 in Clebsch's methods, 76, 78, 209, 225, 229;  
 coextensive with a system of exact equations, 68;  
 number of independent solutions of, 69;  
 construction of solutions by successive derivation, 68; by general method of Mayer, 72; by simplest form of Mayer's method, 74;  
 Mayer's derivation of single solution of, from any integral of subsidiary system, 76.
- Completely** integrable systems of equations, Chap. ii, 300.
- Condition** of consistency, of first subsidiary system when the number of variables is odd, 107;  
 effect of this condition on the first transformation, 109;  
 verification of, from Natani's method, 164.
- Conditioned** expression, transformed by Cauchy substitutions into an unconditioned expression having normal form of same character, 258, 269, 297;  
 normal form of, deduced from that of the transformed unconditioned expression, 259—266.
- Conditions**, for existence of integrals fewer in number than the general minimum, in form of interrupted products, 123;  
 with numerical interpretation, 124—127;  
 for expression in even number of variables, 166, 167, 207;  
 for expression in odd number of variables, 168, 169, 207;  
 given by Lie to determine character of normal form, 253;  
 inferences from, 255;  
 that partial differential equation of second order possess an intermediary integral, 309.
- Consistency** of subsidiary equations when number of variables is odd, condition for, 107;  
 effect of this condition, 109;  
 verification of, from Natani's method, 164.
- Construction**, gradual, of solution and normal form:  
 in Pfaff's reduction, 112;  
 by Jacobi, 120;  
 by Grassmann, 141;  
 by Natani, 151, 157, 160;  
 by Clebsch, 198, 199, 208, 215, 223;  
 by Lie, 257, 268, 270.
- Corollaries** from cylindrical tangential transformation used by Lie, 242, 248;  
 derivable from Clebsch's results, 244, 245.
- Covariant**, bilinear differential, associated with expression, 273;

- no other similarly formed, exists for expression, 274;  
 multilinear, not associated with expression, 274 (note).
- Cylindrical tangential transformation, 239;  
 equations of, 241, 242;  
 Lagrange's examples of, 242;  
 Lie's corollaries from, 242, 248.
- Darboux's investigations, 88;  
 abstract of, Chap. xii, see table of contents.
- Deahna's conditions of exactness of system of equations, 51.
- Definition of tangential transformation, 232.
- Derivation, of exact equation from one integral, 1;  
 of system of exact equations from system of integrals, 37.
- Determinants, in differential transformation, 94, 101, 103, 106, 166, 169, 201, 214, 254;  
 in algebraical transformation of bilinear form, 276 et seq.
- Determination of character of normal form, 253—255;  
 inferences from conditions which lead to, 255.
- Differential covariant, bilinear associated with expression, 273, 294;  
 nonebut bilinear so associated, 274.
- Differential relation characteristic of tangential transformation, 231;  
 can be treated as a Pfaffian equation, 233;  
**Mayer's establishment of relations among functions in, 288—288.**
- Diminution, of number of equations in equivalent of a single expression when conditions are satisfied, 165—169;  
 of number of equations in subsidiary system, when one integral is known, 173; when two integrals are known, 176; when more than two are known, 179.
- Direct construction by Clebsch of equations which determine all the integrals of an expression, 213.
- du Bois-Reymond's integration of exact equation, 31, 32;  
 compared with Natani's, 32.
- Effect, of vanishing determinant on result of first transformation, 102, 105;  
 of integrals already determined, on the remainder, 200, 215;  
 equations determining the, constructed direct, 213.
- Engel on canonical form of two equations in four variables, 326.
- Equations, determining the first integral of an expression, in an even number of variables, when unconditioned, 200—203;  
 when conditioned, 203—207;  
 when normal form involves an odd number of functions, 224;  
 determining all the integrals, 213, 291;  
 some properties of the simultaneous, in Clebsch's method, 215.
- Equivalence of two expressions through normal forms, 252, 285, 289.
- Equivalent, integral, of expression deduced from normal form, 116, 117;  
 of irreducible differential relation, 250 (note);  
 minimum number of equations in, of a non-integrable system, 312—316;  
 of general partial differential equation of second order, 320;  
 reduced forms, 195;  
 relations between elements of, 252;  
 solutions, of an exact equation, 2;  
 of system of exact equations, 38.
- Euler's, integration of an exact equation, 18, 109;  
 compared with Natani's, 30;  
 generalised to system of exact equations, 55;  
 view of inexact equations, 80.
- Even reduction, 105;  
 can always be applied to expression in  $2n$  variables, 105;  
 not unique, 105.
- Exact equation, all solutions of a single, are equivalent, 3;  
 methods of integration of a single:  
 Euler's, 18, 109;  
 by integrating factor, 14, 25;  
 Bertrand's for three variables, 18, 109, 226, 227;  
 Bertrand's generalised for  $n$  variables, 20;  
 Natani's, 27;  
 comparison of Euler's and Natani's, 80;  
 du Bois-Reymond's, 81;  
 inference from Lie's general process, 270.
- Exact equations, system of: any combination of solutions of, is a solution, 38:  
 partial differential equations satisfied by solution of, 38;  
 are coextensive with original system, 67, 69, 71;  
 integrating factors of, 89;  
 multiplier of, 40;  
 methods of integration of:  
 Euler's generalised, 55;  
 Natani's, 56;  
 Mayer's, 59;

- special method for two equations  
in four variables, 62;  
by solution of partial differential  
equations, 67, 71.
- Exact integral** of system of equations,  
what is meant by, 300, 301;  
satisfies a system of partial dif-  
ferential equations, 302;  
conditions that it may exist, 304,  
305;  
used to modify the system, 311.
- Exact ordinary equations**, note on and  
memoirs relating to, 33.
- Exactness**,  
necessary conditions of, for a  
single equation, 4;  
number of necessary conditions  
independent, 6;  
necessary conditions are sufficient,  
7–12;  
necessary conditions of, for a  
system, 44;  
necessary conditions are sufficient,  
45–51;  
Frobenius' form of, 51–54.
- Expression**:  
in  $2n$  variables can always be sub-  
jected to an even reduction, 105;  
completely reduced, 113;  
in  $2n + 1$  variables can always be  
subjected to an odd reduction,  
111;  
completely reduced, 114.
- Extensive equation**, representing single  
equation, 122;  
representing system of equations,  
122, 329;  
subsidiary, transformed, 132, 135;  
solved, 136, 137 (note).
- Extensive variable**, introduction of, 121.
- Factor**, form of, removable from ex-  
pression after transformation:  
when the Pfaffian determinant  
does not vanish, 90, 99;  
when it does vanish, 101, 105,  
203 (note).
- First integral** of expression:  
if unconditioned, 203;  
if conditioned, 207;  
if normal form involve odd num-  
ber of functions, 224.
- First Method**, Clebsch's, of treating  
Pfaff's problem, 199–209.
- Frobenius**, on conditions of exactness of  
system of equations, 51–54.
- Frobenius' method** of treatment of  
Pfaff's problem, Chap. xi, see table of  
contents;  
abstract and general character of,  
87, 272;  
connection of Darboux's method  
with, 88.
- Functions**, number of that can be ob-  
tained without affecting invariance of  
determinantal integer, 282.
- Fundamental problems** for non-inte-  
grable systems, 311;  
two of the three yet unsolved,  
329, 330.
- Generalisation of normal forms**:  
in even number of functions, 197,  
252;  
in odd number of functions, 222,  
253.
- Generalisation of Natani's process** for a  
single Pfaffian:  
to determine number of equations  
in equivalent of system, 313;  
inadequate to determine the equa-  
tions in equivalent of system,  
322.
- Generality of integral equivalent**, cha-  
racteristic properties of, 145, 311.
- Gradual construction of solution and  
normal form**:  
in Pfaff's reduction, 112;  
by Jacobi, 120;  
by Grassmann, 141;  
by Natani, 151, 157, 160;  
by Clebsch, 198, 199, 208, 215, 223;  
by Lie, 257, 268, 270.
- Grassmann**, on system of equations,  
122, 329.
- Grassmann's method** of treatment of  
Pfaff's problem, Chap. v, see table of  
contents;  
abstract and general character of,  
83, 84.
- Hamburger's comparison** of methods of  
Natani and Clebsch, 85.
- Historical summary** of methods of treat-  
ing Pfaff's problem, Chap. iii, see  
table of contents.
- Homogeneous tangential transforma-  
tions**, 244;  
equations characteristic of, 245;  
infinitesimal, a special case of,  
246;  
applied by Lie to Pfaff's problem,  
252.
- Imschenetsky** on intermediary integral  
of partial differential equation of  
second order, 307.
- Incompletely integrable system** of equa-  
tions, 301;  
number of exact integrals of, 304;  
example of, in subsidiary equa-  
tions for partial of second order,  
together with conditions for  
two integrals, 307–310;  
modified by use of exact inte-  
grals, 311.

## INDEX

337

- Independent solutions of a complete system of linear and homogeneous partial differential equations**, number of, 69.
- Inexact equations**, views of, before Pfaff's memoir, 80.
- Infinitesimal homogeneous transformation**, 246;  
equations characteristic of, 247.
- Initial values of variables**, used by Jacobi, 82, 183, 183 (note);  
introduction of, leads to principal integrals, 118.
- Integers associated with determinants in transformation of bilinear quantics are invariante**, 277;  
arithmetic mean of the two, 278.
- Integral, an exact, of system of equations**:  
what is meant by, 300, 301;  
must satisfy partial differential equations, 302;  
conditions for existence, 304, 305;  
used to modify system, 311.
- Integral equivalent, minimum number of equations in**:  
of single equation, 145, 146;  
of system of equations, 312–317.
- Integral equivalent, of Pfaffian expression deduced from normal form**, 116, 117;  
of irreducible differential relation, 250 (note);  
of general partial differential equation of second order, 320, 321.
- Integrals of subsidiary system used to diminish the system**:  
when one is known, 178;  
when two are known, 175; with alternative inferences, 177;  
when more than two are known, 179.
- Integrals of Pfaffian equation, effect of, on subsidiary system**:  
when one is known, 173;  
when two are known, 178.
- Integrating factor of single exact equation**:  
general form of, 4;  
quotient of two is a solution, 4, 16;  
determined, by linear equations, 14–17;  
by De Morgan's partial equation, 20;  
by Collet's equations, 25.
- Integrating factors of system of equations**, 39;  
determinant of sets of, a multiplier, 40;  
special method for determination of, when there are four variables, 62–67.
- Integration of extensive equation**, 136, 137 (note).
- Intermediary integral, conditions that partial differential equation of second order possess an**, 307–310.
- Interpretation, numerical, of Grassmann's forms**, 124, 125, 133.
- Interrupted products**, 123;  
conditions in form of, for existence of integrals fewer than general minimum, 123–127;  
interpretation of, 124–127.
- Invariantive integers in transformation of bilinear quantics**, 277;  
arithmetic mean of two, a single sufficient invariant, 278;  
persistence of, sufficient to ensure equivalence of two expressions, 286, 290.
- Invariantive persistence of character of normal form**, 249–251.
- Irreducible differential relation, integral equivalent of**, 250 (note).
- Jacobi's additions to Pfaff's theory**, 81, 82;  
introduction of initial values of variables, 82;  
simplification of Pfaff's reduction, 117–120;  
simplification of Pfaff's solution of differential equation, 183–187;  
theorem relative to four-termed expression, 83, 115, 226.
- Jordan, on multipliers**, 40;  
on tangential transformations, 298.
- Lagrange's examples of cylindrical tangential transformation**, 242.
- Legendrian transformation is a tangential transformation**, 242.
- Lemmas of transformation in Clebsch's theory**, 210.
- Lie's method of treatment of Pfaff's problem, Chap. x, see table of contents**; abstract and character of, 86, 87;  
connection herewith of Darboux's method, 88.
- Lie's theorem for functions in tangential transformation**, 287;  
modified and simplified by Mayer, 237.
- Lie's corollaries from tangential transformations**, 242–244, 248;  
derivable from Clebsch's results, 244.
- Linear transformations, introduced into treatment of Pfaff's problem**, 274;  
applied to bilinear form give two invariantive integers, 277.
- Mayer's method of integrating system of exact equations**, 59;  
in its simplest form, 61.



- Mayer's method, of solving complete system of linear homogeneous partial differential equations, 72–76;  
of obtaining at least one integral, 76–78.
- Mayer's establishment of equations of tangential transformation, 233, 238;  
will not apply to systems of Pfaffians, 331;  
modification and simplification of Lie's theorem, 237.
- Memoirs on systems of Pfaffians, 299.
- Minimum, conditions that the number of equations in the integral equivalent be less than the general:  
in Grassmann's symbolical form, 123–127;  
obtained by Natani, 166, 168;  
by Clebsch, 207;  
by Lie, 255.
- Minimum number of integrals, general:  
for single equation, 89, 113, 114, 145, 146;  
for non-integrable system, 312–316.
- Modification of incompletely integrable system by use of its exact integrals, 311.
- Monge's view of inexact equations, 80.
- Monomial equation of Grassmann:  
if numerical, includes a single partial differential equation of the first order, 122;  
if extensive, includes a system of simultaneous equations in several dependent variables, 122;  
relation of, to integrals of Pfaffian, 123;  
transformation of, 137, 141.
- Multipliers of system of exact equations, 40;  
quotient of two is a solution, with other properties, 41, 43;  
partial differential equations satisfied by, 42.
- Natani's method of integration, of exact equation, 27;  
compared with Euler's, 30;  
compared with du Bois - Reymond's, 32;  
of system of exact equations, 56;  
developed by Mayer, 59–62.
- Natani's method of treatment of Pfaff's problem, Chap. vi, see table of contents;  
abstract and character of, 85, 86;  
application of, to solution of single partial differential equation, 187, 188;  
to solution of system of partial differential equations, 190;  
generalised to system of Pfaffians to determine number, 313;  
inapplicable to determine integrals, 322.
- Normal form:  
involving even number of functions, 194;  
Clebsch's generalisation of, 196;  
involving odd number of functions, 221;  
Clebsch's generalisation of, 222;  
reduction of expression to  $a$ , by Lie, 248;  
relations between elements of equivalent, 252;  
of conditioned expression deduced from that of transformed unconditioned expression, 259–266;  
invariantive persistence of character of, 249;  
conditions which determine character of, 253;  
construction of (see Gradual construction);  
unaltered in character when Cauchy substitution is applied, 258, 269;  
character of, to be inferred from results of Natani and Frobenius, 278 (note);  
used by Frobenius to make expressions equivalent, 285, 289;  
equations of Frobenius which determine, 291, 292.
- Non-integrable systems, 301;  
three fundamental problems relating to, 311;  
minimum number of equations in integral equivalent of, 312–316;  
number of arbitrary integrals in equivalent of, 316;  
arbitrary integrals used to modify, 319.
- Number of independent solutions of a complete system of linear and homogeneous partial differential equations, 69.
- Number of equations in most general integral equivalent of non-integrable system, 312–316;  
application of result to simultaneous partial equations in several dependent variables, 317, 329.
- Number of exact integrals of incompletely integrable system, 304.
- Odd invariantive integer in Frobenius' theory, 287.
- Odd number of variables, unconditioned expression involving, treated by Natani's method, 157–165;

Cambridge University Press

978-1-107-65024-4 - Theory of Differential Equations: Part I: Exact Equations and Pfaff's Problem

Andrew Russell Forsyth

Index

[More information](#)

## INDEX

339

- by Clebsch's method, 219;
- by Lie's method, 268.
- Odd reduction, 111;
  - can always be applied to expression involving odd number of variables, 111;
  - not unique, 111.
- Osculational transformations, 230.
- Partial differential equations, system of, coextensive with system of exact equations, 67, 69, 71.
- Partial differential equation of first order, a single:
  - special case of Pfaffian equation, 122;
  - only one integration necessary in Natani's method, 154;
  - integration of, by Pfaff, 181—183;
  - Pfaff's integration improved by Jacobi, 183—187;
  - by Natani, 187, 188.
- Partial differential equations, solution of simultaneous system of, by Natani, 190;
  - application to solution of, Chap. vii, see table of contents, 238.
- Partial differential equation of second order:
  - conditions for intermediary integral of, 307;
  - nature of integral equivalent of general, 320.
- Persistence of character of normal form, 249.
- Pfaff's Problem, why so called, 81.
- Pfaff's method of reduction, Chap. iv, see table of contents:
  - general theory, 81, 90;
  - additions to, by Gauss and Jacobi, 81, 82.
- Pfaff's theorem as to transformation of an expression:
  - stated, 89;
  - completion of proof of, 112;
  - applied to deduce a later theorem of Jacobi's, 83, 115;
  - simplified by Jacobi, 117—120.
- Pfaff's solution of partial differential equation of first order, 181;
  - simplified by Jacobi, 183.
- Principal integrals, of system of equations, 118, 151;
  - of subsidiary system used for transformation by literal transcription, 120, 152, 153, 157, 170, 171.
- Reciprocation a particular case of tangential transformation, 230, 232.
- Reduced form, integral equivalent derived from, 116, 117;
  - equivalence of, to another reduced form, 195, 219, 249;
  - containing even number of independent functions, 203, 255;
  - generalisation of, 194;
    - containing odd number of independent functions, 219, 268;
  - generalisation of, 220;
    - normal form of, 222;
- Reduction of expression by Pfaff's method, subsidiary equations for, 89—94;
  - three cases of, 95, 100, 103;
  - procedure in, when number of variables is odd, 106—111;
  - completed by Gauss for all cases, 112—114.
- Reduction, even, 105;
  - odd, 111;
  - neither unique, 105, 111;
  - simplified by use of principal integrals, 117—120.
- Reduction of two equations in four variables to canonical form, 325—327.
- Second Method, Clebsch's, 209—218;
  - applies only to equation in even number of variables, if unconditioned, 210;
  - or if Pfaffian determinant alone vanishes, 214.
- Second order, partial differential equation of:
  - conditions that it may possess intermediary integral, 307—310;
  - nature of integral equivalent of, 320, 321.
- Simplification of Pfaff's reduction, by Jacobi, 117—120;
  - of process when some coefficients vanish, 153.
- Simultaneous partial differential equations of first order,
  - in a single dependent variable:
    - solution of, by Natani, 190—193;
    - conditions of coexistence, 191 and note;
  - in several dependent variables:
    - included in extensive monomial equation, 122;
    - number of integral equations equivalent to, 317, 329.
- Simultaneous variations:
  - used by Binet, 148;
  - used by Natani, 148, 149, 157, 174, 178;
  - used by Frobenius, 273;
  - used by Darboux, 295.
- Single exact equation, Chap. i, see table of contents.
- Single integration of subsidiary system sufficient for partial differential equation of first order, 154.

- Solution of partial differential equation:  
 note on history of, 183 (note);  
 Pfaff's method of, 181–183;  
 improved by Jacobi, 183–187;  
 connected with Natani's general  
 result, 187;  
 Natani's method of, 188–190.
- Solution of system of equations:  
 by Natani's method, 190–193;  
 in Clebsch's method, 215;  
 by Mayer, 72–78.
- Subsidiary equations for conditioned  
 expression an exact system, 54, 170;  
 integration of, 170–172.
- Subsidiary equations for Pfaff's reduc-  
 tion, formation of, 89–94.
- Subsidiary equations, when number of  
 variables is even:  
 alternative cases, 94, 296;  
 solution of, in most general case,  
 96;  
 Cayley's symbolical form of, 100;  
 still valid, if determinant alone  
 vanish, 101;  
 principal integrals of, 118.
- Subsidiary equations, when number of  
 variables is odd:  
 in general inconsistent, 110;  
 Jacobi's condition for consistency  
 of, 107;  
 effect of satisfaction of, 109;  
 verified by Natani's results, 164;  
 formed by Natani, 157–159;  
 solution of, 160, 164;  
 principal integrals of, used to  
 transform expression, 151, 160.
- Subsidiary equation for transformation  
 of Grassmann's monomial form, 128–  
 130;  
 solution of, 130–136;  
 interpreted numerically, 133;  
 integration of solved equation,  
 136, 137 (note).
- Subsidiary equations constructed by use  
 of simultaneous variations, 148.
- Subsidiary system, modified by one of  
 its integrals, 173–175;  
 by two of its integrals, 176, 177;  
 by two integrals of original equa-  
 tion, 178;  
 by more than two integrals, 179.
- Summary of methods of treatment of  
 Pfaff's problem, Chap. iii, see table of  
 contents.
- Symbolical solution by Cayley of subsi-  
 diary system, 100.
- System of exact equations, Chap. ii, see  
 table of contents.
- System of Pfaffians, Chap. xiii, see table  
 of contents;  
 memoirs relating to, 299;  
 represented by a single extensive  
 monomial equation, 122, 329.
- System of equations:  
 completely integrable, 300;  
 incompletely integrable, 301;  
 non-integrable, 301;  
 what is meant by exact integral  
 of, 300, 301.
- Tangential transformations, Chap. ix,  
 see table of contents; 298;  
 note on history of, 230;  
 applied by Lie to Pfaff's problem,  
 252.
- Tanner's symbolical forms similar to  
 Grassmann's, 142.
- Transformation of expression when  
 number of variables is even:  
 three cases of, 99, 100–102, 103  
 –105;  
 completed, 113.
- Transformation of expression when num-  
 ber of variables is odd:  
 two cases of, 107–109, 110–111;  
 completed, 114.
- Transformation of expression to mono-  
 mial form, 122.
- Transformation, gradual, of monomial  
 form, 127–141.
- Transformation of expression, by princi-  
 pal integrals, 118, 152, 157;  
 by Cauchy substitutions when it  
 is conditioned, 258, 269.
- Transformations, tangential (see tan-  
 gential transformations).
- Uniqueness, want of, in reductions even  
 or odd, 105, 111.
- Unsolved problems, connected with non-  
 integrable systems of equations, 330.
- Vanishing coefficients, effect of, in sim-  
 plifying construction of normal form,  
 153.
- Vanishing Pfaffian determinant, effect  
 of, on transformation of expression,  
 102, 105.
- Variable, extensive, 121.