

Cambridge University Press

978-1-107-66014-4 - Theory of Differential Equations : Part IV: Partial
Differential Equations: Vol. V

Andrew Russell Forsyth

Table of Contents

[More information](#)

CONTENTS.

CHAPTER I.

INTRODUCTION : TWO EXISTENCE-THEOREMS.

| ART. | | PAGE |
|-------|--|------|
| 1. | Introductory explanations | 1 |
| 2—4. | The character of the general integral of partial equations not to be inferred either by generalising from ordinary equations nor from a reversal of any process of construction | 2 |
| 5. | Sets of variables: the number of equations must generally (though not universally) be not greater than the number of dependent variables | 6 |
| 6. | Any system of equations, equal in number to the number of dependent variables, can be made an amplified system in an increased number of dependent variables of the same type | 8 |
| 7. | Or it can be replaced by one equation (or several equations) of higher order in one dependent variable | 10 |
| 8. | Cauchy's theorem as to the existence of the integral of a system of m equations in m dependent variables made to depend upon two existence-theorems: the first of these theorems | 10 |
| 9—11. | Madame Kowalevsky's proof of the <i>first theorem</i> | 11 |
| 12. | The integrals thus obtained are the only regular integrals that satisfy the imposed conditions | 17 |
| 13. | Note as to possible non-regular integrals and particular variations of the complex variable | 21 |
| 14. | The <i>second theorem</i> , deduced by Madame Kowalevsky from the first theorem: with example and note | 22 |

Cambridge University Press

978-1-107-66014-4 - Theory of Differential Equations : Part IV: Partial
Differential Equations: Vol. V

Andrew Russell Forsyth

Table of Contents

[More information](#)

CONTENTS

xiii

CHAPTER II.

CAUCHY'S THEOREM.

| ART. | | PAGE |
|------|--|------|
| 15. | Remarks on the limitations imposed by the conditions assigned in the subsidiary existence-theorems | 26 |
| 16. | A single equation of the <i>first order</i> $f(x, y, z, p, q)=0$ | 27 |
| 17. | Existence-theorem for $f=0$ when resolvable with regard to p | 28 |
| 18. | Likewise when $f=0$ is resolvable with regard to q | 31 |
| 19. | Summary of results, with statement of exception | 31 |
| 20. | More general form of existence-theorem for $f=0$: with notes | 32 |
| 21. | Geometrical illustration | 34 |
| 22. | Statement of results for single equation of the <i>first order</i> involving n independent variables | 35 |
| 23. | Application of the subsidiary existence-theorems to the equa- tion $f(x, y, z, p, q, r, s, t)=0$ of the <i>second order</i> : its integral when $f=0$ is resolvable with regard to r or to t | 37 |
| 24. | Summary of results, with statement of exception | 41 |
| 25. | Cauchy's theorem for a general class of m equations in m dependent variables of <i>any orders</i> ; with a remark on the number of arbitrary functions in the integral | 43 |
| 26. | Significance of the limitation defining the preceding class of equations | 48 |
| 27. | Comprehensiveness of the preceding class not universal. | 50 |
| 28. | Recent investigations as to the existence-theorem for more widely comprehensive classes of equations | 53 |

CHAPTER III.

LINEAR EQUATIONS AND COMPLETE LINEAR SYSTEMS.

| | | |
|--------|--|----|
| 29. | Equations $X_1p_1 + \dots + X_np_n = 0$, where z does not occur in X_1, \dots, X_n : its $n-1$ functionally independent integrals; and its most general integral, which includes the Cauchy integral | 55 |
| 30. | Equations $X_1p_1 + \dots + X_np_n = Z$, where z does occur in the coefficients: integrals of the subsidiary system and their relation to the equation: a general integral | 60 |
| 31—33. | This general integral is not completely comprehensive: <i>special</i> integrals may exist that are not included | 63 |
| 34. | Summary of results, with examples of special integrals | 67 |
| 35. | The equation in § 30 can be changed into the equation in § 29: relation of their integrals | 71 |
| 36. | Cauchy's theorem associated with the general integral of the equation in § 30 | 73 |

Cambridge University Press

978-1-107-66014-4 - Theory of Differential Equations : Part IV: Partial
Differential Equations: Vol. V

Andrew Russell Forsyth

Table of Contents

[More information](#)

xiv

CONTENTS

| ART. | | PAGE |
|---------|--|------|
| 37. | Simultaneous equations, homogeneous and linear in the derivatives of one dependent variable, the latter not occurring: <i>complete linear systems</i> | 76 |
| 38. | Two properties of complete linear systems | 79 |
| 39. | Canonical form of complete linear systems: systems <i>in involution</i> : complete Jacobian systems | 81 |
| 40. | A complete Jacobian system of m equations in $m+n$ independent variables possesses n functionally distinct integrals: determined by assigned initial conditions | 83 |
| 41, 42. | <i>Fundamental system</i> of integrals of a complete Jacobian system: any integral expressible in terms of them: and the system is unique as a system of regular integrals | 86 |
| 43. | Mayer's method of constructing a fundamental system of integrals of a complete linear system | 89 |
| 44. | Jacobi's method of integration of complete linear systems by successive reduction | 91 |
| 45, 46. | Complete linear systems that are not homogeneous: <i>special</i> integrals: conditions of completeness | 97 |

CHAPTER IV.

NON-LINEAR EQUATIONS: JACOBI'S SECOND METHOD, WITH
MAYER'S DEVELOPMENTS.

| | | |
|---------|---|-----|
| 47. | Equations, or systems of equations, of the first order and of any degree in the derivatives of one dependent variable: Jacobian conditions of coexistence | 100 |
| 48, 49. | These conditions are necessary and sufficient, when satisfied, (i) in the case of n equations, (ii) in the case of $n+1$ equations, there being n independent variables | 103 |
| 50. | A system of simultaneous equations must be rendered <i>complete</i> : the tests | 108 |
| 51. | The case excepted in § 22: tests of a <i>singular</i> integral | 110 |
| 52. | A property of the combinants (F, G) of three functions of $x_1, \dots, x_n, p_1, \dots, p_n$; with inferences | 112 |
| 53. | A property of the combinants [F, G] of three functions of $x_1, \dots, x_n, p_1, \dots, p_n, z$; with inferences | 114 |
| 54. | Consequent removal of the dependent variable from a complete system of equations | 116 |
| 55. | Mayer's development of Jacobi's second method of integration of a complete system of m equations, by transforming it into a system in involution | 117 |
| 56, 57. | Successive integration of this system in involution, with examples | 121 |
| 58, 59. | Removal of one limitation in Mayer's process of integration, by using a contact-transformation | 127 |

Cambridge University Press

978-1-107-66014-4 - Theory of Differential Equations : Part IV: Partial
Differential Equations: Vol. V

Andrew Russell Forsyth

Table of Contents

[More information](#)

CONTENTS

xv

| ART. | | PAGE |
|--------|--|------|
| 60. | Mayer's method is also applicable when the dependent variable is allowed to remain | 134 |
| 61—64. | Jacobi's <i>second method</i> when the Mayer modifications are inexpedient | 136 |
| 65—67. | The method when the dependent variable occurs explicitly | 146 |
| 68. | Application to $f(x, y, z, p, q)=0$; Charpit's method | 156 |
| 69. | Particular examples of integration, chosen so as to give indications of the character of the integral of an equation when the conditions in Cauchy's theorem are not satisfied | 157 |

CHAPTER V.

CLASSES OF INTEGRALS POSSESSED BY EQUATIONS OF THE
FIRST ORDER: GENERALISATION OF INTEGRALS.

| | | |
|---------|---|-----|
| 70. | Are the different integrals obtained for a non-linear equation connected? | 164 |
| 71, 72. | Variation of parameters applied to an integral that involves n arbitrary constants and n independent variables | 165 |
| 73. | <i>Complete integral, general integrals, singular integral</i> | 171 |
| 74. | But this aggregate is not completely comprehensive: <i>special integrals</i> : tests, with examples | 172 |
| 75. | Conditions that an integral involving the proper number of parameters is complete | 178 |
| 76. | The singular integral, as derived from the complete integral, can be derived from the differential equation: formal tests of its existence | 182 |
| 77. | But these tests are not sufficient: they can be satisfied for integrals that are not singular | 185 |
| 78. | Summary of the preceding results for an equation $f(x, y, z, p, q)=0$, with example of the exception in § 77 | 186 |
| 79. | Construction of equations having exceptional integrals as in § 77 | 190 |
| 80. | An integral of an equation may contain more arbitrary parameters than independent variables: tests that it leads to a single equation | 192 |
| 81. | An equation involving m essential parameters, where $m < n$, leads to a <i>complete system</i> of partial equations; it is a complete integral of that system and, by means of the variation of parameters, leads to other integrals | 193 |
| 82. | The aggregate thus obtained, though widely comprehensive, does not include special integrals of the system | 198 |
| 83. | When a complete system possesses a singular integral, that integral may be derivable without a knowledge of the complete integral | 201 |

Cambridge University Press

978-1-107-66014-4 - Theory of Differential Equations : Part IV: Partial

Differential Equations: Vol. V

Andrew Russell Forsyth

Table of Contents

[More information](#)

xvi

CONTENTS

CHAPTER VI.

THE METHOD OF CHARACTERISTICS FOR EQUATIONS IN TWO INDEPENDENT
VARIABLES: GEOMETRICAL RELATIONS OF THE VARIOUS INTEGRALS.

| ART. | | PAGE |
|---------|---|------|
| 84. | Subsidiary equations obtained by changing the two independent variables | 205 |
| 85. | Effect of the additional conditions thus introduced | 208 |
| 86. | Modes of satisfying these conditions: the different integrals obtained | 210 |
| 87, 88. | Darboux's detailed modification of Cauchy's method and proof | 212 |
| 89. | The singular integral | 216 |
| 90. | Exceptional case: use of Legendrian contact-transformation; examples of the whole theory | 217 |
| 91. | Geometrical interpretation: the complete integral, with a cone T , a reciprocal cone N , and a curve C in any plane | 221 |
| 92. | General integrals as derived from complete integrals: <i>characteristics</i> of families of surfaces and their properties | 224 |
| 93. | <i>Characteristic developable</i> , and its properties | 227 |
| 94. | Differential equations of the characteristics derived from the original equation: they are the subsidiary equations in Cauchy's method | 228 |
| 95. | They can be derived also from the integral equations | 230 |
| 96. | Equations of the characteristic when the integral is given in a form expressing x, y, z in terms of two variables and two parameters | 233 |
| 97. | <i>Edge of regression</i> of the surface representing the general integral: its equations | 236 |
| 98. | <i>Integral curves</i> ; they include characteristics; their contact-relations with the surfaces representing the integrals | 238 |
| 99. | Equations of edge of regression of a selected general surface derived from original equation | 242 |
| 100. | Lie's classification of partial equations of the first order according to the nature of the characteristics on the complete integral: in particular, when they are (i) asymptotic lines, (ii) lines of curvature, (iii) geodesics | 244 |

CHAPTER VII.

SINGULAR INTEGRALS AND THEIR GEOMETRICAL PROPERTIES:
SINGULARITIES OF THE CHARACTERISTICS.

| | | |
|------|---|-----|
| 101. | The Singular Integral (when it exists) is an envelope of the complete integral; its relations to the other surfaces and the curves: Darboux's theorem on <i>conjugate characteristics</i> | 249 |
|------|---|-----|

Cambridge University Press

978-1-107-66014-4 - Theory of Differential Equations : Part IV: Partial
Differential Equations: Vol. V

Andrew Russell Forsyth

Table of Contents

[More information](#)

| | CONTENTS | xvii |
|-----------|--|------|
| ART. | | PAGE |
| 102. | The properties of the singular integral are used to connect it with the original equation | 252 |
| 103. | Relation between the general integral and the singular integral | 254 |
| 104, 105. | Contact between the general integral and singular integral usually is of first order: conditions that the contact may be of the second order | 255 |
| 106. | Order of contact between complete integral and singular integral | 260 |
| 107. | Effect of singularities on the characteristics: the five possible cases | 261 |
| 108. | Case I: an ordinary point | 262 |
| 109. | Case II: when $P=0, Q=0$ | 263 |
| 110. | Case III: when $X+pZ=0, Y+qZ=0$ | 265 |
| 111. | Case IV: when $P=0, Q=0, X+pZ=0, Y+qZ=0$ | 266 |
| 112. | Various Sub-cases of Case IV | 269 |
| 113—115. | Exceptions to Case IV, when the equations are not independent of one another: characteristics and a conditioned singular integral | 271 |
| 116. | Case V: when $P=0, Q=0, X=0, Y=0, Z=0$ | 278 |

CHAPTER VIII.

THE METHOD OF CHARACTERISTICS IN ANY NUMBER OF
INDEPENDENT VARIABLES.

| | | |
|------|---|-----|
| 117. | Equations of characteristics, when there are n independent variables, in Darboux's form | 282 |
| 118. | The additional conditions associated with the new variable | 284 |
| 119. | Modes of satisfying these conditions: kinds of integrals thus deduced | 288 |
| 120. | The singular integral: special integrals | 291 |
| 121. | Exceptional case | 292 |
| 122. | Equations of characteristics derived from the complete integral in two ways | 293 |
| 123. | The various hypersurface integrals touch at a position common to all of them | 297 |
| 124. | Order of this contact | 301 |
| 125. | The investigation in § 124, when $n=2$ | 306 |
| 126. | Examples of the general theory | 310 |
| 127. | Contact of the complete integral and singular integral | 312 |

CHAPTER IX.

LIE'S METHODS APPLIED TO EQUATIONS OF THE FIRST ORDER.

| ART. | | PAGE |
|------|---|------|
| 128. | Contact-transformations: Mayer's form of equations giving the most general transformation defined by Lie . . . | 315 |
| 129. | Equations for infinitesimal contact-transformations . . . | 317 |
| 130. | Also for those which involve the dependent variable in only a single equation, with the special cases, (i) infinitesimal, (ii) homogeneous, (iii) infinitesimal and homogeneous: relation of the last to theoretical dynamics | 318 |
| 131. | The determination of infinitesimal homogeneous contact-transformations, which leave any partial equation of the first order unaltered, is equivalent to the integration of the equation | 323 |
| 132. | Application of finite contact-transformations to the integration of $f(x_1, \dots, x_n, z, p_1, \dots, p_n) = 0$: with the various types of integrals | 324 |
| 133. | Likewise for an equation $f(x_1, \dots, x_n, p_1, \dots, p_n) = 0$. . . | 329 |
| 134. | Likewise for a complete system when the dependent variable occurs explicitly | 333 |
| 135. | And for a complete system when the dependent variable does not occur explicitly | 337 |
| 136. | Relations between the equations of a complete system and the constituents of a contact-transformation . . . | 342 |
| 137. | <i>Groups of functions: sub-groups: systems in involution: a function in involution with a group: simple properties of groups</i> | 344 |
| 138. | Lie's theorem on a group of order r and an associated complete Jacobian system of equations: <i>reciprocal groups</i> | 347 |
| 139. | <i>Indicial functions of a group: their number: two reciprocal groups have the same indicial functions</i> | 350 |
| 140. | <i>Canonical form of a group not in involution, determined by its indicial functions</i> | 354 |
| 141. | When a group of order $m + 2g$, having m indicial functions, is in canonical form, it can be amplified into a group of order $2n$ (where $2n$ is the total number of variables) in canonical form | 361 |
| 142. | When a group of functions is subjected to a contact-transformation, the order of the group and the number of indicial functions are invariants: and conversely . . . | 364 |
| 143. | Highest order of a sub-group in a group, and greatest number of indicial functions that a group can possess . . . | 365 |

Cambridge University Press

978-1-107-66014-4 - Theory of Differential Equations : Part IV: Partial
Differential Equations: Vol. V

Andrew Russell Forsyth

Table of Contents

[More information](#)

CONTENTS

xix

| ART. | | PAGE |
|------|---|------|
| 144. | Construction of the sub-group of greatest order in a group . | 366 |
| 145. | Application of properties of groups to the integration of complete systems of partial equations of first order . | 367 |

CHAPTER X.

THE EQUATIONS OF THEORETICAL DYNAMICS.

| | | |
|-----------|--|-----|
| 146. | Hamilton's canonical form of the equations of general theoretical dynamics: the function H , and the asso- ciated partial equation | 371 |
| 147. | Hamilton's equations connected with the function S . . . | 376 |
| 148. | Relations between the two sets of equations, connected with <i>initial values</i> | 379 |
| 149. | Hamilton's equations connected with the function A . . . | 381 |
| 150. | Extensions of Hamilton's results by Jacobi into a method (called <i>Jacobi's first method</i> or the <i>Hamilton-Jacobi method</i>) for the integration of equations of the first order | 382 |
| 151. | Mayer's generalisation of the complete integral thus ob- tained | 387 |
| 152. | Corresponding results when the dependent variable occurs | 391 |
| 153, 154. | The Poisson-Jacobi <i>combinant</i> and its use in the inte- gration of the canonical system | 392 |
| 155, 156. | A contact-transformation leaves a canonical system un- altered: and every transformation having this property is of the contact type | 398 |
| 157. | Special case of contact-transformation applied to a canoni- cal system | 403 |
| 158. | The canonical equations are the equations of the contact- transformation applied to the energy of the system . . | 404 |

CHAPTER XI.

SIMULTANEOUS EQUATIONS OF THE FIRST ORDER.

| | | |
|------|---|-----|
| 159. | Introductory remarks: differential equations derived from an integral system | 408 |
| 160. | König's system of rm equations in m dependent variables and n independent variables ($r \leq n$) | 411 |
| 161. | Conditions of <i>complete integrability</i> | 413 |
| 162. | Method of integration when the conditions are satisfied . . | 418 |

Cambridge University Press

978-1-107-66014-4 - Theory of Differential Equations : Part IV: Partial
Differential Equations: Vol. V

Andrew Russell Forsyth

Table of Contents

[More information](#)

xx

CONTENTS

| ART. | | PAGE |
|-----------|---|------|
| 163—5. | Different kinds of integrals indicated : how far they can be connected by the variation of parameters : with an example | 419 |
| 166. | Hamburger's sets of m linear equations in m dependent variables : the method ineffective in general when there are more than two independent variables | 428 |
| 167. | Hamburger's method when there are two independent variables | 430 |
| 168. | Extreme case in Hamburger's method : Jacobi's linear equations and their integral | 432 |
| 169. | Equations for which $m = 2$, $n = 2$, with examples | 435 |
| 170, 171. | Hamburger's process in detail when $m > 2$ | 442 |
| 172. | Alternative method, in which the subsidiary equations are partial, not ordinary | 449 |
| 173. | Hamburger's method applied to non-linear simultaneous equations in two independent variables and as many dependent variables as equations | 455 |
| 174. | Proof of the main theorem | 460 |
| 175. | Alternative method which sometimes is effective | 466 |
| 176. | When the Jacobian process for a single equation is generalised, as is the Lagrangian process in Hamburger's method, it leads to subsidiary equations, which are lineo-linear but cannot be integrated by inverse operations of only the first class | 474 |
| 177. | Two equations of the first order in two dependent variables do not, in general, lead to an equation of the second order in either variable alone | 477 |