

Index

2, 4-D, 4, 29

Acacia longifolia, 252, 273

Acacia saligna, Port Jackson willow, 286

acacias, 59

acaricide, 89

acclimitization societies, 19–20

Achilles heel, 25, 259

Acridotheres tristis, mynah bird, 137

activation–inhibition hypothesis, 257

Aculus fockeui, peach silver mite, 6

Acyrtosiphon pisum, pea aphid, 99, 125, 132, 155, 181

Adalia bipunctata, two-spotted lady beetle, 144

additive strategy, 96

Adoxophyes honmai, 222

Aedes aegypti, 384

Aedes vexans, 209

Africa, 55, 64, 68, 157, 237, 369

Ageratum conyzoides, 105

Agrius planipennis, emerald ash borer, 377

Agrobacterium radiobacter, 299

Agrobacterium tumefaciens, 298

agrocin, 299

Agrypon flaveolatum, 119

airplane, 80

alders, 282

alfalfa, 97, 101, 125, 155, 379

alfalfa caterpillar, *Colias eurytheme*, 207

alfalfa weevil, *Hypera postica*, 52, 101

Algarobius prosopis, 64

allee effects, 125

Alliaria petiolata, garlic mustard, 20

amber disease, 213

Amblyseius swirskii, 70

ambushers, nematodes, 193

American chestnut, *Castanea dentata*, 309

Americas, 19

Ampelomyces quisqualis, 300, 319

amphibian, 16

Amsinckia intermedia, fiddleneck, 255

Amyeloides transitella, navel orangeworm, 199

Amylostereum areolatum, 196

Anagrus epos, 103

Anagyrus kamali, 60

Anagyrus sp. near *kivuensis*, 50

Andrés, L. A., 31

Anguina amsinckiae, 255

Anisoplia austriaca, wheat grain beetle, 31

Anoplophora chinensis, citrus longhorned beetle, 239

ant lions (Myrmeleontidae), 141

Anticarsia gemmatilis, velvetbean caterpillar, 101, 221

ants (Formicidae), 34, 36, 93, 140, 141, 151, 154

ants, tending Hemiptera, 179, 267

Aonidiella aurantii, California red scale, 6, 7, 113

aphid flies (Chamaemyiidae), 148

Aphidius ervi, 125

aphids (Aphididae), 6, 18, 47, 62, 67, 79, 94, 100, 101, 124, 143, 305, 330, 343

Aphis glycines, soybean aphid, 382

Aphis gossypii, cotton aphid, 87, 92, 131, 153, 157, 379

Aphis nerii, oleander aphid, 141

Aphthona lacertosa, 273

Aphthona nigricutis, 273

Aphytis chrysomphali, 113

Aphytis lingnanensis, 113

Aphytis melinus, 113

Apis mellifera, honey bee, 12, 329

Apoanagyrus lopezi, 183

apple orchards, 7, 88, 104, 221, 380

apples, 98, 222, 302

aquatic weeds, 13, 246, 249

Argentine stem weevil, *Listronotus bonariensis*, 380

Arizona, 380

Armillaria mellea, 294

armyworm, *Spodoptera exigua*, 379

arsenic, 4

Arthrobotrys, 324

Arthrobotrys anconia, 301

Arthrobotrys dactyloides, 301

Ascension Island, 246

Asclepias spp., milkweed, 256, 381

Ascomycota, 231–233, 311

ash, *Fraxinus* spp., 377

Asia, 64, 330

- Asian corn borer, *Ostrinia furnacalis*, 82–83
 Asian gypsy moth, *Lymantria dispar asiatica*, 28
 aspens, 282
Aspergillus flavus, 311
 Atkinson, P. L., 360
 Atlantic Canada, 259
 attract and reward, 100
 augmentation biological control, 66–83, 379
 comparing inundative vs. inoculative, 71–72
 definitions, 66–69
 greenhouses, 69–71
 host specificity, 73
 inoculation, 68–69
 inundation, 66–67
 macroorganisms, 77
 microorganisms, 77–79
 needing a market, 72–73
 Australia, 19, 20, 30, 43, 50, 51, 59, 64, 96, 137,
 197, 224, 252, 267, 340, 342, 347
 autumnal moth, *Epirrita autumnata*, 119
Azadirachta indica, 368
- Bacillaceae, 204
Bacillus subtilis, 306
Bacillus thuringiensis, 23, 28, 67, 72, 78, 83,
 204–212
 safety, 328–329
Bacillus thuringiensis israelensis, 209, 384
 bacteria, entomopathogenic, 203–214
 Bt resistance, 211–212
 toxins, 23, 203
 bacteria, symbiotic, 190
 Baculoviridae, 215
 Bailey, V. A., 116
 balance of nature, 24
 balanced mortality hypothesis, 174
 banker plants, 70, 94, 105, 186
 Basidiomycota, 284
 basking, 238
 Bassi, Agostino, 29
Bathyplectes curculionis, 52
 beans, 71, 92
Beauveria bassiana, 29, 83, 236
Beauveria brongniartii, 71, 239
 beekeepers, 347
 beetle banks, 93
Bemisia tabaci, 50, 373
Bessa remota, 334
 big-eyed bug, *Geocoris pallens*, 138
 BioControl, 29
 Biocontrol Science and Technology, 30
 biodiversity, 16, 329, 338, 353, 378
 bioherbicides, 67, 279
 biointensive IPM, 362
 biological control
 agents (BCAs), 292
 arthropod pests, history, 30–31
 definition, 22–24
 diversity, 25–28
 history, 30
 host density, 28
 market, 371
 plant pathogens and plant parasitic nematodes,
 32–33
 strategies, 25
 weeds, history, 31–32
 Biological Control: Theory and Application in Pest
 Management, 30
 biological pollution, 20
 biopesticides, 67, 379
 biopiracy, 386
 bioprotectants, 313
 bioresidual control, 374
 biotrophy, 278, 291
 biotypes, 51, 279
 birch trees, 119
 bird cherry-oat aphid, *Rhopalosiphum padi*, 154
 black crust of rubber, 320
 blackberry, *Rubus* spp., 276
 blackberry leafhopper, *Dikrella californica*, 103
 blind releases, 70, 186
 Blossey, B., 274
 bodyguards, 156, 190, 233
 Boettner, G. H., 338
Bombyx mori, silkworm, 29, 329
 Bordeaux mix, 4
Boreioglycaspis melaleucae, 253
 botanical pesticides, 367
Botrytis, 319
Botrytis cinerea, 294
 bottom-up forces, 256
 Braconidae, 162
 Brazil, 20, 32, 221
 British Columbia, 28, 259, 367
 British House of Lords, 15
 British Isles, 20
 brown marmorated stink bug, *Halyomorpha halys*,
 88
 brown planthopper, *Nilaparvata lugens*, 374
 browntail moth, *Euproctis chrysorrhoea*, 344
 bruchids (Bruchidae), 63
 Bt-cotton, 373
Buchnera, 203
 buckwheat, 95, 100
 bumble bees, 16
 bunds, 104
Bupalus piniarius, pine looper moth, 177
 Burdon, J. J., 246
Burkholderia cepacia, 324
 Burma, 28
 Bushland, R. C., 367
 Butler, L., 328

- cabbage, 99
 cabbage looper, *Trichoplusia ni*, 173, 211
 cacao, 104
Cactoblastis cactorum, 252, 262, 266, 342
 calicivirus, 227
 California, 18, 30, 43, 82, 95, 97, 113, 279
 California red scale, *Aonidiella aurantii*, 6, 7, 113
Callosamia promethea, 338
 Canada, 28, 69, 174, 272, 282, 385
Candida oleophila, 312
 cane toad, *Rhinella marina*, 137, 339
 cannibalism, 75, 76, 241, 268
 Carabidae (ground beetles), 149–151
 carbamates, 16, 89
 carbon dioxide, 379
 cardenolide steroids, 141
 cardiac glycosides, 256
Cardiochiles nigriceps, 176
Carduus nutans, musk thistle, 249, 252
 Caribbean, 343
 Carson, Rachel, 13–16, 332
 cassava, 157, 183
 cassava green mite, *Mononychellus tanajoa*, 55, 157
 cassava mealybug, *Phenacoccus manihoti*, 55, 64, 183
Castanea, 377
Castanea dentata, American chestnut, 309
 Cecidomyiidae (gall and wood midges), 148
 cecropia moth, *Hyalophora cecropia*, 338
Centaurea diffusa, diffuse knapweed, 248
Centaurea spp., knapweed, 260
 Central America, 92
 centrifugal phylogenetic method, 345
Ceratitis capitata, Mediterranean fruit fly, 353
 cereal aphids, 34
 cereal cyst nematode, *Heterodera avenae*, 295, 323
 cereal leaf beetle, *Oulema melanopus*, 90
 cereals, 94
 Chalcidoidea, 162
 Chamaemyiidae (aphid flies), 148
 Charudattan, R., 280
 Chateauguay region, 321
 chemical pesticides, 5, 6–12, 22
 side effects, 15–17
 chestnut, 377
 chestnut blight, 309
 China, 4, 5, 28, 82, 92, 96, 104, 105, 236, 265
 chitin, 322
 chlamydospores, 314, 321
Chloris gayana, Rhodes grass, 96
Chondrostereum purpureum, 282
 Chontrol, 282
Choristoneura fumiferana, spruce budworm, 209
Chromobacterium subtsugae, 23
Chrysolina quadrigemina, 261
 Chrysopidae (green lacewings), 131
 cinnabar moth, *Tyria jacobaeae*, 255, 257
 citrus, 7, 18, 28, 30, 96, 105, 113, 276, 281
 citrus black scale, *Saissetia oleae*, 53
 citrus longhorned beetle, *Anoplophora chinensis*, 239
 citrus red mite, *Panonychus citri*, 96, 105
Citrus sinensis, sweet orange, 305
 citrus tristeza, 305
 classical biological control, 41–65, 386
 definition, 41
 economics, 64–65
 foreign exploration, 51–52
 main uses, 45–48
 methods, 55
 new associations, 49
 success, 55–64
 climate change, 378–379
 climate modeling, 51
Coccinella novemnotata, nine-spotted lady beetle, 343
Coccinella septempunctata, seven-spotted lady beetle, 154, 155, 157, 343
 Coccinellidae (lady beetles), 143–144, 156, 330, 343, 379
 cochineal, 31, 267
Cochliomyia hominivorax, screwworm fly, 366
 coconut, 220
 coconut moth, *Levuana iridescens*, 333
 coconut palms, 220, 334
 coconut rhinoceros beetle, *Oryctes rhinoceros*, 64, 220
 Code of Best Practices, 352
 Code of Conduct, 352
 codling moth, *Cydia pomonella*, 222, 367, 380
Coelomomyces, 233
 coevolution, 226, 350
 coevolutionary arms race, 181
 coffee, 5, 49
 coffee mealybug, *Planococcus kenyae*, 49
 Coleoptera, 271
Coleotichus blackburniae, green koa bug, 341
Colias eurytheme, alfalfa caterpillar, 207
 collards, 91, 130, 153
 Collego, *Colletotrichum gloeosporioides* f. sp. *aeschnomene*, 281
Colletotrichum gloeosporioides f. sp. *aeschnomene*, 280
Colletotrichum salsolae, 279
 Colombia, 5
 Colorado potato beetle, *Leptinotarsa decemlineata*, 4, 19, 209
 community ecology, 129–135
 apparent competition, 131
 competitive interactions, 131
 exploitative (or resource) competition, 130

- food webs, 129
- interference competition, 131
- intraguild predation, 131
- companion plants, 102
- competitive displacement, 113
- compost, 322
- Compsilura concinnata*, 337, 344
- Congo, 183
- conservation biological control, 85–105
 - biodiversity, 85–86
 - conserving natural enemies, 87–90
 - definition, 85
 - enhancement, 90–105
 - predatory mites (Acari), 88–89
- Convention on Biological Diversity, 386
- convergent lady beetle, *Hippodamia convergens*, 159
- copepod, 233
- Copidosoma floridanum*, 173
- Copidosomopsis tanytmenus*, 172
- copper sulfate, 4
- corn, 4, 82, 92, 140, 210, 381
- corn borers, 236
- corn leafhopper, *Dalbulus maidis*, 93
- corn rootworm, *Diabrotica* spp., 93
- cost:benefit analysis, 64, 251, 386
- Costelytra zealandica*, grass grub, 213
- Cotesia flavipes*, 237
- Cotesia marginiventris*, 135
- Cotesia sesamiae*, 375
- cotton, 92, 97, 101, 153, 210, 306, 373, 378, 380
- cotton aphid, *Aphis gossypii*, 87, 92, 131, 153, 157, 379
- cotton bollworm or tomato fruitworm, *Helicoverpa zea*, 176, 360
- cottony cushion scale, *Icerya purchasi*, 30, 43
- CpGV, 380
- crab spiders (Thomisidae), 152
- crop rotation, 320, 322
- cross protection, 305
- crown gall, 298
- crucifers, 269
- cruisers, nematodes, 193
- Cryphonectria parasitica*, 309
- Cryptochaetum iceryae*, 30, 44
- Cryptolaemus montrouzieri*, mealybug destroyer, 59
- Ctenopharyngodon idella*, grass carp or white amur, 276, 277
- cucumbers, 97
- Culex pipiens*, 209
- Culex quinquefasciatus*, southern house mosquito, 212
- cultural controls, 4
- Cydia pomonella*, codling moth, 222, 367, 380
- Cylindrosporium concentricum*, 320
- Cyperus esculentus*, yellow nutsedge, 268
- Cyrtobagous salviniae*, 250
- cyst nematodes, 324
- cytoplasmic polyhedroviruses, 216
- Dactylopius ceylonicus*, 32, 273
- Dactylopius opuntiae*, 276
- Dactylopius* spp., 267
- dairy, 362
- Dalbulus maidis*, corn leafhopper, 93
- damping off, 316, 317
- Danaus plexippus*, monarch butterfly, 256, 381
- dates, 28
- DBCP, 16
- DDT, 4, 6, 7, 9, 16, 29
- DeBach, P., 23
- Deladenus siricidicola*, 197
- delayed density dependence, 118, 120
- delta-endotoxins, 206
- deltamethrin, 11
- Dendrolimus* spp., pine caterpillars, 236
- dengue, 384
- Denno, R. F., 155
- density dependence, 118, 267
- density independence, 118
- density vague, 118
- density-dependent control, 68
- Dermolepida albohirtum*, greyback cane beetle, 340
- Desmodium*, 375
- Deuteromycota, 232
- DeVine, *Phytophthora palmivora*, 281
- Diabrotica* spp., corn rootworms, 93
- Diabrotica virgifera virgifera*, western corn rootworm, 135, 190
- Diachasmimorpha tryoni*, 353
- Diadegma semiclausum*, 95
- Dialectica scalariella*, 347
- diamondback moth, *Plutella xylostella*, 99, 211
- Diatraea saccharalis*, sugarcane borer, 237
- Dicyma pulvinata*, 320
- dieldrin, 15
- Dietrick, E. J., 97
- diffuse knapweed, *Centaurea diffusa*, 248
- Dikrella californica*, blackberry leafhopper, 103
- Dipel, 67
- Dixon, B., 329
- domatia, 98
- drift, 332, 378
- earwigs (Dermaptera), 104
- eastern subterranean termite, *Reticulotermes flavipes*, 364
- Echium plantagineum*, 347
- ecological host range, 348
- economic injury level, 26–27, 360

- ecosystem services, 25, 85, 329, 348
ecotypes, 51
ectoparasitoids, 168
education, 71
Edwardsiana prunicola, prune leafhopper, 103
Egypt, 4
Elatobium abietinum, 379
Elkinton, J. S., 338
emerald ash borer, *Agrilus planipennis*, 377
emerging diseases, 384
encapsulation, 181
Encarsia formosa, 31, 70, 76, 97, 185
endoparasitoids, 168
endophytes, 232
endospores, 314
enemies hypothesis, 91
enemy release hypothesis, 41, 126–127
England, 31
enhancement, conservation biological control
 artificial food, 101–102
 cover crops, 96–97
 crop management, 97–98
 crop residue management, 97
 physical environment, 101
 plant characteristics, 97–99
 plant messaging, 100
 polyculture in use, 93
 providing alternate food, 96
 providing refuges, 93–94
 providing shelter, 104
 soil, 100–101
 surrounding areas, 104
 vegetational diversity, 91–92
Entomophaga aulicae, 53
Entomophaga maimaiga, 53, 128, 348
Entomophthorales, 231
environmental safety
 changes to support, 356–357
environmental weed, 265
Eotetranychus sexmaculatus, six-spotted mite, 120
Ephesia kuehniella, Mediterranean flour moth, 172, 207
Epilachna varivestis, Mexican bean beetle, 71
Epirrita autumnata, autumnal moth, 119
epizootics, 101, 127, 233, 329
eradication, 28
Eretmocerus hayati, 51
Erwinia amylovora, 319
Erynnis ello, 224
Erythroneura elegantula, grape leafhopper, 103
Eucalyptus, 96
Euglandina rosea, 333, 340
Euphorbia, 102
Euphorbia esula, leafy spurge, 51, 272
Euproctis chrysorrhoea, browntail moth, 344
Europe, 4, 31, 69, 330
European corn borer, *Ostrinia nubilalis*, 74, 177, 381
European rabbit, *Oryctolagus cuniculus*, 336
European spruce sawfly, *Gilpinia hercyniae*, 220
European Union, 12, 17, 211, 361
Eutreta xanthochaeta, 353
Everglades, 253
evolution of increased competitive ability, 265
experimental methods
 caging, 34–35
 direct observation, 36
 natural enemy activity, 36–37
 prey enrichment, 36
 removal, 35
 sampling, 33–34
extirpation, 333
Fagopyrum esculentum, buckwheat, 95
fall armyworm, *Spodoptera frugiperda*, 93
farmer field schools, 374
fenitrothion, 11
Fennoscandia, 119
fermentation, 78
fertilization, 275
fertilizer, 249
fiddleneck, *Amsinckia intermedia*, 255
Fiji, 334
fire ants, *Solenopsis* spp., 179, 361
fire blight, 319
fish, 276
flea beetles (Chrysomelidae), 273
fleas (Siphonaptera), 225
Fleming, Alexander, 311
Florida, 195, 253, 276, 281, 343
flowers, 94, 102
fluvalinate, 6
Follett, P. A., 341
Food and Agriculture Organization, 17, 54, 352, 374
food webs, 129–135
 bottom-up, 129, 132
 consumer–resource interactions, 130
 guild, 130
 top-down, 129, 132
 tritrophic interactions, 132
 trophic cascade, 129
forest, 24, 344
Formica, 151
Formica animosa, 28
formulation, 78
France, 4, 321
Fraxinus spp., ash, 377
frenchi cane beetle, *Lepidiota frenchi*, 340
fruit borers, 6
fruit flies (Tephritidae), 260
functional redundancy, 86

- functional responses, 112
 fundamental host range, 348
 fungi, entomopathogenic, 229–241
 augmentation biological control, 235–239
 classical biological control, 233–235
 dispersal, 230
 persistence, 231
 Fungi Imperfecti, 232
 fungicides, 359
Fusarium oxysporum, 321
Fusarium oxysporum f. sp. *melonis*, 321
- Gaeumannomyces graminis* var. *tritici*, 322
Galendromus occidentalis, 120
 gall, 245
 gall midge (Cecidomyiidae), 253
Galleria mellonella, wax moth, 36
Gambusia affinis, 139
Gambusia holbrooki, 139
Gambusia, mosquito fish, 139, 384
 garlic mustard, *Alliaria petiolata*, 20
 Gause, G. F., 292
 geese, 276
 genetic bottleneck, 350
 genetic engineering, 210–211
 genetically engineered microbial agents, 78
 genetically modified crops, 5, 23
Geocoris pallens, big-eyed bug, 138
 Germany, 151, 209
 giant African snail, *Lissachatina fulica*, 333, 340
 GiHeNPV, 220
Gilpinia hercyniae, European spruce sawfly, 220
Gliocladium virens, 300
 global change, 376
 global homogenization, 377
 globalization, 377
Glossina austeni, tsetse fly, 367
 glucosides, 269
 glyphosate, 381
Glyptapanteles liparidis, 352
 goats, 245, 276
 Goeden, R. D., 31
 granulovirus, 216, 222, 223
 augmentation, inoculative, 222
 grape leafhopper, *Erythroneura elegantula*, 103
 grapes, 4
 grass carp, *Ctenopharyngodon idella*, 276–277
 grass grub, *Costelytra zealandica*, 213
 grasshoppers, 68, 237
 green fluorescent protein, 296
 green koa bug, *Coleotichus blackburniae*, 341
 green lacewings (Chrysopidae), 75, 76, 97, 101, 131, 158
 Green Muscle, *Metarhizium acridum*, 238
 green peach aphid, *Myzus persicae*, 360
 green rice leafhopper, *Nephotettix cincticeps*, 36
 greenhouse whitefly, *Trialeurodes vaporariorum*, 31, 70, 98, 185
 greenhouses, 69–71, 185, 211, 371, 372, 380
 gregarious parasitoids, 170
 greyback cane beetle, *Dermolepida albohirtum*, 340
 ground beetles (Carabidae), 93, 140, 149–151
Gryllus integer, western trilling cricket, 177
 gypsy moth nucleopolyhedrovirus, 355
 gypsy moth, *Lymantria dispar*, 35, 36, 132, 133, 209, 233, 328, 337, 344, 348, 352, 367
- habitat depletion, 341, 343
 habitat displacement, 329
 Hagen, K. S., 101
 Halloween lady beetle, *Harmonia axyridis*, 329
Halyomorpha halys, brown marmorated stink bug, 88
Hamiltonella, 181, 203
 harlequin ladybird, *Harmonia axyridis*, 18, 329, 343
Harmonia axyridis, multicolored Asian lady beetle, 18, 329, 344
Harpalus pensylvanicus, 155
Harpobittacus nigriceps, 259
 Harris, P., 248, 260
 Hawaii, 19, 63, 332, 340
Helicoverpa zea, cotton bollworm or tomato fruitworm, 176, 360
Heliothis virescens, tobacco budworm, 176
Hemileuca maia maia, 338
 Hemimetabola, 138
 Hemiptera, 58, 131, 145–146, 151, 267
 herbivore-induced plant volatiles, 100, 132
Herpestes javanicus, small Indian mongoose, 137
Heterobasidion annosum, 32, 303
Heterodera avenae, cereal cyst nematode, 295, 323
Heteropan dolens, 334
 Heterorhabditidae, 190
Heterorhabditis, 190–194
Heterorhabditis megidis, 135
 Hindu, 4
Hippodamia convergens, convergent lady beetle, 159
Hirsutella rhossiliensis, 323
 Holling, C. S., 111
 Holometabola, 138, 263
 Homer, 4
Homona magnanima, 222
 honey bee, *Apis mellifera*, 12, 330
 honey bee colonies, 16
 honeydew, 98, 151, 267
 hops, 100
 host feeding, 173, 183, 184
 host-induced plant volatiles, 138
 host range, 53
 native, 344–345
 potential for changes, 351

- host-range testing, 53, 344–345
 - parasitoids and predators, 347–348
 - phytophages, 345–347
- host specificity, 58, 332
- host switching, 333
- house fly, *Musca domestica*, 9, 362
- hoverflies (Syrphidae), 95, 148
- Howarth, F. G., 31, 351
- Huffaker, C. B., 120
- Hunt-Joshi, T. R., 274
- Hyalophora cecropia*, cecropia moth, 338
- hybridization, 351
- Hypera postica*, alfalfa weevil, 52, 101
- Hypericum perforatum*, Klamath weed, 261
- hyperparasitism, plant pathogens, 300
- hyperparasitoids, 53, 171
- hyphal interference, 304
- Hypoaspis miles*, 148
- Hypocreales, 232
- hypovirulence, 305, 310
- Icerya purchasi*, cottony cushion scale, 30, 43
- Ichneumonidae, 162
- idiobionts, 168
- India, 20, 32, 59, 97
- Indian meal moth, *Plodia interpunctella*, 211
- Indonesia, 374
- induced resistance, plant, 304
- induced systemic resistance, 305, 316
- infective juvenile, 189
- inoculative augmentation, 275
- insect growth regulators, 367
- insecticidal check method, 35
- insecticide resistance, 362
- insidious flower bug, *Orius insidiosus*, 76
- integrated pest management, 31, 375
 - autocidal controls, 366–367
 - bio-rational chemicals, 367–368
 - definition, 360
 - ecologically based, 368–369
 - host plant resistance, 365–366
 - mechanical, physical, cultural controls, 365
 - synthetic chemical pesticides, 368
- integrated vector management, 385
- intercropping, 92
- international cooperation, 386–387
- International Federation of Organic Agriculture Movements, 18
- International Organization for Biological Control, 30, 75, 371
- intraguild predation, 153
- invasive plants, 382
- invasive species, 13, 334, 341, 342, 344, 377–378, 387
- invasive species treadmill, 262
- inverse density dependence, 125
- islands, 340
- Ixodes*, 384
- Jacobaea vulgaris*, tansy ragwort, 255, 257, 276
- Japan, 6, 222, 239
- Japanese beetle, *Popillia japonica*, 15, 212
- jumping spiders (Salticidae), 140, 152
- Karner blue butterfly, *Plebejus melissa samuelis*, 355
- Kennedy, J. F., 15
- Kentucky, 101
- Kenya, 49, 375
- Klamath weed, *Hypericum perforatum*, 261
- knawweed, *Centaurea* spp., 260
- Knipling, E. F., 367
- koa, 341
- Koebele, Albert, 43
- koinobionts, 168
- K-selected species, 109, 294
- kudzu, 12
- lace bugs (Tingidae), 247
- lacewings (Neuroptera), 146–147
- lady beetles (Coccinellidae), 67, 143–144, 156, 276, 329, 343, 379
- ladybird (lady beetle) fantasy, 30
- Lagenidium giganteum*, 233
- Lagerlöf, Selma, 24
- landscape design, 380
- landscape scale, 104
- landscape simplification, 104, 380
- Lantana camara*, 246, 275, 350, 353
- large blue butterfly, *Maculinea arion*, 336
- late blight, *Phytophthora infestans*, 359
- leaf beetles (Chrysomelidae), 264
- leaf wax, 99
- leafy spurge, *Euphorbia esula*, 51, 272
- Lecanicillium lecanii*, 101
- Lepidiota frenchi*, frenchi cane beetle, 340
- Lepidoptera, 58
- Leptinotarsa decemlineata*, Colorado potato beetle, 4, 19
- Letourneau, D. K., 86
- lettuce, 95
- lettuce aphid, *Nasonovia ribisnigri*, 95
- Levuana iridescens*, coconut moth, 333
- life tables, 33
- Linnaeus, 29
- Lissachatina fulica*, giant African snail, 333, 340
- Listronotus bonariensis*, Argentine stem weevil, 380
- livestock, 64, 366, 375
- Lixophaga sphenophori*, 102
- Lobularia maritima*, sweet alyssum, 95
- locusts, 237, 369
- Longitarsus aeneus*, 347

- Longitarsus echii*, 347
Longitarsus jacobaeae, 258
Lophodiplosis trifida, 253
 Losey, J. E., 155
 lucerne. *See* alfalfa
 Luckmann, W. H., 18
Lygus hesperus, western tarnished plant bug, 97, 373
Lymantria dispar, gypsy moth, 35, 36, 132, 133, 209, 233, 328, 337, 344, 348, 352, 367
Lymantria dispar asiatica, Asian gypsy moth, 28
Lymantria monacha, nun moth, 24, 218
 lynx, 228
Lysinibacillus sphaericus, 212

Maconellicoccus hirsutus, pink hibiscus mealybug, 59
 macro-beneficials, 74
Macrolophus, 145
Macrolophus pygmaeus, 156
 macroorganisms, 74
 macroorganisms, augmentation
 mass production, 75
 quality control, 75
 releasing, 76
 species/strain, 74–75
 storage and transport, 76
Maculinea arion, large blue butterfly, 336
Mahanarva fimbriolata, 236
 maize. *See* corn
 malathion, 11
 Malaysia, 104
 Marshall, D. R., 246
 Massachusetts, 14
 mathematical models, 33, 111, 115
 mating disruption, 367
 Mauritius, 137
 maximal host range, 348
 McEvoy, P. B., 259
 McFadyen, R. E. C., 63
 mealybug destroyer, *Cryptolaemus montrouzieri*, 59
 mealybugs (Pseudococcidae), 47, 59
 Mediterranean flour moth, *Ephestia kuehniella*, 172, 207
 Mediterranean fruit fly, *Ceratitis capitata*, 353
Megamelus scutellaris, 276
Megarhyssa, 179
Melaleuca quinquenervia, paperbark tea tree, 253
Melanoplus femurrubrum, red-legged grasshopper, 153
Meligethes aeneus, rape pollen beetle, 92, 101
Melinis minutiflora, molasses grass, 375
Meloidogyne spp., root-knot nematodes, 324
 melons, 321
 Mermithidae, 194
 mesquite trees, *Prosopis* spp., 63

 meta-analysis, 86
 metapopulations, 118–121
Metarhizium, 380
Metarhizium acridum, 68, 237, 369
Metarhizium anisopliae, 31
Metarhizium rileyi, 101
 Metcalf, R. L., 18
 Metchnikoff, Elie, 31
Meteorus autographae, 102
 methyl bromide, 12
 Mexican bean beetle, *Epilachna varivestis*, 71
 Mexico, 59, 92
 mice, 36
 microbial pesticides, 67
 microclimate, 35, 101
 microorganisms, augmentation
 formulation, 79
 mass production, 78–79
 quality control, 79
 releasing, 79
 species/strain, 77–78
 storage and transport, 79
 microsporidia, 239
 microtype eggs, 165
 mile-a-minute weed, *Persicaria perfoliata*, 265
 milkweed, *Asclepias* spp., 256, 381
 milky disease, 212
 Mills, N. J., 58
 Miridae (plant bugs), 156
 mites (Acari), 143
 molasses grass, *Melinis minutiflora*, 375
 mole crickets, *Scapteriscus* spp., 195
 monarch butterfly, *Danaus plexippus*, 256, 381
 monocultures, 18, 29, 66, 86, 91, 101, 322, 368
Mononychellus tanajoa, cassava green mite, 55, 157
 monophagy, 331
 Moorea, 333
Morrenia odorata, stranglervine, 281
 mosquito fish, *Gambusia*, 139, 384
 mosquitoes, 16, 194, 209, 225, 233, 384
 multicolored Asian lady beetle, *Harmonia axyridis*, 18, 329, 344
 multiple parasitism, 170
 multiple stress hypothesis, 61, 249
 Murdoch, W. W., 123
Musca domestica, house fly, 9, 362
Muscidifurax raptor, 362
 musk thistle, *Carduus nutans*, 249, 252
 mycoherbicides, 67, 279
 mycoparasitism, 300
 mycorrhizae, 306
 mycorrhizal fungi, 323
 Myers, J. H., 62, 248, 260
Myrmica sabulei, 336
 myxoma virus, 225

- myxomatosis, 225
Myzus persicae, green peach aphid, 360
- Nasonovia ribisnigri*, lettuce aphid, 95
 natural balance, 41
 natural control, 22, 24, 25, 132
 natural enemies, 22
 against invertebrates, 111
 lying in wait, 126
 search and destroy, 125
 navel orangeworm, *Amyelois transitella*, 199
 necrotrophy, 278, 291
 neem, 368
 nematoposition, 189
 nematodes, parasitic, 189–200
 augmentation biological control, 195–200
 classical biological control, 195
 ecological requirements, 190
 entomopathogenic nematodes, 190–194
 finding hosts, 192–193
 nematode-suppressive soils, 323
 nematode-trapping fungi, 323, 324
Nematophthora gynophila, 295
 neoclassical biological control, 331
Neodiprion swainei, Swaine jack pine sawfly, 174
 neonicotinoids, 12, 88
Neoseiulus cucumeris, 67, 70
Neozygites fresenii, 87
Nephotettix cincticeps, green rice leafhopper, 36
 Netherlands, 186, 369, 380
 new associations, 274, 331, 349
 New York State, 7, 360, 362
 New Zealand, 19, 28, 63, 204, 213, 377, 380
Nezara viridula, southern green stink bug, 341
 Nicaragua, 17, 93
 niche, 294
 niche markets, 80, 355
 Nicholson, A. J., 116
Nicotiana attenuata, tobacco, 138
 nicotine, 179
Nilaparvata lugens, brown planthopper, 374
 nine-spotted lady beetle, *Coccinella novemnotata*, 343
 Nixon, R. M., 15
 Nobel Prize in Medicine, 311
Nomadacris septemfasciata, red locust, 137
 nonconsumptive effect, 153
 nonoccluded invertebrate virus, 220
 nonoccluded virus, 64
 nontarget controversy
 extinction, 333–335
 history, 332–337
 indirect effects, 336
 population reduction, 333–335
 nontarget effects, 327–329
 indirect, 343–344
 prevention, 351–357
 nontarget impacts, causes for
 changing priorities, 337–339
 dispersal, 342
 fragile environments, 340–341
 insufficient information, 339–340
 nopales, 343
 North America, 233, 329, 343, 377, 382
 northern jointvetch, *Aeschynomene virginica*, 281
Nosema locustae, 240
 NPV, 220
 nucleopolyhedrovirus, 216
 nuisance, 18, 330, 361, 362
 numerical response, 112
 nun moth, *Lymantria monacha*, 24, 218
- oak trees, *Quercus* spp., 133, 248
 occlusion body, 216
 Oceania, 64
Oecophylla smaragdina, weaver ant, 28
 oil palms, 220
 oilseed rape, 92, 101
 old associations, 331
 oleander aphid, *Aphis nerii*, 141
 oligophagy, 331
 one fungus – one name, 311
 Oomycota, 233, 306, 312
Opuntia, 343
Opuntia corallicola, semaphore cactus, 343
Opuntia spp., prickly pear cacti, 252, 262, 267
 orb weavers (Araneidae), 152
 orchard, 239
 organic amendments, 322
 organic growers, 96
 organic lettuce, 95
 organophosphates, 11, 16
Orgyia thyellina, white-spotted tussock moth, 28
Orius insidiosus, insidious flower bug, 76
Orius spp., 140
Ormia ochracea, 177
 ornamentals, 372
 OrNV, 220
Orthotydeus caudatus, 99
Oryctes rhinoceros, coconut rhinoceros beetle, 64, 220
Oryctolagus cuniculus, European rabbit, 336
Ostrinia furnacalis, Asian corn borer, 82–83
Ostrinia nubilalis, European corn borer, 74, 177, 381
Oulema melanopus, cereal leaf beetle, 90
 outbreaks, 24, 28, 123, 134
 cyclic, 119
 outreach, 386
Oxyops vitiosa, 253
- Paenibacillus popilliae*, 212
 Paine, R. T., 129
 painted apple moth, *Teia anartoides*, 28

- Palmartona catoxantha*, 334
Pandora neoaphidis, 99, 203
Panonychus citri, citrus red mite, 96, 105
 paperbark tea tree, *Melaleuca quinquenervia*, 253
 Papua New Guinea, 249
 paradox of biological control, 123
 paradox of enrichment, 123
 parasite, definition, 161
 parasitic beetles, 165
 parasitic wasps, 162–163
 parasitoids, 154, 161–186
 augmentation biological control, 186
 chemical ecology, 177
 classical biological control, 183–184
 definition, 161
 host defense, 178–179
 life history strategies, 168–174
 offense, 179–181
 polydnavirus, 181
 symbiont, host, 181
 Paris green, 4
 parsnip, *Pastinaca sativa*, 256
Partula, 333
Pasteuria penetrans, 295, 323
Pastinaca sativa, parsnip, 256
 pastures, 226, 274, 283, 380
 Paterson's curse, 347
 pathogens
 chronic infections, 128
 cycles of infection, 128
 definition, 202–203
 epizootics, 127
 fatal disease, 128
 propagules, 127
 reservoirs, 128
 pathotypes, 279
 PCR-based methods, 37
 pea aphid, *Acyrtosiphon pisum*, 99, 125, 132, 155, 181
 peach orchards, 96
 peach silver mite, *Aculus fockeui*, 6
 peach trees, 6
 pears, 222, 302
 peas, 99
Pectinophora gossypiella, pink bollworm, 373
Pediobius foveolatus, 71
 Pemberton, R. W., 350
 penicillin, 311
Penicillium, 29, 294
Penicillium notatum, 311
Penicillium oxalicum, 298
 Pennsylvania, 14, 88
 Pentatomidae (stink bugs), 341
Perillus bioculatus, 145
Peromyscus leucopus, white-footed mouse, 133
Persicaria perfoliata, mile-a-minute weed, 265
 pest complex, 359
 pesticide treadmill, 6–11
 pesticide resistance, 8–11
 secondary pest outbreaks, 6–9
 target pest resurgence, 6–7
 pesticides, 4–6
 drift, 380
 mindset, 376
 phytotoxicity, 73
 resistance, 11, 157
 sublethal effects, 370
 pests, 3, 18–19
 invertebrates, 110
Phasmarhabditis hermaphrodita, 200
Phenacoccus manihoti, cassava mealybug, 55, 64, 183
 pheromones, 367
Phlebiopsis gigantea, 32, 304
Photorhabdus, 190
Phyllachora huberi, 320
 physiological host range, 348
Phytoecia coerulescens, 347
 phytophagous mites (Acari), 36, 96, 98
Phytophthora, 317
Phytophthora cinnamomi, 306
Phytophthora infestans, late blight, 359
Phytophthora palmivora, 283
 Phytoseiidae, 98
Phytoseiulus persimilis, 70, 82, 148
Picea sitchensis, Sitka spruce, 379
Pieris, 176
 pine caterpillars, *Dendrolimus* spp., 236
 pine looper moth, *Bupalus piniarius*, 177
 pine trees, *Pinus* spp., 248, 306
 pink bollworm, *Pectinophora gossypiella*, 373
 pink hibiscus mealybug, *Maconellicoccus hirsutus*, 59
Pinus spp., pine trees, 248, 306
 pistachios, 199
Pisuarina mira, 153
Planococcus kenya, coffee mealybug, 49
 plant bodyguards, 100, 132
 plant growth-promoting fungi (PGPF), 305
 plant growth-promoting rhizobacteria (PGPR), 305
 plant parasitic nematodes, biological control, 323–324
 plant pathogens, biological control
 antagonists, definition, 292
 antagonists, finding, 308–311
 antibiosis, 300
 bacterial antagonists, 312
 bioprotectants, 69
 classical biological control, 319–320
 competition, 301–304
 foliage, 319
 fungal antagonists, 311–312

- plant pathogens, biological control (*cont.*)
 indirect effects, 304–306
 oomycete antagonists, 312
 parasitism, 302
 postharvest fruit, 318–319
 seeds and roots, 317–318
 stems and crowns, 318
 use in control, 313–315
 plant stress, 249
Plebejus melissa samuelis, Karner blue butterfly, 355
 Pliny, 4
Plodia interpunctella, Indian meal moth, 211
Plutella xylostella, diamondback moth, 99, 211
Pochonia chlamydosporia, 78, 295, 300, 323
Poecilia spp., 139
Poecilostictus cothurnatus, 177
 polar filament, 239
 pollen, 101
 polyculture, 92
 polyembryonic, 170
 polyphagy, 332
Popillia japonica, Japanese beetle, 212
 population regulation, 115
 Port Jackson willow, *Acacia saligna*, 286
 post-release monitoring, 355
 potatoes, 18, 359
 Potter, M. C., 29
 powdery mildews, 98, 300, 319
 Poxviridae, 216
 praying mantids (Mantodea), 141, 149, 153, 158
 predators
 diet breadth, 152–153
 gut content analysis, 37
 interactions with natural enemies, 153
 nonconsumptive effects, 153–155
 predators, invertebrate, 156
 active hunting, 140
 ambush, 141
 augmentation biological control, 138–159
 classical biological control, 156–157
 prey defense, 141
 sit and pursue, 141
 sit and wait, 141
 zoophytophagy, 154–156
 predators, vertebrate, 139
 domestic cats, 137
 mosquitofish, *Gambusia*, 137
 mynah bird, *Acridotheres tristis*, 137
 predatory ants (Formicidae), 104
 predatory flies, 148–149
 predatory midges (in Cecidomyiidae), 148
 predatory mites (Acari), 101, 147–148
 predatory true bugs, 131, 145–146
 preventive releases, 70, 71
 prickly pear cacti, *Opuntia* spp., 252, 262, 267
 prickly pear cactus, *Opuntia vulgaris*, 32
 primary parasitoids, 171
 Proctotrupoidea, 162
 propoxur, 11
Propylaea japonica, 379
Prosopis spp., mesquite trees, 63
 Protista, 239
 prune leafhopper, *Edwardsiana prunicola*, 103
Pseudacteon, 177, 179
Pseudomonas, 321, 322
Pseudomonas fluorescens, 295, 312
Pseudomyza flocculosa, 319
 Psyllidae, 253
 public health, 344
Pueraria, 12
Purpureocillium lilacinus, 323
 push-pull, 375
 pyralids (Pylalidae), 267
 pyrethrins, 367
 pyrethroids, 6, 11, 89
Pyrilla perpusilla, sugarcane leafhopper, 97
 pyrrolizidine alkaloids, 347
Pythium, 317
Pythium oligandrum, 312

 quality control, 75, 373
Quaylea whittieri, 53
Quercus spp., oak trees, 133, 248

 rabbit hemorrhagic disease virus, 227
 rabbits, 224
 radioactive isotopes, 37
 rangeland, 13, 63, 240, 272, 283
 rape pollen beetle, *Meligethes aeneus*, 92, 101
 rats, 137
 realized host range, 348
 red imported fire ant, *Solenopsis invicta*, 151, 153
 red locust, *Nomadacris septemfasciata*, 137
 red-legged grasshopper, *Melanoplus femurrubrum*, 153
 refuges, 122–123
 refugia, 212
Regiella, 203
 registration, 77, 355
 Reoviridae, 216
 replacement strategy, 96
 reproductive parasites, 162
 resistance, 223, 226, 279
 resource concentration hypothesis, 91
Reticulotermes flavipes, eastern subterranean termite, 364
 reverse-order method, 346
Rhabdoscelis obscurus, sugarcane weevil, 102
Rhinella marina, cane toad, 137, 339
Rhinocyllus conicus, 252, 351
Rhinoncomimus latipes, 265

- rhizobacteria, 323
Rhizoctonia, 317
 rhizosphere, 296, 297, 301, 323
 competence, 302, 318
 microbiome, 321
 Rhodes grass, *Chloris gayana*, 96
Rhopalosiphum padi, bird-cherry oat aphid, 154
Rhyssa persuasoria, 180
Rhyssomatus marginatus, 248
 rice, 104, 281, 374
 Riley, Charles V., 43
 risk, quantification, 353–355
 Risley, C., 248
 r–K selection, 110
 Roberts, W., 29
 rodents, 4
Rodolia cardinalis, vedalia beetle, 30, 43, 59, 64
 romaine lettuce, 95
Romanomermis culicivorax, 194, 200
Romanomermis iyengari, 200
 Root, R. B., 91
 root and butt rot of conifers, 303
 root-knot nematodes, *Meloidogyne* spp., 324
 rove beetles (Staphylinidae), 151
 r-selected species, 109, 246, 294
 rubber, 320
Rubus spp., blackberry, 276
 Russia, 31, 139
 Russian thistle, *Salsola tragus*, 279
 rust fungi (Pucciniales), 284
 Rutz, D. A., 362
- sabadilla, *Schoenocaulon officinale*, 367
 safeguard considerations, 348
Saissetia oleae, citrus black scale, 53
Salsola tragus, Russian thistle, 279
 Salvation Jane, 347
Salvinia molesta, 246, 249
 sampling, 27, 360
 Saturniidae (silk moths), 338
 scale insects (Coccoidea), 18, 31, 47, 62, 144, 260, 267
Scapteriscus spp., mole crickets, 195
 Schlinger, E. I., 97
Schoenocaulon officinale, sabadilla, 367
Sclerotinia sclerotiorum, 298
 scouting, 360
 screwworm fly, *Cochliomyia hominivorax*, 366
 Scutelleridae (shield bugs), 341
 seasonal inoculative releases, 69, 70, 71, 185
 secondary cycling of infection, 68
 secondary pests, 88
 secondary plant compounds, 269
 seed bank, 62
 selective pesticides, 88
- semaphore cactus, *Opuntia corallicola*, 343
Serratia entomophila, 204, 213
 sesame, 104, 350
Sesbania punicea, 248, 276
 seven-spotted lady beetle, *Coccinella septempunctata*, 154, 155, 157, 343
 sex pheromones, 360
 sheep, 245, 276
 shellfish, 322
 siderophores, 304
Silent Spring, 13–15
 silkmoths (Saturniidae), 338
 silkworm, *Bombyx mori*, 29, 330
 silverleaf whitefly, *Bemisia tabaci*, 50
Sirex noctilio, 195
 Sitka spruce, *Picea sitchensis*, 379
 six-spotted mite, *Eotetranychus sexmaculatus*, 120
 Skeeter Doom, 200
 slugs, 200
 small Indian mongoose, *Herpestes javanicus*, 137
 Smith, H. S., 29
 Smith, R. F., 360
 snails, 200
 soap, 4
 Society Islands, 333
 soil fumigation, 323
Solanum viarum, tropical soda apple, 283
Solenopsis invicta, red imported fire ant, 151, 153
Solenopsis spp., fire ants, 179, 361
Solidago, 153
 solitary parasitoids, 170
 SolviNix, 283
 sorghum, 375
Sorghum vulgare sudanense, Sudan grass, 375
 South Africa, 20, 63, 248, 252, 273, 276, 286, 382
 South America, 305
 South Pacific, 220
 Southeast Asia, 220
 southern green stink bug, *Nezara viridula*, 341
 southern house mosquito, *Culex quinquefasciatus*, 212
 soybean aphid, *Aphis glycines*, 382
 soybeans, 4, 101, 221, 281
 Spain, 105
 spider mites (Tetranychidae), 6, 70, 100, 148, 185
 spiders (Araneae), 36, 93, 104, 138, 143, 152, 153
 spillover, 269, 327, 350
 spittlebugs (Cercopoidea), 236
Spodoptera exigua, armyworm, 379
Spodoptera frugiperda, fall armyworm, 93
Spodoptera littoralis, 135, 140
 spruce, 24
 spruce budworm, *Choristoneura fumiferana*, 209
 squash, 92
 Sri Lanka, 251

- stable equilibrium, 117, 123
 stable fly, *Stomoxys calcitrans*, 362
 stable habitats, 57
 stand-alone strategy, 359
 Staphylinidae (rove beetles), 151
Steinernema, 190–194
Steinernema carpocapsae, 76, 199
 Steinernematidae, 190
 Steinhaus, E. A., 207
 sterile insect technique, 366
Stethorus punctum, 88
 stimulo-deterrent diversion, 375
 Stockholm Convention on Persistent Organic Pollutants, 17
Stomoxys calcitrans, stable fly, 362
 stone fruit, 299
 strangervine, *Morrenia odorata*, 281
 strawberries, 82
Streptomyces griseoviridis, 312
Striga spp., witchweed, 375
 subsistence farmers, 48, 359
 succession, plant pathogens, 295
 Sudan grass, *Sorghum vulgare sudanense*, 375
 sugarcane, 137, 236, 340, 380
 sugarcane borer, *Diatraea saccharalis*, 237
 sugarcane weevil, *Rhabdoscelis obscurus*, 102
 suicidal germination, 375
 summit disease, 218
 superorganisms, 321
 superparasitism, 170
 suppressive soils, 32, 100, 295, 321–322
 sustainability, 368, 379–383
 Swaine jack pine sawfly, *Neodiprion swaini*, 174
 Sweden, 24
 sweet alyssum, *Lobularia maritima*, 95
 sweet orange, *Citrus sinensis*, 305
 sweetpotato whitefly, *Bemisia tabaci*, 50
 symbionts, 203
 symbiosis, 190, 196, 306
 synergism, 154
 Syrphidae (hoverflies), 95, 148
 system stability, 123
 systemic acquired resistance, 316
- Tachinidae, 164, 334, 337
 take-all, 322
 tansy ragwort, *Jacobaea vulgaris*, 255, 257, 276
Taphrocerus schaefferi, 268
 taxonomy, 50
 tea, 222
 teak, 59
Teia anartoides, painted apple moth, 28
Teleonemia scrupulosa, 247, 350
 Tennessee, 15
 Tephritidae (fruit flies), 260, 353
- Tetranychus urticae*, two-spotted spider mite, 82, 156
The Sea Around Us, 14
The Wonderful Adventure of Nils Holgersson, 24
 thistles, 351
 Thomas, M. B., 384
 thrips (Thysanoptera), 67, 70
 ticks (Ixodida), 384
 Tingidae (lace bugs), 350
 tobacco, 4
 tobacco, *Nicotiana attenuata*, 138
 tobacco budworm, *Heliothis virescens*, 176
 tobacco mild green mosaic tobamovirus, 283
 tomato, 105, 156
 tomato fruitworm or cotton bollworm, *Helicoverpa zea*, 176
 tomato pinworm, *Tuta absoluta*, 105
 top-down theory, 256
 total system approach, 368
 toxins, 190, 196, 205, 212
 transgenic crops, 366
Trialeurodes vaporariorum, greenhouse whitefly, 31, 70, 98, 185
Trichapion lativentris, 248, 276
Trichilogaster acaciaelongifoliae, 252, 273
Trichoderma, 311, 316–317
Trichoderma asperellum, 306
Trichoderma atroviride, 306
Trichogramma, 31, 74, 75, 77, 83, 166, 176, 185, 381
Trichogramma dendrolimi, 83
Trichogramma galloi, 237
Trichogramma nubilale, 177
Trichogramma ostrinae, 74, 83
 trichomes, 98
Trichoplusia ni, cabbage looper, 173, 211
Trichopoda pilipes, 341
Trichosirocalus horridus, 252
 Trinidad, 137
Trissolcus basalis, 341
 triangulins, 165
 tropical soda apple, *Solanum viarum*, 283
 tsetse fly, *Glossina austeni*, 367
Tuta absoluta, tomato pinworm, 105
 two-spotted lady beetle, *Adalia bipunctata*, 144
 two-spotted spider mite, *Tetranychus urticae*, 82, 156
 Tydeidae, 98
Typhlodromalus aripo, 157
Typhlodromus pyri, 88, 148
Tyria jacobaeae, cinnabar moth, 255, 257
- Uganda, 50, 350
 UNESCO Science Prize, 251
 United Kingdom, 93, 186, 200, 336, 379
 United Nations, 17

- United Nations Environment Programme, 386
- United States, 4, 15, 17, 18, 20, 74, 151, 159, 211, 240, 265, 272, 281, 283, 309, 328, 337, 343, 366, 367, 373, 381
- urban, 360
- Urophora affinis*, 260
- Urophora quadrifasciata*, 260
- US Congress, 15
- US Environmental Protection Agency, 15
- US Fish and Wildlife Service, 14
- US National Academy of Sciences, 23
- van Leeuwenhoek, Antonie, 29
- van Lenteren, Joop, 56, 80, 353, 371
- vedalia beetle, *Rodolia cardinalis*, 43, 59, 64
- vegetables, 13, 69, 97, 185, 372
- velvetbean caterpillar, *Anticarsia gemmatilis*, 101, 221
- vertebrates, 327
- vertical transmission, 240
- Verticillium*, 323
- Verticillium dahliae*, 306
- Vespa vulgaris*, 158
- vetch, *Vicia*, 102
- Vicia*, vetch, 102
- Vicia faba*, 132
- Vietnam, 104
- virulence, 77
- viruses, general biology, 215
- viruses, invertebrate, 215–224
 - augmentation biological control, 220–224
 - classical biological control, 219–220
 - dispersal, 218–219
 - genetic engineering, 224
- viruses, plant-pathogenic, 305
- viruses, vertebrate, 224–228
- Vitis riparia*, wild river grape, 99
- von Tubeuf, C. F., 29
- Wagner, D. L., 328
- Washington, 322
- water hyacinth, 276
- wax moth, *Galleria mellonella*, 36
- weaver ant, *Oecophylla smaragdina*, 28
- weed biological control
 - agents, 245–246
 - augmentation biological control, 275, 279–283
 - classical biological control, 270–275
 - conservation biological control, 276
 - host plants, 269
 - host range, 269–270
 - impacts, 262
 - plant pathogens, 278–287
 - plant stress, 255
 - vertebrates, 276–277
 - weed injury, 247–251
- weed strips, 102
- weevils (Curculionidae), 248, 252, 253, 267
- West Nile virus, 384
- West Virginia, 328
- western corn rootworm, *Diabrotica virgifera virgifera*, 135, 190
- western tarnished plant bug, *Lygus hesperus*, 97, 373
- western trilling cricket, *Gryllus integer*, 177
- wheat, 92, 255, 322
- wheat grain beetle, *Anisoplia austriaca*, 31
- Whisler, H. C., 233
- white amur, *Ctenopharyngodon idella*, 276
- whiteflies, 97
- whiteflies (Aleyrodidae), 50, 260
- white-footed mouse, *Peromyscus leucopus*, 133
- white-spotted tussock moth, *Orgyia thyellina*, 28
- wild insectary, 102
- wild river grape, *Vitis riparia*, 99
- wildlands, 64, 274, 378
- Wipfelkrankheit, 218
- Wisconsin, 125
- witchweed, *Striga* spp., 375
- Wolbachia*, 162, 203, 384
- wolf spiders (Lycosidae), 141, 152
- woodwasps, 179
- Working for Water, 382
- World Food Prize, 360, 367
- World Health Organization, 17, 140
- World War II, 14
- Xenorhabdus*, 190
- yeasts, 318
- yellow nutsedge, *Cyperus esculentus*, 268
- Yemen, 28
- yield increasing bacteria, 315
- Zanzibar, 367
- Zika, 384
- Zoophthora phytonomi*, 101
- zoophytophagy, 154
- Zyginidia scutellaris*, 140