

Index

ABCD (Airway, Breathing, Circulation, Disability), 328–9 <i>see also</i> resuscitation	imaging studies <i>see</i> imaging studies	STEMI <i>see</i> ST elevation	costs of morbidity/mortality, 44
abdomen	medications, 94	myocardial infarction (STEMI)	fatal, 44
acute mesenteric ischemia	misdiagnosis, 26	treatment, 26, 199–200	incidence, 44
<i>see also</i> acute mesenteric ischemia (AMI)	mortality rate and, 94	percutaneous coronary intervention (PCI), 200	inpatients, 45
aging, 208	physical examination, 94–5	pathology, medications contributing to, 208	nursing homes, 45
aortic aneurysm <i>see</i> abdominal aortic aneurysm, ruptured	pancreatitis <i>see</i> pancreatitis	(AHFS), 201–2	outcomes, 44–5
appendicitis <i>see</i> appendicitis	pathology, medications contributing to, 208	acute heart failure syndrome (AHFS), 201–2	pain management, 69
atypical disease presentation, 26	trauma, 64	acute ischemic stroke (AIS), 170–3	prevention, 48–9
biliary tract disease, 97–8	<i>see also</i> gastrointestinal system	computed tomography, 171, 173	risk factors, 45–6
bowel obstruction <i>see</i> bowel	abdominal aortic aneurysm, ruptured, 95–6	diagnosis, 170–1	unintentional exposures, causes, 46
chest pain and, 83	back pain and, 150	evaluation, 170–1	<i>see also</i> medication
diagnostic testing, 208–9	signs and symptoms, 96	headache and, 141	Affordable Care Act (2010, US) <i>see</i> financial issues
differential diagnosis, 209–16	abscess	magnetic resonance imaging, 171	aging
acute colonic pseudo-obstruction (Ogilvie's syndrome), 214	brain, 260	resuscitation, 35	aged heterogeneity (principle), 41
appendicitis <i>see</i> appendicitis	dental <i>see under</i> dental issues	risk factors and age-related, 170	categories, 8
constipation <i>see</i> constipation	epidural <i>see</i> spine, epidural abscess	treatment, 171	demographics <i>see</i> demographics
diverticulitis <i>see</i> diverticulitis	peritonsillar <i>see</i> peritonsillar abscess (PTA)	thrombolysis with rt-PA, 171	diagnostic tests and, 24
esophageal obstruction and perforation, 212	retropharyngeal <i>see</i> retropharyngeal abscess (RPA)	acute mesenteric ischemia (AMI), 96–7, 214–16	drug metabolism, 43–4
gallbladder disease	abuse, elder <i>see</i> elder mistreatment	arterial embolism, 96, 214	healthy, 20
<i>see</i> gallbladder	ACE inhibitors	arterial thrombosis, 96, 214	immunity and, 23, 208, 254
gastrointestinal bleed (upper and lower)	angioedema due to, 165–6	types, 96	physiologic changes <i>see</i> physiology of aging
<i>see</i> gastrointestinal system	indications and side effects, 165–6	venous thrombosis, 96, 214	polypharmacy <i>see</i> polypharmacy
inflammatory and infectious colitis <i>see</i> colitis	acetaminophen, 70–1	acute otitis externa (AOE), 160	population <i>see</i> population aging
ischemic colitis <i>see</i> colitis	acid–base disorders, 296–7	acute respiratory distress syndrome (ARDS), 34	risks to health, 254
mesenteric ischemia <i>see</i> acute mesenteric ischemia (AMI)	activated partial thromboplastin time (PTT), 193	acute subdural hematoma, 344	vital signs, 254
pancreatitis <i>see</i> pancreatitis	activities of daily living (ADL), 56, 314	acute thoracic aortic syndrome, 202–4	agitation, 103
peptic ulcer disease <i>see</i> peptic ulcer disease (PUD)	hospital at home, 304	atraumatic, 202–3	antipsychotic treatment, 109
small bowel obstruction, 97, 213–14	instrumental <i>see</i> instrumental activities of daily living (IADL)	identification, 203–4	non-pharmacological treatment, 108–9
diverticular disease, 99–100, 210	physical (PADLs), 56	management, 204	airway reactivity, 185
emergencies, 207–18	acute abdomen, atypical disease presentation, 26	traumatic, 202, 203	albuterol, 188
disposition (admission route), 216	Acute Care Elderly (ACE) unit/mobile care elderly unit, 308	acute vestibular syndrome (AVS), 131, 133, 135	alcohol abuse/dependence
physical examination, 207–8	acute coronary syndrome (ACS), 199–200	Addison's disease, 288	abuse, 275–6
history taking, 207	atypical presentation, 24–6	ADH secretion, 293	dependence, 275–6
infections <i>see</i> infections	chest pain, 77	Admission Avoidance Hospital at Home, 304–5	early and late onset, 276
pain, 26, 94–101, 207	dyspnea and, 89	admission route <i>see</i> disposition (admission route)	harmful and hazardous drinking, 275–6
diagnostic tests, 95	identification, 199	adrenal crisis, 288–9	risk factors, 276
differential diagnoses, 95–100	morbidity/mortality, 24	adrenal insufficiency, 288	screening and diagnosis, 276–7, 317–18
disposition (admission route), 100	NSTEMI <i>see</i> non-ST elevation myocardial infarction (NSTEMI)	advance directive (AD), 36, 329	SBIRT (Screening, Brief Intervention, and Referral to Treatment), 277
emergency department (ED)	secondary, 25	capacity and, 330	treatment, 277–8
admission, 94		DNR order, 175	Alcohol Use Disorders Identification Test (Audit-C), 277
extra-abdominal causes, 100		emergency medicine physicians and, 36	allergic conjunctivitis, 157
history taking, 94		living will, 36	altered mental status <i>see</i> mental status, altered
		power of attorney, 36	alternative care, 301–2
		ventilator discontinuation <i>see</i> palliative care	hospital at home, 304–5
		adverse drug reaction (ADR), 11, 42	<i>see also</i> Admission Avoidance Hospital at Home
		abdominal pain, 94	
		admission due to, 44	
		common drugs, 44	

Index

- Alzheimer's dementia, 279  
*see also* dementia
- analgesia  
  non-opioids, 70–1  
  nonsteroidal anti-inflammatory drugs (NSAIDs), 71  
  opioids, 71–2  
  options, 70  
  regional anesthesia, 72–3  
  *see also* pain, management
- anemia, 266–7
- anesthesia, regional, 72–3
- aneurysm, abdominal aortic  
  *see* abdominal aortic aneurysm, ruptured
- aneurysmal subarachnoid hemorrhage (aSAH), 140, 175–6  
  diagnosis and investigations, 175  
  treatment, 175–6  
  *see also* subarachnoid hemorrhage (SAH)
- angioedema, due to ACE inhibitors, 165–6
- angiography, abdomen, 209
- angiotensin-converting enzyme inhibitors *see* ACE inhibitors
- angiotensin receptor blockers (ARBs), 166
- anion gap acidosis, 297
- anterior inferior cerebellar artery (AICA) strokes, 131
- antibiotics  
  chronic obstructive pulmonary disease (COPD), 188  
  community-acquired bacterial meningitis, 178  
  conjunctivitis, 157  
  elderly and, 261  
  Ludwig's angina, 163  
  male genital system infections, 228–9  
  peritonsillar abscess (PTA), 164  
  pneumonia, 91, 256–7  
  urinary tract infections, 233, 257
- anticholinergic medication, 46
- asthma and COPD, 188
- anticoagulants  
  dysrhythmias, 200  
  epistaxis and, 162  
  pulmonary embolism, 193  
  trauma and, 61–2
- antidepressants, 275
- tricyclic (TCAs), 46
- antihistamines, 46
- antiplatelet agents, 61–2
- antipsychotics, 46  
  agitation treatment, 109  
  delirium treatment, 109
- aorta  
  acute thoracic aortic syndromes  
    *see* acute thoracic aortic syndrome  
  penetrating atherosclerotic ulcer (PAU), 203
- aortic aneurysm, 81–2  
  back pain and, 150  
  ruptured abdominal, 95–6  
  signs and symptoms, 96
- aortic dissection (AD), 80–1, 203  
  back pain and, 150  
  chest pain and, 80  
  classification, 203  
  diagnosis and CT, 150  
  diagnostic difficulties, 80  
  prevalence, 150
- surgical repair, 81  
  thoracic, 203
- aorto-enteric fistula (AEF), 96
- appendicitis, 99, 209–10, 258  
  atypical presentation, 26  
  MANTRELS scoring system, 99  
  symptoms, 209  
  treatment, 210
- arrhythmias *see* dysrhythmias
- arthritis  
  osteoarthritis *see* osteoarthritis  
  rheumatoid *see* rheumatoid arthritis  
  septic *see* septic arthritis
- aspiration pneumonia, 190–1
- assessment in ED *see* emergency department (ED)
- asthma, 187  
  dyspnea and, 90, 187  
  emergencies, 187  
  exacerbation, 90  
  presentation, 187–8  
  treatment, 188
- atrial fibrillation, 200
- ATTEST (dizziness timing and triggers paradigm), 129–31, 132
- atypical disease presentation, 24–6
- back pain, 146–52  
  aging and, 146  
  characteristics, 146  
  diagnostic tests, 147–8  
  differential diagnoses, 148–50  
  disposition (admission route), 151  
  emergency department (ED) presentation, 146  
  history taking, 146  
  mechanical causes, 148–9  
  degenerative disk disease, 148  
  disk displacement causing sciatica, 148–9  
  lumbar spine stenosis, 148  
  unstable lumbar spine, 148  
  osteoporotic causes, 149  
  vertebral compression fractures, 149  
  physical examination, 147  
  spinal stenosis *see* spinal stenosis  
  systemic causes, 149–50  
    abdominal aortic aneurysm (AAA), 150  
    aortic dissection (AD), 150  
    cancer, primary or metastatic, 149  
    spinal infections, 149–50  
    visceral diseases unrelated to spine, 150  
  treatment, 150–1
- bacterial meningitis *see* meningitis, bacterial
- bacteriuria, 257
- balance, screening for common conditions, 318
- balanitis, 225–6
- Beers Criteria, 46, 69–70, 71
- analgesic risks, 69–70  
  delirium, 104
- behavioral/psychological symptoms  
  in dementia syndrome (BPSD), 278, 280
- benign paroxysmal positional vertigo (BPPV), 129, 132  
  treatment, 135
- benign prostatic hyperplasia (BPH), 224, 225
- benzodiazepines, 46–8  
  delirium treatment, 109
- beta-adrenergic agonists, 188
- bicarbonate, diabetes, 285
- biliary tract disease, 97–8
- bladder  
  decompression, 224  
  *see also* genitourinary emergencies
- bleeding  
  intracerebral hemorrhage, 174  
  nose (epistaxis) *see* epistaxis  
  vaginal, 222
- blood gas, 187
- blood pressure, 88  
  ambulatory recording in syncope, 120  
  control in aortic aneurysm, 81  
  elevated *see* hypertension  
  intracerebral hemorrhage, 174
- blood transfusion, 267–8
- blood viscosity, 266
- body composition, aging, 21, 43
- Boerhaave's syndrome, 212
- Boston Syncope Criteria, 121
- bowel  
  large, inflammation *see* colitis  
  mesenteric ischemia *see* acute mesenteric ischemia (AMI)
- obstruction, 97  
  large bowel, 97, 214  
  small bowel, 97, 213–14  
  peptic ulcer disease, 97  
  radiography, 214
- brain abscess, 260
- brain injury  
  aneurysmal subarachnoid hemorrhage *see* aneurysmal subarachnoid hemorrhage  
  non-traumatic intracranial hemorrhage, 173–6  
  *see also* intracerebral hemorrhage  
  traumatic (TBI), 176–8  
  epidemiology, 176  
  etiology, 176  
  imaging studies, 176  
  outcomes and treatment, 176–8
- brain tumor, headache and, 141–2
- branch retinal artery occlusion (BRAO), 158
- branch retinal vein occlusion (BRVO), 159
- Brief Alzheimer's Screen (BAS), 106
- BRIGHT (Brief Risk Identification for Geriatric Health Tool), 315
- bronchiectasis, 189–90  
  *see also* chronic obstructive pulmonary disease
- B-type natriuretic peptide (BNP), 187
- burns, 60
- CAGE questionnaire, 277
- calcitonin, 294, 295
- calcium  
  hyperkalemia management, 291  
  normal serum levels, 294
- calcium disorders, 294–6  
  hypercalcemia *see* hypercalcemia  
  hypocalcemia *see* hypocalcemia
- cancer  
  aging and, 263  
  back pain and, 149  
  chest pain and, 82
- hypercalcemia due to, 295  
  lungs, 196–7  
  management and treatment, 263  
  spine, 149  
  *see also* hematologic and oncologic emergencies
- Candida*, 222  
  balanitis, 225
- capacity, decision-making  
  *see* decision-making capacity assessment
- carbon dioxide, end-tidal, 187
- carbon monoxide exposure, 140
- cardiac arrhythmias  
  *see* dysrhythmias
- cardiac catheterization, 26, 120
- cardiac output, 88
- cardiopulmonary resuscitation (CPR)  
  outcomes, 31–2, 323–6  
  prognosis in advanced illness, 323–6  
  *see also* resuscitation
- cardiovascular system  
  aging, 21, 60, 87–8, 204  
  physiological changes, 88  
  structural changes, 88  
  vascular changes, 88
- chest pain *see* chest pain
- congestive heart failure  
  *see* congestive heart failure (CHF)
- emergencies, 199–206  
  acute coronary syndrome  
    *see* acute coronary syndrome (ACS)  
  acute heart failure syndromes  
    *see* acute heart failure syndrome (AHFS)  
  acute thoracic aortic syndrome  
    *see* acute thoracic aortic syndrome  
  dysrhythmias  
    *see* dysrhythmias  
  syncope *see* syncope
- cardioversion, 201
- carotid pulse, 33
- carotid sinus hypersensitivity (CSH)  
  carotid sinus massage and, 119  
  syncope and, 116–17, 119
- carotid sinus massage, syncope and, 119, 120
- case management, 338  
  discharge planning *see* discharge in ED, 337  
  efficacy, 339  
  models, 337  
  monitoring, 338  
  staff team, 337–8
- catheters, indwelling, urinary tract  
  infections, 108, 257–8
- cauda equina syndrome, 149, 243
- central cord syndrome, 343
- central nervous system (CNS)  
  aging, 44, 61  
  infection, 178–9, 259–60  
  bacterial meningitis  
    *see* meningitis  
  brain abscesses, 260  
  headache and, 142  
  microorganisms  
    implicated, 259  
  spinal epidural abscess  
    *see* spine  
  symptoms, 259  
  trauma, 63
- central positional vertigo, 132

- central retinal artery occlusion (CRAO), 157–8
- central retinal vein occlusion (CRVO), 158–9
- central venous pressure (CVP), 33–4
- cerebral herniation, signs, 177
- cerebrospinal fluid (CSF) leak, 289
- cerebrovascular emergencies, headache and, 140–3
- chest pain, 3, 77–85
  - cardiac causes, 77–9
    - acute coronary syndrome *see* acute coronary syndrome (ACS)
    - congestive heart failure, 78
    - demand ischemia, 78
    - non-ST elevation myocardial infarction (NSTEMI), 78
    - pericarditis, 78–9
    - post MI complications, 78
    - ST elevation myocardial infarction (STEMI), 77–8
  - chest wall pathology, 82
  - esophageal obstruction and perforation, 212
  - gastrointestinal causes, 82–3
  - pneumothorax, 80
  - pulmonary causes, 79–80
  - vascular causes, 80–2
- chest wall, compliance loss, 185
- chest X-ray *see* radiographs, chest
- cholecystitis, acute, 97–8, 258–9
- chronic bacterial prostatitis, 227
- chronic obstructive pulmonary disease (COPD)
  - bronchiectasis *see* bronchiectasis
  - dyspnea and, 89–90, 187
  - emergency treatment, 33, 90
  - management, 189
  - presentation, 187–8
  - pulmonary function and, 186
  - treatment, 188–9
- circulation, resuscitation and, 33–4
- claudication, neurogenic, 148
- clinical decision rules (CDR), syncope, 120–1
- Clostridium difficile*
  - diagnostic tests, 211
  - diarrhea, 259
  - see also* colitis
- coagulopathy, 266
  - hyperviscosity syndrome, 266–8
- Cockcroft–Gault formula, 43
- cognitive impairment
  - in altered mental status *see* mental status, altered
  - assessment, 105–6
  - chronic, 104–5
  - dementia, 104–5
  - discharge and, 28
  - ED, implications for, 3
  - physical examination, 27
  - quality indicators, 308
  - suicide and, 271
- colitis, 211
  - Crohn’s disease, 211
  - infectious and inflammatory, 211
  - ischemic, 216
  - ulcerative, 211
- colonic pseudo-obstruction, acute (Ogilvie’s syndrome), 214
- coma, 102
  - diagnostic testing, 108
  - Glasgow Coma Scale, 176
  - myxedema *see* myxedema coma
  - see also* consciousness
- community-acquired bacterial meningitis, symptoms, 178
- community-acquired pneumonia (CAP), 79, 91, 190
  - diagnostic tests, 256
  - ED scoring, 256–7
  - etiology, 190
  - microorganisms implicated, 256
  - symptoms, 256
  - treatment, 191, 256–7
- community-dwelling elders, urinary tract infections, 257
- community resource
  - coordination, 15–16
- complex patients, 2–3
- comprehensive geriatric assessment (CGA), 313
- computed tomography (CT)
  - abdomen, 209
  - acute ischemic stroke, 171, 173
  - back pain and, 148
  - epilepsy, 35
  - intracerebral hemorrhage (ICH), 174
  - non-contrast, aneurysmal subarachnoid hemorrhage, 175
  - pulmonary angiogram (CTPA), pulmonary embolism, 79, 192–3
  - pulmonary diseases, 187
  - subarachnoid hemorrhage (SAH), 141, 175, 177
  - syncope, 120
  - traumatic brain injury (TBI), 176
- congestive heart failure (CHF), 78, 88–9
  - chest pain, 78
  - myocardial infarction complication, 78
  - prevalence, 78
  - types, 88
  - see also* acute heart failure syndrome (AHFS)
- conjunctivitis, 156–7
- consciousness
  - assessment, 105
  - disturbances in, 102–4
    - coma and stupor, 102, 108
    - delirium *see* delirium
  - headache and, 140
  - loss of *see* syncope
  - subarachnoid hemorrhage (SAH) and, 141
- consent, capacity to, 110
- constipation, 213
- continence, 4
- cost of care *see* financial issues
- costochondritis, 82
- C-reactive protein, 55
- creatine kinase (CK), elevated, rhabdomyolysis, 287
- creatinine, 43, 219
- Crohn’s disease, 211
- crossed straight leg test, 147
- CURB-65 (severity score), 256–7
- cytochrome P450 system, 43
- D-dimer, 79–80, 86, 89, 192
- death, causes of, adverse drug reaction (ADR), 44
- decision-making capacity
  - assessment
  - advance directive (AD) and, 330
  - altered mental status and, 110
  - see also* consent
- CURVES, 330
  - goals of care, 330–1
  - palliative care and, 330
  - surrogates, 110, 330
- dehydration, 23, 32, 133, 265
- delirium, 102–4
  - assessment, 107
  - epidemiology and outcomes, 103
  - geriatric psychoses, 278, 279
  - hyperactive, 102
  - hypoactive, 102
  - medications and, 103–4
  - non-pharmacological treatment, 108–9
  - palliative care, 329–30
  - pharmacological treatment, 109
  - risk factors, 103–4, 279
    - dementia, 103
  - subtypes, 102
- demand ischemia, chest pain, 78
- dementia
  - behavioral disturbances, etiology, 279
  - cognition, chronic disturbances in, 104–5
  - delirium and, 103
  - depression and, 273
  - geriatric psychoses, 278, 279
  - with Lewy bodies (DLB), 104
  - secondary/reversible, 104
  - treatment, non-pharmacological, 280
- demographics
  - elderly patients, 1, 2, 86
  - emergency medicine (EM) (for elders), 4–5
  - shift in *see* population aging
- dental issues, 162–3
- depression, 273–5
  - antidepressants *see* antidepressants
  - diagnosis, 274
  - DSM IV-TR criteria, 274
  - emergency department (ED) screening, 274
  - epidemiology, 273
  - headache with, 143
  - in older adults, 23, 274
  - poor recognition of, 274
  - risk factors, 273, 274
  - treatment, 274–5
  - design, of geriatric emergency department, 10–11
- diabetes mellitus
  - bicarbonate, 285
  - gastrointestinal conditions and, 208
  - insulin therapy, 285
  - potassium, 285
  - proliferative diabetic retinopathy, 155
- diabetic ketoacidosis (DKA), 284–5
- diabetic retinopathy, 155
- Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV)
  - alcohol abuse, 275
  - alcohol dependence, 275
  - delirium, 107
  - depression, 274
- diagnostic tests/testing, 25
  - abdominal emergencies, 208–9
  - abdominal pain, 95
  - abnormal laboratory parameters, 25
  - acute kidney injury, 221
  - age and, 24
  - altered mental status, 107–8
  - back pain *see* back pain
  - bacterial meningitis, 259
  - balanitis, 225
  - CNS infections, 259
  - coagulopathy, 266
  - coma, 108
  - community-acquired pneumonia (CAP), 256
  - diabetic ketoacidosis (DKA), 285
  - dizziness, 134–5
  - fever and leukopenia, 264
  - gastrointestinal system, 208–9
  - generalized weakness and fatigue, 52
  - gout, 239
  - hypercalcemia, 295
  - hyperosmolar hyperglycemic state (HHS), 285
  - hyponatremia, 293
  - myxedema coma, 286
  - pelvic fractures, 247
  - pituitary apoplexy, 289–90
  - pneumonia, 256
  - prescribing cascade, 41
  - pulmonary embolism, 89, 192–3
  - pulmonary emergencies, 186–7
  - rhabdomyolysis, 288
  - septic arthritis, 238
  - stupor, 108
  - tuberculosis, 194–5
  - urinary tract infections (UTI), 232–3
  - vertebral fractures, 248
- diarrhea, 208, 286
  - Clostridium difficile*-related diarrhea, 259
  - hyperthyroidism, 298
  - large bowel obstruction, 97, 214
  - watery, 211
- dipyridamole, 48
- discharge (from hospital), 341
  - Acute Care Elderly (ACE) Unit/Mobile Care Elderly Unit, 308
  - cognitive impairment, 28
  - DEED study, 306
  - from ED, 307–10
  - follow-up, 14
  - functional decline, 13–15
  - planning, 28, 338–9
  - transitional care, 307–8
    - Geriatric Floating Interdisciplinary Transition Team (Geri-FITT), 308
    - key elements, 307
- Discharge of the Elderly from the Emergency Department (DEED) study, 306
- disease-modifying anti-rheumatic drugs (DMARDs), 240–1
- disk *see* intervertebral disk
- disposition (admission route)
  - abdominal pain, 100
  - altered mental status, 110
  - back pain, 151
  - diabetic ketoacidosis (DKA), 285
  - distal radius fractures, 251
  - dizziness, 135–6
  - gastrointestinal emergencies, 216
  - gout, 239
  - hematuria, 230
  - hypernatremia, 294
  - hyperosmolar hyperglycemic state (HHS), 285
  - pelvic fractures, 248

## Index

- disposition (admission route) (*cont.*)  
 septic arthritis, 238  
 shoulder dislocations, 250  
 trauma, 64–5  
 urinary tract infection (UTI), 234
- disseminated intravascular  
 coagulation, 267
- distal radius, fractures *see* radius,  
 distal
- diverticular disease, 99–100, 210
- diverticulitis, 210–11, 258  
 diagnosis, 210, 259  
 treatment, 210–11
- Dix–Hallpike test, 134
- dizziness, 128–38  
 acute vestibular syndrome (AVS),  
 131, 133, 135  
 aging and, 128  
 benign paroxysmal positional  
 vertigo *see* benign  
 paroxysmal positional  
 vertigo (BPPV)  
 central positional vertigo, 132  
 diagnostic strategy, 129–31  
 describing difficulties, 129  
 differential diagnosis, 131–2  
 symptom quality, 129  
 timing and triggers paradigm  
 (ATTEST), 129–31  
 diagnostic tests, 134–5  
 disposition (admission  
 route), 135–6  
 episodic vestibular syndrome  
 (EVS), 131–2  
 exacerbated, 132  
 falls, 128–9  
 history taking, 132–3  
 medication issues, 128  
 nystagmus, 133–4  
 orthostatic hypotension, 132  
 physical examination, 133–4  
 gait, 133  
 head impulse test (HIT),  
 133, 134  
 tympanic membrane, 133  
 positional vestibular syndrome  
 (PVS), 132  
 safety of patient, 135  
 skew deviation, 134  
 transient ischemic attack  
 (TIA), 132  
 treatment, 135  
 triggered, 132  
 vestibular physiology, 128
- DNR (Do Not Resuscitate) orders,  
 intracerebral hemorrhage  
 (ICH), 175
- Dressler's syndrome, 78
- driving  
 conditions affecting, 339–40  
 older driver safety, 339–41  
 screening tools, 340  
 syncope and, 124  
*see also* motor vehicle collisions
- drug  
 absorption and binding, 43  
 clearance, 43–4, 69  
 metabolism, 43–4, 69  
 liver, 43–4  
 drug-related problems *see* adverse  
 drug reaction (ADR)
- DSM *see* *Diagnostic and Statistical  
 Manual of Mental Disorders*  
 (DSM-IV)
- dyspnea, 3, 86–93  
 acute coronary syndrome, 89
- asthma and, 90, 187
- cardiovascular changes causing  
*see* cardiovascular system
- chronic obstructive pulmonary  
 disease, 89–90, 187
- congestive heart failure, 88–9  
 etiology, 86, 91  
 evaluation and management, 86  
 history taking, 186  
 palliative care, 329  
 patient assessment, 86  
 pneumonia and, 90–1  
*see also* pneumonia
- pulmonary embolism, 89
- pulmonary hypertension, 195
- respiratory system *see* respiratory  
 system
- dysrhythmias, 200–1
- ears, 160–1  
*see also* hearing impairment
- echocardiography, syncope, 120
- education, 5  
 geriatric emergency department  
 (ED), 13  
 curriculum, 13
- elder mistreatment, 60, 355–9  
 definition, 355–6  
 documentation, 357  
 ED and, 358  
 epidemiology, 356  
 legal/ethical considerations, 357–8  
 preventive services, 358  
 risk factors, 356–7  
 screening, 357, 358  
 social history/historical  
 aspects, 355  
 suspicious presentations, 357–8
- electrocardiography (ECG), 21
- arrhythmic syncope, 202
- chest pain, 89
- hyperkalemia, 290
- NSTEMI, 25
- pulmonary embolism, 89, 192
- syncope, cardiovascular  
 testing, 119
- transient ischemic attack  
 (TIA), 173
- electroencephalography (EEG), 108
- syncope, 120
- electrolyte abnormalities *see specific  
 electrolytes*
- electrophysiologic (EPS) testing,  
 syncope, 120
- emergency department (ED)  
 Affordable Care Act (2010, US), 353
- alcohol abuse screening, 317–18
- assessment, 27–8  
 history taking, 27, 42
- blood transfusions *see* blood  
 transfusion
- cardioversion, 201
- case management *see* case  
 management
- clinical outcomes of elderly  
 patients, 9
- common presentations, 3, 303
- current status, 8
- demographics, 2
- discharge *see* discharge
- Do Not Resuscitate (DNR)  
 orders, 175
- drivers (older), safety of  
*see* driving
- elderly patients and, 1, 8,  
 307, 336–7
- end-of-life care, 322–3  
*see also* palliative care
- environment, 4
- financial issues *see* financial issues
- follow-up care, 14, 15
- functional assessment  
*see* functional assessment
- geriatric *see* geriatric emergency  
 department (ED)
- Glasgow Coma Scale, 176
- last hours of hospice  
 patients, 332–3
- malnutrition screening, 317
- older adults use of, 301
- pain care quality indicators,  
 72, 309
- palliative care in, 5, 17
- primary care providers (PCPs)  
 and, 5
- quality indicators *see* quality
- readmission, 336
- reducing demand, 302–7  
 Admission Avoidance Hospital  
 at Home *see* Admission  
 Avoidance Hospital at Home
- alternative care  
*see* alternative care
- costs of outpatient visits, 304
- DEED study, 306
- financial implications, 351
- immediate readmissions, 305
- off-hours care, access to, 303
- preventing hospital returns  
*see* hospital-based care
- primary care, access to, 303
- telemedicine *see* telemedicine
- see also* geriatric emergency  
 department (ED)
- emergency medical system  
 (EMS), 26–7
- emergency medicine (EM)  
 (geriatric), 1, 9, 20–30
- barriers to, 8–9
- care model, 26–8
- clinical challenge, 2–5  
 complex patients *see* complex  
 patients
- ED environment, 4
- expensive care, 3
- specific demographic  
 groups, 4–5
- context, 1
- current status, 8
- Do Not Resuscitate (DNR)  
 orders, 175
- ED assessment *see* ED
- education *see* education
- in emergency departments  
 (ED), 2
- expenditure *see* expenditure
- Glasgow Coma Scale, 176
- goal of emergency care program, 9
- overview, 1–7
- principles, 20, 21  
*see also* individual conditions
- emergency medicine physicians,  
 advance directive (AD)  
 and, 36
- Emergency Severity Index (ESI), 13
- emotional/psychological  
 abuse, 356
- see also* elder mistreatment
- EMTALA (Emergency Medical  
 Treatment and Active Labor  
 Act), 351
- endocrine system
- aging, 23, 61
- emergencies *see* metabolic and  
 endocrine emergencies
- end-of-life care *see* palliative care
- ENT emergencies; *see* ears; nose;  
 throat
- environment, of geriatric emergency  
 department, 10–11
- epididymitis, 227
- epidural abscess, spinal *see* spine,  
 epidural abscess
- epiglottitis, 164–5
- epilepsy, 35–6  
*see also* seizures
- status epilepticus (SE)
- episodic vestibular syndrome  
 (EVS), 131–2
- epistaxis, 161–2
- erythrocyte sedimentation rate  
 (ESR), giant cell arteritis, 142
- esophagitis, upper gastrointestinal  
 bleed (UGIB) and, 213
- esophagus  
 obstruction and perforation, 212  
 rupture, chest pain and, 82–3
- ethical considerations, elder  
 mistreatment, 357–8
- ethics, 36–7  
 advance directive (AD)  
*see* advance directive (AD)
- end-of-life care, 36
- medical futility, 36–7
- resuscitation, 36
- euvolemic hyponatremia, 292
- expenditure  
 emergency medicine (EM) (for  
 elders), 1–2, 3  
 health care, 336, 351  
*see also* financial issues
- eye emergencies, 153–60  
 acute angle closure glaucoma  
*see* glaucoma, acute
- branch retinal artery occlusion  
 (BRAO), 158
- branch retinal vein occlusion  
 (BRVO), 159
- central retinal artery occlusion  
*see* central retinal artery  
 occlusion (CRAO)
- central retinal vein occlusion  
 (CRVO) *see* central retinal  
 vein occlusion (CRVO)
- conjunctivitis *see* conjunctivitis
- giant cell arteritis *see* giant cell  
 arteritis (GCA)
- glaucoma *see* glaucoma
- retinal detachment *see* retinal  
 detachment
- uveitis *see* uveitis
- vitreous hemorrhage *see* vitreous  
 hemorrhage
- failure to thrive, 52–3
- falls, 54–5, 59, 244, 343–50  
 assessment, 11  
 chest trauma due to, 82  
 discharge after, 65  
 dizziness, 128–9  
 ED role in treatment and  
 prevention, 344, 347–8
- epidemiology, 343
- fear of falling, 54–5
- fractures, 343
- future directions/research, 347–8
- geriatric emergency department  
 (ED), 11



- head injuries, 344  
hip fracture *see* hip injuries, 343  
morbidity and mortality, 343–4  
opioid-associated, 71–2  
orthopedic injuries  
    *see* rheumatologic and orthopedic emergencies  
pelvis fractures *see* pelvic fractures  
prevention, 344–7  
    intervention, 347  
    PROFET (Prevention of Falls in the Elderly Trial), 344–7  
recommendations, 347–8  
recurrent, 65  
risk factors, 54, 59, 344, 345, 346  
    multiple falls, 346  
screening for common conditions, 318
- families, 4  
    as health system, 2  
    home care *see* home health care  
    preparation for ventilator discontinuation, 332  
    resuscitation witnessed by, 37–8, 326–7  
    support for, 37
- fatigue, generalized weakness and  
    *see* generalized weakness
- females *see* gynecologic emergencies
- fever, 254  
    differential diagnosis, 255  
    headache and, 139  
    leukopenia and *see* leukopenia  
    *see also* infections
- financial issues, 351–4  
    Affordable Care Act (2010, US), 352–3  
    implications, 353  
    costs of outpatient visits, 304  
    emergency department (ED), 351–2  
    reducing demand for ED use, 351  
    emergency medicine (EM) (for elders), 1–3  
    cost of care (expensive), 3  
    EMTALA (Emergency Medical Treatment and Active Labor Act), 351  
    financial abuse, 356  
    health care expenditure, 336, 351  
    Medicare, 352  
    *see also* expenditure
- fistula  
    aorto-enteric (AEF), 96  
    vaginal discharge, 222
- fluid resuscitation, 33, 288  
follow-up care, 14, 15
- forced expiratory volume in one second (FEV<sub>1</sub>), 87, 185
- forced vital capacity (FVC), 87, 196
- Fournier's gangrene, 231
- fractures  
    chest pain and, 82  
    distal radius *see* radius, distal  
    falls, 244, 343  
    *see also* falls  
    hip fracture *see* hip orthopedic injuries  
    orthopedic injuries  
        *see* rheumatologic and orthopedic emergencies  
    pelvis *see* pelvic fractures  
    periprosthetic, 64  
    sacral, 149  
    trauma evaluation, 64
- vertebral *see* vertebral fractures
- frailty, 55–6  
Frank–Starling mechanism, 88  
functional assessment, 313–21  
    assessment tools, 314  
    activities of daily living (ADL) *see* activities of daily living (ADL)  
    instrumental ADL  
        *see* instrumental activities of daily living (IADL)  
    performance testing of function *see* performance testing of function  
    comprehensive geriatric assessment (CGA), 313  
    in ED, 318  
    elements, 313  
    post-ED functional decline  
        *see* functional decline  
    screening for common conditions, 316–18  
        alcohol abuse, 317–18  
        balance, gait, and falls, 318  
        hearing impairment *see* hearing impairment  
        malnutrition, 317  
        visual impairment *see* vision  
    trajectories of illness and prognosis, 322
- functional decline, 56–7, 313  
    activities of daily living  
        *see* activities of daily living (ADL)  
    after ED visit, 56  
        two-step geriatric program, 13–15  
    prediction tools (post ED), 57, 314–16  
        BRIGHT (Brief Risk Identification for Geriatric Health Tool), 315  
        HARP (The Hospital Admission Risk Profile), 315  
        ISAR (Identification of Seniors At Risk), 314–15  
        SHERPA (Score Hospitalier d'Evaluation du Risque de Perte d'Autonomie), 315–16  
        SIGNET (The Systematic Intervention for a Geriatric Network of Evaluation and Treatment), 315  
        TRST (The Triage Risk Screening Tool), 315  
    *see also* functional assessment
- gait  
    assessment, 27  
    dizziness and, 133  
    fall risk factor, 54, 314  
    normal pressure hydrocephalus, 104  
    screening for common conditions, 318  
    slow, 52, 314  
    speed and sarcopenia, 52, 53  
    spinal stenosis, 243
- gallbladder  
    acute cholecystitis, 97–8, 258–9  
    disease, 209, 211  
gangrene *see* Fournier's gangrene
- gastrointestinal system  
    aging, 22–3, 61  
    chest pain and *see* chest pain (LGIB), 213  
    peptic ulcer disease (PUD)  
        *see* peptic ulcer disease (PUD)  
    upper gastrointestinal bleed (UGIB), 212–13  
    *see also* abdomen  
    generalized weakness, 51–8  
    diagnoses/causes, 51, 52  
    failure to thrive *see* failure to thrive  
    falls *see* falls  
    fatigue and, 51–2  
    frailty *see* frailty  
    functional decline *see* functional decline  
    geriatric syndromes, 51–4  
    prevalence, 51  
    rehabilitation, 51  
    sarcopenia *see* sarcopenia
- genital system (female)  
    *see* gynecologic emergencies
- genital system (male)  
    aging, 220  
    balanitis *see* balanitis  
    epididymitis, 227  
    Fournier's gangrene *see* Fournier's gangrene  
    infections, 226–9  
        antibiotic use, 228–9  
    orchitis, 226–7  
    pain, 226  
    phimosis/paraphimosis, 230–1  
    physical examination, 220  
    prostatitis *see* prostate  
    *see also* genitourinary emergencies
- genitourinary emergencies, 219–36  
    acute kidney injury *see* kidney  
    aging  
        genital systems, 219–20  
        renal system, 219
- geriatric emergency department (ED), 2, 4, 10–17, 306–7  
    components, 10  
    discharge and unplanned readmission, 3  
    follow-up, 14  
    key considerations, 10, 306  
        community resource coordination, 15–16  
        design, 10–11  
        falls and falls assessment, 11  
        hospital resource coordination, 15  
        medications, 11–13  
        noise reduction, 11  
        observation *see* observation  
        operations *see* triage  
        palliative care *see* palliative care  
        patient satisfaction, 16  
        physical environment/equipment, 10–11  
        quality initiatives, 11  
        staff education *see* education  
        staffing, 16  
    quality, 11, 12  
        falls and fall assessment  
            *see* falls  
        medication *see* medication  
    reasons for specific provision, 9–10  
    triage, 14  
    visit number, 10
- see also* emergency department (ED)
- geriatric emergency medicine  
    *see* emergency medicine (EM) (geriatric)
- Geriatric Floating Interdisciplinary Transition Team (Geri-FITT), 308
- geriatric psychoses, 278–80  
    delirium, 278, 279  
    dementia, 278, 279  
    epidemiology, 278  
    risk factors, 278–9  
    treatment, 279–80  
    *see also* psychiatric emergencies
- geriatric syndromes, 52, 53–4
- giant cell arteritis (GCA), 159–60, 242  
    clinical features, 140, 142, 159  
    diagnosis, 142, 159, 242  
    headache and, 140, 142  
    history taking, 242  
    treatment, 142, 159–60, 242
- Glasgow Coma Scale, 176
- glaucoma  
    acute angle closure (AACG), 153–4  
    emergency medicine (EM) (geriatric), 143  
    headache and, 142–3  
    glomerular filtration rate (GFR), 23, 32, 43, 80, 292  
    glucocorticoids, 288  
        asthma and COPD, 188  
    gout, 238–9  
    gynecologic emergencies, 219–36  
        aging, 219  
        Fournier's gangrene *see* Fournier's gangrene  
        physical examination, 220  
        uterine prolapse, 222–3  
        *see also* uterine prolapse  
        vaginal discharge *see* vaginal discharge
- haloperidol, 109
- HARP (The Hospital Admission Risk Profile), 315
- head impulse test (HIT), 133, 134
- headache, 139–45  
    acute angle closure glaucoma (AACG) and, 140, 153  
    cluster, 143  
    depression and, 143  
    emergency department presentations, 139, 144  
    etiology, 139, 140  
    subarachnoid hemorrhage (SAH), 139, 140  
    high-risk headache features, 139–40  
        new-onset, 139  
        “thunderclap” headache, 139  
    visual disturbances, 140  
    high-risk patients, 139–40  
    hypnic, 143  
    prevalence, 139  
    primary headache disorders, 143  
    secondary headache disorders, 140  
        acute glaucoma, 142–3  
        brain tumor, 141–2  
        central nervous system infection, 142  
        cervicogenic, 143  
        giant cell arteritis, 140, 142

Index

- headache (*cont.*)  
  intracerebral hemorrhage (ICH), 141  
  ischemic stroke, 141  
  meningitis, 142  
  non-emergent secondary headache, 143  
  subarachnoid hemorrhage (SAH), 139, 140  
  subdural hemorrhage, 141  
  trigeminal neuralgia, 143  
  tension-type, 143  
health care expenditure, 336, 351  
  emergency medicine (for elders), 1–3  
health care proxy, 4  
health system, 2  
health care-associated pneumonia, 190–1  
hearing impairment, 316–17  
heart  
  aging, 88  
  *see also* cardiovascular system  
  failure  
    acute *see* acute heart failure syndrome (AHFS)  
    congestive *see* congestive heart failure  
hematologic and oncologic emergencies, 263–9  
  aging and hematology, 23  
  anemia, 266–7  
  coagulopathy *see* coagulopathy  
  hemologic diseases, 263  
  hyperviscosity syndrome, 266–8  
  oncologic diseases, 263  
  specific disease conditions, 263–5  
  transfusion of blood, 267–8  
hematuria, 229–30  
hemoptysis, 194–5  
herbal medications, 42  
herpes simplex meningoencephalitis (HSME), 259–60  
herpes zoster, 260  
  chest pain, 82  
hip  
  dislocation, 245–6  
  fracture, 244–5  
    occult, after trauma, 64  
    periprosthetic, 64  
    trauma evaluation, 64  
history taking, 27, 42, 94  
  *see also individual conditions*  
home health care, 4  
  costs, 2  
  extended home observation, 17  
  telemedicine *see* telemedicine  
  *see also* alternative care  
hospice, 332–3  
hospital-acquired pneumonia  
  *see* health-care associated pneumonia  
hospital at home  
  *see under* alternative care  
hospital-based care  
  admission route *see* disposition (admission route)  
  alternatives *see* alternative care  
  continuation of care, 304  
  DEED study, 306  
  follow-up care, 14, 15  
  hospital at home  
    *see* alternative care  
  preventing hospital returns, 305–6  
  reducing demand for ED use, 302–7  
  resource coordination, 15  
  risks of, 303  
  short-stay geriatric unit, 3  
human immunodeficiency virus (HIV), 260  
humerus, proximal, fractures, 249, 343  
hydrocephalus, normal pressure, 104  
hyperactive delirium, 102  
hypercalcemia, 265, 294–5  
  diagnostic tests, 295  
  etiology, 295  
  malignancy-associated, 295  
  symptoms and diagnosis, 265, 295  
  treatment, 265, 295  
hyperkalemia, 290–1  
  in diabetes, 285  
  tumor lysis syndrome, 264  
hyponatremia, 294  
hyperosmolar hyperglycemic state (HHS), 284–5  
hypertension, 21, 88  
  alcohol consumption and, 276  
  atrial fibrillation and, 200  
  chronic, intracerebral hemorrhage, 174  
  epistaxis and, 161  
  malignant, 229  
  prevalence, 94  
  pulmonary *see* pulmonary hypertension  
  subarachnoid hemorrhage and stroke, 139, 175  
  supine, 118, 202  
hyperthyroidism, emergency  
  *see* thyroid emergencies  
hyperviscosity syndrome, 266–8  
hypoactive delirium, 102  
hypocalcemia, 296  
hypokalemia  
  in diabetes, 285  
  disposition (admission route), 292  
  etiology, 291  
  symptoms and examination, 291  
  treatment, 285, 292  
hyponatremia, 292–4  
  diagnostic tests, 293  
  etiology, 292  
  euvolemic, 292  
  hypovolemic, 292, 293  
  symptoms and examination, 293  
  treatment, 293  
hypoparathyroidism, 296  
hypotension, traumatic brain injury, 177  
hypothyroidism, complication  
  *see* myxedema coma  
hypovolemic hyponatremia, 292, 293  
idiopathic thrombocytopenic purpura, 267  
imaging studies  
  abdominal pain, 95, 208–9  
  angiography, 95  
  computed tomography, 95, 97, 98  
  plain radiographs, 95, 99  
  ultrasound, 95  
  pelvic fractures, 247  
  shoulder dislocations, 250  
  traumatic brain injury (TBI), 176  
  urinary tract infections (UTI), 233  
immunology, aging, 23, 208, 254  
inattention, consciousness and, 105  
infections  
  abdominal, 258–9  
    acute cholecystitis, 97–8, 258–9  
  appendicitis *see* appendicitis  
  *Clostridium difficile*  
    *see Clostridium difficile*  
  diverticulitis *see* diverticulitis  
  atypical presentations, 24  
  brain abscesses, 260  
  CNS *see* central nervous system (CNS)  
  colitis *see* colitis  
  fever, 254, 264  
  differential diagnosis, 255  
  Fournier's gangrene *see* Fournier's gangrene  
  history taking, 255  
  human immunodeficiency virus (HIV), 260  
  intervertebral disk, 149–50  
  physical examination, 255  
  pneumonia *see* pneumonia  
  skin and soft tissue, 260  
  special geriatric populations, 260–1  
  suspected, emergency department (ED), 255  
  urinary tract infections  
    *see* urologic emergencies  
  vital signs, 254  
  white blood cell count (WBC), 255  
    leukopenia *see* leukopenia  
  infectious diseases, 254–62  
    aging and immunity, 254  
  inferior vena cava, collapse, testing, 33–4  
  Inouye et al. (prediction tool for functional decline), 316  
  instrumental activities of daily living (IADL), 56, 314  
  hospital at home, 304  
  insulin, 284–5  
  interleukin-6 (IL-6), 53, 55  
  intervertebral disk  
    degenerative disease, 148  
    displacement causing sciatica, 148–9  
    herniation, 149, 243  
    infections, 149–50  
  intestine *see* bowel  
  intracerebral hemorrhage (ICH), 35, 173–5  
    blood pressure, 174  
    diagnosis and investigations, 174  
    headache and, 141  
    treatment, 174–5  
  intubation, 32–3  
    initial evaluation of trauma, 62  
  ISAR (Identification of Seniors At Risk), 314–15  
  ischemic colitis *see* colitis  
  ischemic heart disease *see* acute coronary syndrome (ACS)  
  ischemic stroke *see* acute ischemic stroke (AIS)  
  Kayexalate, 291  
  ketorolac, 71  
  kidney  
    acute injury, 220–1  
    aging, 219  
    function *see* renal function  
  laboratory parameters, normal/abnormal, 25  
  labyrinthitis, 131  
  lactic acidosis, 297  
  large bowel obstruction *see* bowel  
  legal considerations, elder mistreatment, 357–8  
  leukopenia, fever and, 263–4  
  life expectancy, screening for conditions, 2  
  liver, drug metabolism, 43–4  
  living will, 36  
    *see also* advance directive (AD)  
  loop diuretics, 201  
  low-molecular-weight heparin (LMWH), 79  
  Ludwig's angina, 163–4  
lungs  
  aging and, 87, 185–6  
  cancer, 196–7  
  residual volume (RV), 185  
  restrictive lung disease, 196  
    pulmonary function and, 186  
  smoking and, 185–6  
  structural changes with age, 87  
  *see also* pulmonary function  
magnesium, asthma and COPD, 188  
magnetic resonance imaging (MRI)  
  acute ischemic stroke, 171  
  back pain, 148  
  brain, 108  
  pulmonary embolism, 193  
  spinal epidural abscess (SEA), 179  
major depressive disorder  
  *see* depression  
malignancies *see* cancer  
malignant otitis externa (MOE), 160–1  
Mallory–Weiss tears, 212  
malnutrition, screening, 317  
mannitol, 288  
median nerve, distal radius fractures and, 251  
medical futility, 36–7  
Medicare *see* financial issues  
medication  
  adverse drug reaction (ADR) *see* adverse drug reaction (ADR)  
  age-related changes, resuscitation, 34–5  
  appropriate prescribing, 42–3  
  Beers Criteria *see* Beers Criteria  
  delirium and, 103–4  
  dizziness and, 128  
  geriatric emergency department, use in, 11–13  
  noncompliance, 45  
  overuse, primary headache disorders, 143  
  polypharmacy, 3, 11  
  prescription, quality indicators, 309–10  
  resuscitation, 34–5  
  safe prescribing, 48–9  
  thrombocytopenia due to, 267  
  trauma and, 61–2  
    anticoagulants  
      *see* anticoagulants  
  urinary retention/stricture and, 223  
  *see also* pharmacology  
  *individual drugs and drug groups*  
meningitis, bacterial  
  community-acquired, symptoms, 178  
  diagnostic tests, 259  
  headache and, 142

- herpes simplex
  - meningoencephalitis versus, 259–60
- treatment, 259
- mental health
  - antipsychotics, 46
  - benzodiazepines, 46
  - depression *see* depression
  - psychiatric emergencies
    - see* psychiatric emergencies
  - schizophrenia, 278
  - see also* geriatric psychoses
- mental status, altered, 102–13
  - assessment of patient, 103, 105–8
  - diagnostic testing, 107–8
  - history taking, 105
  - neurological examination, 106
  - physical, 105–8
- cognition
  - dementia, 104–5
  - see also* cognitive impairment
- consciousness, disturbances in
  - see* consciousness
- decision-making capacity
  - assessment, 110
- prevalence, 103
- quality indicators and, 309
- treatment, 108–9, 110
- mesenteric venous thrombosis (MVT), 96
- metabolic and endocrine emergencies, 284–99
  - acid–base disorders *see* acid–base disorders
  - adrenal crisis *see* adrenal crisis
  - calcium disorders
    - see* hypercalcemia;
    - hypocalcemia
  - diabetic ketoacidosis *see* diabetic ketoacidosis (DKA)
  - hyperosmolar hyperglycemic state *see* hyperosmolar hyperglycemic state (HHS)
  - metabolic acidosis *see* acid–base disorders
  - metabolic alkalosis, 296, 297
  - pituitary apoplexy *see* pituitary apoplexy
  - potassium disorders
    - see* hyperkalemia
    - hypokalemia
  - rhabdomyolysis
    - see* rhabdomyolysis
  - sodium disorders *see* sodium disorders
  - thyroid emergencies *see* thyroid emergencies
- metabolism, drugs, 43–4
- methicillin-resistant *Staphylococcus aureus* (MRSA), 91, 256
- methylxanthines, 188
- migraine, 143
- Mini-Mental State Exam (MMSE), 105
- modified Richmond Agitation and Sedation Scale (mRASS), 103
- mortality rate, abdominal pain and, 94
- motor vehicle collisions, 60
  - older driver safety *see* driving
  - pedestrians struck by automobiles, 60
  - pelvic fractures *see* pelvic fractures
  - syncope and, 124
  - traumatic rupture of the aorta (TRA), 202
- mucosa, aging, 21
- Multinational Association for Supportive Care in Cancer (MASCC) Score, 264
- muscle, 54
  - creatinine levels and, 219
  - sarcopenia *see* sarcopenia
- musculoskeletal aging, 23, 61
  - see also* muscle
- Mycobacterium tuberculosis*
  - see* tuberculosis
- myocardial infarction
  - abdominal symptoms, 95
  - complications after, 78
  - NSTEMI *see* non-ST elevation myocardial infarction (NSTEMI)
  - STEMI *see* ST elevation myocardial infarction (STEMI)
- myoglobin, 287
- myxedema coma, 286
- National Institutes of Health Stroke Scale (NIHSS), 171, 172
- necrotizing fasciitis, 260
- necrotizing skin and soft tissue infections (NSTI), 260
- neglect, 356
  - see also* elder mistreatment
- NEISS-CADES database, 44
- nephrotoxic drugs, 221
- neurological emergencies, 170–84
  - acute ischemic stroke *see* acute ischemic stroke (AIS)
  - CNS infections *see* central nervous system (CNS)
  - non-traumatic intracranial hemorrhage *see* brain injury
  - seizures *see* seizures
  - traumatic brain injury (TBI)
    - see* brain injury
- neurology, aging, 23
- neuromuscular blocker, non-depolarizing, 33
- non-anion gap acidosis, 297
- noncompliance, medication, 45
- non-convulsive status epilepticus (NCSE), 181
- noninvasive positive pressure ventilation (NIPPV), 33, 90
- noninvasive ventilation
  - asthma and COPD, 189
  - pneumonia, 191
- nonocclusive mesenteric ischemia (NOMI), 96, 215
- non-ST elevation myocardial infarction (NSTEMI), 24, 25
  - chest pain, 78
  - see also* acute coronary syndrome (ACS)
- nonsteroidal anti-inflammatory drugs (NSAIDs), 48
  - analgesia, 71
  - gastrointestinal pathology, 208
  - gout, 239
- normal pressure hydrocephalus (NPH), 104
- nose, 161–2
  - bleeding *see* epistaxis
- nuchal rigidity, 140
- nursing homes
  - adverse drug reaction (ADR), 45
  - pulmonary infections, 79
- nutrition/diet
  - malnutrition screening, 317
  - supplements (vitamins), 42
- nystagmus, dizziness and, 133–4
- observation
  - within ED, 16
  - extended home observation, 17
- OESIL score, 121, 122
- Ogilvie’s syndrome (acute colonic pseudo-obstruction), 214
- oncologic diseases, 263
- oncologic emergencies
  - see* hematologic and oncologic emergencies
- opioids
  - altered mental status, 108
  - analgesia, 71–2
  - falls/adverse events
    - associated, 71–2
  - specific, risks, 72
- orchitis, 226–7
- orthopedic emergencies
  - see* rheumatologic and orthopedic emergencies
- orthostatic hypotension, 132
  - syncope and, 114, 117, 118
- osteoarthritis, 241–2
- osteomyelitis, vertebral, 150
- osteoporosis, 23, 244
  - back pain and *see* back pain
  - corticosteroid side effect, 240
  - fall-associated fractures, 244
  - pelvic fractures *see* pelvic fractures
  - proximal humeral fractures, 249, 343
  - rib fractures, 22
  - sacral fractures, 149
  - thoracic cage, lung volume and, 87
  - vertebral fractures, 149, 248
- otitis externa, 160
- pain
  - abdominal *see* abdomen
  - assessment and scores, 72
  - back *see* back pain
  - chest *see* chest pain
  - male genital system, 226
  - management, 69–75
    - adverse drug events, 69
    - analgesics *see* analgesia
    - Beers Criteria, 69–70
    - drug sensitivity and clearance, 69
    - goals, 72–3
    - quality indicators (ED), 72, 309
    - undertreatment, 69
    - see also* analgesia
  - palliative care, 329–30
  - scores, 72–3
- palliative care, 4, 17, 322–35
  - advance directive *see* advance directive (AD)
  - aim, 322
  - better symptom care, 329–30
  - capacity, 330
  - CPR and prognosis in advanced illness, 323–6
  - death disclosure, 328
  - emergency assessment, 327–9
    - ABCD (Airway, Breathing, Circulation, Disability), 328–9
    - SPEED (Screen for Palliative and End of Life Care Needs), 328
  - emergency department (ED), 5, 17, 322–3
  - end-of-life care, 1, 322–35
    - DNR (Do Not Resuscitate) orders, 175
- ethics, 36
  - resuscitation, 36
- ethics, 36
  - last hours, 332–3
  - pre-hospital setting, 326
  - resuscitation, 326–7
  - trajectories of illness and prognosis, 322–3
  - ventilator discontinuation, 331–2
- pancreatitis, 98–9, 210
  - symptoms and diagnosis, 210
  - treatment, 210
- papilledema, 140
- paraphimosis, 230–1
  - description and risks, 230
- parathyroid hormone (PTH), 294, 295
- patient (geriatric), 20–30
  - atypical disease presentation, 24–6
  - diagnostic testing *see* diagnostic tests/testing
  - geriatric emergency medicine
    - see* emergency medicine (EM) (geriatric)
  - history taking, 27, 42, 94
    - see also* individual conditions
  - medication *see* medication
  - prescribing cascade, 41
  - satisfaction, 16, 28–9
  - supplements (vitamins), 42
  - treating delirium (non-pharmacological), 108–9
- pedestrians struck by automobiles, 60
  - see also* motor vehicle collisions
- pelvic fractures, 246–8, 343
  - diagnostic tests, 247
  - disposition (admission route), 248
  - history taking, 247
  - physical examination, 247
  - trauma causing, 64
  - treatment, 247
- penetrating atherosclerotic ulcer (PAU), 203
- penis
  - glans, inflammation *see* balanitis
  - see also* genital system (male)
- pentazocine, 48
- peptic ulcer disease (PUD), 26, 97, 211–12
  - diagnosis, 212
  - emergency department (ED), 211
  - treatment, 212
- PERC decision rule, 192
- percutaneous coronary intervention (PCI)
  - acute coronary syndrome, 200
  - ST elevation myocardial infarction (STEMI), 77
- performance testing of function, 314
- pericarditis, chest pain, 78–9
- peritonsillar abscess (PTA), 164
- pharmacology, 41–50
  - adverse drug reaction *see* adverse drug reaction (ADR)
  - aged heterogeneity, 41
  - aging and *see* aging
  - definitions, 42–3
  - epidemiology, 41–2
  - herbal medicines, 42
  - medication history taking, 42
  - pharmacodynamics, 44
  - pharmacokinetics, 43–4
  - poisoning (therapeutic errors), 46
  - polypharmacy *see* polypharmacy
  - prescribing cascade, 41



Index

- pharmacology (*cont.*)  
  preventability, 48–9  
  safe medication for elderly patients, 46–9  
  supplements (vitamins), 42  
  *see also* medication  
phimosi/s/paraphimosis, 230–1  
physical abuse, 356  
  *see also* elder mistreatment  
physical activities of daily living (PADLs), 56  
physical environment, of  
  geriatric emergency department, 10–11  
physical examination, emergency department (ED), 27  
physiology of aging, 20–3, 32  
  back pain, 146  
  body composition, 21, 43  
  cancer, 263  
  cardiovascular system, 21, 60, 87–8, 204  
  central nervous system (CNS), 44, 61  
  creatinine, 219  
  dental issues, 162  
  dizziness, 128  
  endocrine system, 23, 61  
  gastrointestinal system, 22–3, 61  
  genitourinary system, 23, 219–20  
  hearing impairment, 316  
  heart, 88  
  hematologic, 23  
  immunologic, 23  
  lung function, 32, 61, 185–6  
  lungs, 87, 185–6  
  male genital system, 220  
  musculoskeletal, 23, 61  
  neurologic, 23  
  psychological, 23  
  renal system, 32, 61, 219  
  respiratory, 22, 87, 185–6  
  resuscitation and *see* resuscitation  
  skin/mucosa, 21, 61, 260  
  syncope, 114–15  
  trauma, 60–1  
  vital signs, 21  
  *see also* aging  
pituitary apoplexy, 289–90  
platelet transfusion, 267  
pneumonia, 24, 90–1, 190–1, 255–7  
  aspiration *see* aspiration pneumonia  
  atypical presentation, 24  
  causative organisms, 190  
  chest pain and, 79  
  community-acquired *see* community-acquired pneumonia (CAP)  
  diagnostic tests, 256  
  dyspnea and, 90–1  
  etiology, 91, 190–1  
  health care-associated pneumonia *see* health care-associated pneumonia  
  pathogenesis, 190  
  severity scores, 191, 192  
    CURB-65, 256–7  
  symptoms, 190, 256  
  treatment, 91, 191  
pneumothorax, 193–4  
  chest pain, 80  
  definition, 193  
  spontaneous, 193–4  
  tension, 194  
poisoning (therapeutic errors), 46  
polymyalgia rheumatica, 142, 159, 242  
  diagnosis, 242  
  history taking, 242  
  treatment, 242  
polypharmacy, 41  
  adverse drug reaction (ADR), 45, 69  
    *see also* adverse drug reaction (ADR)  
  elders and, 23–4  
  geriatric emergency department (ED), 11  
  quality indicators, 309–10  
    *see also* medication  
population aging, 1–2  
  health care provision and, 8  
    *see also* demographics  
PORT study, 191  
positional vestibular syndrome (PVS), 132  
  Dix–Hallpike test, 134  
potassium  
  diabetes and, 285  
  disorders, 290–2  
    hyperkalemia *see* hyperkalemia  
    hypokalemia *see* hypokalemia  
potassium iodide, 287  
power of attorney, 36  
  *see also* advance directive (AD)  
Pradaxa, 62  
pre-hospital care, 26–7  
pre-hospital setting, palliative care, 326  
presbycusis *see* hearing impairment  
prescribing, appropriate, 42–3  
  “prescribing cascade”, 41  
prescription (of medication), 49  
  prescribing cascade, 41  
  quality indicators, 309–10  
  safe, 46–9  
preventive care, emergency department (ED), 28  
primary care, 303  
primary care providers (PCPs), EDs and, 5  
PROFET (Prevention of Falls in the Elderly Trial), 344–7  
propylthiouracil (PTU), 287  
prostheses, septic arthritis and, 237  
prostate, 220  
  benign prostatic hyperplasia (BPH), 224, 225  
  examination, 224  
  prostatitis, 227–8  
prothrombin complex concentrate (PCC), 62  
pseudoclaudication, 148  
pseudogout, 239–40  
psychiatric emergencies, 270–83  
  alcohol use and, 276  
  depression *see* depression  
  psychoses *see* geriatric psychoses  
  suicide *see* suicide  
  psychological (changes with age), 23  
  psychoses *see* geriatric psychoses  
  pulmonary embolism, 79–80, 89, 191–3  
    diagnosis and difficulties, 79–80  
    diagnostic tests, 89, 192–3  
      CT pulmonary angiogram (CTPA), 79, 80  
      ECG, 192  
    ventilation–perfusion scan, 192  
  outcomes, 80  
  risk factors, 79  
  treatment, 79, 193  
    anticoagulation, 193  
    surgical procedures, 193  
    thrombolytic therapy, 193  
    vena cava filter, 193  
pulmonary emergencies, 185–98  
  abdominal examination, 186  
  asthma *see* asthma  
  bronchiectasis *see* bronchiectasis  
  cardiac examination, 186  
  diagnostic tests, 186–7  
  history taking, 186  
  hypertension, pulmonary (PH), 195–6  
  infections, nursing homes, 79  
  physical examination, 186  
  pneumonia *see* pneumonia  
  pneumothorax *see* pneumothorax  
  restrictive lung disease *see* lungs  
  smoking and, 185–6  
  tuberculosis *see* tuberculosis  
pulmonary function  
  aging, 32, 61, 186  
  COPD *see* chronic obstructive pulmonary disease (COPD)  
  dyspnea, 186  
  pulmonary hypertension, 195–6  
  pulmonary vascular resistance (PVR), 87  
purified protein derivative (PPD), tuberculosis diagnosis, 194  
pyelonephritis, 233–4  
quality  
  geriatric ED *see* geriatric emergency department (ED)  
  indicators, 308–10  
  mental impairment, 308  
  mental status screening, 309  
  pain management, 309  
  polypharmacy, 309–10  
radiographs  
  abdomen, 208–9  
  bowel, 214  
  chest, 187  
    pneumonia, 191  
    pulmonary embolism, 192  
  in TB, 195  
  spinal, 148  
radius, distal, fractures, 251, 343  
  disposition (admission route), 251  
  median nerve and, 251  
  symptoms and diagnosis, 251  
  treatment, 251  
recombinant tissue plasminogen activator (rt-PA), 171  
rectum, back pain and, 147  
regional anesthesia, 72–3  
renal function  
  acid–base disorders *see* acid–base disorders  
  acute renal failure *see* kidney  
  aging effect, 32, 61  
  assessment, 43  
  respiratory acidosis, 296, 297  
  respiratory alkalosis, 296, 297  
  respiratory muscle changes, 185  
respiratory system  
  aging, 22  
  asthma *see* asthma  
  breathing assessment, 33  
  bronchiectasis *see* bronchiectasis  
  distress *see* dyspnea  
  failure, respiratory, 63  
    *see also* resuscitation  
intubation, 32–3  
lungs *see* lungs  
noninvasive positive pressure ventilation (NIPPV), 33, 90  
structural mechanics, 86–9  
  aging, 87  
  cardiovascular changes, 87–8  
  lung function *see* lungs  
throat emergencies *see* throat  
resuscitation, 31–40  
  advance directive (AD)  
    *see* advance directive (AD)  
  airway, 32–3  
  anatomic age-related changes, 32  
  breathing assessment, 33  
  circulation, 33–4  
  end-of-life care, 36  
  epidemiology and outcomes, 31–2  
  epilepsy, 35–6  
  ethics, 36  
  family-witnessed, 37–8, 326–7  
  fluids, 33, 288  
  intubation, 32–3  
  management, 32–6  
  medical futility, 36–7  
  medications, 34–5  
  out-of-hospital, outcomes, 31  
  palliative care, 326–7  
  pathophysiology, 32  
  sepsis, 34  
  stroke and, 35  
  traumatic brain injury, 177  
  under-resuscitation, 63  
  ventilation *see* ventilation  
    *see also* cardiopulmonary resuscitation (CPR)  
retinal artery occlusion  
  branch (BRAO), 158  
  central *see* central retinal artery occlusion (CRAO)  
retinal detachment, 155  
retinal vein occlusion 158–9  
retinopathy, diabetic, 155  
retropharyngeal abscess (RPA), 165  
rhabdomyolysis, 287–8  
rheumatoid arthritis, 240–1  
rheumatologic and orthopedic emergencies, 237–53  
  giant cell arteritis *see* giant cell arteritis (GCA)  
  gout *see* gout  
  hip dislocation, or fracture *see* hip  
  humeral, proximal fractures  
    *see* humerus, proximal  
  orthopedic injuries, 243–4  
  osteoarthritis *see* osteoarthritis  
  pelvic fractures *see* pelvic fractures  
  polymyalgia rheumatica  
    *see* polymyalgia rheumatica  
  pseudogout *see* gout  
  rheumatoid arthritis  
    *see* rheumatoid arthritis  
  rheumatologic emergencies, 237  
  septic arthritis *see* septic arthritis  
  shoulder dislocations *see* shoulder  
  spinal stenosis *see* spine  
  vertebral fractures *see* vertebral fractures  
rib fractures, 22, 82, 343  
Risk Stratification of Syncope in the Emergency Department (ROSE), 121  
sacral fractures, 149  
sarcopenia, 53  
satisfaction, patient, 16, 28–9



- SBIRT (Screening, Brief Intervention, and Referral to Treatment), 277
- schizophrenia, 278  
*see also* geriatric psychoses
- sciatica, disk displacement causing, 148–9  
*see also* back pain
- screening, conditions reducing active life expectancy, 2
- seizures, 179–81  
  complex partial, 180  
  new, 179–80  
  status epilepticus *see* status epilepticus (SE)  
  syncope, differential diagnosis, 117–18
- self-neglect, 356  
*see also* elder mistreatment
- sepsis, 34
- septic arthritis, 237–8
- sexual abuse, 356  
*see also* elder mistreatment
- sexual activity, 222
- SHERPA (Score Hospitalier d’Evaluation du Risque de Perte d’Autonomie), 315–16
- shingles *see* herpes zoster
- shock, 34  
  management, 33, 34
- Short Blessed Test (SBT), 106
- Short Michigan Alcoholism Screening Test, 277
- shoulder dislocations, 249–51  
  disposition (admission route), 250  
  history taking and symptoms, 250  
  imaging, 250  
  neurovascular complications, 249  
  treatment, 250
- SIGNET (Systematic Intervention for a Geriatric Network of Evaluation and Treatment), 315
- Six-Item Screener (SIS), 106
- skew deviation, dizziness, 134
- Skilled Nursing Facility (SNF), 352
- skin  
  aging, 21, 61, 260  
  infections, 260
- Slone Survey, 42
- small bowel obstruction, 97, 213–14
- smoking, 185–6
- social services  
  and case management, 336–42  
  ED social worker, 338  
*see also* case management
- sodium disorders, 292–4
- soft tissue  
  infections, microorganisms implicated, 260  
  necrotizing skin and soft tissue infections (NSTI), 260
- SPEED (Screen for Palliative and End of Life Care Needs), 328
- spinal nerve roots, 147
- spinal stenosis, 242–3
- spine  
  back pain, 148–50  
    *see also* back pain  
  cancer, primary or metastatic, 149  
  disk disorders  
    *see* intervertebral disk  
  epidural abscess (SEA), 179  
  fractures, 343  
  infections, 149–50  
  lumbar spine, 148  
  trauma, 63  
  vertebrae *see* vertebral fractures
- spontaneous pneumothorax, 193–4
- ST elevation myocardial infarction (STEMI)  
  chest pain, 77–8  
  percutaneous coronary intervention (PCI), 77  
*see also* acute coronary syndrome (ACS)
- staffing  
  education *see* education  
  geriatric emergency department (ED), 16
- statins, rhabdomyolysis due to, 288
- status epilepticus (SE), 180–1  
  resuscitation, 35–6  
  treatment, 35–6, 180–1
- straight leg test, 147
- stroke, 172  
  acute ischemic *see* acute ischemic stroke (AIS)  
  resuscitation and, 35  
  transient ischemic attack *see* transient ischemic attack (TIA)  
  treatment, 35
- stupor, 102  
  diagnostic testing, 108
- subarachnoid hemorrhage (SAH), 140–1  
  aneurysmal *see* aneurysmal subarachnoid hemorrhage (aSAH)  
  clinical features, 140–1  
  computed tomography (CT), 141, 175, 177  
  consciousness and, 141  
  epidemiology, 140  
  headache and, 139, 140  
  misdiagnosis, 140  
  “thunderclap” headache, 139
- subclavian steal, 117
- subdural hemorrhage (SDH), 177, 344  
  headache, 141
- substance use/abuse, 275–8  
  alcohol *see* alcohol abuse/dependence  
  epidemiology, 275  
  suicide, 271
- suicide, 23, 270–3  
  epidemiology, 23, 270  
  prevention, 272–3  
  risk factors, 270–2  
    access to means, 271  
    cognitive deficits, 271  
    medical illness, 271  
    psychiatric illness, 271  
    psychosocial, 271  
    sociodemographic, 271  
    substance use disorders, 271
- signs and assessment, 271–2
- treatment, 272
- superior mesenteric artery (SMA)  
  embolus, 96  
  thrombosis, 96
- superior vena cava syndrome, 265
- supplements (vitamins), 42
- surrogates (decision making), 110, 330
- symptoms and diagnostic tests, 266
- syncope, 114–27, 202  
  ancillary studies, 119–20  
  cardiovascular testing, 119–20  
  carotid sinus massage and, 119  
  definition, 114  
  diagnosis, 115, 124  
  driving and, 124  
  emergency evaluation, 114, 115, 118, 119, 123  
  epidemiology, 114, 115  
  etiology, 115–17, 122  
    age-related, 114–15  
    cardiac, 117, 118  
    carotid sinus hypersensitivity, 116–17, 119  
    neurally mediated, 115–17  
    orthostatic hypotension, 117, 118  
  transient ischemic attack (TIA), 117
- history taking, 117–18  
  autonomic failure, 118  
  sequence of events, 117  
  specific data points, 117
- identification, 202
- laboratory testing, 120
- management, 202
- morbidity, 115
- neurologic testing, 120
- physical examination, 118–19
- prognosis/risk stratification, 122
- recurrent, 115
- risk factors, 122
- seizure, differential diagnosis, 117–18
- systems of evaluation, 120–2
- syndrome of inappropriate ADH secretion (SIADH), 293
- systolic blood pressure, 88
- telemedicine, 306
- telemetry monitoring, syncope, 119–20
- temporal arteritis *see* giant cell arteritis (GCA)
- tension pneumothorax, 194
- theophylline, 188
- thorax, trauma, 63–4
- throat, 162–6  
  angioedema due to ACE inhibitors *see* ACE inhibitors  
  dental abscess *see* dental issues  
  epiglottitis *see* epiglottitis  
  esophagus *see* esophagus  
  Ludwig’s angina *see* Ludwig’s angina  
  peritonsillar abscess *see* peritonsillar abscess  
  retropharyngeal abscess *see* retropharyngeal abscess
- thrombocytopenia, 266–8  
  aging and, 267  
  etiology, 267
- thrombolytic therapy  
  ischemic stroke, 171  
  pulmonary embolism, 193  
  STEMI, 77
- thrombotic thrombocytopenic purpura, 267
- thyroid emergencies, 286–7  
  thyroid storm, 286–7  
  examination and diagnosis, 287  
  symptoms, 286–7  
  treatment, 287
- thyroid hormones, 286
- tonsil, peritonsillar abscess *see* peritonsillar abscess
- transient ischemic attack (TIA), 171–3  
  dizziness and, 132  
  management, 173  
  syncope and, 117
- transitional care *see* discharge
- transport, elders in emergency, 26–7
- trauma, 59–68  
  brain *see* brain injury  
  chest, 82  
  disposition (admission route), 64–5  
  epidemiology, 59  
  evaluation and management, 62–4  
  abdomen, 64  
  central nervous system, 63  
  initial evaluation, 62–3  
  musculoskeletal, 64  
  prognosis, 64  
  spine, 63  
  thorax, 63–4
- falls *see* falls
- injury patterns in the elderly, 59
- intubation, 62
- mechanisms of injury, 59–60
- physiology and *see* physiology of aging
- pre-existing medicines, 61–2
- pre-hospital considerations, 62
- rheumatoid arthritis and, 241
- traumatic brain injury (TBI)  
  *see* brain injury
- triage, geriatric emergency department (ED), 13–15
- Emergency Severity Index (ESI), 13, 14, 27
- two-step geriatric program, 13–15
- tricyclic antidepressants (TCAs), 46
- trigeminal neuralgia, 143
- TRST (The Triage Risk Screening Tool), 315
- tuberculosis, 194–5
- tumor lysis syndrome, 264–5
- tumor necrosis factor (TNF- $\alpha$ ), 53, 55
- Two-step Geriatric Program, 13–15
- tympanic membrane, dizziness and, 133
- ultrasound  
  abdomen, 209  
  bedside  
    pulmonary diseases, 187  
    resuscitation, 34
- under-resuscitation, 63
- undertreatment, pain management, 69
- urinary retention/stricture, 223–5  
  medications and, 223  
  phimosis/paraphimosis, 230  
  treatment, 224–5
- urinary tract infections (UTI), 24, 231–4, 257–8  
  antibiotic use, 233, 257  
  bacteriuria *see* bacteriuria  
  community-dwelling elders, 257  
  definition, 258  
  diagnostic testing, 232–3  
  disposition (admission route), 234  
  imaging studies, 233  
  indwelling catheters, elders with, 257–8  
  long-term care facilities, elders in, 257, 258  
  microorganisms implicated, 257  
  pyelonephritis, 233–4  
  symptoms and examination, 232, 258  
  treatment, 233

Index

urine alkalinization, 288	vasoreactivity test, 196	vertebral osteomyelitis, 150	in shock, 21
urine flow, phimosis/paraphimosis and, 230	vasovagal syncope, 116	vertigo	vitamin D, hypocalcemia and, 296
urologic emergencies	vena cava	benign paroxysmal positional	vitamin K
hematuria <i>see</i> hematuria	filter, pulmonary embolism, 193	<i>see</i> benign paroxysmal	administration, 268
kidney <i>see</i> kidney	superior vena cava	positional vertigo (BPPV)	antagonists, 266
nephrotoxic drugs, 221	syndrome, 265	central positional, 132	vitreous hemorrhage, 154–5
paraphimosis <i>see</i> phimosis/	venous thromboembolic disease	vestibular neuritis, 133, 135	VO <sub>2max</sub> , 55
paraphimosis	(VTE), 191–2	vestibular physiology	volvulus, 97
renal function <i>see</i> renal function	ventilation, 33, 90	acute vestibular syndrome (AVS),	
urinary retention/stricture	assessment of breathing, 33	131, 133	warfarin
<i>see</i> urinary retention/	discontinuation <i>see</i> palliative care	dizziness, 128	epistaxis and, 162
stricture	intubation, 32–3	episodic vestibular syndrome	trauma and, 62
UTIs <i>see</i> urinary tract	noninvasive positive pressure	(EVS), 131–2	<i>see also</i> anticoagulants
infections (UTI)	ventilation (NIPPV), 33, 90	positional vestibular syndrome	weakness, generalized
<i>see also</i> genitourinary emergencies	ventilation–perfusion scan,	(PVS) <i>see</i> positional	<i>see</i> generalized weakness
uterine prolapse, 222–3	pulmonary embolism, 192	vestibular syndrome (PVS)	white blood cell count (WBC)
uveitis, 155–6	ventilator discontinuation		infections, 255
	<i>see</i> palliative care	vision	<i>see also</i> leukopenia, fever and
vaginal bleeding, 222	vertebral fractures, 248–9	disturbances	withholding treatment, 4, 37
vaginal discharge, 221–2	compression, 149	headache and, 140	wrist fractures, 343
varicella zoster virus <i>see</i> herpes	diagnostic testing, 248	<i>see also</i> eye emergencies	<i>see also</i> radius, distal
zoster	history taking, 248	impairment, screening, 316	
vascular steal syndromes, 117	osteoporosis and, 149, 248	loss, giant cell arteritis (GCA),	
vasodilators, 201	physical examination, 248	159, 160	Xarelto, 62
vasopressin secretion, 293	trauma causing, 64	vital capacity (VC), 185	X-rays <i>see</i> radiographs
	treatment, 248	vital signs	
		aging, 21, 187	zoster <i>see</i> herpes zoster