

Index

Note: page numbers in *italics* refer to figures, tables and flow charts

- adverse outcomes, minimising 111–23
- anaesthesia
 - concerns following birth 136
 - general 136
 - local anaesthetic infiltration 127
 - regional blockade 131
 - concerns following birth 136
 - contraindications 133
 - spinal single-shot 131, 132, 135
- analgesia 19
 - caesarean section at full dilatation 98
 - combined spinal–epidural 135
 - Kielland's forceps 89
 - manual rotation 76
 - nonrotational forceps birth 65
 - regional 132
 - contraindications 133
 - see *also* epidural analgesia
 - antibiotic prophylaxis, caesarean section at full dilatation 99, 105
 - anticipation 35
 - asynclitism 85, 89, 91
- biparietal diameter 20
- Bird GC 5–7
- birthing partner, professional behaviour with 39–40
- bladder catheterisation 19
 - nonrotational forceps birth 65
- bladder, in caesarean section at full dilatation
 - adherence 100
 - damage 105
 - reflection 100
- blood transfusion, caesarean section at full dilatation 105
- breech birth, reverse 103, 103
- caesarean section
 - developed/developing countries
 - comparison 9
 - epidemiology 95
 - incidence 95
 - low resource countries 7–8
 - operative vaginal birth in theatre 26
 - proceeding to from forceps birth 79
 - rates
 - in developed countries 6
 - in developing countries 5, 6
- caesarean section at full dilatation 95–108
 - advanced resuscitation 106
 - analgesia 98
 - antibiotic prophylaxis 99, 105
 - bladder adherence 100
 - bladder damage 105
 - bladder reflection 100
 - blood transfusion 105
 - cephalopelvic disproportion 96
 - cervical laceration 104
 - complication prevention 106
 - consent 98
 - debriefing 107
 - documentation 107
 - epidural analgesia 97
 - examination in theatre 98
 - fetal complications 106
 - fetal distress 97

Index

fetal head delivery 101–3	Clinical Negligence Scheme for Trusts (CNST) 123
fetal head disimpaction 101	cognitive skills 32, 33, 34
difficulties 101–3	combined spinal–epidural (CSE)
fetal pillow use 98–9, 102	analgesia 135
fetal skull fracture due to impacted head 106	communication 37–9
further surgery 105	with mother 39–40
high maternal body mass index 97	caesarean section at full dilatation 107
high vaginal laceration 104	operative vaginal birth 120–1
ICU admission 105	consent <i>see</i> maternal consent
labour early immobilisation 97	decision making 35–7, 41
labour ward supervision 106	Desperate Debra simulated training 106
lower uterine segment identification 100	documentation
maternal complications 103–5	caesarean section at full dilatation 107
perinatal asphyxia risk 106	nonrotational forceps birth 75
postpartum haemorrhage 105	operative vaginal birth 121–3
predisposing factors 96–7	electronic fetal monitoring 21
preparation 98–9	fetal distress identification 97
previous abdominal surgery 100	fetal status review 20
reasons for 96	epidural anaesthesia 132
reverse breech birth 103, 103	epidural analgesia 133
safety comparison with OVB 113, 115	avoidance 111
sepsis 105	caesarean section at full dilatation 97
surgical technique 99–103, 103	delayed pushing 112
ureter damage 105	operative vaginal birth risk 134
uterine angle extension 103–4	oxytocin use 112
uterine cavity opening 100	procedure 134
uterine incision 99	episiotomy 56, 72, 73
WHO checklist 98	Kielland's forceps use 92
cardiotocography	perineal tears 118
fetal compromise 113	face presentation 77
fetal status review 20	fetal assessment
cephalopelvic disproportion 20, 26	abdominal examination 20–1
caesarean section at full dilatation 96	for Kielland's forceps use 89
deep transverse arrest 84–5	fetal compromise
predisposing factors 96	operative vaginal birth indication
trial of operative vaginal birth in theatre 78	113, 122
clinical history taking 19–20	suspected 16

Index

- fetal head
 - assessment for Kielland's forceps
 - use 89
 - autorotation 51–2
 - biparietal diameter 20
 - completing birth 72, 73
 - delivery in caesarean section at full dilatation 101–3
 - disimpaction in caesarean section at full dilatation 101
 - difficulties 101–3
 - engagement 20
 - flexion point 44–5
 - location 46
 - vacuum cups manoeuvring toward 46–7
 - malposition 16, 26
 - midpoint 44–5
 - position 13, 21–4, 23
 - malpresentations 16, 26
 - skull fracture due to impaction 106
- fetal pillow 98–9, 102
- fetal status review 20, 21
- fetus
 - accidental local anaesthetic injection 131
 - caesarean section at full dilatation
 - complications 106
 - delivery of head 101–3
 - disimpaction of head 101–3
 - distress 97
 - limb fractures 106
 - clinically large 20
 - general anaesthetic drug effects 136
- forceps 2–3
 - instrument choice 25, 25, 117
 - see *also* Kielland's forceps, nonrotational forceps
- forceps birth 4
 - after vacuum-assisted birth failure 54
 - consent 64–5
 - consultant presence 79
 - injuries 60
 - manual rotation 75–6
 - risks 25
 - simulation role in training 79–80
 - success rate 60
 - supervision 79
 - trends
 - developed world 3–4
 - developing world 4–5
 - see *also* nonrotational forceps birth, rotational forceps birth
- general anaesthesia 136
- glyceryl trinitrate (GTN) 102
- information exchange 38
- information gathering 34–5
- information understanding/analysing 35
- intrapubic translabial ultrasound (ITU) 22
- instrumental delivery see operative vaginal birth (OVB)
- intensive care unit (ICU) admission 105
- interpersonal skills 41
- intracranial haemorrhage 78
- Joel Cohen incision 99
- Kielland's forceps 83–4, 86–93
 - application 90–2
 - preparation for 89
 - episiotomy 92
 - features 87–8
 - maternal trauma 92–3
 - prerequisites 88–9
 - rotation 92
 - safety 118
 - traction 92
 - wandering application 91
 - see *also* rotational forceps birth

Index

labour <ul style="list-style-type: none">delay in second stage 14, 113early immobilisation 97see <i>also</i> caesarean section at full dilatation	litigation potential areas 111
labour ward supervision 106	operative vaginal birth rate reduction 111–12
leadership, key skills 40–1	mother
lidocaine 127	after vacuum-assisted birth 52
litigation <ul style="list-style-type: none">communication in reduction 121perineal tears 120potential areas 111	communication with 39–40
Litzmann's obliquity 85, 86	caesarean section at full dilatation 107
local anaesthetic <ul style="list-style-type: none">pharmacology 127toxicity 128	operative vaginal birth 120–1
local anaesthetic infiltration 127	debriefing after nonrotational forceps birth 75
complications 131	dignity maintenance 40
perineal 128	distress/fatigue 16
pudendal block 128	vacuum-assisted birth impact 55–6
	see <i>under</i> maternal entries
malposition/malpresentations 16, 84	Naegele's obliquity 85
maternal body mass index	National Institute for Health and Care Excellence (NICE)
fetal complications 106	antibiotic prophylaxis guidelines 99
high 97	intrapartum guideline 113, 122
maternal complications	neonates
caesarean section at full dilatation 103–5	operative vaginal birth
operative vaginal birth 26–7	complications 27
Kielland's forceps 92–3	trauma
maternal consent 16, 18	baby in occiput posterior position 86
caesarean section at full dilatation 98	with operative vaginal birth 118
forceps birth 64–5	risk with sequential instruments 119
maternal trauma 27	nonrotational forceps 61, 62, 63
assessment for post-birth 75	application 66–9, 70
Kielland's forceps use 92–3	assembly 66
occiput posterior position 86	blade locking 69, 69
operative vaginal birth 119	nonrotational forceps birth 59–80, 70
medico-legal matters 110–23	abandoning procedure 77–8
adverse outcome minimising 111–23	analgesia 65
	assessment before 62–5
	catheterisation 65
	clinical assessment 64
	completing birth 75

Index

- nonrotational forceps birth (cont.)
 - of head 72, 73
 - consent 64–5
 - debriefing woman 75
 - episiotomy 72, 73
 - face presentation 77
 - forceps application 66–9
 - forceps removal 72, 74
 - indications for 62, 64
 - patient follow-up 75
 - post-birth trauma assessment 75
 - preparation 65
 - record keeping 75
 - sequential instrument use 77–8
 - special circumstances 77–9
 - technique 65–75, 72
 - traction 71–5, 73, 74
 - low OP position 77
 - ultrasound assessment 64
- nulliparous women, delay in second stage of labour 113
- obstetric forceps *see* forceps
- occiput posterior position 85–6, 91–2
- operative vaginal birth (OVB)
 - abandoning 119
 - abdominal examination 19–21
 - aim 60
 - alternatives 18–19
 - analgesia 19
 - assessment before 19–24, 23
 - bladder catheterisation 19
 - classification 13, 14
 - coaching model 10
 - communication 121
 - complications 26–7
 - contraindications 15, 16–18, 115
 - documentation 121–3
 - flow chart 28
 - historical perspectives 2
 - indications 13–16, 15
 - informed consent 16, 18
 - instrument choice 24–6, 37
 - instrument use 117
 - intervention indications 112–13, 122
 - on the job training 10
 - low resource countries 7–8, 9
 - mannequins 10
 - maternal trauma 27, 119
 - neonatal trauma 118
 - observation period 24
 - Papua New Guinea experience 5–7
 - perineal injuries 119
 - identification 119–20
 - prerequisites 17, 18–19
 - rates 60, 110
 - reduction 111–12
 - RCOG guidelines 115, 116, 119–20, 123
 - risk with epidural analgesia 134
 - risk/benefit balance 110
 - safest option 113, 115
 - safety 110
 - sequential 77–8
 - neonatal trauma risk 119
 - simulation role in training 79–80
 - skills
 - attainment 8–10
 - training 10
 - supervision 118
 - technique use 118
 - in theatre 26
 - training 8–10, 13, 118
 - trends
 - developed world 3–4
 - developing world 4–5
 - trial of in theatre 78, 115, 117
 - vaginal examination 19, 21–4, 23
 - venous thromboembolism risk 136
- overweight/obesity, maternal 97
 - fetal complications 106
- oxytocin 112
- Pajot's manoeuvre 71, 71, 72
- Papua New Guinea 5–7

Index

- parous women, delay in second stage of labour 113
- partogram 111
- patient follow-up, nonrotational forceps birth 75
- pelvic mannequins 79
- perinatal asphyxia, caesarean section at full dilatation 106
- perineal infiltration 128
- perineal injuries
 - litigation 120
 - operative vaginal birth 119
 - identification 119–20
- Pfannenstiel incision 99
- postpartum haemorrhage 105
- preterm birth, vacuum birth
 - contraindication 18
- private practice, in developing world 5
- professional behaviour
 - maintaining 40–1
- pudendal block 128
- record keeping
 - caesarean section at full dilatation 107
 - nonrotational forceps birth 75
 - operative vaginal birth 121–3
- regional anaesthesia 131
 - contraindications 133
- regional analgesia 132
 - contraindications 133
- regional blockade 131
 - postdelivery concerns 136
- resources, identification 39
- resuscitation, advanced 106
- rotational forceps birth 83–93
 - analgesia 89
 - asynclitism 85
 - complications 92–3
 - contraindications 88–9
 - occiput posterior position 85–6, 91–2
 - transverse position 84–5, 90–1
 - vaginal examination 89
 - see also Kielland's forceps
- Royal College of Obstetricians and Gynaecologists (RCOG)
 - guidelines for operative vaginal birth 115, 116, 119–20, 123
 - guidelines for repair of third- and fourth-degree perineal tears 119–20
- sepsis, caesarean section at full dilatation 105
- shoulder dystocia 26
 - traction force 80
- situational awareness 34–5
- skills
 - attainment 8–10
 - interpersonal 41
 - leadership 40–1
 - nontechnical 31–42
 - categories 34–41
 - classification 32–4
 - definition 32
 - elements 33
 - simulation training 80
 - training 10
 - see also cognitive skills, social skills
- small-for-gestational age fetus 20
- social skills 32, 33, 34, 41
- spinal anaesthesia, single-shot 131, 132, 135
- spinal–epidural analgesia, combined 135
- StationMaster tool 24
- subgaleal haemorrhage 55
- supervision
 - caesarean section at full dilatation 106

Index

- supervision (cont.)
 - forceps birth 79
 - operative vaginal birth 118
- teamwork 37–9
 - respect for team members 38–9
 - team capabilities 39
- traction
 - force 80
 - Kielland's forceps 92
 - nonrotational forceps birth 71–5, 73
 - low OP position 77
 - simulation training 80
- training 8–10, 13
 - complication prevention in second-stage caesarean section 106
 - operative vaginal birth 118
 - perineal tear diagnosis 120
 - simulation role in 79–80
- transverse position/transverse arrest 84–5, 90–1
- trauma see maternal trauma, neonates, trauma
- ultrasound
 - assessment before nonrotational forceps birth 64
 - infrapubic translabial 22
- ureter, damage in caesarean section at full dilatation 105
- uterine angle, extension 103–4
- uterine cavity, opening for caesarean section at full dilatation 100
- uterine relaxant 102
- vacuum cups 25–6
 - detachment 53–4
 - holding 46
 - insertion 46
 - distance 46, 47
 - manoeuvring toward flexion point 46–7
 - newborn infant effects 55
 - selection 45–6, 117–18
 - traction 48
 - finger–thumb position of nonpulling hand 49–50
 - finger-tip position of pulling hand 48–9
 - method 48–50
 - negative 51
 - two-handed technique 49
 - vacuum induction/maintenance 48
 - vacuum release 51
- vacuum extractors 2
 - developed world 4
 - failure rate 119
 - instrument choice 25, 25–6, 117
- vacuum-assisted birth 44–56
 - after the birth 51–2
 - baby and mother on day after 52
 - descent phase 50–1
 - difficulties 53–4
 - failure
 - rate 119
 - sequential forceps use 54
 - five steps 46–50
 - instrument choice 24
 - maternal effects 55–6
 - newborn infant impact 54–5
 - pelvic floor phase 50–1
 - principles 44–6
 - risks 25–6
 - rotational 52
 - signs of progress 51
 - three pulls rule 51
 - trends
 - developed world 3–4
 - developing world 4–5
- venous thromboembolism 136
- ventouse see vacuum extractors
- WHO checklist, caesarean section at full dilatation 98