

Cambridge University Press

978-1-107-69274-9 - Theory of Differential Equations: Part IV: Partial
Differential Equations

Andrew Russell Forsyth

Table of Contents

[More information](#)

CONTENTS.

CHAPTER XII.

GENERAL INTEGRALS OF EQUATIONS OF ORDERS HIGHER
THAN THE FIRST.

ART.	PAGE
178. Equations of the second order: the Cauchy integral, and the singular integral (if any)	1
179. The <i>general integral</i> : Ampère's definition: Darboux's definition: which is the more comprehensive?	4
180. Different kinds of integrals	8
181. Modes of occurrence of the arbitrary elements in a general integral in finite form	13
182. Two classes of equations, discriminated by the occurrence or non-occurrence of partial quadratures in the integral	16
183. Ampère's test as to whether an equation can have an integral free from partial quadratures	17
184. The number of independent arbitrary functions in the general integral, of finite form and without partial quadratures, of an equation of any order is equal to the order of the equation: their arguments	21
185. The result of § 184 in relation to Cauchy's theorem	28
186. Equation satisfied by the argument of an arbitrary function in the general integral of $f=0$, supposed free from partial quadratures: various cases	29
187. Likewise for an equation of order n	35
188. Likewise for an equation of any order in any number of independent variables	35

CHAPTER XIII.

LINEAR EQUATIONS OF THE SECOND ORDER IN TWO INDEPENDENT
VARIABLES : THE LAPLACE-TRANSFORMATIONS.

ART.		PAGE
189.	Linear equation of the second order in two independent variables is transformable to one of a couple of forms	39
190.	<i>Elliptic</i> , <i>hyperbolic</i> , and <i>parabolic</i> cases when all the variables are real	42
191.	The equation $s+ap+bq+cz=0$: its two <i>invariants</i> for the substitutions $z=\lambda z'$; effect of vanishing invariants	44
192.	Canonical forms of the equation	46
193.	The two Laplace-transformations $q+az=\sigma(z)=z_1, \quad p+bz=\Sigma(z)=Z_1,$ lead to linear equations of the second order for z_1 and for Z_1 ; they are inverse to one another	49
194.	Invariants of the equations in successive transformations	52
195.	General integral of the original equation is derivable from that of any transformed equation	56
196.	Form of an integral when a finite series of transformations leads to a vanishing invariant: <i>rank</i> of an integral	59
197.	Reducibility of rank; when possible	61
198.	Construction of equations having an integral of finite rank in either transformation	64
199.	Equations having an integral of finite rank in both transformations: characteristic number	69
200.	Simplification of the form of the results in § 199	74
201.	Darboux's alternative expressions when the integral is of finite rank in either transformation	82
202.	Application to the construction of the equations	85
203.	Also to the construction of equations having integrals of doubly-finite rank	87
204.	Goursat's theorem on the rank of an equation when $n+1$ integrals are connected by a linear relation the coefficients in which involve only one variable	90
205.	Lévy's transformation of the equation $s+ap+bq+cz=0$	94
206.	The equation $r+2ap+2\beta q+\gamma z=0$ and its invariants under transformations that conserve its form	97
207.	Forms of the equation as affected by the invariants	100
208.	The general integral of the equation in § 206, when expressed free from partial quadratures, cannot be finite in form	103
209.	Borel's expression, by means of partial quadratures, of regular integrals of an equation that are not finite in form: with examples	104

Cambridge University Press

978-1-107-69274-9 - Theory of Differential Equations: Part IV: Partial
Differential Equations

Andrew Russell Forsyth

Table of Contents

[More information](#)

CONTENTS

vii

CHAPTER XIV.

ADJOINT EQUATIONS: LINEAR EQUATIONS HAVING EQUAL INVARIANTS.

ART.		PAGE
210.	The equation adjoint to $s+ap+bg+cz=0$	111
211.	Relations between the invariants of adjoint equations in the system resulting from the Laplace-transformations	114
212.	Construction, <i>à priori</i> , of adjoint equations: and their integrals	117
213, 214.	Riemann's use of the adjoint equation to construct the Cauchy integral of the original equation: with examples	119
215.	Linear equations with equal invariants: they are self-adjoint	131
216.	Construction of equations with equal invariants	133
217.	And of their integrals, when of finite rank	136
218.	Moutard's method for equations having equal invariants	139
219.	Another form of the construction of these equations	146
220.	The process of obtaining their integrals: with examples	147

CHAPTER XV.

FORMS OF EQUATIONS OF THE SECOND ORDER IN TWO INDEPENDENT
VARIABLES HAVING THEIR GENERAL INTEGRALS IN EXPLICIT FINITE
FORM.

221.	Moutard's theorem on the classes of equations of the second order in two independent variables having general integrals represented by a single equation of finite and specified form	159
222.	Cosserat's proof of Moutard's theorem: preliminary simplifications; the equation can be transformed to $s+ap+bg+c=0$, where a, b, c are functions of x, y, z	161
223.	The coefficients a and b belong, each of them, to one of three types	164
224.	When $a=\mu+\lambda e^{2p}$, two forms are possible for b	166
225.	One form of b is $\mu'+\lambda'e^{-2p}$; the resulting equation, and its general integral, with examples	167
226.	The other form of b , when a has the value in § 224, is $b=\mu'$; the resulting equation is only a special form of the equation in § 225	178
227, 228.	When $a=\mu+\lambda z$, then b is μ' : consequent form of equation	186
229, 230.	When $a=\mu$, then $b=\mu'$: the resulting equation can be transformed either to Liouville's equation $s=e^z$, or to Laplace's linear equation	191
231.	Summary of results	195

Cambridge University Press

978-1-107-69274-9 - Theory of Differential Equations: Part IV: Partial
Differential Equations

Andrew Russell Forsyth

Table of Contents

[More information](#)

viii

CONTENTS

CHAPTER XVI.

EQUATIONS OF THE SECOND ORDER IN TWO INDEPENDENT VARIABLES
HAVING AN INTERMEDIATE INTEGRAL.

ART.		PAGE
232.	Equations of the second order in two independent variables possessing an intermediate integral	199
233.	The equation $Rr + 2Ss + Tt + U(rt - s^2) = V$: Monge's method; subsidiary equations when U is not zero	202
234.	Subsidiary equations in Monge's method when U is zero	206
235.	Completion of the primitive from the intermediate integral	208
236.	Boole's method for Monge's equations: subsidiary equations	208
237.	The subsidiary equations in Monge's method, involving differential elements, and those in Boole's method, involving differential coefficients, are equivalent	212
238.	More general method for the construction of an intermediate integral, if it exists: application to an equation of the second order of any form; examples not of Monge's form	220
239, 240.	Application of the method of § 238 to the equations in §§ 233, 234: deduction of Boole's subsidiary equations	226
241.	Conditions that the system of two subsidiary equations in § 240 should have <i>three</i> independent integrals: the primitive of the original equation	230
242.	Construction of equations satisfying conditions of § 241	236
243.	Conditions that the system of two subsidiary equations in § 240 should have <i>two</i> independent integrals: the intermediate integral, as in the methods of Monge and Boole; the possible intermediate integrals for each of the two sets of subsidiary equations	242
244.	Ampère's theorem on the equations subsidiary to the integration of an intermediate integral of an equation of Monge's type	248
245.	Conditions that the system of two subsidiary equations should possess only a single integral; in that case, the intermediate integral leads to two equations of the second order with which the given equation is consistent	253
246.	Inference when the system of two subsidiary equations possesses no integral	259
	<i>Supplementary Note</i> : the comprehensiveness of the aggregate of integrals retained in Chapter XII, illustrated by reference to an equation of Monge's type	261

Cambridge University Press

978-1-107-69274-9 - Theory of Differential Equations: Part IV: Partial
Differential Equations

Andrew Russell Forsyth

Table of Contents

[More information](#)

CONTENTS

ix

CHAPTER XVII.

AMPÈRE'S METHOD APPLIED TO EQUATIONS OF THE SECOND
ORDER IN TWO INDEPENDENT VARIABLES.

ART.		PAGE
247.	Equations of the second order in two independent variables not necessarily having any intermediate integral, but with a primitive equivalent of specified type . . .	266
248.	Ampère's method of procedure by transformation of the independent variables; selection of these in connection with the primitive: subsidiary equations . . .	267
249.	Tests as to whether the equation has a primitive of the specified type, with examples and notes . . .	269
250.	Application to the equations discussed (in Chap. xvi) by Monge's method: the subsidiary equations agree with Monge's but now have a wider significance . . .	281
251, 252.	Equations subsidiary to the construction of the primitive .	284
253.	Significance and use of the subsidiary equations in Ampère's method	289
254.	Lie's theorem on equations of the second order which possess two general intermediate integrals, with examples	295

CHAPTER XVIII.

DARBOUX'S METHOD AND OTHER METHODS FOR EQUATIONS OF THE
SECOND ORDER IN TWO INDEPENDENT VARIABLES.

	Preliminary note	302
255.	General remarks on processes of integration of partial equations	303
256, 257.	The Cauchy method, as restated by Darboux and applied to Monge's equations	305
258.	The number of subsidiary equations is less by one than the number of variables to be determined	309
259.	Why the method sometimes succeeds in providing integrals, by admitting integrable combinations	313
260.	Darboux's method of obtaining integrable combinations (if any) of higher orders	314
261.	Application of the method to obtain integrals of $f=0$ which are of the second order: two subsidiary systems: simple properties of these systems: an example	316
262.	Vályi's integration of simultaneous equations of the second order, with a note on its relation to Darboux's method	325

Cambridge University Press

978-1-107-69274-9 - Theory of Differential Equations: Part IV: Partial
Differential Equations

Andrew Russell Forsyth

Table of Contents

[More information](#)

x

CONTENTS

ART.		PAGE
263.	Lie's systems in involution	329
264.	Hamburger's application, to $f=0$, of his process of integrating simultaneous equations of the first order; the subsidiary equations	336
265.	Equations obtained by any of these methods can be combined with $f=0$; mode of using such equations	339
266.	Equations of the form $f(r, s, t)=0$ amenable to Darboux's method: the conditions	344
267.	Discussion of the conditions: examples	348
268.	When $f=0$ admits no compatible equation of the second order, Darboux's method can be applied so as to deduce compatible equations (if any) of the third order; subsidiary equations in two forms	353

CHAPTER XIX.

GENERALISATION OF INTEGRALS.

269.	Lagrange's application of the variation of parameters to generalise integrals of equations of the second order; usually it leads to a problem more difficult than the integration of the original equation	361
270.	Imschenetsky's use of Lagrange's method, keeping only three parameters; application to Monge's equation; generalising equation, and mode of procedure	366
271.	Limitations upon the form of the generalising equation when the integral to be generalised has been obtained by one or other of Ampère's subsidiary systems	374
272.	Generalisation of an intermediate integral involving two parameters	377
273.	Generalisation of a primitive of $s+ap+bg+cz=0$ which contains three parameters	379
274.	The relation between the two equations in § 273 effectively is a contact-transformation	382
275.	Alternative method of proceeding	384

CHAPTER XX.

CHARACTERISTICS OF EQUATIONS OF SECOND ORDER:

INTERMEDIATE INTEGRALS.

276.	Cauchy's problem, as ordinarily stated for $f=0$, is completely solved except along a curve $Rdy^2 - Sdx dy + Tdx^2 = 0$	388
277.	Relation of this curve to the integral surface: equations satisfied along the curve: <i>characteristics</i>	391

Cambridge University Press

978-1-107-69274-9 - Theory of Differential Equations: Part IV: Partial
Differential Equations

Andrew Russell Forsyth

Table of Contents

[More information](#)

CONTENTS

xi

ART.		PAGE
278.	Characteristics of Monge's equation; <i>orders</i> of characteristics: those of first order and of second order	393
279.	Equations of characteristics in general; different cases	396
280.	Goursat's geometrical interpretation	397
281.	Classification of equations of the second order according to their characteristics	400
282.	Characteristics of the first order and intermediate integrals	401
283.	Construction of equations having intermediate integrals in the simplest case	403
284.	Derivation of their primitive	406
285.	The critical equation: particular classes of solutions	409
286.	Two remaining important cases	413
287.	Case when $B=0$; examples	414
288.	Case when B does not vanish; examples	420
289, 290.	Unimportant cases	422

CHAPTER XXI.

GENERAL TRANSFORMATION OF EQUATIONS OF THE
SECOND ORDER.

291.	Transformation of surface-elements into one another	425
292.	Transformation of surfaces by equations for the transformation of elements: critical relation	427
293.	First case: the dependent variable satisfies two equations of the third order: consequent correspondences	429
294.	Second case: the dependent variable satisfies a single equation of the second order: consequent correspondences	430
295, 296.	Bäcklund transformations: Clairin's classification	432
297.	Transformation of equation of the second order: the critical condition for simplified relations of transformation	434
298.	When the critical condition is satisfied identically, the transformation is a contact-transformation.	436
299.	When the critical condition is not an identity, it is an equation of the Monge-Ampère type: but it is not the most general equation of that type	438
300.	Bäcklund transformations of the linear equation $s + ap + \beta q + \gamma z = 0$: the condition limiting the coefficients obtained	441
301.	Goursat's method of treating the linear equation	447
302, 303.	Two simultaneous equations of the first order in two dependent variables: classes of such equations	450

Cambridge University Press

978-1-107-69274-9 - Theory of Differential Equations: Part IV: Partial
Differential Equations

Andrew Russell Forsyth

Table of Contents

[More information](#)

xii

CONTENTS

CHAPTER XXII.

EQUATIONS OF THE THIRD AND HIGHER ORDERS, IN
TWO INDEPENDENT VARIABLES.

ART.		PAGE
304.	Equations of the third order having an intermediate integral in the form of an equation of the second order: possible cases	456
305.	Application to the equation $E(a\gamma - \beta^2) + F(a\delta - \beta\gamma) + G(\beta\delta - \gamma^2)$ $+ Aa + B\beta + C\gamma + D\delta + H = 0,$ with particular examples	459
306, 307.	Equations for which $EG - F^2 = 0$, without E, F, G all vanishing	462
308.	The linear equation $Aa + B\beta + C\gamma + D\delta + H = 0$: examples .	466
309, 310.	Equations in general having intermediate integrals . . .	470
311.	When an equation of the third order does not possess an intermediate integral, it may be soluble by an extension of Ampère's method: examples	474
312.	Darboux's method, extended and applied to equations of the third order	478
313.	Hamburger's method applied	481
314.	Connection of the subsidiary equations in §§ 312, 313 . . .	483
315.	Statement of some results for equations of order n . . .	487

CHAPTER XXIII.

EQUATIONS OF THE SECOND ORDER IN MORE THAN TWO INDEPENDENT
VARIABLES, HAVING AN INTERMEDIATE INTEGRAL.

316.	Equations, involving more than two independent variables and of order higher than the first, to be considered: sufficient to have three independent variables: notation	490
317.	Equations of second order in three independent variables having an intermediate integral: subsidiary equations: various cases	492
318, 319.	Cases when there are three subsidiary equations; the differential equation in the simplest case, and its primitives: examples	495
320.	Case next in simplicity to those in §§ 318, 319	509
321.	Equations having an intermediate integral $F(\theta, \phi, \psi) = 0$.	511
322, 323.	Construction of the intermediate integral for the equations in § 321	513
324.	The linear equation of the second order	520
325.	Equations in differential elements subsidiary to the con- struction of an intermediate integral	522

Cambridge University Press

978-1-107-69274-9 - Theory of Differential Equations: Part IV: Partial
Differential Equations

Andrew Russell Forsyth

Table of Contents

[More information](#)

CONTENTS

xiii

CHAPTER XXIV.

EQUATIONS OF THE SECOND ORDER IN THREE INDEPENDENT VARIABLES,
NOT NECESSARILY HAVING AN INTERMEDIATE INTEGRAL.

ART.		PAGE
326.	Equations of the second order in three independent variables: Cauchy's problem, as ordinarily stated for an equation $\phi=0$, is completely resolved except along the surface $Ap^2 + Hpq + Bq^2 - Gp - Fq + C = 0$	528
327, 328.	Extension of Ampère's method to equations in three independent variables: the <i>characteristic invariant</i>	530
329.	When the subsidiary equations provide any integrable combination, this is compatible with the original equation	533
330, 331.	Extension of Darboux's method to equations in three independent variables	539
332.	The aggregate of subsidiary equations is made up of two sets (i) those which are common to all equations (ii) those which belong to a particular equation	543
333.	Relation between the resolubility of the characteristic invariant and the character of the general integral	550
334, 335.	Equations having a resolvable characteristic invariant: subsidiary equations connected with each of the two resolved linear equivalents of the invariant	553
336.	The use that is made of any integral of the subsidiary systems: the alternative when they possess no integral	560
337.	Darboux's method applied to construct a compatible equation of the third order	562
338.	The subsidiary equations in the investigation of § 337 when the characteristic invariant is resolvable	567
339.	Limitations in practice when the characteristic invariant is not resolvable.	571
340.	Application to Laplace's equation $a+b+c=0$: with other examples from mathematical physics	571
341.	Whittaker's integral of Laplace's equation: its relation to the Cauchy integral	576
	Concluding remarks	582
	INDEX TO PART IV.	585