

Cambridge University Press
 978-1-108-00525-8 - The Effects of Cross and Self Fertilisation in the Vegetable Kingdom
 Charles Darwin
 Index
[More information](#)

(471)

INDEX.

ABUTILON.

A.

- Abutilon darwinii*, self-sterile in Brazil, 333, 358; moderately self-fertile in England, 344; fertilised by birds, 371
Acacia spherocephala, 404
Acanthaceæ, 96
Aconitum napellus, 428
Adlumia cirrhosa, 366
Adonis vernalis, 128; measurements, 128; relative heights of crossed and self-fertilised plants, 277; self-fertile, 365
Ajuga reptans, 368
Allium cepa (blood-red var.), 369
Anagallis collina (var. *grandiflora*), 217, 267; measurements, 218; seeds, 316, 323, 325
 Anderson, J., on the *Calceolaria*, 87; removing the corollas, 420
Anemone, 394
 Anemophilous plants, 400; often diclinous, 408
Antirrhinum majus (red var.), 363; perforated corolla, 429
 — (white var.), 363
 — (peloric var.), 363
Apium petroselinum, 172; result of experiments, 277
Argemone ochroleuca, 366
 Aristotle on bees frequenting flowers of the same species, 415
Aristolochia, 417
Arum maculatum, 417

B.

- Bailey, Mr., perforation of corolla, 427

BENNETT.

- Bartonia aurea*, 170; measurements, 170, 171; result of experiments, 277
Bartsia odontites, 369
 Beal, W. J., sterility of *Kalmia latifolia*, 359; on nectar in *Ribes aureum*, 432
 Beau, the common, 432
 Bees distinguish colours, 373; frequent the flowers of the same species, 415, 420; guided by coloured corolla, 420; powers of vision and discrimination, 422; memory, 423; unattracted by odour of certain flowers, 423; industry, 424; profit by the corolla perforated by humble-bees, 427; skill in working, 428; habit, 431; foresight, 433.
 —, humble, recognise varieties as of one species, 416; colour not the sole guide, 421; rate of flying, 424; number of flowers visited, 425; corolla perforated by, 426, 433; skill and judgment, 429
 Belt, Mr., the hair of *Digitalis purpurea*, 82; *Phaseolus multiflorus*, 151; not visited by bees in Nicaragua, 360; hummingbirds carrying pollen, 371; secretion of nectar, 403; in *Acacia spherocephalus* and passion-flower, 404; perforation of corolla, 430.
 Bennett, A. W., on *Viola tricolor*, 123; structure of *Impatiens fulva*, 367; plants flowering in winter, 384; bees frequenting flowers of same species, 416

BENTHAM.

- Bentham, on protection of the stigma in *Synaphea*, 412
Beta vulgaris, 228; measurements, 229, 230; crossed not exceeded by self-fertilised, 289, 367; prepotency of other pollen, 397
Bignonia, 363
 Birds, means of fertilisation, 371
 Blackley, Mr., on anthers of rye, 376; pollen carried by wind, experiments with a kite, 406
Boraginaceæ, 185
Borago officinalis, 185, 276; measurements, 186; early flowering of crossed, 293; seeds, 323; partially self-sterile, 362
 Brackenridge, Mr., organism of animals affected by temperature and food, 446; different effect of changed conditions, 455
Brassica oleracea, 98; measurements, 100; weight, 101, 102; remarks on experiments, 262; superiority of crossed, 288; period of flowering, 292; seeds, 322; self-fertile, 365
 —— *napus*, 393
 —— *rapa*, 393
 Brisout, M., insects frequenting flowers of same species, 419
 Broom, 163
Brugmansia, 371; humming-birds boring the flower, 432
 Bulrush, weight of pollen produced by one plant, 405
 Bundy, Mr., *Ribes* perforated by bees, 432
 Bütschli, O., sexual relations, 409

C.

- Cabbage, 98; affected by pollen of purple bastard, 378; prepotency of other pollen, 393, 397
 ——, Ragged Jack, 395
Calceolaria, 87, 369
Calluna vulgaris, 421
Campanula carpatica, 174, 364
Campanulaceæ, 174
 Candolle, A. de, on ascending a

CORYDALIS.

- mountain; the flowers of the same species disappear abruptly, 389
Canna variegata, 230; result of crossed and self-fertilised, 278; period of flowering, 294; seeds, 323, 325; highly self-fertile, 369
Cannaceæ, 230
Carduus arctoides, 402
 Carnation, 132
 Carrière, relative period of the maturity of the sexual elements on same flower, 441
Caryophyllaceæ, 130
 Caspary, Professor, on *Corydalis cava*, 331; *Nymphæaceæ*, 358; *Euryale ferox*, 365
Cecropia, food-bodies of, 404
Centradenia floribunda, 364
 Cereals, grains of, 354
 Cheeseman, Mr., on Orchids in New Zealand, 390
Chenopodiaceæ, 228
Cineraria, 335
Clarkia elegans, 169; measurements, 170; early flowering of self-fertilised, 294, 295; seeds, 316
 Cleistogene flowers, 90
 Coe, Mr., crossing *Phaseolus vulgaris*, 153
 Colgate, R., red clover never sucked by hive-bees in New Zealand, 361
 Colour, uniform, of flowers on plants self-fertilised and grown under similar conditions for several generations, 306, 307
 Colours of flowers attractive to insects, 372; not the sole guide to bees, 421
Compositæ, 173
Coniferæ, 401
Convolvulus major, 28
 —— *tricolor*, 55
 Corolla, removal of, 420; perforation by bees, 425
Coronilla, 405
Corydalis cava, 331, 358
 —— *halleri*, 331
 —— *intermedia*, 331
 —— *lutea*, 359
 —— *ochroleuca*, 359

Cambridge University Press

978-1-108-00525-8 - The Effects of Cross and Self Fertilisation in the Vegetable Kingdom

Charles Darwin

Index

[More information](#)

INDEX.

473

CORYDALIS.

- Corydalis solida*, 358
Corylus avellana, 390
 Cowslip, 219
Crinum, 395
 Crossed plants, greater constitutional vigour of, 285
 Cross-fertilisation, 370 : see Fertilisation.
 Crossing flowers on same plant, effects of, 297
Cruciferæ, 98
 Crüger, Dr., secretion of sweet fluid in *Maregraviaceæ*, 405
Cuphea purpurea, 323, 362
Cycadæ, 401
Cyclamen persicum, 215; measurements, 216; early flowering of crossed, 293; seeds, 317, 323; self-sterile, 362
 — *repandum*, 215
Cytisus laburnum, 362

D.

- Dandelion, number of pollen grains, 376
 Darwin, C., self-fertilisation in *Pisum sativum*, 161; sexual affinities, 209; on *Primula*, 219; bud variation, 298; constitutional vigour from cross parentage in common pea, 305; hybrids of *Gladiolus* and *Cistus*, 306; *Phaseolus multiflorus*, 360; nectar in orchids, 405; on cross-fertilisation, 438, 439, 457; inheritance of acquired modifications, 446; change in the conditions of life beneficial to plants and animals, 454
 —, F., structure of *Phaseolus multiflorus*, 150: *Pteris aquilina*, 404; perforation of *Lathyrus sylvestris*, 429
 —, G., on marriages with first cousins, 460
 Decaisne on *Delphinium consolida*, 129
 De Candolle, nectar as an excretion, 402

DYER.

- Delphinium consolida*, 129; measurements, 130; seeds, 322; partially sterile, 358; corolla removed, 420
 Delpino, Professor, *Viola tricolor*, 123; *Phaseolus multiflorus*, 150; intercrossing of sweet-pea, 156; *Lobelia ramosa*, 176; structure of the *Cannaceæ*, 230; wind and water carrying pollen, 371; *Juglans regia*, 390; anemophilous plants, 400; fertilisation of *Plantago*, 401; excretion of nectar, 402, 405; secretion of nectar to defend the plant, 403, 404; anemophilous and entomophilous plants, 408; dioecious plants, 414
 Denny, *Pelargonium zonale*, 142
 Diagram showing mean height of *Ipomoea purpurea*, 53
Dianthus caryophyllus, 132; crossed and self-fertilised, 133–136; measurements, 135–138; cross with fresh stock, 136; weight of seed, 139; colour of flowers, 139; remarks on experiments, 263, 274; early flowering of crossed, 292; uniform colour of self-fertilised, 309; seeds, 316, 319, 323, 325; few capsules, 360
 Dickie, Dr., self-fertilisation in *Cannaceæ*, 230
Dictamnus fraxinella, 416
Digitalis purpurea, 81; measurements, 84–87; effects of intercrossing, 85, 299; superiority of crossed, 288, 447; self-sterile, 363
Dipsacæ, 172
 Dobbs, bees frequenting flowers of same species, 416
 Dodel, Dr. A., sexual reproduction, 409
 Duhamel on *Raphanus sativus*, 393
 Dunal, nectar as an excretion, 402
 Dyer, Mr., on *Lobelia ramosa*, 176; on *Cineraria*, 335

Cambridge University Press

978-1-108-00525-8 - The Effects of Cross and Self Fertilisation in the Vegetable Kingdom

Charles Darwin

Index

[More information](#)

EARLEY.

E.

- Earley, W., self-fertilisation of *Lathyrus odoratus*, 153
 Eaton, Rev. A. E., on *Pringlea*, 408
 Engelmann, development of sexual forms, 409
 Engler, Dr., on dichogamous *Saxifraga*, 440
 Entomophilous plants, 408
Epipactis latifolia, attractive only to wasps, 375, 423
Erica tetralix, 421; perforated corolla, 426, 434
Erythrina, 360
Eschscholtzia californica, 109; measurements, 110; plants raised from Brazilian seed, 111; weight, 113; seeds, 115, 116, 315, 319, 322; experiments on, 263, 275; superiority of self-fertilised over crossed, 290; early flowering, 292, 294; artificially self-fertilised, 332; pollen from other flowers more effective, 340; self-sterile in Brazil, 343, 358
Euphrasia officinalis, 368
Euryale amazonica, 358
 — *ferox*, 365

F.

- Fabricius on *Aristolochia*, 417
Fagopyrum esculentum, 228; early flowering of crossed plant, 293
 Faivre, Professor, self-fertilisation of *Cannaceæ*, 230
 Farrer, T. H., papilionaceous flowers, 5; *Lupinus luteus*, 147; *Phaseolus multiflorus*, 150, 431; *Pisum sativum*, 150; cross-fertilisation of *Lobelia ramosa*, 176; on *Coronilla*, 405
 Fermond, M., *Phaseolus multiflorus*, 151; *P. coccineus hybridus*, 151
 Fertilisation, means of, 356; plants sterile, or partially so without insect-aid, 357-364; plants fer-

FUMARIA.

- tile without insect-aid, 365-369; means of cross-fertilisation, 370; humming-birds, 371; Australian flowers fertilised by honey-sucking birds, 371; in New Zealand by the *Anthornis melanura*, 371; attraction of bright colours, 372; of odours, 374; flowers adapted to certain kinds of insects, 375; large amount of pollen-grains, 376, 377; transport of pollen by insects, 377-379; structure and conspicuousness of flowers, 381; pollen from a distinct plant, 388; prepotent pollen, 393-400
 Fertility, heights and weights, relative, of plants crossed by a fresh stock, self-fertilised, or intercrossed (Table C), 245-252
 Fertility of plants as influenced by cross and self-fertilisation (Table D), 312; relative, of crossed and self-fertilised parents (Table E), 314-319; innate, from a cross with fresh stock (Table F), 319; relative, of flowers crossed with pollen from a distinct plant and their own pollen (Table G), 320; of crossed and self-fertilised flowers, 324, 325

Flowering, period of, superiority of crossed over self-fertilised, 291-297

Flowers, white, larger proportion smelling sweetly, 374; structure and conspicuousness of, 381; conspicuous and inconspicuous, 385; papilionaceous, 385; fertilised with pollen from a distinct plant, 388

- Forsythia viridissima*, 341
 Foxglove, 81
 Frankland, Dr., chemical affinity, 456
Fraxinus ornus, 402
Fumaria capreolata, 366
 — *officinalis*, 366

Cambridge University Press

978-1-108-00525-8 - The Effects of Cross and Self Fertilisation in the Vegetable Kingdom

Charles Darwin

Index

[More information](#)

INDEX.

475

GALIUM.

G.

- Galium aparine*, 369
Gallesio, spontaneous crossing of oranges, 394
 Galton, Mr., *Limnanthes douglasii*, 146; report on the tables of measurements, 16-19, 146, 234; self-fertilised plants, 290, 291; superior vigour of crossed seedlings in *Lathyrus odoratus*, 353, 355
 Gärtner, excess of pollen injurious, 24; plants fertilising one another at a considerable distance, 152; *Lobelia fulgens*, 179, 330; sterility of *Verbascum nigrum*, 330; number of pollen-grains to fertilise *Geum urbanum*, 377; experiments with pollen, 379
 Gentry, Mr., perforation of corolla, 427
Geraniaceæ, 142
Geranium phænum, 420
Gerardia pedicularia, 427
 Germination, period of, and relative weight of seeds from crossed and self-fertilised flowers, 352-355
Gesneria pendulina, 92; measurements, 92; seeds, 322
Gesneriaceæ, 92
Geum urbanum, number of pollen-grains for fertilisation, 377
Glaucium luteum, 366
 Godron, intercrossing of carrot, 172; *Primula grandiflora* affected by pollen of *P. officinalis*, 378; tulips, 394
 Gould, humming-birds frequenting *Impatiens*, 371
Graminaceæ, 233, 440
 Grant, Mr., bees of different hives visiting different kinds of flowers, 423
 Gray, Asa, sexual relations of trees in United States, 412; on sexual reproduction, 439

H.

- Hallet, Major, on selection of grains of cereals, 354

HOFFMANN.

- Hassall, Mr., number of pollen-grains in Peony and Dandelion, 376; weight of pollen produced by one plant of Bulrush, 405
 Heartsease, 123
Hedychium, 364
Hedysarum onobrychis, 361
 Heights, relative, of crossed and self-fertilised plants (Table A), 240-243
 Heights, weights, and fertility, summary, 238-284
 Henschel's experiments with pollen, 379
 Henslow, Rev. G., cross fertilisation in *Sarrothamnus scoparius*, 164
 Herbert on cross-fertilisation, 7; pollen brought from distant plants, 378; spontaneous crossing of rhododendrons, 394
 Hero, descendants of the plant, 47-51, 238; its self-fertilisation, 349
Heterocentron mexicanum, 361
Hibiscus africanus, 140; measurements, 140; result of experiments, 277; early flowering of crossed plant, f., 296; number of pollen-grains for fertilisation, 377
 Hildebrand on pollen of *Digitalis purpurea*, 82; *Thunbergia alata*, 96; experiments on *Eschscholtzia californica*, 110; *Viola tricolor*, 123; *Lobelia ramosa*, 176; *Fago-pyrum esculentum*, 228; self-fertilisation of *Zea mays*, 233; *Corydalis cava*, 331; *Hypecoum grandiflorum*, 331, 354; and *H. procumbens*, 331, 365; sterility of *Eschscholtzia*, 332; experiments on self-fertilisation, 340; *Corydalis lutea*, 359; spontaneously self-fertilised flowers, 366; various mechanical structure to check self-fertilisation, 382; early separation of the sexes, 400; on *Aristolochia*, 417; fertilisation of the *Gramineæ*, 440; wide dissemination of seeds, 450
 Hoffmann, Prof. H., self-fertilised capsules of *Papaver somniferum*, 108, 366; *Adonis aestivalis*, 129,

HONEY-DEW.

- 365; spontaneous variability of *Phaseolus multiflorus*, 151; self-fertilisation of kidney-bean, 152; *Papaver alpinum*, 331; sterility of *Corydalis solida*, 358; *Linum usitatissimum*, 366
 Honey-dew, 402
 Hooker, Dr., *Euryale ferox* and *Victoria regia*, each producing several flowers at once, 365; on sexual relation of trees in New Zealand, 411
 Horse-chestnut, 399
 Humble-bees, 416: *see* Bees
 Humboldt, on the grains of cereals, 354
 Humming-birds a means of cross-fertilisation, 371
 Hyacinth, 394
 Hybrid plants, tendency to revert to their parent forms, 379
Hypecomum grandiflorum, 331, 359
 — *procumbens*, 331, 366

I.

- Iberis umbellata* (var. *kermesiana*), 103; measurement, 104–106; cross by fresh stocks, 105; remarks on experiments, 262; superiority of crossed over self-fertilised seedlings, 289; early flowering, 292; number of seeds, 315; highly self-fertile, 365; prepotency of other pollen, 392
 — *amara*, 365
Impatiens frequented by humming-birds, 371
 — *barbigera*, 366
 — *fulva*, 341, 367
 — *noli-me-tangere*, 367
 — *pallida*, 341
 Inheritance, force of, in plants, 305
 Insects, means of cross-fertilisation, 371; attracted by bright colours, 372; by odours, 374; by conspicuous flowers, 383; dark streaks and marks as guides for, 372; flowers adapted to certain kinds, 375

KURR.

- Ipomoea purpurea*, 28; measurements, 29–49; flowers on same plant crossed, 41–44; cross with fresh stock, 45–47; descendants of Hero, 47–51; summary of measurements, 52; diagram showing mean heights, 53; summary of observations, 53–62; of experiments, 257–259; superiority of crossed, 289; early flowering, 291, 297; effects of intercrossing, 300; uniform colour of self-fertilised, 308; seeds, 314, 322, 324; highly self-fertile, 368; prepotency of other pollen, 397
Iris, secretion of saccharine matter from calyx, 402
Isotoma, 176, 364

J.

- Juglans regia*, 390

K.

- Kalmia latifolia*, 359
 Kerner, on protection of the pollen, 376; on the single daily flower of *Villarsia parnassifolia*, 390; pollen carried by wind, 405, 406, 412
 Kidney-bean, 152
 Kitchener, Mr., on the action of the stigma, 64; on *Viola tricolor*, 123
 Knight, A., on the sexual intercourse of plants, 7; crossing varieties of peas, 163; sexual reproduction, 439
 Kohl-rabi, prepotency of pollen, 393
 Kölreuter on cross-fertilisation, 7; number of pollen-grains necessary for fertilisation, 24; sexual affinities of *Nicotiana*, 210; *Verbascum phoeniceum*, 330; experiments with pollen of *Hibiscus vestitus*, 377
 Kuhn adopts the term cleistogene, 90
 Kurr, on excretion of nectar, 402; removal of corolla, 420

Cambridge University Press

978-1-108-00525-8 - The Effects of Cross and Self Fertilisation in the Vegetable Kingdom

Charles Darwin

Index

[More information](#)

INDEX.

477

LABIATÆ.

L.

- Labiatae*, 93
Lactuca sativa, 173, 369; measurement, 174; prepotency of other pollen, 397
Lamium album, 389, 416
 — *purpureum*, 416
Lathyrus odoratus, 153–160; measurements, 157–160; remarks on experiments, 265; period of flowering, 295; cross-fertilisation, 304; seeds, 316, 325; self-fertile, 367
 — *grandiflorus*, 155, 360
 — *nissolia*, 367
 — *sylvestris*, perforation of corolla, 429
 Lawes and Gilbert, Messrs., consumption of inorganic matter by plants, 448
 Laxton, Mr., crossing varieties of peas, 163, 305
 Lecop, *Cyclamen repandum*, 215; on *Fumariaceæ*, 359; annual plants rarely dioecious, 413
Leersia oryzoides, 350
Leguminosæ, 147; summary on the, 168
 Leighton, Rev. W. A., on *Phaseolus multiflorus*, 151; *Acacia magnifica*, 405
Leptosiphon androsaceus, 368
Leschenaultia formosa, 364
 Lettuce, 173
Lilium auratum, 341
Limnanthes douglasii, 145; measurements, 146; early flowering of crossed, 293; seeds, 316, 323; highly self-fertile, 367; prepotency of other pollen, 397
Linaria vulgaris, 9, 88; seeds, 322; self-sterile, 363
 — *cymbalaria*, 384, 422
 Lindley on *Fumariaceæ*, 359
 Link, hypopetalous nectary in *Chironia decussata*, 403
Linum grandiflorum, 343
 — *usitatissimum*, 366
Loasaceæ, 170

MEEHAN.

- Lobelia erinus*, 176; secretion of nectar in sunshine, 403; experiments with bees, 420
Lobelia fulgens, 179; measurements, 180–182; summary of experiments, 274; early flowering of self-fertilised, 291, 294, 295; seeds, 323; sterile unless visited by humble-bees, 364
 — *ramosa*, 176; measurements, 177, 178; early flowering of crossed, 293, 295; seeds, 325; self-sterile, 364
 — *tenuior*, 176
 Loiseleur-Deslongchamp, on the grains of cereals, 354
Lotus corniculatus, 361
 Lubbock, Sir J., cross-fertilisation of flowers, 6; on *Viola tricolor*, 123; bees distinguishing colours, 373; instinct of bees and insects sucking nectar, 415
Lupinus luteus, 147; measurements, 148; early flowering of self-fertilised, 294, 296; self-fertile, 367; prepotency of other pollen, 397
 — *pilosus*, 149; self-fertile, 367
Lychnis dioica, 411

M.

- Macnab, Mr., on the shorter or longer stamens of rhododendrons, 298
Mahonia aquifolium, 395
 — *repens*, 395
Malvaceæ, 140
Maregraviaceæ, 404
 Masters, Mr., cross-fertilisation in *Pisum sativum*, 161; cabbages affected by pollen at a distance, 378
 —, Dr. Maxwell; on honey-dew, 402
 Measurements, summary of, 241; Table A, 240–243; Table B, 244; Table C, 245–252
Medicago lupulina, 368
 Meehan, Mr., fertilising *Petunia violacea* by night moth, 188

Cambridge University Press

978-1-108-00525-8 - The Effects of Cross and Self Fertilisation in the Vegetable Kingdom

Charles Darwin

Index

[More information](#)

MELASTOMACEÆ.

- Melastomaceæ*, 298
Melilotus officinalis, 360
Mercurialis annua, 418
 Miller, Professor, on chemical affinity, 456
Mimulus luteus, effects of crossing, 10; crossed and self-fertilised plants, 64–70; measurements, 70–78; cross with a distinct stock, 72–75; intercrossed on same plant, 75–78; summary of observations, 78–81; of experiments, 259–261; superiority of crossed plants, 286; simultaneous flowering, 294, 296; effects of intercrossing, 301; uniform colour of self-fertilised, 307; seeds, 315, 319, 322, 324; highly self-fertile, 348, 369; potency of other pollen, 392, 397
 — *roseus*, 63
 Miner, Mr., red clover never sucked by hive-bees in the United States, 361
Mirabilis, dwarfed plants raised by using too few pollen-grains, 298; number of grains necessary for fertilisation, 377
 Mitchell, Dr., on first-cousins intermarrying, 460
Monochætum ensiferum, 364
 Moore, Mr., on Cinerarias, 335
 Müller, Fritz, on *Posoqueria fragrans*, 5, 391; experiments on hybrid *Abutilons* and *Bignonias*, 305, 306; large number of orchidaceous genera sterile in their native home, also *Bignonia* and *Tabernæmontana echinata*, 331; sterility of *Eschscholtzia californica*, 332, 342; *Abutilon darwinii*, 334; experiments in self-fertilisation, 340; self-sterile plants, 341; incapacity of pollen-tubes to penetrate the stigma, 342; cross-fertilisation by means of birds, 371; imperfectly developed male and female Termites, 380; food-bodies in *Cecropia*, 404
 Müller, Hermann, fertilisation of flowers by insects, 6, 7; on *Digitalis purpurea*, 82; *Calceo-*

NAUDIN.

- laria*, 87; *Linaria vulgaris*, 88; *Verbascum nigrum*, 89; the common cabbage, 98; *Papaver dubium*, 107; *Viola tricolor*, 123, 124; structure of *Delphinium consolida*, 129; of *Lupinus lutea*, 147; flowers of *Pisum sativum*, 160, 161; on *Sarrothamnus scoparius* not secreting nectar, 164; *Apium petroselinum*, 172; *Borago officinalis*, 185; red clover visited by hive-bees in Germany, 361; insects rarely visiting *Fumaria officinalis*, 366; comparison of lowland and alpine species, 375; structure of plants adapted to cross and self-fertilisation, 380; large conspicuous flowers more frequently visited by insects than small inconspicuous ones, 383; *Solanum* generally unattractive to insects, 387; *Lamium album*, 389; on anemophilous plants, 400; fertilisation of *Plantago*, 401; secretion of nectar, 405; instinct of bees sucking nectar, 415; bees frequenting flowers of the same species, 416; cause of it, 419; powers of vision and discrimination of bees, 422
 Müller, Dr. H., hive-bees occasionally perforate the flower of *Erica tetralix*, 427; calyx and corolla of *Rhinanthus alestrolophus* bored by *Bombus mastrucatus*, 432
 Munro, Mr., some species of *Oncidium* and *Maxillaria* sterile with own pollen, 331
Myrtaceæ, 412

N.

- Nägeli on odours attracting insects, 374; sexual relations, 409
 Natural selection, effect upon self-sterility and self-fertilisation, 345, 346
 Naudin on number of pollen-grains necessary for fertilisation, 24; *Petunia violacea*, 188

Cambridge University Press

978-1-108-00525-8 - The Effects of Cross and Self Fertilisation in the Vegetable Kingdom

Charles Darwin

Index

[More information](#)

INDEX.

479

NECTAR.

- Nectar regarded as an excretion, 402
Nemophila insignis, 182; measurements, 183–185; early flowering of crossed plant, 293; effects of cross and self-fertilisation, 303; seeds, 316, 323
Nepeta glechoma, 416
Nicotiana glutinosa, 210
— *tabacum*, 203; measurements, 205–208; cross with fresh stock, 210; measurements, 212–215; summary of experiments, 266, 267, 279; superiority of crossed plants, 288–290; early flowering, 293–295; seeds, 323, 325; experiments on, 349; self-fertile, 368
Nolana prostrata, 186; measurements, 187; crossed and self-fertilised plants, 277; number of capsules and seeds, 321, 323; self-fertile, 368
Nolanaceæ, 186
Nymphaea, 358, 365

O.

- Odours emitted by flowers attractive to insects, 374
Ogle, Dr., on *Digitalis purpurea*, 82; *Gesneria*, 92; *Phaseolus multiflorus*, 151, 360, 431; perforation of corolla, 426; case of the Monkshood, 428
Onagraceæ, 169
Onion, prepotency of other pollen, 393
Ononis minutissima, 167; measurements, 168; seeds, 323; self-fertile, 367
Ophrys apifera, 350, 369, 439
— *muscifera*, 384, 406
Oranges, spontaneous crossing, 394
Orchideæ, 364, 369; excretion of saccharine matter, 402
Orchis, fly, 406
Origanum vulgare, 94; measurements, 95; early flowering of crossed plant, 292; effects of intercrossing, 301

PHASEOLUS.

P.

- Paeony, number of pollen-grains, 376
Papaveraceæ, 107
Papaver alpinum, 331, 358
— *argemonoides*, 366
— *bracteatum*, 108
— *dubium*, 107
— *orientale*, 108
— *rhæas*, 107
— *somniferum*, 108, 331, 365
— *vagum*, 107; measurements, 109; number of capsules, 315; seeds, 358; prepotency of other pollen, 397
Papillæ of the *Viola tricolor* attractive to insects, 124
Parsley, 172
Passiflora alata, 330, 334
— *gracilis*, 171; measurements, 171; crossed and self-fertilised, 276; seeds, 323; self-fertile, 365
Passifloraceæ, 171, 357
Pea, common, 160, 351
Pelargonium zonale, 142; measurements, 143; effects of intercrossing, 301; almost self-sterile, 359
Pentstemon argutus, perforated corolla, 426, 428, 433
Petunia violacea, 188; measurements, 189–203; weight of seed, 196; cross with fresh stock, 196–201; relative fertility, 201–203; colour, 203; summary of experiments, 265, 274; superiority of crossed over self-fertilised, 289; early flowering, 293, 294; uniform colour of self-fertilised, 309; seeds, 316, 319, 323, 325; self-sterile, 362
Phalaris canariensis, 235; measurements, 236, 237; early flowering of crossed, 293
Phaseolus coccineus, 150
— *multiflorus*, 150; measurement, 152; partially sterile, 168, 360; crossed and self-fertilised, 276;

Cambridge University Press

978-1-108-00525-8 - The Effects of Cross and Self Fertilisation in the Vegetable Kingdom

Charles Darwin

Index

[More information](#)

PHASEOLUS.

- early flowering of crossed, 293; seeds, 316; perforated by humble-bees, 430
- Phaseolus vulgaris*, 153; self-fertile, 168, 367
- Pisum sativum*, 160; measurements, 162; seldom intercross, 169; summary of experiments, 264, 278; self-fertile, 367
- Plants, crossed, greater constitutional vigour, 285
- Pleroma*, 364
- Polemoniaceæ*, 182
- Pollen, relative fertility of flowers crossed from a distinct plant, or with their own, 320; difference of results in *Nolana prostrata*, 321, 323; crossed and self-fertilised plants, again crossed from a distinct plant and their own pollen, 324; sterile with their own, 330–338; semi-self-sterile, 338–340; loss of, 376; number of grains in Dandelion, Peony, and *Wistaria sinensis*, 376; number necessary for fertilisation, 377; transported from flower to flower, 377; prepotency, 391–400; aboriginally the sole attraction to insects, 402; quantity produced by anemophilous plants, 405
- Polyanthus, prepotency over cow-slip, 396
- Polygonæ*, 228
- Poqueria fragrans*, 5, 391
- Potato, 387
- Poterium sanguisorba*, 407
- Potts, heads of *Anthornis melanura* covered with pollen, 371
- Primrose, Chinese, 225
- Primula elatior*, 422, 427
- *grandiflora*, 378
- *mollis*, 368
- *officinalis*, 378
- *scotica*, 362
- *sinensis*, 225, 279; measurements, 227; early flowering of crossed, 293, 296
- *veris* (var. *officinalis*), 219; measurements, 221; result of experiments, 267, 268; early flower-

SALVIA.

- ing of crossed, 293; seeds, 317; self-fertility, 351; prepotency of dark red polyanthus, 396
- Primulaceæ*, 215
- Pringlea*, 408
- Proteaceæ* of Australia, 412
- Prunus avium*, 403
- *laurocerasus*, 404
- Pteris aquilina*, 404

R.

- Radish, 393
- Ranunculaceæ*, 128
- Ranunculus acris*, 365
- Raphanus sativus*, 365, 393
- Reinke, nectar-secreting glands of *Prunus avium*, 403
- Reseda lutea*, 117; measurements, 118, 119; result of experiments, 339; self-fertile, 365
- *odorata*, 119; measurements, 120–123; self-fertilised scarcely exceeded by crossed, 289; seeds, 316; want of correspondence between seeds and vigour of offspring, 328; result of experiments, 336; sterile and self-fertile, 358, 365
- Resedaceæ*, 117
- Rheum rhaboticum*, 401
- Rhexia glandulosa*, 364
- Rhododendron, spontaneous crossing, 394
- Rhododendron azaleoides*, 432
- Rhubarb, 394, 401
- Ribes aureum*, 432

- Riley, Mr., pollen carried by wind, 405; *Yucca* moth, 418
- Rodgers, Mr., secretion of nectar in *Vanilla*, 408
- Rye, experiment on pollen cf, 376

S.

- Salvia coccinea*, 93; measurements, 93; early flowering of crossed, 292; seeds, 315, 322; partially self-sterile, 363
- *glutinosa*, 427

Cambridge University Press

978-1-108-00525-8 - The Effects of Cross and Self Fertilisation in the Vegetable Kingdom

Charles Darwin

Index

[More information](#)

INDEX.

481

SALVIA.

- Salvia grahami*, 426, 428, 433
— tenori, 362
Sarothamnus scoparius, 163; measurements, 165–167; superiority of crossed seedlings, 285, 289; seeds, 323; self-sterile, 360
Scabiosa atro-purpurea, 172; measurements, 172, 173
Scarlet-runner, 150
Scott, J., *Papaver somniferum*, 108; sterility of *Verbascum*, 330; *Oncidium* and *Maxillaria*, 331; on *Primula scotica* and *Cortusa matthioli*, 362
Scrophulariaceæ, 63
Self-fertile varieties, appearance of, 347–351
Self-fertilisation, mechanical structure to check, 382
Self-sterile plants, 329–347; wide distribution throughout the vegetable kingdom, 341; difference in plants, 342; cause of self-sterility, 343; affected by changed conditions, 344–346; necessity of differentiation in the sexual elements, 347
Senecio cruentus, 335, 364
— heritieri, 335
— maderensis, 335
— populifolius, 335
— tussilaginis, 335
Sharpe, Messrs., precautions against intercrossing, 395
Snow-flake, 176
Solanaceæ, 188
Solanum tuberosum, 362, 387
Specularia perfoliata, 174
— speculum, 174; measurements, 175, 176; crossed and self-fertilised, 276; early flowering of crossed, 293; seeds, 323; self-fertile, 369
Spencer, Herbert, chemical affinity, 457
Spiranthes autumnalis, 391, 421
Sprengel, C. K., fertilisation of flowers by insects, 5, 6; *Viola tricolor*, 123; colours in flowers attract and guide insects, 372, 373; on *Aristolochia*, 417; *Aconi-*

URBAN.

- tum napellus*, 428; importance of insects in fertilising flowers, 455
Stachys coccinea, 426, 428, 433
Stellaria media, 367
Strachey, General, perforated flowers in the Himalaya, 434
Strawberry, 394
Strelitzia fertilised by the Nectarinidae, 371
Structure of plants adapted to cross and self fertilisation, 380
Swale, Mr., garden lupine not visited by bees in New Zealand, 150
Sweet-pea, 153

T.

- Tabernæmontana echinata*, 331, 362
Tables of measurements of heights, weights, and fertility of plants, 240–270
Termites, imperfectly developed males and females, 380
Thunbergia alata, 96, 277, 331
Thyme, 419
Tinzmann, on *Solanum tuberosum*, 362, 388
Tobacco, 203
Transmission of the good effects of a cross to later generations, 303
Trees, separated sexes, 411
Trifolium arvense, 367, 384
— incarnatum, 361
— minus, 368
— pratense, 361, 426, 434
— procumbens, 368
— repens, 361
Tropaeolum minus, 144; measurements, 145; early flowering of crossed, 293; seeds, 316, 323
— tricolor, 427; seeds, 323
Tulips, 394
Typha, 405

U.

- Umbelliferæ*, 172
Urban, Ig., fertilisation of *Medicago lupulina*, 368

2 i

Cambridge University Press

978-1-108-00525-8 - The Effects of Cross and Self Fertilisation in the Vegetable Kingdom

Charles Darwin

Index

[More information](#)

VANDELLIA.

V.

- Vandellia nummularifolia*, 90, 278; seeds, 315, 322; self-fertile, 369
Vanilla, secretion of nectar, 403
Verbascum lychnitis, 89, 341, 369
 —— *nigrum*, 89, 330, 341
 —— *phænicum*, 330, 341, 364
 —— *thapsus*, 89; measurements, 90; self-fertile, 341, 369
 Verlot on *Convolvulus tricolor*, 55; intercrossing of *Nemophila*, 183; of *Leptosiphon*, 394
Veronica agrestis, 369
 —— *chamædrys*, 369
 —— *hederifolia*, 369
Vicia faba, 360, 404
 —— *hirsuta*, 367
 —— *sativa*, 367, 403, 404
Victoria regia, 365
Villarsia parnassifolia, 390
 Vilmorin on transmitting character to offspring, 451
Vinca major, 362
 —— *rosea*, 362
Viola canina, 357
 —— *tricolor*, 123; measurements, 126, 127; superiority of crossed plants, 286, 289; period of flowering, 292, 296; effects of cross-fertilisation, 304; seeds, 316, 325; partially sterile, 358; corolla removed, 420
Violaceæ, 123
Viscaria oculata, 130; measurement, 132; average height of

ZEA.

- crossed and self-fertilised, 276; simultaneous flowering, 295; seeds, 316, 323; self-fertile, 367

W.

- Wallace, Mr., the beaks and faces of brush-tongued lories covered with pollen, 371
 Wasps attracted by *Epipactis latifolia*, 375
 Weights, relative, of crossed and self-fertilised plants, 244, 283; and period of germination of seeds, 352–355
 Wilder, Mr., fertilisation of flowers with their own pollen, 341
 Wilson, A. J., superior vigour of crossed seedlings in *Brassica campestris ruta baga*, 353
Wistaria sinensis, 376, 427

Y.

- Yucca* moth, 448

Z.

- Zea mays*, 16, 233; measurements, 16–18, 234; difference of height between crossed and self-fertilised, 288; early flowering of crossed, 293; self-fertile, 369; prepotency of other pollen, 397