

INDEX

- Abnormalities, 318
 Abortion of basal pole, 346
 Accessory strands, 156
 Acrostichoid derivatives, 232; species, 232; state, 231, 233, 238
Acrostichum aureum, 146, 239
Adiantum, embryo of, 20 (Fig. 25)
 Adult plant of *Dryopteris Filix-mas*, Rich., 26 (Fig. 31)
 Adventitious buds, 72, 79; buds of *Cystopteris bulbifera*, 72 (Fig. 66); roots, 159
Alsophila blechnoides, pinna of, 227 (Fig. 223)
A. crinita, lenticel-like pneumathodes of, 203
 Alternating generations, 21
Amphicosmia Walkeræ, 68 (Fig. 60)
 Amphiphloic solenostely, 141, 145
 Anadromic helicoid branching, 88
 Analysis, morphological, 66
 Anatomical memoirs, list of, 193
 Anatomy, vascular, 192
Anemia adiantifolia, juvenile leaves of, 85 (Fig. 77)
A. phyllitidis, young plant of, 125 (Fig. 118)
Angiopteris, stipules of, 100 (Fig. 96); wings of leaves of, 112 (Fig. 109)
A. evecta, apex of stem of, 113 (Fig. 112); sporangium of, 250 (Fig. 244)
 Angular leaf, 73
Ankyropteris Grayi, 68 (Fig. 59); stele of, 123 (Fig. 117), 182
 Annulus, 13, 14 (Fig. 17), 242, 249, 251, 255 (Fig. 250); many-rowed, 252; of *Loxosoma*, 255; position of, 254 (Fig. 250); single-rowed, 252; vestigial, 255
Anogramme chaerophylla, perennation in, 276
A. leptophylla, 33
 Antheridia (male organs), 15, 17 (Fig. 20); of various Ferns, differences in number of spermatocytes in, 291 (Fig. 281); of various Ferns seen in surface view, 293 (Fig. 283); of *Woodsia ilvensis*, 287 (Fig. 277); parallel between, and sporangia, 296; position of, 278; segmentation of, 293 (Fig. 282)
 Antheridium of *Marattia Douglasii*, 288 (Fig. 278); of *Matteuccia Struthiopteris*, 285 (Fig. 276); of *Ophioglossum pendulum*, 288 (Fig. 278)
 Apex of leaf of *Osmunda*, 111 (Fig. 108); of root of *Pteridium*, 10 (Fig. 12); of stem of *Angiopteris evecta*, 113 (Fig. 112); of stem of *Osmunda regalis*, 9 (Fig. 10); of stem of *Trichomanes radicans*, 107 (Fig. 101)
 Aphlebiae, 99 (Fig. 95)
 Apical cell, 106, 107; cell, segments of, 107, 111 (Fig. 107); meristem, 9; meristem of leaf of *Zygopteris corrugata*, 109 (Fig. 104); segmentation, 115
 Apices of pinnules of *Osmunda regalis*, var. *cristata*, 333 (Fig. 307)
 Apogamous growth in *Nephrodium pseudo-mas*, var. *polydactylum*, 325 (Fig. 302)
 Apogamy (apomixis), 24, 320, 323; in *Nephrodium pseudo-mas*, var. *cristatum*, 322 (Fig. 297); in *Pteris cretica*, 319 (Fig. 295); in *Scolopendrium vulgare*, 322 (Fig. 298)
 Apomixis (apogamy), 24, 320
 Apospory, 24, 320 (Fig. 296), 321 (Fig. 296 *conf.*), 323; in *Nephrodium pseudo-mas*, var. *cristatum*, 322 (Fig. 297)
 Appendages, dermal, 61
 Archegonia (female organs), 15; bisexual, of *Mnium cuspidatum*, 290 (Fig. 280); of *Marattia Douglasii*, 287, 288 (Fig. 278); of *Mnium*, 289 (Fig. 280); of *Nephrodium dilatatum*, 323 (Fig. 299); of *Ophioglossum pendulum*, 287, 288 (Fig. 278); of *Polypodium vulgare*, 18 (Fig. 21); position of, 278
 Archegoniophore of *Trichomanes pyxidiferum*, 282 (Fig. 272)
 Archegonium of *Matteuccia Struthiopteris*, 288 (Fig. 279); standardised, 296
 Arches, commissural, 228 (Fig. 225)
 Archesporium, 242, 257
 Archetype for Class, 336, 339
 Architecture of leaf, 61, 81
 Armature, 202
Aspidium anomalum, sori of, 225
A. trifoliatum, spores of, 261 (Fig. 257)
Asplenium nidus, epiphytic Nest Fern, 45 (Fig. 52)
A. Trichomanes, sporangia of, 245 (Fig. 239)
Asterocalamites scrobiculatus, 103 (Fig. 98)
Asterochlaena laxa, stele of, 183
Athyrium (Lady Fern), 26
A. Filix-foemina, var. *clarissima*, apospory of, 320 (Fig. 296); diploid throughout, 326

- Axial system, polycyclic, of *Saccoloma elegans*, 153 (Fig. 146)
- Axillary bud, 76; bud of *Trichomanes radicans*, 70 (Fig. 63); position, 70
- Axis, 344; arrangement of leaves on, 67; relation of leaf to, 315; vascular system of, 140
- Azolla*, 48, 261; massula of, in young state, showing glochidia, 269 (Fig. 264)
- A. filiculoides*, megasporangium of, 268 (Fig. 263)
- Basal indusium, 237; pole, abortion of, 346; wall, 19, 301
- Basipetal sorus, 212
- Bifurcated trunk of *Cyathea medullaris*, 158 (Fig. 152)
- Bisexual archegonia of *Mnium cuspidatum*, 290 (Fig. 280)
- Blechnoid fusion-sorus, disintegration of, 230 (Fig. 228)
- Blechnum*, flange in species of, 239 (Fig. 237); indusium of, 238; mucilaginous hairs of, 197 (Fig. 185)
- B. longifolium*, fusion-sorus of, 229 (Fig. 227)
- B. punctulatum*, Sw. var. *Krebsii*, 230 (Figs. 228, 229), 231 (Fig. 229)
- B. tabulare*, 29 (Fig. 35)
- Blepharoplast, 286
- Botrychioxylon*, 137
- Botrychium*, 131; young plant of, 184
- B. daucifolium*, sporangium of, 247 (Fig. 242)
- B. Lunaria*, embryo of, 306 (Fig. 292); medullation of, 125 (Fig. 119); old embryo of, 307 (Fig. 293); reconstruction of stelar structure of, 131 (Fig. 124); seedling of, 307 (Fig. 293); sporangia of, 208 (Fig. 198)
- B. obliquum*, 306; suspensor of, 305; young sporophyte of, 305 (Fig. 289)
- B. virginianum*, stele of, 137 (Fig. 129)
- Botryopteris cylindrica*, 121 (Fig. 114), 161 (Fig. 153); transverse section of stem of, 182 (Fig. 174)
- B. forensis*, equisetoid hairs of, 199 (Fig. 187); solid xylem-core of, 127
- Bracken (*Pteridium aquilinum*), meristele of, 5 (Fig. 4), 6 (Fig. 5); rhizome of, 4 (Fig. 3)
- Bramble Ferns, 39 (Fig. 44), 171
- Branching, 69, 79; anadromic helicoid, 88; catadromic helicoid, 88; dichopodial, 89; helicoid, 88; monopodial, 91; terminal, dichotomous, 337
- Bud, axillary, 76
- Budding, gametophytic, 15, 320; sporophytic, 206, 320
- Buds, 69, 70; adventitious, 72, 79; adventitious, of *Cystopteris bulbifera*, 72 (Fig. 66); extra-axillary, 70
- Bulk, proportion of surface to, 177, 181
- Cambial increase, 178
- Canal-cell, 17, 18 (Fig. 21)
- Cap-cells, 293
- Capsule of sporangium, 242
- Caspary-band, 185
- Catadromic helicoid branching, 88
- Cauline stele, 139; vascular system, 139
- Cell, apical, 106, 107; apical, segments of, 107, 111 (Fig. 107); embryonic, 300; opercular, 287 (Fig. 277); opercular, single, 293
- Cell-cleavages, embryology based upon, 299
- Cells, initial, 105, 107, 109; marginal, 109; of endodermis, 186 (Fig. 177); sporogenous, 257
- Cellular construction, 105; segmentation of embryo, 314
- Ceratopteris*, young leaf of, 9 (Fig. 11)
- C. thalictroides*, 324
- Chantransia* filaments, 342
- Cheiropleuria*, 138, 233 (Fig. 231); four-rowed stalk of, 254 (Fig. 251); proto-stele of, 122 (Fig. 115); sporophyll of, 233 (Fig. 231); two-rowed segmentation of, 246
- C. bicuspis*, 71 (fig. 65)
- Christensenia*, scales of, 201; synangia of, 249
- Chromosome-cycle, 22; normal, 324
- Chromosome-distinction between generations, 324
- Chromosomes, 21
- Cibotium Barometz*, solenostelic rhizome of, 151 (Fig. 142); vascular system of, 170 (Fig. 163)
- Cilia, 286
- Classification, early methods of, 57; phyletic, 53
- Cleavages, segmental, in embryo, 299
- Climbing leaf, 38 (Fig. 43)
- Coaxial scheme, 114 (Fig. 113)
- Coenopterideae, 162
- Coenosori, 228
- Commissural arches, 228 (Fig. 225)
- Commissures, 229
- Common vascular system, 139
- Comparison, criteria of, 59; organographic, 336
- Compensation strand, 152
- Cone, 179
- Conical form of stele, 180; form of young stem, 179; stem of *Polypodium vulgare*, 179 (Fig. 171); stem of *Pteris podophylla*, 187 (Fig. 178)
- Conjunctive parenchyma, 5
- Cordate type of prothallus, 273 (Fig. 265)
- Corrugation, 151; of solenostele of *Histiopteris incisa*, 151 (Fig. 143)

- Corynepteris Essenghi*, sorus of, 250 (Fig. 245)
 Cotyledon, 311, 344
 Cover-cell, 286
 Covering of scales, 42 (Figs. 47, 48)
 Creeping habit, 32, 33
 Crested state related to sexual irregularity, 333; varieties, 331
 Criteria of comparison, 59
 Cushion, 273
 Cutting of leaf-blade, 83
Cyathea dealbata, double-headed specimen of, Frontispiece
 C. Inrayana, 156 (Fig. 150)
 C. medullaris, bifurcated trunk of, 158 (Fig. 152)
 C. Schansin, 34 (Fig. 40)
 Cyatheaceae, 54
 Cylindrical processes, 284 (Fig. 275)
Cystopteris bulbifera, adventitious buds of, 72 (Fig. 66)
 C. fragilis, scale or ramentum of, 201 (Fig. 189)
Danaea, scales of, 201; syngangia of, 249
 D. alata, 226 (Fig. 221)
Davallia Griffithiana, 215 (Fig. 208)
D. speluncae, petiole of, 166 (Fig. 159)
 Dehiscence, slit of, shifting of, 256
 Delay of certain parts, 314
 Dendroid habit, 30; types, 29
Dennstaedtia, rhizomes of, 152 (Fig. 144)
 D. apifolia, sorus of, 255 (Fig. 252)
 D. dissecta, 221 (Fig. 215)
 D. rubiginosa, sorus of, 215 (Fig. 208)
Deparia Moorei, sori of, 225; starved, 179 (Fig. 172)
 Dermal appendages, 61; tissues, 195
 Dichopodial branching, 89
 Dichopodium, 76, 337; scorpionid, 88
 Dichotomous terminal branching, 337
 Dichotomy, 69, 79, 86, 158 (Fig. 152); equal, 90; of meristele of *Schizaea dichotoma*, 172 (Fig. 166); of rhizome, 69 (Fig. 61); sympodial, 91
Dicksonia Scheidei, young sorus of, 218 (Fig. 211), 219 (Fig. 212)
 Dictyostelic stem of Ferns, 155 (Fig. 149)
 Dictyostely, 147
 Dictyo-xylic ring, 133
 Differentiation of leaves, 49
 Diffusing surface, area of, 178
 Dimorphic leaves, 46 (Fig. 53)
 Diplodesmic condition, 233
 Diploid generation, 323; sporophyte, 21
Diplolabis Römeri, 82 (Fig. 73)
 Dipterid-derivatives, 232
Dipteris, two-rowed segmentation of, 246
 D. conjugata, young sori of, 216 (Fig. 209)
 D. Lobbiana, fertile pinna of, 227 (Fig. 222)
 Disintegration of fusion-sorus, 230 (Fig. 228); of stele, 135, 157, 189 (Fig. 180), 191
 Distal region of wall, 252
 Divergence, two-fifths, 68
 Divergent unit, 168
 Dorsiventral and dichotomous rhizome of *Lygodium scandens*, 67 (Fig. 58); rhizome of *Polypodium vulgare*, 67 (Fig. 57 bis)
 Dorsiventrality of shoot, 67
Drymoglossum subcordatum, 42 (Fig. 49)
Drynaria, 46 (Fig. 53)
 D. Filix-mas, Rich. (Male Shield Fern), 1, 26; adult plant of, 26 (Fig. 31)
 D. Linnæana, rhizome of, 27 (Fig. 32)
 D. vivipara, extra-marginal type of pinna-supply in, 172 (Fig. 168)
 Early methods of classification, 57
 Ectophloic condition, 141
 Embryo, 19 (Fig. 24), 298; as a living whole, 300; cellular segmentation of, 314; encapsulation of, 343; of *Adiantum*, 20 (Fig. 25); of *Botrychium Lunaria*, 306 (Fig. 292); of *Helminthostachys*, 304 (Fig. 288); of *Macroglossum*, 303 (Fig. 287); of Marattiaceae, 303 (Fig. 286); old, of *Botrychium Lunaria*, 307 (Fig. 293); parts of, origin of, 311; parthenogenetic, of *Marsilia Drummondii*, 325 (Fig. 303); polarity of, 300, 313; rotation of, 312; segmental cleavages in, 299; segmentation of, 301 (Fig. 284)
 Embryological method, 299
 Embryology based upon cell-cleavages, 299; endoscopic, 309; exoscopic, 309; of sporophyte, 63; theoretical problem in, 298
 Embryonic cell, 300
 Encapsulation of embryo, 343
 Endodermis, 5, 138, 180, 185; as physiological barrier, 181; cells of, 186 (Fig. 177); continuity of, 143; internal, 130, 131, 133; involution of, 145; of *Helminthostachys zeylanica*, 181 (Fig. 173)
 Endoscopic embryology, 309
 Epibasal hemisphere, 19, 311; tier, 301
 Epiphytic Ferns, 43, 44; habit, 35, 45, 46 (Fig. 53); Nest Fern, 45 (Fig. 52)
 Equal dichotomy, 90
 Equisetoid hairs of *Botryopteris forensis*, 199 (Fig. 187)
 Eusporangiate Ferns, 242, 244; Ferns, segmentation in, 111; sporangium, 270
 Exoscopic embryology, 309
 Exospore, 261 (Fig. 257)
 Extra-axillary buds, 70
 Extra-marginal pinna-trace, 173; type of pinna-supply, in *Dryopteris vivipara*, 172 (Fig. 168)
 Extrastelar pith, 188

- Factor, limiting, 181
 Female organs (archegonia), 15
 Fern, life-cycle of, 20, 21 (Fig. 28); starved by unfavourable culture, 179 (Fig. 172)
 Fern-leaf, primitive type of, 337
 Fern-Plant, 1
 Ferns, branching in, 79; epiphytic, 43, 44; Eusporangiate, 242, 244; Filmy, 41, 54; Leptosporangiate, 244; segmentation in Eusporangiate, 111; solenostelic stem of, 155 (Fig. 149); stems of, 189 (Fig. 180); differences of number of spermatocytes in, 291 (Fig. 281); antheridia of, seen in surface view, 293 (Fig. 283)
 Fern-sporophyte, primitive, 337
 Fertile leaf of *Leptochilus tricuspis*, 234 (Fig. 233); pinna of *Dipteris Lobbiana*, 227 (Fig. 222); pinnule of *Klukia exilis*, 252 (Fig. 248)
 Fertilisation in *Onoclea sensibilis*, 18 (Fig. 22)
 Filament, simple, 342; transversely septate, 301
 Filamentous construction, 342; origin, 346; type of prothallus, 275 (Fig. 266), 278, 280 (Fig. 270)
 Filmy Ferns, 41, 54
 First leaf, 311; root, 312, 345
 Fission of sorus, 227
 Flange, 239 (Fig. 237); indusial, 238
 Floating habit, 48
 Foliar gap, 142, 145
 Foot, 19, 310, 345
 Fossil Osmundaceae, 129; actual size of steles of, 184 (Fig. 176)
 Frond, 81
 Fructification of *Ptychocarpus unitus*, 210 (Fig. 201)
 Fungus, symbiotic, 284 (Fig. 274)
 Fusion of sori, 227; of veins, 64
 Fusion-sori of *Lindsaya lancea*, 228 (Fig. 225)
 Fusion-sorus, 228, 229 (Fig. 227); disintegration of, 230 (Fig. 228); of *Blechnum longifolium*, 229 (Fig. 227)
 Gametangia, 273, 285, 295; analogies between, and sporangia, 285, 289; arrangement of, 278; homology of, 289; male and female, 285
 Gametophyte, 21, 273; comparison based on, 295; haploid, 21
 Gametophytic budding, 15, 320
 Gaps, foliar, 142, 145
 Gemmae, 15; of *Trichomanes*, 280; of *Trichomanes alatum*, 281 (Fig. 271)
 Generations, alternating, 21
 Germination of spores, 15 (Fig. 18)
 Glands, water, of *Polypodium vulgare*, 205 (Fig. 196)
 Glandular hairs, 198; of *Notholaena trichomanoides*, 199 (Fig. 186)
Gleichenia, sorus of, 208
G. dicarpa, base of petiole of, 170 (Fig. 164)
G. flabellata, 142 (Fig. 132); protostele of rhizome of, 141 (Fig. 131); sori of, 209 (Fig. 200)
G. pectinata, juvenile plant of, 188 (Fig. 179); plan of stelar construction, 144 (Fig. 134); solenostelic structure of, 143 (Fig. 133); sorus of, 212 (Fig. 203); stiff hairs of, 202
 Gleicheniaceae, 54
 Glochidia, 261, 268, 269 (Fig. 264)
 Gradatae, 213
 Gradate sorus, 212
Gymnogramme japonica, vascular system of, 167 (Fig. 160)
 Habit, 26; creeping, 32, 33; dendroid, 30; epiphytic, 35, 45, 46 (Fig. 53); floating, 48; monophyllous, 31; nest, 44; scandent, 35; upright, 29; xerophytic, 41
 Habitat, 26
 Hairs, 195, 197; equisetoid, of *Botryopteris forensis*, 199 (Fig. 187); glandular, 198; glandular, of *Notholaena trichomanoides*, 199 (Fig. 186); indusial, 239; mucilaginous, of *Blechnum*, 197 (Fig. 185); on prothalli, 277 (Fig. 268); peltate, 239; simple linear, 198; stellate, of *Polypodium lingua*, 200 (Fig. 188); stiff, of *Gleichenia pectinata*, 202
 Haploid gametophyte, 21; generation, 323; sporophyte of *Lastraea pseudo-mas*, var. *crinata*, 326
 Haustorium, 310, 345
 Helicoid branching, 88
Helminthostachys, 131; embryo of, 304 (Fig. 288); root of, 175; saprophytic prothallus of, 283; suspensor of, 304; young plant of, 184
H. zeylanica, 31 (Fig. 37); endodermis of, 181 (Fig. 173); prothalli and sexual organs of, 283 (Fig. 274)
 Hemisphere, epibasal, 19, 311; hypobasal, 312
Hemitelia capensis, aplebiae of, 99 (Fig. 95)
H. grandifolia, scales of, 202 (Fig. 191)
 Heterospory, 267
Histiopteris incisa, corrugation of solenostele of, 151 (Fig. 143); young sori of, 224 (Fig. 219)
 Homology of gametangia, 289
 Homoplasia, 343
Hornea Lignieri, reconstruction of, by Kidston and Lang, 340 (Fig. 308)
 Hymenophyllaceae, 54, 115
Hymenophyllum dilatatum, 282 (Fig. 273)
H. Wilsoni, sorus of, 212 (Fig. 204)
 Hypobasal hemisphere, 312; tier, 19, 301
Hypolepis repens, pinnule of, 221 (Fig. 216); young sorus of, 222 (Fig. 217)

INDEX

353

- Increase by spores, 206; cambial, 178; in size, 188; primary, 178
- Indusial flange, 238; hairs, 239; protections, 62
- Indusium, 234, 236 (Fig. 236), 237, 238, 240; basal, 237; of *Blechnum*, 238; of *Pteris*, 238; peltate, 238; prototype of, 235; reniform, 238; types of, in Polypodiaceae (Fig. 236); vestigial lower, 222
- Initial cells, 105, 107; cells of leaves, 109
- Intercellular spaces, 180, 184
- Internal endodermis, 130, 131, 133; phloem, 145
- Intrastelar pith, 127, 188
- Inversicatenales, 162
- Involution of endodermis, 145
- Isolated tracheides, 126
- Juvenile leaf of *Todea superba*, 85 (Fig. 76); leaves of *Anemia adiantifolia*, 85 (Fig. 77); leaves of *Osmunda regalis*, 87 (Fig. 80); plant of *Gleichenia pectinata*, 188 (Fig. 179)
- Klukia exilis*, fertile pinnule of, 252 (Fig. 248)
- Lady Fern (*Athyrium*), 26
- Lastraea pseudo-mas*, var. *crispata apospora*, haploid sporophyte of, 326
- Lattice-work, 149
- Leaf, 7; angular, 73; apex of, of *Osmunda*, 111 (Fig. 108); apical meristem of, 109 (Fig. 104); architecture of, 61, 81; climbing, 38 (Fig. 43); definition of, 81; fertile, of *Leptochilus tricuspis*, 234 (Fig. 233); first, 311; juvenile, of *Todea superba*, 85 (Fig. 76); relation of, to axis, 315; vascular tissue of, 61; venation of, 61; young, of *Ceratopteris*, 9 (Fig. 11)
- Leaf-architecture, 81
- Leaf-blade, cutting of, 83; venation of, 83
- Leaf-fall, 33
- Leaf-margin, overlapping of, in *Matteuccia intermedia*, 238
- Leaf-trace, 160, 161, 162, 163 (Fig. 156); in *Thamnopteris Schlechtendalii*, 162 (Fig. 155); origin of, *Lophosoria quadrifinnata*, 163 (Fig. 156)
- Leaves, arrangement of, on axis, 67; differentiation of, 49; dimorphic, 46 (Fig. 53); initial cells of, 109; juvenile, of *Anemia adiantifolia*, 85 (Fig. 77); juvenile, of *Osmunda regalis*, 87 (Fig. 80); segmentation at margin of, 110 (Fig. 105); of Zygopterideae, 81; wings of, of *Angiopteris*, 112 (Fig. 109); young, of *Schizaea rupestris*, 220 (Fig. 213)
- Lecanopteris carnosus*, 43 (Fig. 50)
- Leptochilus tricuspis*, fertile leaf of, 234 (Fig. 233); soral region of, 233 (Fig. 232)
- Leptosporangiate Ferns, 244; sporangium, 270
- Life-history of a Fern, 1
- Limiting factor, 181; surfaces, 178
- Lindsaya lancea*, fusion-sori of, 228 (Fig. 225)
- L. linearis*, stele of, 146 (Fig. 136)
- L. scandens*, sections through node of, 147 (Fig. 137)
- Lindsaya*-condition, 146
- Linear hairs, simple, 198
- Liverworts, sporogonia of, 117
- Lonchitis pubescens*, 173 (Fig. 169)
- Lophosoria*, pinna-trace in, 174 (Fig. 170)
- L. quadrifinnata*, origin of leaf-trace in, 163 (Fig. 156)
- Loxsona*, 145; annulus of, 255; meristele of, 165 (Fig. 158)
- L. Cunninghamii*, mature sorus of, 214 (Fig. 206); plan of stelar construction of, 145 (Fig. 135); vascular system at node of (Fig. 130)
- Lygodium*, monangial sorus of, 235 (Fig. 235)
- L. circinatum*, sporangium of, 265 (Fig. 260)
- L. scandens*, dorsiventral and dichotomous rhizome of, 67 (Fig. 58)
- Macroglossum*, embryo of, 303 (Fig. 287)
- Male organs (antheridia), 15
- Male Shield Fern (*Dryopteris Filix-mas*, Rich.), 1
- Marattia*, synangia of, 249
- M. Douglasii*, antheridium of, 288 (Fig. 278); archegonia of, 287, 288 (Fig. 278)
- M. fraxinea*, sporogenous tissue of, 258 (Fig. 254)
- Marattiaceae, 55, 135; embryos of, 303 (Fig. 286); sori of, 211 (Fig. 202); suspensors of, 303
- Margin of leaves, segmentation at, 110 (Fig. 105)
- Marginal cells, 109; origin of sorus of *Pteridium*, 222; pinna-trace, 173; position of sorus, 217, 219; type of pinna-supply in *Pteris umbrosa*, 172 (Fig. 167)
- Marginales, 140
- Marsilia*, 48
- M. Drummondii*, parthenogenetic embryo of, 325 (Fig. 303)
- Marsiliaceae, 55
- Massulae, 261, 268, 269 (Fig. 264)
- Matonia*, 154
- M. pectinata*, stelar system of, 154 (Fig. 148)
- Matoniaceae, 54
- Matteuccia intermedia*, overlapping of leaf-margin in, 238; sorus of, 229 (Fig. 226)
- M. Struthiopteris*, antheridium of, 285 (Fig. 276); archegonium of, 288 (Fig. 279)

- Mature sorus of *Loxsona Cunninghamii*, 214 (Fig. 206)
 Mechanisms, opening, 253
 Medulla, 124
 Medullation, 124, 183; of *Botrychium Lunaria*, 125 (Fig. 119)
 Mega-gametangia, 290
 Megasporangium, 268 (Fig. 263), 290
 Megaspores, 267, 268 (Fig. 263)
 Memoirs, anatomical, list of, 193
 Meristele, 4; dichotomy of, of *Schizaea dichotoma*, 172 (Fig. 166); of Bracken, 5 (Fig. 4), 6 (Fig. 5); of *Loxsona*, 165 (Fig. 158)
 Meristemes in petioles, 171 (Fig. 165)
 Meristem, apical, 9; of leaf of *Zygopteris corrugata*, 109 (Fig. 104)
Metaclepsydropsis duplex, mixed pith of, 127; stele of, 128 (Fig. 122)
Metaxya rostrata = *Alsphila blechnoides*
 Micro-gametangia, 290
 Microsporangia, 290; of Salviniaceae, 267 (Fig. 262)
 Microspores, 267
 Mitosis, vegetative, 22 (Fig. 29)
 Mixed pith, 126, 184; of *Metaclepsydropsis duplex*, 127; of *Osmunda regalis*, 127 (Fig. 120)
 Mixtae, 214, 216
Mnium, archegonia of, 289 (Fig. 280)
M. cuspidatum, bisexual archegonia of, 290 (Fig. 280)
Mohria, scales of, 201
M. caffrorum, position of sporangium of, 220 (Fig. 214)
 Monangial sorus, 208, 241; of *Lygodium*, 235 (Fig. 235)
 Monophyllous habit, 31
 Monopodial branching, 91
 Morphological analysis, 66
 Morphology, stelar, 177
 Mosaics, theory of, 299
 Mucilaginous hairs of *Blechnum*, 197 (Fig. 185)
 Mycorrhizic habit, 47; prothalli of primitive Ophioglossaceae, 295; type, 283
 Neck of archegonium, 295
 Nectary of *Pteridium aquilinum*, 205 (Fig. 195)
Nephrodium dilatatum, prothalloid cylindrical process of, bearing archegonia, 323 (Fig. 299)
N. Filix-mas, *multi-cristatum*, 332 (Fig. 305)
N. Filix-mas, var. *cristatum*, raised apogamously, 332 (Fig. 306)
N. pseudo-mas, var. *cristatum*, apogamy and apospory of, 322 (Fig. 297)
N. pseudo-mas, var. *polydactylum*, apogamous growth of, 325 (Fig. 302)
Nephrolepis, stolons of, 190 (Fig. 182); tubers of, 43
Nervatio Anaxeti, 98 (Fig. 94); *Caenopteridites*, 97 (Fig. 93); *Cyrtophlebii*, 98 (Fig. 94); *Doodyae*, 98 (Fig. 94); *Drynariaceae*, 98 (Fig. 94); *Eupteridites*, 97 (Fig. 93); *Goniophlebii*, 97 (Fig. 93); *Goniopteridites*, 97 (Fig. 93); *Marginariaceae*, 98 (Fig. 94); *Neuropteridites*, 97 (Fig. 93); *Pecopteridites*, 97 (Fig. 93); *Sageniaceae*, 98 (Fig. 94); *Sphenopteridites*, 97 (Fig. 93); *Taeniopteridites*, 97 (Fig. 93)
 Nest Ferns, 45 (Fig. 52), 49
 Nest-habit, 44
 Node of *Lindsaya scandens*, sections through, 147 (Fig. 137); of *Loxsona Cunninghamii*, vascular system at, 141 (Fig. 130)
Notholaena trichomanoides, glandular hairs of, 198, 199 (Fig. 186)
 Octant-division, 300
Onoclea sensibilis, fertilisation in, 18 (Fig. 22)
 Ontogeny, 120
 Opening mechanisms, 253
 Opercular cell, 287 (Fig. 277); single, 293
 Ophioglossaceae, 55; primitive, mycorrhizic prothalli of, 295
Ophioglossum Bergianum, 136 (Fig. 128)
O. palmatum, storage-stock of, 30 (Fig. 36)
O. pedunculatum, view of Campbell on, 316
O. pendulum, antheridium of, 288 (Fig. 278); archegonia of, 287, 288 (Fig. 278)
O. simplex, 48 (Fig. 55)
O. vulgatum, sporogenous tissue of, 260 (Fig. 255)
 Organographic comparisons, 336
Osmunda, apex of leaf of, 111 (Fig. 108)
O. cinnamomea, 133 (Fig. 126)
O. regalis, apex of stem of, 9 (Fig. 10); juvenile leaves of, 87 (Fig. 80); mixed pith of, 127 (Fig. 120); root-tip of, 112 (Fig. 110), 113 (Fig. 111); sporangium of, 265 (Fig. 259); spore of, 260 (Fig. 256)
O. regalis, var. *cristata*, apices of pinnules of, 333 (Fig. 307)
 Osmundaceae, 55; fossil, 129; living and fossil, actual size of steles of, 184 (Fig. 176)
 Osmundaceous series, 128
Osmundites Carnieri, 134, 135 (Fig. 127)
O. Kidstoni, 129, 137
O. skiegatensis, 134
 Output per sorus, 266
Ouvirandra, 149
 Ovum, 17; uniformity of, 298
 Parenchyma, 195; conjunctive, 5
 Parkeriaceae, 54

INDEX

355

- Parthenogenetic embryo of *Marsilia Drummondii*, 325 (Fig. 303)
- Pellaea*, root of, 175
- P. rotundifolia*, vascular system of, 148 (Fig. 139)
- Peltate hairs, 239; indusium, 238; scales of *Polypodium incanum*, 200 (Fig. 188)
- Perennation in *Anogramme chaerophylla*, 276
- Perennials, 33
- Perforation, 149, 169
- Pericycle, 5
- Perispore, 259, 261 (Fig. 257)
- Petiole, base of, of *Gleichenia dicarpa*, 170 (Fig. 164); of *Davallia speluncae*, 166 (Fig. 159); of *Saccoloma elegans*, 169 (Fig. 162)
- Petioles, meristemes in, 171 (Fig. 165); sections of, 168 (Fig. 161); transverse sections of, 189 (Fig. 181)
- Phlebodium aureum*, 42 (Fig. 48); scales of, 202 (Fig. 190); sporangium of, 245 (Fig. 240)
- Phloem, 5; internal, 145
- Phloeoterma, 138
- Phyletic classification, 53; drift, 270; slide, 221, 222
- Phyllopodium, 82, 160, 165, 337
- Phyllorhize, 315
- Pilularia*, 48
- P. americana*, sporangium of, 256 (Fig. 253)
- P. globulifera*, 78 (Fig. 72)
- Pinna, fertile, of *Dipteris Lobbiana*, 227 (Fig. 222); of *Alsophila blechnoides*, 227 (Fig. 223)
- Pinna-supply, extra-marginal type of, in *Dryopteris vivipara*, 172 (Fig. 168); marginal type of, in *Pteris umbrosa*, 172 (Fig. 167)
- Pinna-trace, 160, 172; extra-marginal, 173; in *Lophosoria*, 174 (Fig. 170); marginal, 173
- Pinnule, fertile, of *Klukia exilis*, 252 (Fig. 248); of *Hypolepis repens*, 221 (Fig. 216)
- Pinnules, apices of, of *Osmunda regalis*, var. *cristata*, 333 (Fig. 307)
- Pith, 124; extrastelar, 188; intrastelar, 127, 188; mixed, 126, 184; mixed, of *Metaclepsydropsis duplex*, 127; mixed, of *Osmunda regalis*, 127 (Fig. 120)
- Plagiogyria*, 150; pneumatophores of, 204 (Fig. 194)
- P. pycnophylla*, rhizomes of, 148 (Fig. 138)
- Plant-body, constitution of, 60
- Plasmodium, 259
- Platyserium*, sporangium of, 254 (Fig. 251); three-rowed segmentation of, 247
- P. angolense*, sporophyll of, 233 (Fig. 231)
- Platyzoma*, 131; spores of, 265 (Fig. 258); stele of small plant of, 132 (Fig. 125)
- Pneumathode-areas, 169
- Pneumathodes, 169; lenticel-like, of *Alsophila crinita*, 203
- Pneumatophores, 203; of *Plagiogyria*, 204 (Fig. 194)
- Polarity of embryo, 300, 313; of spindle, 313
- Pole, basal, abortion of, 346
- Polybotrya*, sori of, 234 (Fig. 234)
- Polycyclic axial system of *Saccoloma elegans*, 153 (Fig. 146)
- Polycyly, 151, 155
- Polyphyletic, 343
- Polypodiaceae, 54; types of indusium in, 236 (Fig. 236)
- Polypodium bifrons*, 44 (Fig. 51)
- P. incanum*, peltate scales of, 200 (Fig. 188)
- P. lingua*, stellate hairs of, 200 (Fig. 188)
- P. vulgare*, 33 (Fig. 39); archegonia of, 18 (Fig. 21); conical stem of, 179 (Fig. 171); dorsiventral rhizome of, 67 (Fig. 57 bis); spores of, 260 (Fig. 256); water-glands of, 205 (Fig. 196)
- Polystichum angulare*, var. *pulcherrimum*, 321 (Fig. 296 cont.)
- Primary increase, 178
- Primitive Fern-sporophyte, 337; Ophioglossaceae, mycorrhizic prothalli of, 295; spindle, 313, 344; type of fern-leaf, 337; vascular construction, 338
- Primordial organs, relations of, 314
- Principal walls, 107, 112
- Principle of similar structures, 177, 191
- Procambial destination, 144, 145
- Progressive simplification of structure, 117
- Propagation, vegetative, 10
- Protections, indusial, 62
- Prothalli, hairs on, 277 (Fig. 268); mycorrhizic, of primitive Ophioglossaceae, 295; of *Helminthostachys zeylanica*, 283 (Fig. 274); of *Pteris longifolia*, 276 (Fig. 267); of *Scolopendrium*, 284 (Fig. 275)
- Prothalloid cylindrical process of *Nephrodium dilatatum*, bearing archegonia, 323 (Fig. 299)
- Prothallus, 1, 14, 16 (Fig. 19); cordate type of, 273 (Fig. 265); filamentous type of, 278, 280 (Fig. 270); form and physiology of, 62; of *Schizaea pusilla*, 279 (Fig. 269); of *Trichomanes alatum*, 281 (Fig. 271); saprophytic, of *Helminthostachys*, 283
- Protocorm, 341
- Protostele, 121, 181; of *Cheiropleuria*, 122 (Fig. 115); of rhizome of *Gleichenia flabellata*, 141 (Fig. 131); primitive vascular construction, 338; upgrade development from, 133; varying form of, 183
- Protoxylem, 6
- Protoxylem-groups, 164

- Proximal region of wall, 253
Pteridium, apex of root of, 10 (Fig. 12); marginal origin of sorus of, 222
P. aquilinum (Bracken), nectary of, 205 (Fig. 195); origin of sorus of, 223 (Fig. 218); rhizome of, 4 (Fig. 3); young plant of, 75 (Fig. 69)
Pteris, indusium of, 238
P. cretica, apogamy in, 319 (Fig. 295)
P. elata, vascular system of, 152 (Fig. 145)
P. longifolia, prothalli of, 276 (Fig. 267)
P. podophylla, 153 (Fig. 147); conical stem of, 187 (Fig. 178)
P. serrulata, young sori of, 224 (Fig. 220)
P. umbrosa, marginal type of pinna-supply in, 172 (Fig. 167)
Ptychocarpus unitus, 208 (Fig. 201); fructification of, 210 (Fig. 201)
- Radial symmetry, 66
Ramentum, 200; of *Cystopteris fragilis*, 201 (Fig. 189)
Receptacle, origin of, 218
Reniform indusium, 238
Reserve materials, 35
Rhizome, dichotomy of, 69 (Fig. 61); dorsiventral, of *Polypodium vulgare*, 67 (Fig. 57 bis); dorsiventral and dichotomous, of *Lygodium scandens*, 67 (Fig. 58); of Bracken (*Pteridium aquilinum*), 4 (Fig. 3), 34; of *Dennstaedtia*, 152 (Fig. 144); of *Dryopteris Linnæana*, 27 (Fig. 32); of *Plagiogyria pycnophylla*, 148 (Fig. 138); of *Pteridium aquilinum*, 4 (Fig. 3); protosteles of, in *Gleichenia flabellata*, 141 (Fig. 131); solenostelic, in *Cibotium Barometz*, 151 (Fig. 142)
Ring, dictyo-xylic, 133
Ring-cells, 286, 287 (Fig. 277)
Root, 175; apex of, in *Pteridium*, 10 (Fig. 12); first, 312, 345; in *Helminthostachys*, 175; of *Pellaea*, 175
Roots, adventitious, 159; of Ferns, 8; of *Pellaea*, 8 (Fig. 9)
Root-tip, 108 (Figs. 102, 103); of *Osmunda regalis*, 112 (Fig. 110), 113 (Fig. 111)
Rotation of embryo, 312
- Saccoloma elegans*, petiole of, 169 (Fig. 162); polycyclic axial system of, 153 (Fig. 146)
Sac-shaped urns, 44 (Fig. 51)
Salvinia, 48
Salviniaceae, 55; microsporangia of, 267 (Fig. 262)
Saprophytic prothallus of *Helminthostachys*, 283
Scalariform tracheides, 7
Scale, flattened, 199; or ramentum of *Cystopteris fragilis*, 201 (Fig. 189)
Scales, 195, 197; covering of, 42 (Figs. 47, 48); of *Christensekia*, 201; of *Danaea*, 201; of *Hemitelia grandiflora*, 202 (Fig. 191); of *Mokria*, 201; of *Phlebodium aureum*, 202 (Fig. 190); peltate, of *Polypodium incanum*, 200 (Fig. 188)
Scandent habit, 35
Schizaea dichotoma, dichotomy of meristele of, 172 (Fig. 166)
S. pusilla, prothallus of, 279 (Fig. 269)
S. rupestris, young leaves of, 220 (Fig. 213)
Schizaeaceae, 55
Sclerenchyma, 195, 196
Scolopendrium, 230 (Fig. 228); prothalli of, 284 (Fig. 275)
S. vulgare, 323 (Fig. 300); apogamy of, 322 (Fig. 298)
Scorpioid dichopodium, 88
Seedling of *Botrychium Lunaria*, 307 (Fig. 293)
Segmental cleavages in embryo, 299
Segmentation, apical, 115; at margin of leaves, 110 (Fig. 105); cellular, of embryo, 314; in Eusporangiate Ferns, 111; of antheridia, 293 (Fig. 282); of embryo, 301 (Fig. 284), 314; of sporangia, 110 (Fig. 106), 246, 247; primary, 301; seriation according to, 115; sporangial, 115; three-rowed, of *Platyserium*, 247; two-rowed, of *Cheiropleuria*, 246
Segments of apical cell, 111 (Fig. 107)
Selaginella, 117
Senftenbergia elegans, sporangium of, 252 (Fig. 247)
Seriation according to segmentation, 115
Sextant walls, 107, 112
Sexual irregularity, crested state related to, 333; organs, 15, 273; organs of *Helminthostachys zeylanica*, 283 (Fig. 274); organs, structure and position of, 63
Shade-form, 38
Shoot, dorsiventrality of, 67; form of, 27, 60; proportions of, 60; simple, 336; symmetry of, 67
Sieve-tubes, 5, 7; of *Pteridium*, 7 (Fig. 7)
Similar structures, principle of, 177, 191
Simple shoot, 336; sorus, 208, 209, 210, 211
Simplices, 212
Simplification of structure, progressive, 117
Siphonostele, 140
Size, 177; increase in, 188
Slide, phyletic, 221, 222
Slit of dehiscence, shifting of, 256
Solenostele, 140; corrugation of, in *Histiopteris incisa*, 151 (Fig. 143)
Solenostelic rhizome of *Cibotium Barometz*, 151 (Fig. 142); stem of Ferns, 155 (Fig. 149); structure of *Gleichenia pectinata*, 143 (Fig. 133)
Solenostely, 140, 141; amphiphloic, 141, 145
Soleno-xylic stele, 124

- Solitary sporangia, 207
- Soral region of *Leptochilus tricuspis*, 233 (Fig. 232)
- Sori, fusion of, 227; of *Aspidium anomalum*, 225; of *Deparia Moorei*, 225; of *Gleichenia flabellata*, 209 (Fig. 200); of Marattiaceae, 211 (Fig. 202); of *Polybotrya*, 234 (Fig. 234); uniseriate, 210; young, of *Dipteris conjugata*, 216 (Fig. 209); young, of *Histiopteris incisa*, 224 (Fig. 219); young, of *Pteris serrulata*, 224 (Fig. 220)
- Sorus, 12 (Fig. 14), 206, 207; basipetal, 212; constitution of, 240; fission of, 227; gradate, 212; individuality of, 240; loss of individuality of, 225, 227; marginal, 219; marginal origin of, in *Pteridium*, 222; marginal position of, 217; mature, of *Loxsonia Cunninghamii*, 214 (Fig. 206); mixed, 214; monangial, 208, 241; monangial, of *Lygodium*, 235 (Fig. 235); of *Corynepteris Essenghi*, 250 (Fig. 245); of *Dennstaedtia apiifolia*, 255 (Fig. 252); of *Dennstaedtia rubiginosa*, 215 (Fig. 208); of *Gleichenia*, 208; of *Gleichenia pectinata*, 212 (Fig. 203); of *Hymenophyllum Wilsoni*, 212 (Fig. 204); of *Matteuccia intermedia*, 229 (Fig. 226); of *Thyrsopteris elegans*, 213 (Fig. 205); origin of, in *Pteridium aquilinum*, 223 (Fig. 218); output per, 266; position of, 62, 216, 240; primitive marginal position of, 339; protection of, 234, 237; relations of, 62; simple, 208, 209, 210, 211; superficial position of, 217; young, of *Dicksonia Scheidei*, 218 (Fig. 211), 219 (Fig. 212); young, of *Hypolepis repens*, 222 (Fig. 217)
- Spaces, intercellular, 180, 184
- Spermatocytes, 16, 18 (Fig. 21), 286; differences in number of, in antheridia of various Ferns, 291 (Fig. 281); diminishing output of, 292; number of, 292
- Spermatozoid, 16, 17 (Fig. 17), 286, 298
- Sperm-cells, 286
- Sperm-numbers, 296
- Sphenophylleae, 103 (Fig. 99)
- Sphenopteris crenata*, aphanogamiae of, 99 (Fig. 95)
- Spindle, polarity of, 313; primitive, 313; primitive, theory of, 344
- Sporangia, 209 (Fig. 199); of *Botrychium Lunaria*, 208 (Fig. 198); analogies between, and gametangia, 285, 289; brevicidal, 255; formed simultaneously, 339; longicidal, 255; of *Asplenium Trichomanes*, 245 (Fig. 239); of *Thyrsopteris elegans*, 213 (Fig. 205); of *Zygopteris*, 209 (Fig. 199); parallel between, and antheridia, 296; protection of, 240; segmentation of, 110 (Fig. 106), 244 (Fig. 238); solitary, 207; two-sided initial cell of, 246; two-sided segmentation of, 246 (Fig. 241)
- Sporangial segmentation, 115; stalks, sections of, 248 (Fig. 243)
- Sporangium, 207 (Fig. 197), 214 (Fig. 207), 242; capsule of, 242; definition of, 243; development of, 13 (Fig. 16); Eusporangiate, 270; Leptosporangiate, 270; of *Angiopteris evecta*, 250 (Fig. 244); of *Botrychium daucifolium*, 247 (Fig. 242); of *Lygodium circinatum*, 265 (Fig. 260); of *Osmunda regalis*, 265 (Fig. 259); of *Phlebodium aureum*, 245 (Fig. 240); of *Pilularia americana*, 256 (Fig. 253); of *Platyterium*, 254 (Fig. 251); of *Senftenbergia elegans*, 252 (Fig. 247); of *Trichomanes speciosum*, 214 (Fig. 207); ontogenetic origin of, 243; origin of, 62; position of, in *Mohria caffrorum*, 220 (Fig. 214); segmentation of, 12 (Fig. 15); stalk of, 247; wall of, 249
- Spore of *Osmunda regalis*, 260 (Fig. 256)
- Sporeling, stele of, 180
- Spore-mother-cells, 13, 258; number of, 292; of sporangium, 242
- Spore-output, 261; actual, 270; large, importance of, 271; numerical, 62; per sporangium, 264
- Spore-producing organs, 206
- Spores, 11, 13, 20; germination of, 15 (Fig. 18); increase by, 206; of *Aspidium trifoliatum*, 261 (Fig. 257); of *Platyxoma*, 265 (Fig. 258); of *Polypodium vulgare*, 260 (Fig. 256); of sporangium, 242
- Spore-tetrad, 14
- Sporogenous cells, 257; tissue of *Marattia fraxinea*, 258 (Fig. 254); tissue of *Ophioglossum vulgatum*, 260 (Fig. 255)
- Sporogonia of Liverworts, 117
- Sporophyll of *Cheiropleuria*, 233 (Fig. 231); of *Platyterium angolense*, 233 (Fig. 231)
- Sporophylls, 11
- Sporophyte, 21; adaptation of, 49; diploid, 21; embryology of, 63; haploid, of *Lastraea pseudo-mas*, var. *crinata*, 326; primitive, 337; young, of *Botrychium obliquum*, 305 (Fig. 289); young, of *Tmesipteris*, 341 (Fig. 309)
- Sporophytic budding, 206, 320
- Stalk, four-rowed, of *Cheiropleuria*, 254 (Fig. 251); of sporangium, 242
- Stauropteris*, sporangia of, 249
- S. oldhamia*, sporangium of, 207 (Fig. 197)
- Stelar construction, plan of, of *Gleichenia pectinata*, 144 (Fig. 134); construction, plan of, of *Loxsonia Cunninghamii*, 145 (Fig. 135); morphology, 177; structure,

- reconstruction of, of *Botrychium Lunaria*, 131 (Fig. 124); system of *Matonia pectinata*, 154 (Fig. 148); system of *Stenochlaena tenuifolia*, showing perforations, 150 (Fig. 141); theory, 138
- Stele, 57, 139; cauline, 139; conical form of, 180; disintegration of, 135, 157, 189 (Fig. 180), 191; of *Ankyropteris Grayi*, 123 (Fig. 117), 182; of *Asterochlaena laxa*, 183; of *Botrychium virginianum*, 137 (Fig. 129); of *Lindsaya linearis*, 146 (Fig. 136); of *Metaclepsydropsis duplex*, 128 (Fig. 122); of small plant of *Platyzoma*, 132 (Fig. 125); of sporeling, 180; of *Trichomanes scandens*, 123 (Fig. 116); soleno-xylic, 124
- Steles, actual size of, in living and fossil Osmundaceae, 184 (Fig. 176); xylem of, outlines of, 183 (Fig. 175)
- Stellate hairs of *Polypodium lingua*, 200 (Fig. 188)
- Stem, apex of, of *Angiopteris evecta*, 113 (Fig. 112); apex of, in *Osmunda regalis*, 9 (Fig. 10); apex of, in *Trichomanes radicans*, 107 (Fig. 101); conical, of *Polypodium vulgare*, 179 (Fig. 171); conical, of *Pteris podophylla*, 187 (Fig. 178); dictyostelic, 155 (Fig. 149); solenostelic, 155 (Fig. 149); transverse section of, in *Botryopteris cylindrica*, 182 (Fig. 174); vascular tissue of, 61; young, conical form of, 179
- Stem-apices, 106 (Fig. 100)
- Stems of Ferns, 189 (Fig. 180)
- Stenochlaena sorbifolia*, fertile pinnae of, 232 (Fig. 230)
- S. tenuifolia*, stelar system of, showing perforations, 150 (Fig. 141)
- Stiff hairs of *Gleichenia pectinata*, 202
- Stipular growths, 100
- Stipules of *Angiopteris*, 100 (Fig. 96)
- Stock of *Dryopteris*, 149 (Fig. 140)
- Stolons of *Nephrolepis*, 190 (Fig. 182)
- Stomium, 13, 251, 256; of sporangium, 242; position of, 254 (Fig. 250)
- Storage and climbing, 35
- Storage-pith, 130
- Storage-stock of *Ophioglossum palmatum*, 30 (Fig. 36)
- Strand, compensation, 152
- Strands, accessory, 156
- Structure, progressive simplification of, 117
- Structures, similar, principle of, 177, 191
- Sun-form, 38
- Superficial position of sorus, 217
- Superficiales, 140
- Suppression of certain parts, 314
- Surface, diffusing, area of, 178; proportion of, to bulk of organ, 177, 181
- Surfaces, limiting, 178
- Suspensor, 300, 342; elimination of, 302; liable to be suppressed, 308; of *Botrychium obliquum*, 305; of *Helminthostachys*, 304; of Marattiaceae, 303
- Symbiotic fungus, 284 (Fig. 274)
- Symmetry of shoot, 67; radial, 66
- Sympodial dichotomy, 91
- Synangia, 234; of *Christensenia*, 249; of *Danaea*, 249; of *Marattia*, 249
- Synangium, 208, 242
- Synapsis, 22
- Syngamy, 298
- Tapetum, 13, 258; of sporangium, 242
- Terminal branching, dichotomous, 337
- Tetrad-division, 23 (Fig. 30), 258
- Thamnopteris Schlechtendalii*, 130; leaf-trace in, 162 (Fig. 155)
- Thyrsopteris elegans*, 153; sorus of, 213 (Fig. 205); sporangia of, 213 (Fig. 205)
- Tier, epibasal, 301; hypobasal, 19, 301
- Tissue, sporogenous, of *Ophioglossum vulgatum*, 260 (Fig. 255)
- Tissues, dermal, 195
- Tmesipteris*, young sporophyte of, 341 (Fig. 309)
- Todea barbara*, Frontispiece; sporangium of, 251 (Fig. 246)
- T. superba*, juvenile leaf of, 85 (Fig. 76)
- Tracheides, 6; isolated, 126; of *Pteridium*, 7 (Fig. 6); scalariform, 7
- Tree-Ferns, 54
- Trichomanes*, gemmae of, 280
- T. alatum*, prothallus and gemmae of, 281 (Fig. 271)
- T. pyxidiferum*, archegoniophore of, 282 (Fig. 272)
- T. radicans*, apex of, 107 (Fig. 101); axillary bud of, 70 (Fig. 63)
- T. reniforme*, 86 (Fig. 78)
- T. rigidum*, filamentous prothallus of, 280 (Fig. 270)
- T. scandens*, stele of, 123 (Fig. 116)
- T. spectosum*, sporangium of, 214 (Fig. 207)
- T. venosum*, Frontispiece
- Trunk, bifurcated, of *Cyathea medullaris*, 158 (Fig. 152)
- Tubers of *Nephrolepis*, 43
- Tubicaulis solenites*, 128 (Fig. 121), 161 (Fig. 154)
- Two-fifths divergence, 68
- Typical number, 262
- Uniseriate sori, 210
- Upgrade development from protostele, 133
- Urns, sac-shaped, 44 (Fig. 51)
- Variability, 271
- Vascular anatomy, 192; construction, primitive, 338; skeleton (Fig. 1), 2; system, 120; system, cauline, 139; system,

INDEX

359

- common, 139; system at node of *Loxsona Cunninghamii*, 141 (Fig. 130); system of *Cibotium* (*Dicksonia*) *Barometz*, 170 (Fig. 163); system of *Gymnogramme japonica*, 167 (Fig. 160); system of axis, 140; system of *Pellaea rotundifolia*, 148 (Fig. 139); system of *Pteris elata*, 152 (Fig. 145); tissue of leaf, 61; tissue of stem, 61
- Vegetative mitosis, 22 (Fig. 29); propagation, 10
- Veins, fusion of, 64
- Venation, 160, 174; of leaf, 61; of leaf-blade, 83
- Venter, 295
- Ventilating areas, 203; system, 184
- Ventilation, 130
- Ventral-canal-cell, 17
- Vestigial annulus, 255; lower indusium, 222
- Vestiture, 57
- Wall, basal, 19, 301; distal region of, 252; proximal region of, 253
- Walls, principal, 107, 112; sextant, 107, 112
- Water-glands of *Polypodium vulgare*, 205 (Fig. 196)
- Water-storage, 43
- Webbing, 83, 102; progressive, 85
- Wings of leaves of *Angiopteris*, 112 (Fig. 109)
- Woodsia ilvensis*, antheridia of, 287 (Fig. 277); filamentous type of, 275 (Fig. 266)
- Woodwardia*, 11 (Fig. 13)
- Xerophytic habit, 41
- Xylem, 6; of steles, outlines of, 183 (Fig. 175)
- Xylem-core, solid, of *Botryopteris forensis*, 127
- Xylic-gaps, 143
- Xylic-perforations, 149
- Young leaf of *Ceratopteris*, 9 (Fig. 11); leaves of *Schizaea rupestris*, 220 (Fig. 213); plant of *Anemia phyllitidis*, 125 (Fig. 118); plant of *Botrychium*, 184; plant of *Helminthostachys*, 184; plant of *Pteridium aquilinum*, 75 (Fig. 69); sori of *Dipteris conjugata*, 216 (Fig. 209); sori of *Histiopteris incisa*, 224 (Fig. 219); sori of *Pteris serrulata*, 224 (Fig. 220); sorus of *Dicksonia Scheidei*, 218 (Fig. 211), 219 (Fig. 212); sorus of *Hypolepis repens*, 222 (Fig. 217); sporophyte of *Botrychium obliquum*, 305 (Fig. 289); sporophyte of *Tmesipteris*, 341 (Fig. 309); stem, conical form of, 179
- Zaleskia*, 129
- Zygopterideae, leaves of, 81
- Zygopteris*, sporangia of, 209 (Fig. 199)
- Z. corrugata*, apical meristem of leaf of, 109 (Fig. 104)
- Zygote, 19, 20, 298