

Cambridge University Press

978-1-108-01732-9 - Water Plants: A Study of Aquatic Angiosperms

Agnes Arber

Index

[More information](#)

[422]

INDEX

[The names of authors which occur in the bibliography are not included in the following index, since page references are given in connexion with the titles in the bibliography, which thus also serves as an index of authors' names]

- Acacia* phyllode, 340
Achillea ptarmica, 5, 199
 Acquired characters, inheritance of, 333, 334
 Adaptation, 171, 332-335
Aedemone mirabilis. See *Herminiera elaphroxylon*
 Aerating system, in tissues of hydrophytes, **183-194**, **256-259**; of root, 185-187; of stem, primary, 183-185; of stem, secondary, 187-194
Aerenchyma, 187-194; from cambium, 191-192; from phellogen, 187-191, 193-194
Aeschynomene, aerenchyma, 191, 192
Aeschynomene aspera, 191
Aeschynomene hispidula, 191-192
 Affinities of hydrophytes, **308-321**
 Africa, 213, 295, 298, 305
 "Age and Area" in plant distribution, **305-307**
 Air spaces, lysigenous, 184, 185
 Air spaces, schizogenous, 184, 185
Aldrovandia, affinities, 310; carnivorous habit, 110-111, 270; embryo, 110; fruit ripening under water, 239; roots, absence of, 109, 110, 204, 244; seed, 110; sensitive leaves, 110, 111 (Fig. 75); shade plant, 289; stem anatomy, 175; turions, 110, 219
Aldrovandia vesiculosa, 8, **109-111** (Fig. 75), 289, 310
 Algae, 113, 114, 123, 124, 142, 155, 172
 Aliens, 303
Alisma, effect of freezing on fruit, 243; fruit, 242; germination and rupture of seed coats, 244; heterophylly, 19, 20; land and water plants, 153 (Figs. 101 and 102); ranalean features, 320
Alisma graminifolium, 19, 20, 23, 157, 280
Alisma natans, 234
Alisma Plantago, 19, 20, 23, 151, 153 (Figs. 101, 102), 156, 169, 242, 243, 244, 289, 297
Alisma ranunculoides. See *Echinodorus ranunculoides*
Alismaceae, 5, **9-23**, 24, 33, 151, 156, 195, 224, 248, 297, 313, 314, 319, 337, 346
Alocasia, 303
 Aloe, Water. See *Stratiotes aloides*
 Alps, 290
Alihena, bracts, 316; in brackish water, 134; perigonium, 316; reduced stem anatomy, 63, 173
Alihena filiformis, 173
 Altitude above sea-level, 289-291
 Amazons, 31, 99, 113, 229
 "Ambatsch," 192
Ambulia, affinities, 313; heterophylly, 151
Ambulia hottonoides, 151
 America, 61, 108, 120, 190, 193, 210, 216, 286, 290, 295, 298, 312, 313
 American Indians, 17, 118
Ammania, 303
 Amphibious plants, effect of water upon, 201, 202
 "*Amphibolis zosteraefolia*," 123
 "*An Idea of a Phytological History*," 230
Anacharis. See *Elodea*
 Andes, 291
 Anemophily. See Pollination, anemophilous
 Anemophytes, 143
 Angiosperms, Marine. See Marine Angiosperms
 Anthocyanin, 15, 17, 113, **276-278**
 Apical openings in leaves, of *Callitriche*, 268 (Fig. 163); *Heteranthera*, 268; *Littorella*, 269; *Pistia*, 82 (Fig. 53); *Potamogeton*, 167 (Fig. 108), 268, 269; Potamogetonaceae, 133; *Zostera*, 269
Aponogeton, affinities, 314; disarticulation of primary root, 244; distribution, 305; fenestration, 142 (Fig. 91), 143; geotropism, 281; heliotropism, 281; heterophylly, 154; undulated leaf, 62
Aponogeton angustifolius, 143
Aponogeton Bernerianus, 142
Aponogeton distachyus, 215, 244, 281
Aponogeton fenestratis, 142 (Fig. 91), 143, 281, 314
Aponogeton ulvaceus, 62
 Aponogetonaceae, 239, 248, 305, 313, 314, 315
Aquilegia, 314
 Araceae, 74, 82, 314, 315

INDEX

423

- Araguay, River, 295
 Argentine, 55
 Aroideae, 316
 Arrowgrass. See *Aponogeton*
 Arrowhead. See *Sagittaria sagittifolia*
 Asia, 295, 298
 Astrakhan, 303
Auricula, polystely in, 180, 181, 182
 Australia, 295, 305
 Awlwort. See *Subularia aquatica*
 Azores, 295, 333
- Bacteria, 142
 Baltic, 123
 Baltimore, 253
 Bananas, 143
 Band leaves, 11 (Fig. 3), 12, 13 (Fig. 4), 14 (Fig. 5), 19, 20, 22, 23, 140, 141 (Fig. 90), etc.
Barclaya, 33
 Bateson, W., on evolution, 334
 Batrachian Ranuncul. See *Ranunculus* sect. *Batrachium*, *Ranunculus aquatilis*, etc.
Batrachospermum, 155
 Bean, 249
 Beetles, as pollinators of Lemnaceae, 80; in utricle of *Utricularia*, 93
Begonia hydrocotylifolia, 256
 Belgium, 303
Bellis perennis, 165
 Bengal, 110
 Bermudas, 298
Bidens Beckii, 151, 313
 Biological classification of hydrophytes, 4–8, 42
 Birds and dispersal, 35, 298–302
 Bittersweet. See *Solanum dulcamara*
 “Bitter-sweet,” Grew on heterophyllly in, 155
 Black Sea, 302
 Bladderwort. See *Utricularia*
 Bladderwort, Common. See *Utricularia vulgaris*
 Blue Nile, 192
 Bodensee, 322
Boottia, 57
Bostrychia Moritziana, 114
Brasenia peltata. See *B. Schreberi*
Brasenia Schreberi, 38 (Fig. 20), 205, 272
 Brazil, 206, 207, 243, 295
 Brent Geese, 302
 Broads, 288
Brocchinia cordylinoides, 109
 Bromeliaceae, 108, 109
 Bruch-Eicheln, 17
 Brunfels, Otto, 27
 Bull Nut. See *Trapa natans*
Bulliarda (Tillaea), affinities, 310; aquatic with xerophilous ancestry, 310; cleistogamy, 234
Bulliarda (Tillaea) aquatica, 234, 310
- Burton-on-Trent, 211
 Butler, Samuel, 347
 Butomaceae, 157, 248, 313
Butomus, 314
 Buttercup, Water. See *Ranunculus aquatilis*, *Ranunculus* sect. *Batrachium*, etc.
- Cabomba*, anatomy, 37, 38; heterophyly, 29 (Fig. 14), 146; polystely, 37; reduced leaves, 338
Cabomba caroliniana, 338
 Cabomboideae, 38, 309
 Caddice worms, 217
 Calcareous substratum, 286, 287
Caldesia, heterophyly, 23; turions, 22, 225 (Figs. 148, 149)
Caldesia parnassifolia, 22, 23, 224, 225 (Figs. 148, 149)
 California, 123
Calla palustris, 167 (Fig. 107)
 Callitricheaceae, 134, 311, 318
Callitriche, affinities, 311, 312; altitude, 290; annual and perennial forms, 215, 216; as coloniser, 299; chlorophyll, absence in epidermis, 164; distribution, 306, 307; flowers, 237 (Fig. 154); fruit, 242, 243; germination, 280; heterophyly, 146, 147 (Fig. 94); land form, 170 (Fig. 111), 195; leaf anatomy, 163, 169, 170 (Fig. 111); local races, 330; mucilage trichomes, 271; pollination, 236, 237; roots, air spaces in, 187; root anatomy, 208, 209 (Fig. 138); seeds, 297; stomates, 166; vascular strand of axis, 175, 176 (Fig. 114); vegetative reproduction, 216; water pores, 267, 268 (Fig. 163)
Callitriche autumnalis, 6, 134, 169, 237, 268 (Fig. 163), 307
Callitriche stagnalis, 176 (Fig. 114), 208, 209 (Fig. 138), 271
Callitriche verna, 6, 146, 147 (Fig. 94), 163, 166, 169, 170 (Fig. 111), 187, 236, 237 (Fig. 154), 306
Caltha palustris, 198, 199 (Fig. 129)
 Cam, River, 150, 211, 263
 Cambridge, 150
 Cambridge Botanic Garden, 211
 “Camichi,” 300
 “Cammomill,” 144
 Campanulaceae, 313
 Canadian Waterweed. See *Elodea canadensis*
 Canary Islands, 295
Canna, 244
 Carbon dioxide, derived from substratum, 254; excess of, available for hydrophytes, 254; proportion of, in free and dissolved air, 253
Cardamine, adventitious budding from leaves, 216, 217 (Fig. 141); land and water forms, 201, 202 (Fig. 133)

- Cardamine pratensis*, 201, 202 (Fig. 133), 216, 217 (Fig. 141), 309
- Carnation, 155
- Carolina, 286
- Carrot, 249
- Caryophyllaceae, 234, 310, 311
- Caspian Sea, 302
- Castalia*, air leaves, 32; effect of frost on seeds, 243; floating leaves, 30, 146, 159; geophytic habit, 217, 323; length of peduncle and petiole, 31 (Fig. 15), 40; pigmented variety, 276; rhizome, 24–26 (Fig. 11), 39, 217; seedlings, 28 (Fig. 13), 29; seeds, 302; stipules, 25, 26 (Fig. 11); submerged leaves, 29, 146, 159; terrestrial form, 32
- Castalia alba*, 24, 25, 26 (Fig. 11), 27, 28 (Fig. 13), 29–31 (Fig. 15), 32, 243, 276, 302
- Castalia flava*, 37
- Castalia Lotus*, 34, 36, 37 (Fig. 19), 225
- Castalia pygmaea*, 33
- Castelnavia*, 117, 295, 306
- Caucasus, 302
- Ceratophyllaceae, 84–90, 318, 320
- Ceratophyllum demersum*, 84–90; affinities, 84, 309, 312; cuticularisation, 86; distribution, 295, 297, 298; effect of strong illumination, 279; epiphytic fauna, 88; flowers, 84, 85 (Fig. 54); freezing, effect of, 88, 89; hairs containing mucilage, 86, 272; high temperatures necessary for fruiting, 88, 275; in biological classification, 8; in deep water, 86, 288; leaf, anatomy of, 168, dimensions of, 140, juvenile, 86 (Fig. 55); luxuriance, 87; monoecism, 84, 85; movements, 90, 281; mucilage hairs, 86, 272; perenniality, 215; pollination, hydrophilous, 84, 85, 134, 237, 238; rhizoid branches, 88, 89 (Figs. 57 and 58), 98, 336, 337; roots, absence of, 85, 204, 244; seedling, 85, 86 (Fig. 55); stem anatomy, 86, 87 (Fig. 56); vegetative reproduction, 87, 216, 219; water absorption, 269, 270; water content, 86
- Ceylon, 112, 216
- Chalky incrustation on leaves of aquatics, 51
- Chantransia*, 155
- Chaya*, 288
- Chili, 181
- Chimborazo, 290
- China, 17
- Chlorophyll, in epidermis of aquatics, 164, 168 (Fig. 109), 169, 254; in epidermis of terrestrial plants, 164, 165
- Chydorus sphaericus*, as food of *Utricularia*, 94
- Cirsium anglicum*, 198, 199 (Fig. 130)
- Classification, biological, of hydrophytes, 4–8, 42
- Cleistogamy, 233, 234; in *Alisma*, 234; *Bulliarda* (*Tillaea*), 234; *Echinodorus*, 234; *Euryale*, 34, 234; *Heteranthera*, 234 (Fig. 153); *Hydrothrix*, 234; *Illecebrum*, 234; *Limosella*, 233, 235; *Nesaea*, 234; *Pephis*, 234; *Podostemon*, 121 (Fig. 82), 234; *Ranunculus*, 233, 234; *Rotala*, 234; *Tillaea*, 234; *Trappella*, 234
- Climate and life-cycles, 275
- Cnicus arvensis*, 200
- Cnicus pratensis*, 198, 199, 200
- Codium tomentosum*, 123
- “Collet,” development of root-hairs from, 245
- Colocasia*, 303
- Colonisation of waters, 289, 298, 299
- Commissioners’ Pits, Upware, 217
- Competition, aquatic life as a refuge from, 324, 325
- Compositae, 151, 313, 320, 321
- Conifers, “Youth forms” of, 155
- Copepods, as food of *Utricularia*, 94
- Cotton, aliens accompanying, 303
- Cotula myriophylloides*, 313
- Crassulaceae, 234, 310
- Cruciferae, 216, 309
- Crustacea, as food of *Utricularia*, 93, 94
- Cuscuta alba*, 5, 199 (Fig. 131)
- Cuticle, slight development of, in hydrophytes, 163, 254, 260
- Cuyuni River, 119
- Cyclamen*, 240
- Cymodocea*, absence of apical openings in leaves, 269; anatomy, 125 (Fig. 84), 131, 331; chlorophyll in epidermis, 164; flowers, 126; grappling apparatus of fruit, 127, 245; habit, 124 (Fig. 83), 125; leaf form, 124; life-history, 124 (Fig. 83)–127; pollination, hydrophilous, 126, 237; regarded as Alga, 123; spiral roots, 205; *squamulae intravaginales*, 126; vivipary, 127, 246
- Cymodocea aequorea*, 124 (Fig. 83), 125 (Fig. 84), 126, 127, 246
- Cymodocea antarctica*, 123, 127, 205, 245
- Cymodocea isoetifolia*, 124
- Cyperaceae, 154, 317, 326
- Cypris*, as food of *Utricularia*, 94
- Damasonium stellatum*, 23
- Danube, 212
- Daphnidae, as food of *Utricularia*, 94
- Darwin, Erasmus, on evolution, 334
- Delayed germination in aquatics, 36, 71, 72, 243–244
- Delesseria Leprieurii*, 114
- “Dents nageoires,” 133, 314, 315
- Depth to which plants can grow, 86, 123, 275
- Desmanthus natans*. See *Neptunia oleracea*

Cambridge University Press

978-1-108-01732-9 - Water Plants: A Study of Aquatic Angiosperms

Agnes Arber

Index

[More information](#)

INDEX

425

- Diaphragms, 18, 19 (Fig. 8), 183, 184 (Figs. 118, 119), 257
- Dicotyledonous families, proportion of among aquatics, 322
- Dicraea elongata*, 115 (Fig. 77)
- Dicraea stylosa*, 114, 115 (Fig. 78), 116 (Fig. 79), 118
- Dictyota dichotoma*, 123
- "Differentiation Theory" of plant distribution, 304–307
- Dionaea*, 111
- Diplanthera*, 123
- Distribution, geographical, 73, 112, 295–307
- Dock, 271
- Dodder. See *Cuscuta alba*
- Dodoens' *Histoire des Plantes*, 144
- Dollo's "Law of Irreversibility," 336, 347
- Droseraceae, 109, 111, 310
- Duckweed. See *Lemna*, Lemnaceae, *Spirodela*, *Wolffia*
- Duckweed, Rootless. See *Wolffia*
- East Anglia, flood of 1912 in, 296
- Echinodorus ranunculoides*, cleistogamy, 234; heterophylly, 23; inflorescence and vegetative shoots, 224 (Fig. 147); land and water forms, 21 (Fig. 9)
- Ecology, 285–292
- Egypt, 332
- Egyptian cotton, aliens accompanying, 303
- Eichhornia*, affinities, 317; air tissue in petioles, 154; heterophylly, 154, 160, 161; phylloclad interpretation of leaf, 341 (Fig. 169), 342 (Fig. 170), 343, 344; vegetative multiplication, 213
- Eichhornia azurea*, 160, 161
- Eichhornia crassipes*, 154
- Eichhornia speciosa*, 213, 341 (Fig. 169), 342 (Fig. 170), 343, 344
- Elatinaceae, 310, 311
- Elatine*, affinities, 311; annual species, 215; conveyance by birds, 301; root-hairs from "collet," 245 (Fig. 158)
- Elatine hexandra*, 245 (Fig. 158), 311
- Elatine hydrophyper*, 301, 311
- Eleocharis*, 286
- Elisma*, heterophylly, 23; relation of inflorescence and vegetative shoot, 20
- Elisma natans*, 20, 23
- Ellis, Lake, 286
- Elodea*, aerating system, 256; circulation of protoplasm, 212; experimental cultures, 265; history in Britain, 55, 210–213; leaf anatomy, 165 (Fig. 106), 169; leaf form, 141; pollination mechanism, 55, 56 (Fig. 35), 57, 134, 236; root-hairs, 208; vegetative reproduction, 55, 210–213; wintering shoots, 55 (Fig. 34), 219
- Elodea callitrichoides*, 55, 56, 236
- Elodea canadensis*, 6, 7, 55 (Fig. 34), 57, 165 (Fig. 106), 169, 173, 210–213, 219, 253, 254, 265, 266
- Elodea densa*, 57, 236
- Elodea ioensis*, 55, 56 (Fig. 35)
- Embryo, macropodous, 246 (Fig. 159), 248 (Fig. 161), 249, 319 (Fig. 166), 326; protection of, 242, 243; viviparous, 127
- Engadine, 290
- Enhalus*, 57, 123, 124, 131, 236
- Enteromorpha*, 123
- Entomophily. See Pollination, entomophilous
- Epidermis of aquatics, chlorophyll in, 164, 168, 169, 254; form of cells of, 163, 164
- Epilobium hirsutum*, 188
- Epithem, 267
- Eranthis hiemalis*, 319
- Eriocaulon*, 286
- "Eu-anthostrobilus," 315
- Eu-callitriche*, 236, 237, 306, 307, 330
- Eucalyptus Preissiana*, 256
- Eupatorium cannabinum*, 188
- Euphorbiaceae, 311
- Europe, 290, 295
- Euryale*, affinities, 38; cleistogamy, 34, 234
- Euryale ferox*, 34, 234
- Farinosae, 316, 317, 341
- Farmesia metzgerioides*, 114, 248
- Fenestration of leaves, 142 (Fig. 91)
- Fijis, 298, 303
- Flagellates, 142
- "Floating-leaf association," 288
- Floating leaves, 30–32, 44–46, *et passim*
- Floating Sensitive Plant. See *Nepenthes oleracea*
- Floating wood of *Herminiera*, 192
- Florida, 213
- "Flossenzähne," 133, 314
- Flowers of water plants, 227–238, etc.
- Fluviales. See Helobieae
- Fly pollinating *Sagittaria*, 9
- "Flying germinators," Water-fowl as, 301
- Fontinalis*, 225
- Food plants, Alismaceae, 17; *Alocasia*, 303; *Colocasia*, 303; Nymphaeaceae, 24; *Trapa*, 302
- Forest of Dean, 276
- Freezing, effect of, 220, 243, 278
- Frogbit. See *Hydrocharis morsus-vanae*
- Fruits of water plants, 239–249, etc.
- Fruits, ripening under water, 239, 240
- Fungi, 172
- Gaseous exchange, 253–259
- Geneva, Lake of, 278
- Gentianaceae, 39, 205, 312
- Geographical distribution, 295–307, etc.

- Geophytes, aquatic, 217, 323
 Geotropism, 281, 282
 Germany, 17, 53, 290, 299
 Germination, delayed, 36, 71, 72, **243**, **244**; *in situ*, 80, 127, 246, 248; of various genera, see heading, Seedlings
 Giant Waterlily. See *Victoria regia*
 Gill-tufts of *Oenone*, 118, 119 (Fig. 81), 255
Glechoma hederacea, 5
Glyceria, aquatic grass, 317, 318; distributed by birds, 301
Glyceria aquatica, 317, 318
Glyceria fluitans, 301, 317
 "Gramen bulbosum aquaticum." 11
 Gramineae, 317, 318, 326
 Grass-wrack. See *Zostera*
 Grasses in Lake Ellis, 286
 Grew, Nehemiah, on flower colour of aquatics, 230; on heterophylly, 154, 155
Griffithiella Hookeriana, 114
 Ground Ivy. See *Glechoma hederacea*
 Guiana, British, 109, 118, 119, 300
 Guiana, French, 113
Gunnera, gigantic herb, 181; polystely, 180-182, 346
Gunnera scabra, 181
 Gunnereae, 312
- Hairs, absence in submerged leaves, 165, 166; absence in water forms of amphibious plants, 151, 152 (Fig. 99); change in character of hairs of *Rubus* when submerged, 200; growth of hairs due to wounding of Waterlily petiole, 258; loss of hairs of *Mentha* when submerged, 201; mucilage-containing hairs of *Ceratophyllum*, 86, 87, 272; mucilage-secreting hairs of Nymphaeaceae, 38 (Fig. 20), 272; *Myriophyllum* trichomes, 168, 169, 170 (Fig. 110); *Polygonum amphibium*, hairs on air leaves only, 151, 152 (Fig. 99); *Utricularia*, hairs of bladders, 92 (Fig. 60), 93 (Fig. 61), 94, 95 (Fig. 62); *Utricularia*, protective hairs of turions, 101, 102, 220 (Fig. 143 A); sensitive hairs of *Aldrovandia*, 111; stipular hairs of *Nymphaea lutea*, 26; see also under Root-hairs and *Squamulae intravaginales*
Halodule, anatomy of vegetative organs, 132 (Fig. 88), 331; marine Angiosperm, 123
Halodule uninervis, 132 (Fig. 88)
Halophila, anatomy, 131, 169; in biological classification, 6; leaf form, 124; marine Angiosperm, 57, 123; pollen-grains in strings, 130; pollination, hydrophilous, 130, 236; structure and life-history, **129**, **130** (Fig. 87); styles filiform, 130
Halophila ovalis, 129, 130 (Fig. 87)
Halophila ovata, 129
Halophila stipulacea, 129, 130
 Haloragaceae, 180, 205, 311, 312
 Haloragideae, 312
 Haptera of Podostemaceae, 114; of Tristichaceae, 113
 Hawthorn, pigmented variety, 276
 Heliotropism, 281
 Helobieae, 52, 123, 124, 245, 248, 313, 314, 318, 319, 320, 321, 325, 326
 Herbarium material, use in anatomical work, 331
Herbarum vivae eicones of Brunfels, 27 and Frontispiece
Herminiera elaphroxylon, 192
Heteranthera, cleistogamy, 234 (Fig. 153); phylloclad leaf anatomy, **342** (Fig. **170**), **343**, **344**; root differentiation, 207; support of inflorescence, 228; water pores and apical opening, 268
Heteranthera dubia, 234 (Fig. 153)
Heteranthera reniformis, 342 (Fig. 170), 343, 344
Heteranthera zosteraefolia, 207, 228, 268, 342 (Fig. 170), 343, 344
 Heterophylly, **143-162**, *et passim*
Himantalia lorea, 114
 Hippuridaceae, 312
Hippuris vulgaris, affinities, 311, 312; altitude, 290; anemophily, 230, 232; diaphragms of stem, 184 (Fig. 119), 257; flowers, 230, 231 (Fig. 151); fruit dispersal, 297; heterophylly, 141, **146**, **147** (Fig. **95**), **148** (Fig. **96**), **231** (Fig. **151**); in biological classification, 6; nutlets, 242; perennation, 215; reduction of primary root, 244; rhizome, 173 (Fig. 112); root-hairs from "collet," 245; stem, 172, 173; stem anatomy, **175-178** (Fig. **115**), **181**, **184** (Fig. **119**), **185** (Fig. **120**); stomates, 166; tenderness of leaves, 163
 Holland, 303
 Horned Pondweed. See *Zannichellia*
 Hornwort. See *Ceratophyllum demersum*
 Horse Chestnut, submerged germination, 199
Hottonia palustris, affinities, 312, 318; in biological classification, 6, 7; land and water forms, 197 (Fig. 127); leaf anatomy, 169; non-cleistogamic, 233; polystely, 181; ripening of fruit in air, 239; sinking of seeds, 297; support of inflorescence, 228; vegetative reproduction, 216
 Hyacinth, Water. See *Eichhornia speciosa*
Hydrilla, in Britain, 54, 55; leaves, 57; spathe, 315; tendrill roots, 205 (Fig. 136)
Hydrilla verticillata, 54, 55, 205 (Fig. 136)
 Hydrilleae, 175
Hydrobryum, 114, 115 (Fig. 76)
 Hydrocaryaceae, 311

INDEX

427

- Hydrocharis*, anthocyanin, 276; buds, summer, 43 (Fig. 24); buds, winter, 47 (Fig. 29), 48, 49 (Fig. 30); dehiscence, 47, 241; dioecism or monoecism, 46; flowers, 46; fruits, 46, 47, 241; freezing, 220; germination of turions, 48, 49 (Fig. 30), 280; in biological classification, 7; inverted bundles of leaf, 46 (Fig. 28); land form, 42, 49; leaf anatomy, 44 (Fig. 25), 45 (Figs. 26, 27), 46 (Fig. 28); light, effect of, 280; petiole length, experiments on, 283, 284; pollination, 236; root-hairs, 42, 43; roots, 42, 43, 244; stipules, 43 (Fig. 24), 44; stomates, 45 (Fig. 26); submerged form, 45; turions, 47 (Fig. 29), 48, 49 (Fig. 30); winter-buds, 47 (Fig. 29), 48, 49 (Fig. 30)
- Hydrocharis astatica*, 42
- Hydrocharis Morsus-ranae*, 7, 32, 42, 43 (Fig. 24), 44 (Fig. 25), 45 (Figs. 26, 27), 46 (Fig. 28), 47 (Fig. 29), 48, 49 (Fig. 30), 53, 54, 57, 166, 195, 215, 219, 220, 236, 241, 280, 283
- Hydrocharis parnassifolia*, 42
- Hydrocharitaceae, fresh-water, 42–57, 84, 151, 157, 248; marine, 123, 129–131, 133, 134; other references, 169, 205, 235, 236, 239, 314, 340
- Hydrocleis*, apical cavity of leaf, 269, 270 (Fig. 164); heterophylly, 157
- Hydrocleis nymphoides*, 157, 269, 270 (Fig. 164)
- Hydrocotyle vulgaris*, 200, 201 (Fig. 132)
- Hydromystris*, 57
- Hydrophilous pollination. See Pollination, hydrophilous
- Hydrothrix*, cleistogamy, 234; phyllode leaf, 344
- Hydrothrix Gardneri*, 234, 344
- Hydrotriche*, affinities, 313, 318; heterophylly, 151
- Hydrotriche hottoniaefolia*, 151
- Illecebraceae, 311
- Illecebrum*, affinities, 311; cleistogamy, 234
- Illecebrum verticillatum*, 234
- Illumination, 157, 278–280
- “Imbibition theory,” 174
- Indehiscent fruits of aquatics, 241–244
- India, 112, 191, 291, 305
- Infusoria, as food of *Utricularia*, 94
- Inn, River, 228
- Ireland, 210
- Iridaceae, 326
- Iris*, phylloid anatomy, 340
- Iris Pseudacorus*, 199
- Isoetes*, altitude, 291; in peaty water, 287; in mountain lochs and heath pools, 290; ousted by *Potamogeton*, 333; replacement of sporangia by plantlets, 225
- Isoetes amazonica*, 291
- Isoetes echinospora*, 225
- Isoetes lacustris*, 225
- Italy, 303
- Ivy-leaved Duckweed. See *Lemna trisulca*
- Japan, 17
- Juncaginaceae, 248, 313, 314
- Juncus*, 299, 309
- Juncus conglomeratus*, 299
- Jura lakes, 279, 287, 290, 323
- Jussiaea*, aerenchyma from phellogen, 189, 190 (Fig. 122); affinities, 311, 318; breathing roots, 189; replacement of cork by aerenchyma, 188, 189; roots not floats, 192, 193; submerged leaves whorled, 230
- Jussiaea amazonica*, 230
- Jussiaea grandiflora*, 189
- Jussiaea peruviana*, 189, 190 (Fig. 122)
- Jussiaea repens*, 189, 193
- Kaiteur, 109
- Kerguelen’s Land, 233
- “Kiemenbüschel” (gill-tufts) of *Oenone*, 118, 119 (Fig. 81), 255
- Kingston, 273
- Kurdistan, 303
- Laboul, 291
- Lace-plant of Madagascar. See *Aponogeton fenestralis*
- Lacis alata*, 120
- “Lady-Smocks,” Grew on heterophylly in, 155
- Lady’s Smock. See *Cardamine pratensis*
- Lake dwellings, 302
- Lake Ellis, North Carolina, 286
- Lake George, Florida, 213
- Lake St Clair, Michigan, 288
- Lakenheath Lode, 216
- Land forms, of water plants, 195–198; of Alismaceae, 20, 21 (Fig. 9), 153 (Fig. 101), 195; *Callitriche*, 195; *Cardamine*, 202 (Fig. 133); *Hottoma*, 197 (Fig. 127); *Hydrocharis*, 42, 49, 195; Lemnaceae, 77, 78; *Limnanthemum*, 195; *Limosella*, 198; *Littorella*, 198 (Fig. 128); *Myriophyllum*, 195, 223 (Fig. 146); Nymphaeaceae, 32, 195; *Polygonum*, 152 (Figs. 99 and 100), 197, 198; *Potamogeton*, 195, 196 (Fig. 125); *Ranunculus*, 195, 196 (Fig. 126), 203 (Fig. 134)
- Land plants, effect of water upon, 200, 201 (Fig. 132)
- “Law of Age and Area,” 305–307
- “Law of Irreversibility,” 336, 347
- “Law of Loss,” 182, 336–347
- Lawia*, germination, 117; mucilaginous seeds, 300; shoot thallus, 117; special-

- ised Podostemad, 306; starch storage, 120; vegetative reproduction, 216
- Lawia foliosa*, 117
- Lawia zeylanica*, 117, 120, 216, 300
- Leaves, band or ribbon, 11 (Fig. 3), 12, 13 (Fig. 4), 14 (Fig. 5), 19, 20, 22, 23, 140, 141, 343, 344, etc.
- Leaves, floating, 30–32, 44–46, *et passim*
- Leaves, submerged, 139–143, 163–171; “adaptation” in, 171; aerating system, 167; diaphragms, 167; epidermal cells, form of, 163, 164; epidermis, chlorophyll in, 164; fenestration, 142 (Fig. 91); mesophyll for storage, 168 (Fig. 109); non-radial anatomy, 165 (Fig. 106), 169; radial anatomy, 168 (Fig. 109); ratio of surface to volume, 140; reduction of cuticle, 163; stomates and hairs, 165–167 (Fig. 107); undifferentiated mesophyll, 167, 168 (Fig. 109); water pores, 167
- Leguminosae, 188, 189, 191, 192
- Lemna gibba*, 7, 76 (Fig. 48), 77, 78, 81, 275, 297
- Lemna minor*, 7, 76, 77, 78, 80, 291, 295, 297, 301, 307
- Lemna trisulca*, 8, 78, 79 (Figs. 49, 50, 51), 80, 81 (Fig. 52), 208, 215, 295
- Lemnaceae, 73–82; aerenchyma, 76 (Fig. 48); affinities, 74, 82, 314, 316; altitude, 290, 291; anatomy, 78, 79 (Fig. 51); anthocyanin, 276, 277; cotyledon as float, 81 (Fig. 52), 248; dispersed by water birds, 300, 301; distribution, 73, 112, 295, 307; entirely aquatic, 318; entomophily, 80, 230; flowers, 74 (Fig. 47), 79 (Fig. 50), 80; land forms, 77, 78; life in impure water, 81, 287; number of genera, 84; protandry, 80; range, 73, 112, 295, 307; rarity of seeds, 75; reduced inflorescences, 73, 74 (Fig. 47); root-caps, 74, 76 (Fig. 48); roots for equilibrium, 74, 207; seedlings, 80, 81 (Fig. 52); seeds, 80, 297; turions, 74–77; vegetative morphology, 73–74; vigour of vegetative growth, 77, 81, 83
- Length, of axes, of *Oenanthe*, 150, of *Polygonum*, 215, of *Ranunculus*, 214, of *Utricularia*, 215; of petiole, peduncle, etc., in Nymphaeaceae, 28 (Fig. 13), 31 (Fig. 15), 283; of petiole in *Hydrocharis*, *Marsilea*, *Ranunculus*, 283, 284; of submerged leaves of *Sagittaria*, 12, of *Vallisneria*, 140
- Lentibulariaceae, 91, 104, 313
- Lenticels, effect of submergence on, 187
- Lesser Water Plantain. See *Echinodorus ranunculoides*
- Lignification, poor, in aquatics, 260
- Liliaceae, 326
- Limnanthemum*, 39–41; affinities, 313; anthocyanin, 276; dehiscence, 240 (Fig. 156), 241, 242 (Fig. 157); distribution, 304, 305; effect of drying on seeds, 243; fruit ripening in water, 239; geophytic habit, 41 (Figs. 22 and 23), 217, 323; germination, 248; land form, 195; mucilage, 271; rhizome, 39, 41 (Figs. 22, 23), 217, 323; seeds, 240 (Fig. 156), 241, 297; support of inflorescence, 228
- Limnanthemum Humboldtianum*, 239
- Limnanthemum indicum*, 40, 304
- Limnanthemum nymphoides*, 39–41 (Figs. 22, 23), 195, 228, 240 (Fig. 156), 241, 242 (Fig. 157), 243, 248, 271
- Limnobium*, geotropic curvature of fruit-stalk, 239, 282; heterophylly, 157
- Limnobium Boscii*, 157, 282
- Limnocharis Humboldtii*, 166
- Limnophila*, heterophylly, 151, 161; sleep movements, 281; systematic position, 313
- Limnophila heterophylla*, 161, 281
- Limnophila hottonoides*, 151
- Limnosipanea*, heterophylly, 151; affinities, 313
- Limnosipanea Spruceana*, 151
- Limosella*, affinities, 313, 318; cleistogamy, 233, 235; distributed by birds, 301; flowers, 313; land and water forms, 198
- Limosella aquatica*, 198, 233, 301, 313
- Linaria Cymbalaria*, 240
- “Little Bell,” Grew on heterophylly in, 155
- Littorella*, aerating system of leaf, 167; affinities, 313; anemophily, 232; apical openings of leaves, 269; flowers and fruit, 313; fruit, 241, 242; funicular plug in fruit wall, 242; geophytic habit, 323; germination *in situ*, 246; in peaty water, 287, 290; land and water forms, 198 (Fig. 128); ousted by *Potamogeton*, 333; radial leaf, 168; runners, 217, 218 (Fig. 142)
- Littorella lacustris*, 7, 141, 198 (Fig. 128), 217, 218 (Fig. 142), 232, 241, 242, 246
- Liverworts, 327
- Lobelia Dortmanna*, aerating system of leaves, 167; affinities, 313; fruit ripening in air, 239; in biological classification, 7; in peaty water, 287; in sandy pools and mountain lochs, 290; leaves, 141; root, 245; sinking of seeds, 297; stomates, 166
- Lobelia, Water. See *Lobelia Dortmanna*
- Loess alluvium, 287
- Loosestrife, Water. See *Lythrum Salicaria*
- Lotus*, 188
- Lotus*, Sacred. See *Nelumbo Nelumbo*
- Low countries, 9
- Ludwigia*, 311, 318
- Lupinus*, 200

INDEX

429

- Luxuriance of vegetative growth, **210–215**
- Lycopus europaeus*, 188
- Lysimachia*, 188
- Lyte's *Herbal*, 144
- Lythraceae, 175, 188, 193, 234, 303, 311
- Lythrum Salicaria*, 188, 311
- McLean, R. C., on treatment of herbarium material, 331
- Macropodous embryo, 246 (Fig. 159), **248** (Fig. 161), **249, 319** (Fig. 166), 326
- Madagascar, 305
- Madeira, 295
- "Major plant individual," **211–213**
- Manchester, 275, 303
- Marathrum utile*, 113
- Marburg, 189
- Mare's-tail. See *Hippuris vulgaris*
- Marine Angiosperms, **123–135**; affinities, 123, 320; association with Algae, 123; flowers, 126, 127, 129, 130 (Fig. 87); fruits, 126, 127, 248 (Fig. 161); leaf anatomy, **125** (Fig. 84), **128** (Figs. **85, 86**), **130, 131, 132** (Figs. **88, 89**); leaves, 124 (Fig. 83), 130 (Fig. 87), 133; origin of the group, 133–135; pollen-grains, *Conserva*-like, 124, 125, 126; pollen-grains in strings, 130; pollination, hydrophilous, 124–127, 129; seed-coats, 130; vegetative habit, 124 (Fig. 83), 130 (Fig. 87); vivipary, 127
- Marsilea*, 284
- Mauritius, 129, 295
- Mayaca fluviatilis*, 243
- Mayacaceae, 317
- Mediterranean, 123, 125
- Melilotus Taurica*, sleep habits of, 161
- Memory, unconscious, 333
- Mentha aquatica*, 201
- Menyanthes*, affinities, 313; submerged form, 199; tendril roots, 205
- Menyanthes trifoliata*, 199, 313
- Mercurialis*, 311
- Michigan, 288
- Milfoil, Water. See *Myriophyllum*
- Mimosa lacustris*. See *Neptunia oleracea*
- Mirabilis*, 121
- Monocotyledonous families, proportion of, among aquatics, 322
- Monocotyledons, aquatic origin of, **322–326**
- Monstera*, 142, 314
- Montia*, affinities, 310; biennial and perennial forms, 216; submerged xerophyte, 310
- Montia fontana*, 216, 310
- Mosses, 113
- Mourea*, anthocyanin, 113; flowering, 120; haptera, 114
- Mourea fluviatilis*, 113, 114, 120
- Mucilage, 38 (Fig. 20), 47, **271–272**, 300
- Mucilage-secreting trichomes, 13, 15, 38 (Fig. 20)
- Myriophyllum*, affinities, 311, 312; air spaces in roots, 187; in stem, 179 (Fig. 116), 256; altitude, 290; anemophily, 230, 232; effect of freezing on fruits, 243; on turions, 220; fruits, 242; germination of turions, 222 (Fig. 145); growth in still or moving water, 283; habit, 172, 221 (Fig. 144); in biological classification, 6; land form, 195, 223 (Fig. 146); on sandy substratum, 286; relation of turions to inflorescence, 224; sleep movements, 281; stem anatomy, 178, 179 (Figs. 116, 117), 181; submerged leaves, form, 255, structure, 168 (Fig. 109); trichomes, 168–170 (Fig. 110); turions, 219, 220, 221 (Fig. 144), 222 (Fig. 145), 223 (Fig. 146), 224; wave motion, 289
- Myriophyllum alternifolium*, 168
- Myriophyllum proserpinacoides*, 281
- Myriophyllum spicatum*, 134, 140, 168 (Fig. 109), 179 (Figs. 116, 117), 195, 197, 232, 242, 243
- Myriophyllum verticillatum*, 6, 168–170 (Fig. 110), 219, 220, 221 (Fig. 144), 222 (Fig. 145), 223 (Fig. 146)
- Naiadaceae, 248, 313, 315
- Naias*, alien weed with rice and cotton, 303; annual, 215; distribution, 304, 305; effect of depth, 279; flower, 315, 316, 320, 346; in biological classification, 6; in deep water, 279, 288; pollination, hydrophilous, 237; reduction of primary root, 244; root anatomy, 208, 209 (Fig. 140); specific differences, 331, 332; stem anatomy, 175; submerged vegetative organs, 134
- Naias flexilis*, 215
- Naias graminea*, 237, 303
- Naias graminea*, var. *Delilei*, 332
- Naias major*, 209 (Fig. 140)
- Naias marina*, 304, 305
- Naias minor*, 209 (Fig. 140), 215
- Nasturtium*, affinities, 309; air and water shoots, 201; budding from leaves, 216
- Nasturtium amphibium*, 201, 309
- Nasturtium lacustre*, 216
- Nelumbium*. See *Nelumbo*
- Nelumbo*, absence of mucilage, 257, 272; affinities and structure, 38, 39; geologic distribution, 38, 39 (Fig. 21); movements of gases, 257, 258; possible case of reversion from aquatic to terrestrial life, 39; stability of seedling, 245
- Nelumbo Nelumbo*, 38, 39 (Fig. 21)
- Nelumboideae, 38

- Nepenthes*, 93, 310
Nepitunia oleracea, 189–191 (Fig. 123)
Nesaea, aerenchyma, 193 (Fig. 124), 194; cleistogamy, 234; distribution, 295
Nesaea verticillata, 193 (Fig. 124), 194
 Nile, 113
 Nile, Blue, 192
Nitella, 288
 Nitrogen, proportion in free and dissolved air, 253
Nuphar luteum. See *Nymphaea lutea*
Nuphar minima, 28
Nuphar pumilum, 32
Nymphaea, dehiscence, 35, 36; effect of frost on seeds, 243; etiolation with depth, 279; floating leaves, 30, 31, 146, 159; fruit, 34 (Fig. 17), 35, 36, 240; general habit, Frontispiece; geophytism, 217, 323; length of petiole and peduncle, 31 (Fig. 15), 40; operculum of seed, 35 (Fig. 18), 36; pigmented variety, 276; reduction of primary root, 244, 281; rhizome, 24, 25 (Fig. 10), 26, 27 (Fig. 12), 36, 39, 217; roots, 25 (Fig. 10), 204, 281; seedlings, 34, 35 (Fig. 18), 36, 280; submerged leaves, 27 (Fig. 12), 28, 29, 146, 159, 279; trichome diaphragms, 272
Nymphaea alba. See *Castalia alba*
Nymphaea lutea, 6, 24, 25 (Fig. 10), 26, 27 (Fig. 12), 28–31, 34 (Fig. 17), 35 (Fig. 18), 36, 159, 243, 244, 272, 279, 280, 281, 288, see also Frontispiece
Nymphaea lutea, var. *rubropetala*, 276
Nymphaea pumila, 32
 Nymphaeaceae, 24–39; affinities, 309, 314, 318–320; air system in petioles and peduncles, 37, 257; anatomy, 36–38, 182; ancient aquatic habit, 321; and water fowl, 299, 300; anthocyanin, 276, 277; cleistogamy, 234; geophytic habit, 217; heterophylly, 27 (Fig. 12), 28, 29 (Fig. 14), 146; in Lake Ellis, 286; land forms, 32, 195; leaf and flower, 40; leaf, floating, 30, 31; leaf, submerged, 27 (Fig. 12), 28 (Fig. 13), 29 (Fig. 14), 255; mucilage, 35, 36, 38 (Fig. 20), 272; wound effects, 258
 Nymphaeoidae, 32, 38
*Oenanth*e, aerenchyma, 188; heterophylly, 150; perenniation, 215; root system, 204, 205, 229; stomates, 166; submergence, 312
Oenanthe Phellandrium, 150, 204, 205, 229
Oenanthe Phellandrium, var. *fluviatilis*, 150, 166, 215, 312
Oenone, gill-tufts, or Kiemenbüschel, 119 (Fig. 81), 255
Oenone multibranchiata, 119 (Fig. 81)
 Onagraceae, 188, 189, 311, 318
 “Open reed-swamp,” 288
 Organ Mountains, 108
Origin of Species, *The*, 260
 Osmotic pressure of sap in leaves and roots, 266
Ottelia, geotropism of peduncle, 239; heterophylly, 57
Owivirandra. See *Aponogeton fenestralis*
 Oxygen, proportion in free and dissolved air, 253; scarcity in water life, 255
Pacu myletes, 118
Padina pavonia, 123
 Palms, 143
 “Pancy,” 155
 Pandanaceae, 317
 Parallel veining of Monocotyledonous leaves, 338
Parra jacana, 300
 Parsnip, Water. See *Sium latifolium*
 Pea, 249
 Peat-bog lakes, 275
 Peaty substratum, 287
 Pedaliaceae, 151, 234
Peplis, aerating system, 185, 259; affinities, 311; anthocyanin, 276, 277; cleistogamy, 234; detached shoots, 216, 276, 277; flowers and fruit, 230, 232 (Fig. 152); pollination, 230; vascular anatomy, 175; vegetative reproduction, 216; winter state, 216
Peplis Portula, 175, 185, 216, 230, 232 (Fig. 152), 259, 276, 277, 311
 Perenniation among hydrophytes, 215
 Perthshire, 288
Phaseolus, 207
 Phellogen, air-containing, 187–191
 Phellogen producing aerenchyma, 187–191
Philodendron, 206
Phragmites, in Jura Lakes and White Moss Loch, 287, 288 (Fig. 165); “major plant unit,” 212; root differentiation, 207
Phragmites communis, 207, 212
 “Phragmitum,” 288
 “*Phucagrostis major*,” 125
 “*Phucagrostis minor*,” 125
Phyllanthus fluitans, 311
 Phyllode theory of Monocotyledonous leaf, 52, 161, 162, 337–345
Phyllospadix, 123, 124
 Pico, 333
 Piliferous layer, cuticularised, 208; death of, before death of root-hairs, 264
Pilularia, 225
Pinguicula, insectivorous habit and relation to *Utricularia*, 111; polystely, 181
Pinguicula vulgaris, 181

INDEX

431

- Pistia Stratiotes*, air tissue of leaves, 82, 154, 256; comparison with Lemnaceae, 74, 82, 316; hairs, 82, 83; vigour of vegetative growth, 83, 213, 214; water pores, 82 (Fig. 53), 83, 167, 267
 "Pith" helmets, 191
Plantago, 233, 313
 "Plantago aquatica," 20
Plantago major, 241
 Plantain, Lesser Water. See *Echinodorus ranunculoides*
 Plantain, Water. See *Alisma Plantago*
 Podostemaceae (including *Tristichaceae*), 112–122, 327–333; affinities, 310, 319; anatomy, 117, 118 (Fig. 80); ancient aquatics, 321; and natural selection, 327–333; and wading birds, 300; anemophily, 120, 121; anthocyanin, 112, 113, 276, 277; cleistogamy, 121 (Fig. 82); dependence on aeration, 257; distribution, 295, 306; dorsiventrality, 121, 122, 327–329; entirely aquatic, 318; flowers, 120, 121 (Fig. 82); germination *in situ*, 248; "gill-tufts," 118, 119 (Fig. 81), 255; habit, 114, 115 (Figs. 76, 77), 116 (Fig. 79), 117; haptera, 113, 114, 121; in biological classification, 7; inhabit rapids, 112, 113, 119, 257; "Kiemenbüschel," 118, 119 (Fig. 81), 255; lack of adaptation, 328–333; lack of intercellular spaces, 118 (Fig. 80), 257; morphology, 73, 121, 122; mucilaginous seeds, 300; polymorphism of thallus, 114–117; rarity outside tropics, 112, 113; reduction of primary root, 244; root thallus, 114, 115 (Figs. 76, 77, 78), 116 (Fig. 79), 117, 208; secondary shoots, 114; seedling, 114, 115 (Fig. 78), 117; seeds, 121, 300; shoot thallus, 117; silica, 117; simulation of lower plants, 114–117; vegetative reproduction, 216; water reservoir in nucellus, 121
Podostemon, Alga-like form, 114; cleistogamy, 121 (Fig. 82), 234; distribution, 306
Podostemon Barberi, 121 (Fig. 82), 234
Podostemon subulatus, 114
 Pollination, anemophilous, 57, 120, 121, 230, 232, 233
 Pollination, aquatic. See Pollination, hydrophilous
 Pollination, cleistogamic. See Cleistogamy
 Pollination, entomophilous, 9, 57, 80, 230
 Pollination, hydrophilous, 1, 6, 8, 55–57, 70, 71, 84, 85, 124, 127, 129, 130, 134, 235–238, 345, 346
 Polygonaceae, 311
Polygonum, affinities, 311, 318; colonising new waters, 289; land and water forms, 150, 151, 152 (Figs. 99, 100), 197, 198; heterophylly, 150, 151, 152 (Figs. 99, 100); length of shoot system, 215; mucilage, 271; vegetative reproduction, 225
Polygonum amphibium, 150–152 (Figs. 99, 100), 197, 198, 215, 271, 289, 311, 318
Polygonum viviparum, 225
 Polypetalae, 309–312, 319
 Polystely, 37, 180–182, 346
 Pondweed. See *Potamogeton*
 Pondweed, Horned. See *Zannichellia*
Pontederia, geotropic curvature of fruit stalk, 239, 240 (Fig. 155); phylloclad leaf anatomy, 341 (Fig. 169), 342 (Fig. 170), 343, 344; specialised genus, 317; stomates, 166
Pontederia cordata, 166, 341 (Fig. 169), 342 (Fig. 170), 343, 344
Pontederia rotundifolia, 239, 240
 Pontederiaceae, affinities, 316, 317; cleistogamy, 234 (Fig. 153); entirely aquatic, 318; geotropic curvature of fruit stalk, 239, 240 (Fig. 155), 282; heterophylly, 154, 160, 161; phylloclad leaf structure, 337, 341 (Fig. 169), 342 (Fig. 170)–344; vegetative multiplication, 213
 Poplar, 187, 316
 Portulacaceae, 310
Posidonia, apical openings absent, 269; chlorophyll in epidermis, 164; fibres in leaf sheath, 133; habit, 124; leaf anatomy, 132 (Fig. 89); marine Angiosperm, 123; pollen thread-like, 124, 125
Posidonia Caulini, 125, 132 (Fig. 89)
Potamogeton, 58–72; air tissue in fruit wall, 71, 72 (Fig. 46); altitude, 290, 291; anatomy of inflorescence axis, 65; anatomy of root, 65 (Fig. 41), 66, 208; anatomy of stem, 62 (Fig. 39), 63, 64 (Fig. 40), 65, 175; apical openings of leaves, 167 (Fig. 108), 268; chlorophyll in epidermis, 164; choking mill sluices, 210; cortical bundles, 65; cuticle, waxy, 254; delayed germination, 71, 72; diaphragms, 65, 184 (Fig. 118); dispersal by ducks, 301, 302; distribution, 295, 297, 298; dwarfing due to heat, 275; exudation of water drops, 269; fibres in leaves, 61 (Fig. 38), 169; flowers, 69–71; fruits, 71, 72 (Fig. 46), 297; germination, 280; heterophylly, 151, 153, 154 (Fig. 103), 157, 158 (Fig. 104), 159 (Fig. 105), 339 (Figs. 167 and 168); in biological classification, 6; land forms, 61, 195, 196 (Fig. 125); leaf forms, 61, 339 (Figs. 167 and 168); oil drops, 62; perennation, 215; phylloclad interpretation of leaf, 339 (Figs. 167, 168), 340

- Potamogeton crispus*, 61, 62 (Fig. 39)–64 (Fig. 40), 67 (Fig. 42), 68 (Fig. 43), 69, 71, 164, 269, 275, 295
- Potamogeton densus*, 65 (Fig. 41), 71, 107 (Fig. 108), 206 (Fig. 137), 268, 298
- Potamogeton fluitans*, 69, 151, 157, 158 (Fig. 104)
- Potamogeton heterophyllus*, 61, 195
- Potamogeton lucens*, 61–64 (Fig. 40), 164, 173, 262, 330, 339 (Fig. 167)
- Potamogeton natans*, 6, 31, 32, 61, 62 (Fig. 39), 63, 65 (Fig. 41), 66, 70, 72, 151, 157, 159 (Fig. 105), 166, 167 (Fig. 107), 184 (Fig. 118), 195, 196 (Fig. 125), 272, 280, 289, 301, 339 (Fig. 168)
- Potamogeton obtusifolius*, 206
- Potamogeton pectinatus*, 62–64 (Fig. 40), 65 (Fig. 41), 66, 70, 134, 262, 282, 291
- Potamogeton pennsylvanicus*, 303
- Potamogeton perfoliatus*, 58, 59 (Fig. 36), 61, 63, 69, 71, 140, 195
- Potamogeton polygonifolius*, 196 (Fig. 125), 333
- Potamogeton praelongus*, 62
- Potamogeton pulcher*, 61, 62 (Fig. 39), 63, 65
- Potamogeton pusillus*, 63, 64 (Fig. 40), 66, 71
- Potamogeton rufescens*, 69 (Fig. 44)
- Potamogeton trichoides*, 62, 66, 71
- Potamogeton varians*, 195
- Potamogeton zosterifolius*, 61 (Fig. 38)
- Potamogetonaceae, fresh-water, **58–72**;
marine, **123–129, 131–135**, 237, 331;
other references, 205, 248, 314–316, 337
- Primulaceae, 180, 312, 318
- Proserpinaca palustris*, heterophylly, 159–161
- Protection of embryo in aquatics, 242, 243
- Pseudo-callitriche*, 134, 237, 306, 307
- Pseudo-lamina, 339, 340, 341
- Quercus*, 207
- Rafts of wood of *Herminiera*, 192
- Ranales, 146, 238, 308, 319, 320, 321, 346
- Range, wide, of aquatics, 295
- Ranunculaceae, 309, 313, 314, 320
- Ranunculus*, affinities, 318; air spaces, 176 (Fig. 113), 185; altitude, 290; amphibious, aquatic and terrestrial types, 200, 309, 320; dimensions of dissected leaves, 140; geotropic curvature of peduncle, 145 (Fig. 93), 239; germination, 280; heterophylly, 144 (Fig. 92), 145, 146, 155; in biological classification, 6; land and water forms, 195, 196 (Fig. 126), 198, 203 (Figs. 134, 135); luxuriance, 214; non-heterophyllous form, 145 (Fig. 93); root-system, 204, 264; sinking of seeds, 297; stem anatomy, 175, 176 (Fig. 113); sub-aquatic flowering, 233, 234; submerged leaves, 29, 140, 142; submergence of inflorescence, 228; support of inflorescence, 228; toleration of salt, 134; winter-buds, 219
- Ranunculus aquatilis*, 144, 145, 155, 196 (Fig. 126), 204, 234, 280, 297
- Ranunculus Baudotii*, 134
- Ranunculus carinatus*, 228
- Ranunculus circinatus*, 145
- Ranunculus confusus*, 228
- Ranunculus divaricatus*, 234
- Ranunculus Flammula*, 145, 198, 203 (Figs. 134, 135), 309, 320
- Ranunculus hederaceus*, 145 (Fig. 93)
- Ranunculus heterophyllus*, 145
- Ranunculus Lingua*, 146, 219
- Ranunculus Puschii*, 144 (Fig. 92)
- Ranunculus repens*, 200
- Ranunculus sceleratus*, 146, 284, 320
- Ranunculus trichophyllus*, 140, 176 (Fig. 113), 228, 290
- Ranunculus* sect. *Batrachium*, 6, 29, 142, 144–145, 175, 185, 195, 228, 239, 264, 309, 318, 320
- Rheotropism, 282
- Rhizomatous plants of Jura Lakes, 323
- Rhizome, and polystely, 182; rhizome of *Castalia*, 24–26 (Fig. 11), 217; *Gunnera*, 182; *Hippuris*, 173 (Fig. 112); *Hottonia*, 197 (Fig. 127); *Limnanthemum*, 39–41 (Figs. 22, 23), 217; *Nymphaea*, 24, 25 (Fig. 10), 27 (Fig. 12), 217; *Potamogeton*, 58–60 (Fig. 37)
- Rhizopods, as food of *Utricularia*, 94
- Rhyncolacis macrocarpa*, 120, 121
- Ribbon leaves, 11 (Fig. 3), 12, 13 (Fig. 4), 14 (Fig. 5), 19, 20, 22, 23, 140, 141, etc.
- Riccia*, 225
- Rice, aliens accompanying, 303
- Ricinus*, 200
- River-basins, isolation of, 296
- Rodriguez, 129
- Root-caps, of *Brasenia*, 205; of Lemnaceae, 74
- Root-hairs, absence of, in *Lemna trisulca*, 208, in water roots of *Elodea*, 208; in *Hydrocharis*, length of, 42, 43, 208; protoplasmic rotation, 43
- Rootless Duckweed. See *Wolffia*
- Roots of water plants, **204–209**; aerenchyma, secondary, 188, etc.; air tissue, 185, 186 (Fig. 121), 187; anatomy, 65 (Fig. 41), 208, 209 (Figs. 138–140); assimilation, 207; differentiation, 207; equilibrium, 207; importance in life of aquatics, 264–266; reduction of, 208, 244; spiral or tendril, 127, 205 (Fig. 136), 206 (Fig. 137)

INDEX

433

- Roraima, 109
 Rosales, 310, 319
 Roslyn Pits, Ely, 147, 215, 241
Rotala, alien accompanying rice, 303;
 cleistogamy, 234
Rotala indica, 303
 Rotifers, 94, 142
 Rubiaceae, 151, 313
Rubus fruticosus, 200
Ruppia, in brackish water, 134; macro-
 podous embryo, 319 (Fig. 166);
 pollination, 70; reduction of primary
 root, 244
Ruppia brachypus, 319 (Fig. 166)
- Sacred Lotus. See *Nelumbo Nelumbo*
- Sagittaria*, band or ribbon leaves, 9, 11
 (Fig. 3), 12, 13 (Fig. 4), 140, 141 (Fig.
 90); diaphragms, 19 (Fig. 8), 167;
 effect of freezing on fruit, 243; floating
 of fruit, 297; flowers, 10 (Fig. 1);
 fruits, 10 (Fig. 2); heterophylly, 9,
 11 (Fig. 3), 12, 13 (Fig. 4), 14 (Fig. 5),
 16 (Fig. 6), 22, 23; in biological classifica-
 tion, 5; leaf anatomy, 344, 345 (Fig.
 171); mericarps and seeds, 17, 18;
 phylloclad interpretation of leaf, 161,
 162, 339, 340, 344, 345 (Fig. 171);
squamulae intravaginales, 13, 15;
 stolons, 15, 16 (Fig. 6), 18 (Fig. 7);
 tubers, 11 (Fig. 3), 13 (Fig. 4), 15, 16
 (Fig. 6), 17, 18 (Fig. 7), 217, 223, 224
- Sagittaria montevidensis*, 344, 345 (Fig.
 171)
- Sagittaria natans*, 12, 156
- Sagittaria sagittifolia*, 5, 9, 10 (Figs. 1,
 2), 11 (Fig. 3), 12, 13 (Fig. 4), 14
 (Fig. 5), 15, 16 (Fig. 6), 17, 18 (Fig.
 7), 19 (Fig. 8), 23, 32, 34, 61, 141
 (Fig. 90), 156, 157, 217, 223, 224, 297,
 344, 345 (Fig. 171)
- Sagittaria sagittifolia* f. *vallisnerifolia*,
 11, 12
- Sagittaria teres*, 7, 22
- Sagittaria variabilis*, 17
- St John's River, Florida, 213
- Salix*, development of aerenchyma, 188;
 effect of submergence, 200
- Salix viminalis*, 188
- Salt, toleration of, by certain hydro-
 phytes, 133, 134
- Salvinia*, 311, 337
- Santalaceae, 312
- Santarem, 291
- Sarracenia*, 93
- Sarraceniales, 310, 319
- Saxifragaceae, 310
- Scania, 303
- Scirpus*, affinities, 317; heterophylly,
 154; in Jura lakes, 287, 288
- Scirpus fluitans*, 317
- Scirpus lacustris*, 154, 288, 317
- Scotland, 290
- Screw Pine, 317
- Scrophulariaceae, 151, 313, 318
- Scutellaria*, 188
- Sedges in Lake Ellis, 286
- Seed dispersal, 17, 18, 35, 71, 72 (Fig.
 46), 297-303
- Seedlings, 243-249; of *Alisma*, 151, 153
 (Fig. 101); *Castalia*, 28 (Fig. 13), 29;
Ceratophyllum, 85, 86 (Fig. 55);
Dicraea, 114, 115 (Fig. 78); *Elatine*,
 245 (Fig. 158); *Hippuris*, 245; *Lawia*,
 117; *Lemna*, 80, 81 (Fig. 52); *Limnan-
 themum*, 248; *Nymphaea*, 35 (Fig. 18),
 36; *Rhyncolaxis*, 121; *Trapa*, 245-
 247 (Fig. 160); *Utricularia*, 100 (Figs.
 67, 68); *Victoria*, 32, 33 (Fig. 16), 34;
Zannichellia, 245, 246 (Fig. 159)
- Seeds, 239-249; of *Aldrovandia*, 110;
Alisma, 242, 297; *Callitriche*, 297;
Castalia, 243; *Hippuris*, 242, 297;
Hottonia, 297; *Hydrocharis*, 241;
Lemnaceae, 80, 297; *Limnanthemum*,
 240 (Fig. 156), 241, 243, 297; *Litto-
 rella*, 242; *Lobelia*, 297; *Mayaca*, 243;
Myriophyllum, 242; *Nymphaea*, 243;
Potamogeton, 297; *Ranunculus*, 297;
Sagittaria, 17, 18, 297; *Stratiotes*,
 241; *Utricularia*, 99, 100
- Sensitive Plant, Floating. See *Neptunia
 oleracea*
- Sesbania*, 191
- "Shade leaf" characters of submerged
 leaf, 45 (Fig. 27), 164, 165, 171, 279
- Siberia, 302
- "Sifting" in evolution, 162, 203
- Sium latifolium*, heterophylly, 147, 149
 (Fig. 97), 150 (Fig. 98); in biological
 classification, 5
- Sleep movements, 281
- Snails, as food of *Utricularia*, 93
- Snails, water, absence in peaty water,
 287; as pollinators of *Sagittaria*, 9;
 unable to eat turions of *Utricularia*,
 101
- Sneezewort. See *Achillea ptarmica*
- Solanum Dulcamara*, 198
- Soldier, Water. See *Stratiotes aloides*
- Sparganium*, fruits eaten by wild ducks,
 301; resemblance to Pandanaceae,
 317
- Sparganium natans*, 317
- Spathicarpa*, 315
- Spathiflorae, 314
- Spearwort, Greater. See *Ranunculus
 Lingua*
- Spearwort, Lesser. See *Ranunculus
 Flammula*
- Specific characters, constancy of ana-
 tomical, 131, 331; lack of utility of,
 131, 330, 331
- Specific vital energy, 212
- Speedwell, Water. See *Veronica Ana-
 gallis*

- Spirodela polyrrhiza*, 7, 74 (Fig. 47), 75, 76, 78, 80, 215
 Spurwing, 300
 "Squamulae intravaginales," 15, 126, 271
 "Starch-leaved" plants, 164
 Starwort, Water. See *Callitriche*
 Stems, aquatic, condensation of vascular tissue, 174–180; morphology and anatomy, 172–185, 187–194; polystely, 180–182; reduction of xylem, 172, 173; sympodial growth, 172, 173
 Stomates, on submerged leaves, 165, 166, 167 (Fig. 107); water stomates, 30, 267, 268 (Fig. 163), 269
Stratiotes, calcophil., 287; chalky incrustation on leaves, 51; dehiscence, 241; dioecism, 54; flowers, 54 (Fig. 33); fruit, 239; geographical distribution of sexes, 54; habit, 53 (Fig. 32); in biological classification, 7; inversely orientated leaf bundles, 52; leaves, 49–53, 57, 141, 157, 169; mechanism of rising and sinking, 50, 51; Pleiocene and Pleistocene records, 54; roots, 50, 185, 186 (Fig. 121), 187, 207, 244; stem, 49 (Fig. 31), 172; stomates, 51, 52; winter-buds, 53 (Fig. 32), 54
Stratiotes aloides, 7, 49 (Fig. 31), 50–53 (Fig. 32), 54 (Fig. 33), 185–187, 215, 219, 241
 Submerged leaves. See Leaves, submerged
 Substratum, influence of, 286–287
Subularia, affinities, 309; in biological classification, 7; submerged flowering 233
Subularia aquatica, 7, 309
 "Sugar-leaved" plants, 164
 Sundew, 111
 Surface-heating of ponds, 274
 "Swan's Potatoes," 17
 Sweden, 46, 303
 Sweet William, 155
 Switzerland, 302, 303
 Sympetalae, 312, 313, 320
 Tanqui, 181
 Temperature, 273–275
 Tendril roots, of Gentianaceae, 205; Haloragaceae, 205; Hydrocharitaceae, 205 (Fig. 136); *Philodendron*, 206; Potamogetonaceae, 205, 206 (Fig. 137)
 Terrestrial forms. See Land forms
Thalassia, leaf anatomy, 131, 169; marine Angiosperm, 57, 123
 Thallus of Podostemads, 7, 114, 115 (Figs. 76, 77, 78), 116 (Fig. 79), 117
 Thames, 210, 273
 Theophrastus, on ecology of aquatics, 285; on *Trapa*, 207
 Thyme, Water. See *Eloëa canadensis*
 Tibet, 291
Tillaea. See *Bulliardia*
Tillandsia, 108, 109
Tornelia, 142
 Transcaucasia, 303
 Transpiration current, 260–272
Trapa natans, affinities, 311; assimilatory roots, 207, 255; buoyancy due to lacunae, 192; changes in distribution, 302, 303; cotyledons, 245; fixation of seedling, 245; fossil records, 303; hypocotyl, negatively geotropic, 245; reduction of primary root, 244; seedling, 247 (Fig. 160); transverse heliotropism, 281; use in magic and medicine, 302
Trapella, cleistogamy, 234; heterophylly, 151
Trapella sinensis, 234
Trianea, 45
Trifolium, sleep habits of, 161
Trifolium resupinatum, 199 (Fig. 131)
Tristicha, dorsiventral root, 113; primitive Podostemad, 306
Tristicha ramosissima, 113
 Tristichaceae (see also Podostemaceae), 7, 112, 113, 120, 306
 Turions, 217–225; and unfavourable conditions, 222–224; in relation to inflorescences, 224, 225; of *Aldrovandia*, 110, 219; of *Caldesia*, 22, 225 (Figs. 148, 149); of *Hydrocharis*, 47 (Fig. 29), 48, 49 (Fig. 30), 219; of Lemnaceae, 75–77, 219; of *Myriophyllum*, 219, 220, 221 (Fig. 144), 222 (Fig. 145), 223 (Fig. 146); of *Potamogeton*, 66, 67 (Fig. 42), 68 (Fig. 43), 69 (Fig. 44); of *Utricularia*, 101, 102 (Fig. 69), 103, 219, 220 (Fig. 143)
Ulva, 27, 146
 Umbelliferae, 5, 147, 312
 Unconscious memory, 333, 347
 Upware, 216
Utricularia, 91–109; absence of roots, 91, 204, 244; affinities, 313; "air shoots," 96–98 (Fig. 65); anatomy, 107, 108 (Fig. 74), 227; apical development, 106 (Fig. 72), 107 (Fig. 73); benzoic acid in utricles, 96; capsule, 239; carnivorous habit, 93–96, 111, 270; comparison with *Aldrovandia*, 110; dispersal, 299; "earth shoots," 96 (Figs. 63, 64), 97, 270, 336, 337; effect of strong light, 279; epiphytism, 108, 109; floats on inflorescence axis, 99, 229 (Fig. 150); freezing, 220; hairs inside utricle, 92 (Fig. 60), 93 (Fig. 61), 95; hairs on valve of utricle, 92 (Fig. 60), 94, 95; land forms, 91; morphology, 73, 103–107; position in biological classification, 8; regeneration, 104 (Fig. 70), 105 (Fig. 71),

INDEX

435

- 106; relation to *Pinguicula*, 181; "rhizoids," 96, 98, 99 (Fig. 66); seedlings, 100 (Figs. 67, 68); seeds, 99, 100; turions, 101–103, 219, 220; utricles, 91, 92 (Figs. 59, 60), 93 (Fig. 61), 94, 95 (Fig. 62), 96 (Figs. 63, 64), 97, 98 (Fig. 65); vegetative luxuriance, 215; water absorption, 270
- Utricularia Bremii*, 93 (Fig. 61), 95 (Fig. 62), 97, 299
- Utricularia exoleta*, 100 (Fig. 68)
- Utricularia flexuosa*, 92 (Fig. 60), 93
- Utricularia Hookeri*, 93
- Utricularia Humboldtii*, 109
- Utricularia inflata*, 99, 229 (Fig. 150)
- Utricularia inflexa*, 99
- Utricularia intermedia*, 91, 94, 97, 101, 219, 220 (Fig. 143)
- Utricularia minor*, 91, 96 (Figs. 63, 64), 97, 100, 102 (Fig. 69), 108 (Fig. 74), 168, 299
- Utricularia neglecta*, 92 (Fig. 59), 93, 97, 99 (Fig. 66), 104
- Utricularia nelumbifolia*, 108, 109
- Utricularia ochroleuca*, 97
- Utricularia quinqueradiata*, 99
- Utricularia stellaris*, 99
- Utricularia vulgaris*, 91, 93, 94, 97, 98 (Fig. 65), 100 (Fig. 67), 101, 102, 104 (Fig. 70), 105 (Fig. 71), 106 (Fig. 72), 107 (Fig. 73), 215, 220, 227
- Valerian, Grew on heterophylly in, 155
- Vallisneria*, contraction of peduncle, 235, 239; experimental cultures, 265; leaves, 57, 140, 169; ovules, 314; pollination, 57, 235, 236; root anatomy, 208, 209 (Fig. 139); toleration of salt, 134
- Vallisneria spiralis*, 134, 140, 209 (Fig. 139), 235, 236
- Vegetative reproduction, 210–226, etc.
- Venezuela, 113, 122, 191, 291
- Veronica*, anatomy of submerged and air shoots, 201, 259
- Veronica Anagallis*, 201, 259
- Vicia*, effect of water, 200; nutation of roots, 207
- Vicia sativa*, 200
- Victoria regia*, affinities, 38; evolution of heat from flower, 34; in shallow water, 31; leaf succession, 32–34; peltate leaf, 30; rate of growth, 214; ripening of fruit under water, 300; seedlings, 32, 33 (Fig. 16), 34; size, 80; Spurwing nesting on leaf, 300
- Villarsia*. See *Limnanthemum*
- Villarsia ovata*, 166
- Violet, Water. See *Hottonia palustris*
- Vorticellidae*, 237
- Vosges, 225, 287
- "Wapatoo," 17
- Water Aloe. See *Stratiotes aloides*
- Water Buttercup. See *Ranunculus* sect. *Batrachium*, *R. aquatilis*, etc.
- Water Chestnut. See *Trapa natans*
- Water Crowfoot. See *Ranunculus* sect. *Batrachium*, *Ranunculus aquatilis*, etc.
- Water forms, of *Achillea*, 199; *Caltha*, 198, 199 (Fig. 129); *Cirsium* (*Cnicus*), 198, 199 (Fig. 130), 200; *Cuscuta*, 199 (Fig. 131); *Hydrocotyle*, 200, 201 (Fig. 132); *Menyanthes*, 199; *Ranunculus*, 198, 200, 203 (Figs. 134, 135); *Trifolium*, 199 (Fig. 131), etc.
- Water-fowl and dispersal of hydrophytes, 35, 298–302
- Water Hyacinth. See *Eichhornia speciosa*
- Water Lobelia. See *Lobelia Dortmanna*
- Water Loosestrife. See *Lythrum Salicaria*
- Water Milfoil. See *Myriophyllum*
- Water Parsnip. See *Sium latifolium*
- Water Plantain. See *Alisma Plantago*
- Water Plantain, Lesser. See *Echinodorus ranunculoides*
- Water pores, 267–269, etc.
- Water-shield. See *Cabomba*
- Water-snails. See Snails, water
- Water Soldier. See *Stratiotes aloides*
- Water Speedwell. See *Veronica Anagallis*
- Water Starwort. See *Callitriche*
- Water stomates. See Stomates, water
- Water Thyme. See *Elodea canadensis*
- Water Violet. See *Hottonia palustris*
- Waterlily, Giant. See *Victoria regia*
- Waterlily, White. See *Castalia alba*
- Waterlily, Yellow. See *Nymphaea lutea*
- Waterweed, American, or Canadian. See *Elodea canadensis*
- Weddellina squamulosa*, 113
- Whitchurch Weir, 210
- White Moss Loch, Perthshire, 288 (Fig. 165), 289
- White Waterlily. See *Castalia alba*
- Wicken Fen, 20
- Willow Herb, 311
- Wolffia*, conveyance by birds, 300; in biological classification, 7; member of Lemnaceae, 74; size, structure, wintering habits, 80
- Wolffia brasiliensis*, 80, 300
- Wolffia Micheli*, 80
- Xerophyte, aquatic, 310
- Xylem and water conduction, 173, 174
- Yellow Waterlily. See *Nymphaea lutea*
- Yew-tree, 316
- Yorkshire, 303
- "Youth forms" of Conifers, 155
- Zannichellia*, anatomy, 62, 63, 173;

Cambridge University Press

978-1-108-01732-9 - Water Plants: A Study of Aquatic Angiosperms

Agnes Arber

Index

[More information](#)

436

INDEX

- flowers, 70 (Fig. 45), 71, 215, 315, 316;
fruit and seedling, 246 (Fig. 159),
248; hydrophilous pollination, 70, 71,
134, 346; in biological classification,
6; rheotropism, 282; root-hairs from
"collet," 245; stability of seedling,
245; twining roots, 205, 206 (Fig. 137)
- Zannichellia palustris*, 134, 173, 205,
206 (Fig. 137), 215, 282
- Zannichellia polycarpa*, 70 (Fig. 45), 71,
246 (Fig. 159)
- Zonation, 287-289
- Zostera*, affinities, 315; apical openings
of leaves, 269; association with Algae,
123; distribution, 302; food of Brent
Geese, 302; fruit, 248 (Fig. 161);
habit, 124, 140; in biological classi-
fication, 6; leaf anatomy, 128 (Figs
85, 86), 131, 164; macropodous
embryo, 248 (Fig. 161); pollen,
thread-like, 125; pollination, 127,
129, 237
- Zostera marina*, 123, **127, 128** (Figs. **85,**
86), **129**, 248
- Zostera nana*, 125, 302
- "*Zostera oceanica*," 125