

TABLE OF CONTENTS

CHAPTER I. INTRODUCTION TO THE THEORY
OF QUADRIC SURFACES

	PAGES
Preliminary remark. Equation of a quadric surface	1
Definition of a quadric surface by means of its lines	1—3
Definition of a quadric cone	4
The duality of the definition of a quadric	4
Application of the symbols; equation of a quadric	5—7
Quadric defined by two conics having two points in common, and a line meeting these conics	7—10
Three conics of which every two have two points in common lie on a quadric	10
Two conics which have two points in common lie on two quadric cones	10
Quadric defined by two dually related central systems	11—13
Quadric representing a plane with two fundamental points . . .	13, 14
General homogeneous quadric polynomial in four variables . . .	15, 16
Equation of a quadric cone	17
Condition for quadric form to represent a cone	18
Conditions for quadric form to represent two planes	19
Conditions for quadric form to represent a repeated plane . . .	20
General properties of a quadric given by its equation	20—26
Tangent cone, or enveloping cone, of a quadric	26
Polar plane, and tangent plane, descriptively defined	27
Polar line of a line in regard to a quadric	28
Polar point of a plane in regard to a quadric	29, 30
Polar point of a plane in regard to a quadric, expressed by the symbols	31
Examples of the general properties of quadrics (Exx. 1—19) . . .	31—56
Ex. 1. <i>The self-dual character of a quadric, in particular form</i> . . .	31
Ex. 2. <i>The same in general</i>	32
Ex. 3. <i>Two conjugate sections have tangent lines at a common point harmonic in regard to the generators; the dual of this</i>	32, 33
Ex. 4. <i>Self-polar tetrad in regard to a quadric</i>	33, 34
Ex. 5. <i>Tetrad of points whose opposite joins are conjugate lines; self-conjugate pentad; construction from two tetrads in perspec- tive; analytical treatment; expression of a quadric by five squares</i>	34—40
Ex. 6. <i>A general self-polar figure in space of any number of dimen- sions</i>	40, 41

	PAGES
Ex. 7. Joins of two reciprocal tetrads are generators of a quadric	41, 42
Ex. 8. Symmetry of two particular quadrics arising in preceding example	42—44
Ex. 9. Dandelin's figure of generators; its application to Steiner points in Pascal's figure	44—47
Ex. 10. Construction of a self-conjugate hexad. Expression of a quadric by six squares	47—49
Ex. 11. Construction of an incomplete tetrad, or pentad, or hexad, of points of a quadric	49, 50
Ex. 12. Condition for one quadric to be outpolar to another	50—52
Ex. 13. A property of two self-polar tetrads	52
Ex. 14. Quadric for which two proper tetrads are reciprocal	53
Ex. 15. A theorem of Chasles for the intersections of a quadric with the joins of four points	53, 54
Ex. 16. A further theorem for the intersections of a quadric with the joins of four points	54
Ex. 17. Bipunctual quadric, the harmonic conjugate of two planes	54
Ex. 18. The equation of a quadric touching the four planes of reference	55
Ex. 19. A birational transformation of a quadric into itself. Its geometrical meaning	55, 56
The coordinates and equations of a line in three dimensions. Elements of the theory of a linear complex	56—69
Ex. 1. The analytical conditions for a plane to contain a line	57
Ex. 2. Common element of a line and point, or of a line and plane	57, 58
Ex. 3. The condition of intersection of two lines	58
Ex. 4. Lines of a quadric as a linear system	58
Ex. 5. The tangent planes from a line to a quadric	58
Ex. 6. Coordinates of the polar line of a line in regard to a quadric	58, 59
Ex. 7. Condition a line should touch a quadric	59
Ex. 8. Change of line coordinates by change of reference points	59, 60
Ex. 9. Generators of a quadric, of one system, belong to three linear complexes	60
Ex. 10. First properties of a focal system and linear complex. Analytical definition. Derivation from generators of a quadric. Sylvester's derivation. Polar lines. Linear complex consists of those lines of a focal system which intersect, and coincide with, their polars. Invariant of two linear complexes	60—66
Ex. 11. Pole of a plane in a focal system	66
Exx. 12—14. Geometrical theorems. A theorem of Sylvester's	66
Ex. 15. Chasles' generation of a linear complex	66, 67
Exx. 16—22. The focal system and Moebius tetrads	67, 68
Ex. 23. The two lines common to a linear complex and a quadric	68
Ex. 24. Theorem for the mutual perpendiculars of lines	68

Contents xi

	PAGES
Ex. 25. <i>The tangent lines of a cubic curve belong to a linear complex</i>	68
Ex. 26. <i>A skew pentagon determines a focal system</i>	68
Ex. 27. <i>The polar line of a line in regard to a general quadric</i>	69
Ex. 28. <i>The tangent lines of the curve of intersection of two quadrics; eight of these meet an arbitrary line</i>	69
CHAPTER II. RELATIONS WITH A FIXED CONIC. SPHERES, CONFOCAL SURFACES; QUADRICS THROUGH THE INTERSECTION OF TWO GENERAL QUADRICS	
Parallel lines and planes. Middle points. Lines and planes at right angles	70, 71
Exx. 1—3	71
Spheres and Circles	72, 73
Ex. 1—5. <i>Plane sections of a sphere</i>	73
Relations of two spheres. Coaxial spheres	74—76
Ex. 1. <i>Two circles with two common points lie on a sphere</i>	77
Ex. 2. <i>The radical planes of the pairs of three spheres meet in a line</i>	77
Ex. 3. <i>Any three circles are met in two points by another properly taken circle</i>	77
Ex. 4. <i>Analytic treatment of the relations of two spheres. Centres of similitude</i>	77, 78
Ex. 5. <i>A sphere can be constructed of given centre to pass through a given point</i>	78
Ex. 6. <i>Sphere determined by perpendiculars from a fixed point to planes through another point</i>	78
Ex. 7. <i>Theorems for an orthogonal tetrad</i>	78
Relations of a quadric in general with the absolute conic. The circular sections	79
Umbilici of a quadric. Generators through these	80
Relations of a quadric cone with the absolute conic. Cone of revolution; or right circular cone	81
Ex. <i>Analytical conditions for a right circular cone</i>	82
Quadric touching the absolute plane	82, 83
Ex. 1. <i>Umbilici of a paraboloid</i>	83
Ex. 2. <i>Two circular sections of a quadric not in parallel planes lie on a sphere</i>	83
Ex. 3. <i>Condition a plane should give a circular section</i>	83
Ex. 4. <i>Centres of plane sections of a quadric</i>	83, 84
Ex. 5. <i>Locus of line of intersection of two perpendicular tangent planes of a quadric</i>	84
Ex. 6. <i>Locus of normals of a quadric at points of a generator</i>	84

	PAGES
Line of striction on a quadric	84
Ex. 1—6. <i>Properties of the line of striction</i>	85, 86
The director sphere of a quadric. Analogue of Gaskin's theorem .	86—89
Ex. 1. <i>The intersection of the director sphere with the quadric</i> .	89, 90
Ex. 2. <i>Condition two generators should be at right angles</i> . . .	90
Ex. 3. <i>Point on quadric and director sphere at which three con-</i> <i>focals coincide</i>	90
Ex. 4. <i>Generalisation of director sphere</i>	90, 91
Ex. 5. <i>Locus of intersection of three perpendicular tangent lines of</i> <i>a quadric</i>	91
Confocal quadrics, defined algebraically	91, 92
Points through which confocals are not distinct	92
Ex. 1. <i>The umbilical generators lie on a quartic surface</i>	92, 93
Ex. 2. <i>Polar reciprocal of a confocal in regard to original</i>	93
Axis of a plane in regard to a system of confocals. Plane touching all the confocals	93
Alternative ways of initiating the system of confocal quadrics. Dual of quadrics through common curve of two quadrics. By common tangent planes of two quadrics	94
Ex. <i>Confocal paraboloids</i>	94, 95
The relations of the four focal conics. Locus of vertex of right circular cone circumscribing a quadric. Section of quadric by plane at right angles to a focal conic has a focus on this conic .	95
Ex. 1. <i>Equation of enveloping cone of quadric drawn from point</i> <i>of focal conic</i>	96
Ex. 2. <i>Generalisation of focal distance theorem for a conic. The</i> <i>geometrical meaning</i>	96
Ex. 3. <i>Planes through a point whose sections with the quadric have</i> <i>the point as focus</i>	96
Ex. 4. <i>A system of right circular cones with given vertex</i>	96
The axes of a confocal system of quadrics	97
Ex. <i>The condition for a line to be one of these axes</i>	97, 98
The axes of the confocal system lying in a plane touch a conic ; those through a point lie on a quadric cone	98
Ex. <i>The axes are the lines which are met by the principal planes in</i> <i>related ranges</i>	99
The tetrahedral complex. The associated conics and cones	99
Various geometrical conditions for an axis	99, 100
Ex. 1. <i>Condition two polar lines should be at right angles</i>	100
Ex. 2. <i>Axes of the confocals lying in a plane</i>	100
Ex. 3. <i>The parabola touched by the axes lying in a plane is in-polar</i> <i>to the section of a confocal by this plane</i>	101
Ex. 4. <i>An axis of the confocals meets one of the surfaces in points</i> <i>whereat the normals intersect</i>	101

Contents xiii

	PAGES
Ex. 5. <i>Axis of the confocals as axis of a plane section of one of the confocals</i>	101
Ex. 6. <i>Tangent planes of a surface where polar line of an axis meets it</i>	102
Ex. 7. <i>Plane of which a line of the tetrahedral complex is the axis</i>	102
Ex. 8. <i>Normal of one confocal meeting another of the surfaces</i>	102
Ex. 9. <i>The locus of the conics which are touched by the axes intersecting a line is a rational quartic surface</i>	102
The normals of a quadric, and of confocal quadrics, which pass through a point	102—104
The polar planes of a point in regard to the confocals determine a cubic developable	103
Ex. 1. <i>The cubic curve defining the six normals from a point to a quadric</i>	104
Ex. 2. <i>General definition of a cubic curve in space illustrated in this case</i>	104
Exx. 3, 4. <i>The planes containing two triads of feet of normals from a point to a quadric</i>	104, 105
Ex. 5. <i>Six points determining the cubic curve</i>	105
Ex. 6. <i>A property of the six feet of normals</i>	105
Ex. 7. <i>The normals of a quadric at the points of a plane section determine a ruled quartic surface with a cubic curve as double curve</i>	105
The normals of the confocals at right angles to a given line; and the polar lines of this line	106, 107
Ex. 1. <i>Equation of locus of polar lines of a given line in regard to the confocals</i>	107
Ex. 2. <i>Equations of the cubic curve which is the locus of points of contact of tangent planes to the confocals from a given line</i>	107
Special positions for the line	107, 108
Ex. <i>Points of contact of parallel tangent planes of the confocals lie on a conic</i>	108
The enveloping cones drawn from a point to the surfaces of the confocal system; and the tangent planes drawn from a line. Two of the confocals touch a line, with their tangent planes at right angles. The generators of three intersecting confocals are focal lines of enveloping cones	108, 109
Ex. 1. <i>Equation of enveloping cone to a quadric, and of the quadric itself, referred to normals at vertex. Normals lie in threes on four planes</i>	110
Ex. 2. <i>Chasles' theorem of correspondence between two systems of confocals</i>	110, 111
Ex. 3. <i>A property for a line, touching two confocals, leading to the definition of geodesics</i>	111, 112

	PAGES
Ex. 4. <i>The angle between two geodesic tangents of a line of curvature reduced to the angle between two tangents of a conic . . .</i>	112, 113
Ex. 5. <i>Quadric of rigid generators transformable into a confocal . . .</i>	113—116
Ex. 6. <i>The generators, of one system, of confocal surfaces, at points of curve of intersection of two other fixed confocals, generate a conic on a principal plane. Proof of Poncelet's porism for conics . . .</i>	116—118
Ex. 7. <i>Ivory's theorem, in generalised form</i>	118
Ex. 8. <i>The developable of common tangent planes of the confocals . . .</i>	118, 119
Dual method of developing the theory of confocals	119, 120
Ex. 1. <i>Equation of quadric locus of polar lines of a fixed line in regard to quadrics through the intersection of two</i>	121
Ex. 2. <i>Equation of cone touched by tangent planes of the family at points where surfaces are met by a line in a principal plane . . .</i>	121

CHAPTER III. CUBIC CURVES IN SPACE. THE INTERSECTION OF TWO OR MORE QUADRICS

The curve of intersection of two quadrics. Enumeration of possible cases	122, 123
Exx. 1—9. <i>Equations of quadrics having the respective intersections</i>	123—125
The cubic curve in three dimensions	125
Definition by two quadric cones. Tangent line and osculating plane of curve	126, 127
Definition by related axial pencils of planes	127
General quadric containing the curve	128
Cubic curve in space determined by six points	129
Cubic curves, lying on a quadric, passing through five points	129
Intersections with generators, of cubic curve lying on a quadric	129
Cubic developable dual to a cubic curve	130
The cubic curve as a self-dual construct	131
Surface formed by tangent lines of cubic curve	132
Four tangents of cubic curve meet any line	132
Concrete illustration of cuspidal edge of a developable	132
Cubic developable considered initially	133
Examples in regard to the cubic curve in space (Exx. 1—31)	134—149
Ex. 1. <i>Tangent line and osculating plane of curve at a point belong to cone projecting curve from this point</i>	134
Ex. 2. <i>Points and osculating planes of curve define a focal system</i>	134
Ex. 3. <i>Tetrahedral complex of chords depending upon four points of the curve</i>	134
Ex. 4. <i>Equation of general quadric containing the curve. Coordinates of chord of the curve through an arbitrary point</i>	134, 135
Ex. 5. <i>A (1, 2) correspondence of generators of a quadric determining a cubic curve on the quadric</i>	135

Cambridge University Press

978-1-108-01779-4 - Principles of Geometry, Volume 3

H. F. Baker

Table of Contents

[More information](#)

<i>Contents</i>	XV
	PAGES
Ex. 6. <i>Quadrics through six points of the curve determine an involution on any chord</i>	135
Ex. 7. <i>Various generations of a cubic developable</i>	135
Ex. 8. <i>Cone touched by planes joining a point of the curve to the chords which meet an arbitrary line</i>	135, 136
Ex. 9. <i>Planes meeting six lines and a common transversal of these in points lying on a conic belong to a cubic developable</i>	136
Ex. 10. <i>The form apolar to any system of binary forms of the same order</i>	136
Ex. 11. <i>The Hessian pair of three points of the curve</i>	136, 137
Exx. 12, 13. <i>Four mutually inscribed tetrads arising for the curve</i>	137, 138
Exx. 14, 15. <i>General case of four mutually inscribed tetrads</i>	138
Ex. 16. <i>The general involution of sets of four points of the curve</i>	138, 139
Ex. 17. <i>Number of common points of two cubic curves lying on a quadric</i>	139
Ex. 18. <i>A cubic curve to pass through five points and have a given chord</i>	139
Ex. 19. <i>A cubic curve to pass through four points and have two given chords when possible</i>	139, 140
Ex. 20. <i>A cubic curve to pass through three points and have three given chords</i>	140
Ex. 21. <i>A cubic curve to pass through two points and have four given chords</i>	140
Ex. 22. <i>A cubic curve to pass through one point and have five given chords</i>	140, 141
Ex. 23. <i>Two general cubic curves in space have ten common chords</i>	141, 142
Ex. 24. <i>There are six cubic curves having six given lines, of general position, as chords</i>	142
Ex. 25. <i>A cubic transformation giving another proof of preceding results. The condition that four lines should be tangents of a cubic curve</i>	142—145
Ex. 26. <i>A cubic curve to meet each of four lines in a pair of points of an involution given thereon</i>	145
Ex. 27. <i>The condition for a cubic curve to be outpolar to a quadric. Self-polar tetrads, self-conjugate pentads and hexads thereon</i>	145—147
Ex. 28. <i>Cubic developable inpolar to a quadric</i>	147
Ex. 29. <i>Cubic curve containing six points of two self-polar tetrads has join of other two for chord. Three independent quadrics contain the eight points. Associated points</i>	147, 148
Ex. 30. <i>Squares of eight points of a cubic curve linearly related</i>	148
Ex. 31. <i>A theorem of Lie, for complex curves, in generalised form</i>	148, 149
The rational curve of order n in space of n dimensions	149, 150
Examples thereon (Exx. 1—4). <i>Number of conics meeting eight lines</i>	150, 151

	PAGES
The curve of intersection of two general quadric surfaces; its irrationality	151
Examples (Exx. 1—3). <i>The common self-polar tetrad</i>	151, 152
The determination of a quadric by given conditions	152, 153
Ex. 1. <i>The condition for two quadrics to intersect in two conics</i>	153
Ex. 2. <i>Three quadrics intersect in eight points</i>	153
Ex. 3. <i>A detailed analytical consideration</i>	153, 154
Ex. 4. <i>Number of remaining common points of three quadrics with a common curve</i>	154
Ex. 5. <i>Weddle surface as locus of vertex of quadric cone containing six points</i>	154
Ex. 6. <i>Hesse's sextic curve, locus of vertex of quadric cone containing seven points</i>	154
Theorems for eight associated points	154—158
Exx. 1—3. <i>Four lines, determined by the eight points, which lie on a quadric (in analogy with Pascal's theorem)</i>	154, 155
Ex. 4. <i>A second construction for eight associated points</i>	155
Ex. 5. <i>Two determinations of the eighth point when seven are given</i>	155, 156
Ex. 6. <i>Determination of the ninth intersection of two plane cubic curves when eight are given</i>	156, 157
Ex. 7. <i>Associated points interpreted in space of four dimensions</i>	157, 158

CHAPTER IV. THE GENERAL CUBIC SURFACE; INTRODUCTORY THEOREMS

The theorem of a double-six of lines	159—161
Reciprocity of the figure. Schur's quadric	161—163
Another determination of the double-six, by a porismatic relation	163, 164
The Steiner trihedral pairs	164—166
Examples of the figure of twenty-seven lines (Exx. 1—5)	166—170
Ex. 1. <i>Numbers of sets of lines of which no two meet</i>	166
Ex. 2. <i>The sets of three coplanar lines</i>	166
Ex. 3. <i>The number of constants on which the figure depends, and the general cubic polynomial in four coordinates</i>	166—168
Ex. 4. <i>Another proof of the reciprocity of the figure</i>	168
Ex. 5. <i>A double-six determined by five arbitrary points</i>	168—170
Geometrical definitions of a cubic surface	170—174
Cubic curves having as chords five lines with a common transversal	172
Deduction of a double-six of lines	173
Ex. 1. <i>Pairs of ranges, of four points, intersections of lines, which are in perspective from a point of intersection of two lines</i>	174
Ex. 2. <i>Deduction of a pair of Steiner trihedrals</i>	174
Ex. 3. <i>The number of such pairs is one hundred and twenty</i>	174

Cambridge University Press

978-1-108-01779-4 - Principles of Geometry, Volume 3

H. F. Baker

Table of Contents

[More information](#)

<i>Contents</i>	xvii
	PAGES
The definition of a cubic surface in general. The equation of the surface	174, 175
The cubic curves of the surface which belong to a particular double-six	175—177
Each curve is determined by two points. Two curves of the same system have one common point	175
The cubic curves determined by quadrics through three (suitable) lines	176
A curve of either system lies on a quadric containing a curve of the other system	176
Two cubic curves of complementary systems meet in five points	176
The quadrics containing the cubic curves are outpolar to the Schur quadric associated with the double-six	177
The five common points of two complementary cubics form a self-conjugate pentad in regard to the Schur quadric	177
Ex. 1. <i>Consideration of two cubic curves of the surface with one common point</i>	177, 178
Ex. 2. <i>A property of plane cubic curves: if conics be drawn through four given points of the curve, the line joining the two remaining intersections passes through a fixed point of the curve. General theorem of coresiduation for a plane curve</i>	178, 179
Ex. 3. <i>Illustration of the theorem of coresiduation for curves on the cubic surface</i>	179, 180
Ex. 4. <i>Statement of the theorem in general</i>	181
Ex. 5. <i>The conics lying in planes through a line of the surface meet the line in pairs of points of an involution</i>	181
Ex. 6. <i>All polar quadrics meet a line of the surface in pairs of points of an involution</i>	182
Ex. 7. <i>Definition of the cubic surface from two cubic curves with a common point. Number of lines meeting four curves of given orders</i>	183, 184
Definition of a cubic surface by three related star-figures, or central-systems	184—186
Meaning of related star-figures	184
Resulting equation of the cubic surface	186
Ex. 1. <i>The cubic curves, and the quadrics containing these, arising from this form of the equation</i>	186
Ex. 2. <i>All Schur quadrics are inpolar to the polar quadrics; there are six linearly independent Schur quadrics in tangential form; they may be expressed by six point-pairs of the Hessian</i>	187
Ex. 3. <i>The equation of the Hessian for a certain form of the equation of the cubic surface</i>	187, 188
The representation of the cubic surface on a plane. Representation of the systems of curves of the surface by means of seven systems	189—192

	PAGES
Ex. 1. <i>A further way of obtaining the representation upon a plane</i>	192, 193
Ex. 2. <i>Representation of complementary systems of cubic curves</i>	193
Ex. 3. <i>Representation of conics of the cubic surface</i>	194
Ex. 4. <i>Representation of all cubic curves of the cubic surface</i>	194
Ex. 5. <i>Number of intersections, freedom and genus, for systems of curves of the cubic surface</i>	194
Ex. 6. <i>Converse derivation of the cubic surface from six arbitrary points of the plane</i>	194, 195
Ex. 7. <i>Given seven lines with a common transversal, there is a point from which the eight lines are projected by planes touching a quadric cone. Corresponding theorems given six lines, or five lines, with a common transversal</i>	195, 196
Derivation of a plane quartic curve from a cubic surface	196—201
The inflexional lines at a point of the cubic surface. They are the generators of the polar quadric of the point	197
The sextic curve of contact of tangent planes from a point of the cubic surface. Projects into a plane quartic curve	198
The sixty-three systems of conics, each touching the plane quartic curve in four points; each system representable by a pair of double tangents of the curve	198
Statement of the facts for a degenerate contact cone containing the tangent plane at the point of projection. Steiner system of bitangents of a plane quartic curve	199, 200
Proof of the results for the case of planes drawn to two skew lines of the cubic surface. The contact cones are those containing the cubic curves, through the point of projection, associated with the two lines. Relation with the Schur quadric. The transversals from a point of the surface to the six pairs of a double-six of lines lie on a quadric cone	200, 201
Ex. 1. <i>The quadric cone containing the points of contact of five tritangent planes of the cubic surface through a line. A property of cubic curves through these five points</i>	201, 202
Ex. 2. <i>The algebraic treatment of the case of two skew lines of the surface</i>	202, 203
The equation of the cubic surface referred to two Steiner trihedrals. Deduction of the lines of the surface	203—206
The Cremona form of the equation of the cubic surface	206
The Sylvester form of the equation of a general cubic surface	206—208
The polars in regard to a cubic surface. The Hessian surface	208—210
Ex. 1. <i>The vertices of a pair of Steiner trihedrals are corresponding points of the Hessian</i>	210
Ex. 2. <i>The joins of the eight points of which a given plane is the polar plane</i>	210

Contents

xix

	PAGES
Ex. 3. <i>The remaining intersections with the Hessian of the join of a pair of corresponding points</i>	211
Ex. 4. <i>The cubic surface enveloped by the polar planes, in regard to the cubic surface, of the points of a plane, touches the Hessian along a sextic curve</i>	211
Ex. 5. <i>A construction for the expression of a cubic surface by six cubes</i>	211
Ex. 6. <i>Every line of the cubic surface touches the Hessian in two points. The twenty-four intersections with the Hessian of the lines of a double six</i>	211
Ex. 7. <i>The equation of the Hessian when the cubic surface is expressed by six cubes. The question of the generality of Sylvester's form</i>	211, 212
The representation of a general cubic surface upon a quadric surface. The twenty-seven elements on the quadric corresponding to the twenty-seven lines of the cubic surface	212—214
Note on the cubic surface with four double points	214—218
The representation upon a plane. The fundamental cubic curves	215
The tangent cone from any point of the surface breaks up into two quadric cones	216
The inflexional curves of the surface. One touches every fundamental cubic curve	216, 217
Ex. <i>The envelope sextic curve associated with a point. Proof of a theorem leading to a generalisation of the theorem of the nine-point circle</i>	218—221
The Steiner quartic surface. Representation upon a plane	221, 222
Ex. 1. <i>Envelope of polar planes, in regard to a cubic surface, of points of a plane in connexion with Steiner's surface</i>	222
Ex. 2. <i>The inflexional curves of Steiner's surface</i>	222
Ex. 3. <i>These curves are rational quartics. A property of these</i>	222, 223
Ex. 4. <i>A formula for inflexional curves which includes the cases of the cubic surface with four double points and of Steiner's surface</i>	223
Ex. 5. <i>Veronese's surface</i>	223
Ex. 6. <i>The general cubic surface derived from two tetrads in perspective. The cubic surface with four double points as a pedal locus. Kronecker and Castelnuovo's theorem</i>	223
CORRECTIONS FOR VOLUMES I AND II	224
INDEX	225—228