

INDEX

- Anchor Ring, or Tore, and Dupin's Cyclide, 193, 199
- Angle between two circles, in projection from a quadric, 7; unaltered by inversion, 14
- Apolar, or conjugate, linear complexes, 42; apolar relation for point of contact of Hart section, 76
- Apollonius's problem for circles, 66
- Associated lines, five, in fourfold space, 114; all met by planes, 120; planes meeting these through any point, 122; associated system of five planes, 118
- Bauer, on a quartic surface, 20
- Bennett, theorem on circles through threes of four points, 17; Hart section, equations for, 86
- Bertini, on Veronese's surface, 54
- Bitangents of Kummer's surface, from lines of Segre's cubic locus, 156
- Bobillier, theorem for pedal circles, 22
- Burkhardt, the group of the lines of a cubic surface, 105
- Burnside: a figure in fourfold space, 105; the group of the lines of a cubic surface, 105; invariants of a linear group, 210
- Caporali, tetrahedral complexes containing a quadratic congruence, 242
- Cardioid, 99
- Cartesian, 99
- Castelnuovo and Kronecker, surface with double infinity of reducible plane sections, 55
- Caustic, or focal, surface of a quadratic congruence, 225
- Cayley, representation of a line by a point of fivefold space, 40; on Veronese's surface, 54; algebraic solution of Malfatti's problem, 68
- Centre of a circle, in projection from a quadric, 7
- Chords common to two curves, 51
- Circle, obtained by projection of plane section of a quadric, 1; circle and two inverse points changed into same by inversion, 14; circle of similitude, obtained by projection, 15; circles of similitude of the pairs of four circles, 15; circles in a plane, chain of theorems for, 29; circle represented by a line of fourfold space, 39; Hart and Feuerbach circles, Chapter II, p. 65; circles meeting given circles at equal, or at given angles, 67; three circles with three concurrent common tangents, 67
- Clifford, a theorem for circles, 1, 31, 64, 105
- Complex, tetrahedral, determined by planes in fourfold space, 32; linear, determined by planes in fourfold space, 34; linear, represented in fivefold space, 42; six linear complexes mutually conjugate, 42; lines common to three, or to four linear complexes, 43, 44; linear, point and polar plane represented in fivefold space, 48; linear, six conjugate, constructed from six points, 139; linear, six conjugate, from lines of Segre's cubic locus, 155; linear, six conjugate, poles of a plane lie on a conic, 204; quadratic, represented in fivefold space, 213; quadratic, condition for singular point, 214; quadratic, rationality of, 234; tetrahedral, in fivefold space, 239. See also *Focal System*
- Confocal Cyclides, cutting at right angles, 181; equation for, 183
- Congruence of lines, its order and class, 49; for chords of a curve, 50; number of lines common to two, 51; quadratic, in tangent solid of Segre's quartic locus, 137; six congruences associated with a Kummer surface, 219, 223; quadratic, caustic or focal system of, 225; six quadratic, equations for common singular points, 226; quadratic, singular points and planes, 240; quadratic, lies in

- forty tetrahedral complexes, 242; quadratic, and Moebius tetrads, 241
- Conic: conics touching fives of six arbitrary lines, 1; triangularly circumscribed to six conics, 3, 5; in an arbitrary plane, determined by the joins of five points, 11; two conics in fivefold space which are mutually conjugate, 43; conics through a point of Segre's quartic locus, 127; conics lying on quartic surface in fourfold space, 171
- Conjugate, or apolar, linear complexes, 42
- Cremona, triangularly circumscribed conic and line coordinates, 6
- Cubic curve: conic touching seven lines associated with, 4; five arbitrary points on, 11; three tetrads of points on, 28; tetrads of points determined by two points, 29; chords of, represented in fivefold space, 52; in plane, through two points, inverse of quartic with two double points, 97
- Cubic surface, with four double points, theorem for, 28; group of lines of, 105; cubic envelope, by projection of Segre's figure, 149; by projection of quartic surface in fourfold space, 173
- Curvature, lines of; upon a Cyclide, 186; Dupin's theorem, 188; general theorems for, 189; differential equation for, 190
- Curve, in n -fold space, intersection of $(n-1)$ quadrics, 185
- Cyclide: arising by projection from fourfold space, 174; enveloped by spheres, 175; as locus of limiting points of fixed sphere and variable tangent plane of a quadric, 176; tangent planes of, at a point of double conic, 177; four focal conics of principal quadric surfaces, 178; angle between enveloping spheres of different systems, 179; is inverted into itself from five centres, 179; four points of double curve where tangent planes coincide, 180; three confocal through given point, 181; equations for confocal cyclides, 183; normal of, by projection from fourfold space, 184; lines of curvature upon, 186; spheres having stationary contact with, 187; Dupin's Cyclide, 190, 193, 199; with cuspidal conic, 192; by inversion of a quadric, 192; with planes of symmetry, 193; general, irrational form of equation, 194; with two double points, inversion of, 194; with two double points, equations for, 195; with four double points, equations for, 196; a particular surface, Kummer's equation for, 197; Dupin's, treated tangentially, 201; Dupin's, points expressed by two parameters, 202; lines of curvature correspond to inflexional curves on Kummer's surface, 231
- Darboux: surface containing a doubly infinite system of conics, 54; on plane quartic curves, and Cyclides, 172; generalisation of lines of curvature and of geodesic lines, 190; a generalisation of Dupin's Cyclide, 198; inversion of Dupin's Cyclide, 200; on the existence of space of four dimensions, 200; congruences of lines, 225
- Del Pezzo, on Veronese's surface, 54, 55
- Dickson, the group of the lines of a cubic surface, 105
- Director conics, for plane quartic curve with two double points, are confocals, 91
- Double four: of lines on quartic surface in fourfold space, twenty such, 170; of planes, 238
- Double-six theorem, proved in fivefold space, 63
- Double tangents of Kummer's surface, 225
- Dupin: theorem on lines of curvature, 188; Dupin's Cyclide, 190, 193, 199, 201, 202. See *Cyclide*
- Elte, on a six-dimensional polytope, 105
- Feuerbach's theorem, 65; particular case of Hart's theorem, 78
- Figure: of twentyseven points in fourfold space, 104; of fifteen lines and points in fourfold space, 113; arising from four arbitrary planes, 117

- Focal conics of a confocal system of quadrics, 92; descriptive form of theorem of intervals, 103
- Focal system: point and polar plane represented in fivefold space, 48; six conjugate, scheme for, 133; arising in tangent solid of Segre's quartic locus, 135. See *Complex, linear*
- Foci of plane quartic curve with two double points, 92
- Focus, and focal line, of a singular plane in fivefold space, 219
- Fontené, theorem for pedal circles, 22
- Fresnel's wave surface, 217
- Gergonne, solution of Apollonius's problem, 66
- Göpel, equation of Kummer's surface, 216
- Grace: generalisation of a theorem for circles, 8; theorem for pedal circles, 23; theorem for spheres touching planes, 58; Wallace's theorem for space of even dimensions, 59; theorem for six lines with a common transversal, 60; on systems of spheres, 105, 112
- Group, linear, in four homogeneous variables, 208
- Harmonic inversion, with fundamental point and solid, 117; with two fundamental planes, 118; with fundamental line and plane, 119
- Harmonically conjugate quadric, 72
- Hart: theorem for circles, 65; proof of Steiner's solution of Malfatti's problem, 68; Hart circle of three circles, orthogonal to their orthogonal circle, 71; general case, 72, 75; apolar relation for point of contact, 76; Hart circle for spherical triangle, 77; Feuerbach circle as particular case, 78; radius of Hart section of a quadric, 81; Hart section of a quadric, sufficient condition is necessary, 82; Hart circle of three circles, defined by angles of intersection, 88
- Haskell, generalisation of a theorem for circles, 8
- Hudson, on Kummer's quartic surface, 208, 227, 233
- Humbert, a curve on the Weddle surface, 234
- Inflexional lines of a surface, 186; differential equation for, 190; of Kummer's surface, 222, 230; correspond to lines of curvature on Cyclide, 231
- Invariants, representative, for an involution in threefold space, 209
- Inversion in a plane, deduced from projection in space, 12; of plane quartic curve with two double points, 96; of Cyclide, 179; of Cyclide with two double points, 194; of Dupin's Cyclide, to quadric cone of revolution, 201. See also *Harmonic inversion*
- Involution of sets of sixteen points in threefold space, 208
- Jessop, theorem for angles of intersection of tangent circles of a quartic curve, 99
- Jordan, the group of the lines of a cubic surface, 105
- Klein: representation of a line by a point of fivefold space, 40; Lie's correspondence between sphere and lines, 55; six conjugate linear complexes, 202, 208; inflexional lines of Kummer's surface, 233; rationality of quadratic complex, 235
- Kronecker and Castelnuovo, surface with double infinity of reducible plane sections, 55
- Kühne, generalisation of Wallace's theorem, 59
- Kummer, equation for a particular Cyclide, 197
- Kummer's quartic surface: arising by section of Segre's quartic locus with tangent solid, 139; bitangents of, from lines of Segre's cubic locus, 156; from profile of Segre's cubic locus, 156; and Weddle's surface, from four quadrics in threefold space, 160; obtained by transformation from fourfold space, 211; as locus of singular points of a quadratic complex, 212; equation expressed as sum of five squares, 215; range of six points associated with, 215; equation shewing double

- points, 216; dual of, 217; represented in fivefold space, 218; six congruences associated with, 219, 223; tangent lines at a point, 221; identical with envelope of singular planes, 222; inflexional lines of, 222, 230; bitangents of, 224; from confocal complexes, 227; related range of points and pencil of planes, 228; intersections with a line of fundamental complex, 229; inflexional curves correspond to lines of curvature on Cyclide, 231
- Lachlan, Hart circle defined by angles of intersection, 88
- Larmor, A., circles arising from the intersection of three circles, 82
- Leathem, theorem for four concyclic points of a sphere, 101
- Lexell, theorem for four concyclic points of a sphere, 101
- Lie, correspondence of a sphere with two lines, 55; applied to Kummer's surface and Cyclide, 233
- Limaçon, 99
- Line, in threefold space, represented by a point of fivefold space, 41
- Line-cone, quadric, and point-cone, in fourfold space, 121
- Linear complex. See *Complex, linear*
- Malfatti's problem, 68
- Mannheim, inversion of Dupin's Cyclide to Anchor Ring, 200
- Mantel, symmetry of five points, for pedal property, 25
- Maxwell, Dupin's Cyclide, 193, 198
- Meridian surface, Plücker's, 142
- Miquel, a theorem for circles, 1; generalisation of, 8
- Moebius's figure of two tetrads and Wallace's theorem, 18; Moebius tetrads and quadratic congruence, 241
- Morley, theorem for Feuerbach circle, 79
- Noether, rationality of quadratic complex, 235
- Normal of Cyclide, by projection from fourfold space, 184
- Orr, theorem for four concyclic points of a sphere, 101
- Orthocentric, Absolute conic for which five points are, 11
- Pedal circles orthogonal to a fixed circle, 23; pedal property of circum-circle generalised to rational curve, 26; properties in threefold space, 24, 25, 27
- Picard, surface whose plane sections are rational, 55
- Plücker's Meridian surface, 142
- Point-cone, quadric, and line-cone, in fourfold space, 121
- Polar system, by plane section of joins of five points, 10; polar line of a line, in regard to a focal system, or linear complex, 42; polar of a line in regard to a quadric, represented in fivefold space, 49; polar lines, in a focal system, corresponding to the same sphere, 55; polar quadric of point of Segre's cubic locus, 156
- Poncelet, on circles touching three circles, 66
- Quadratic complex. See *Complex, quadratic*
- Quadric: in fourfold space, lines of, through a point, 37; in fivefold space, representing lines, 41; quadric surface, generators of, represented by a conic in fivefold space, 43; in fivefold space, planes lying on, 45; each its own polar plane, 47; quadric surface, number of common points of two curves on, 51; quadrics in n -fold space, through given points, 61; quadric surface, sections of, touching three sections, 65; equations of, 69; condition four sections of quadric surface be touched by four planes, 72; quadric surface, harmonically conjugate, 72; equation of, 73; quadric surface, radius of section of, with respect to a point, 80; four sections and four tangent sections, mutual relations, 86; angles between four sections, 100; quadric surfaces with ring contact, theorem of intervals, 103; quadric, in fourfold space, in regard to which Segre's figure is self-polar, 148, 149; quadric surfaces through five points give rise

- to Segre's cubic locus, 152; system derived from four general quadrics, 160; quadric surfaces, ten, associated with six conjugate linear complexes, 206; sets of four with two common generators, 207; quadric surface, generators touching curve of intersection with another quadric surface, 94, 170, 220, 223, 224; quadric in fivefold space, generation of, 237
- Quartic curve, in a plane, with two double points, 89; as envelope of circles, 90; foci of, 92; number of tangents from a double point, 93; relation between distances to three coneyclic foci, 94; inversion into itself, or another such curve, 96; with four foci in a line, 98; angles of two tangent circles with variable tangent circle, 99; general form of theorem for angles between tangent circles, 102
- Quartic curve, in fourfold space, 120, 147
- Quartic surface, with five double points, from quartic surface with fifteen double points, 160
- Quartic surface, in fourfold space: intersection of two quadrics, 161; algebraic treatment, 162; sixteen lines of, 164; representation on a plane, 165; descriptive theory of lines on, 167, 169; self-polar pentad for, 168; double-fours of lines on, 170; notation for lines, 170; conics upon, five pairs through each point of surface, 171; other equations for lines, 185; lines of, by intersection with a quartic locus, 189
- Rao, co-residuation on quartic surface in fourfold space, 172
- Related planes, three, variable plane containing corresponding points, 145
- Reye, triply infinite system of quadric surfaces, 160
- Richmond, pedal property for general rational curve, 25; figure arising from six points in fourfold space, 116
- Roberts, generalisation of a theorem for circles. 8
- Rosenhain, equation of Kummer's surface, 216
- Salmon, radius of Hart circle on a sphere, 80
- Schoute, on a six-dimensional polytope, 105
- Segre, tetrahedral and linear complex, from planes in fourfold space, 36; representation of conics in fivefold space; on Veronese's surface, 54; loci of third order in fourfold space, 116; theorem of five planes meeting six lines, 143; theorem for a Cyclide, 180; general investigation of Cyclides, 172; degenerate forms, 190
- Segre's figure, of fifteen points and lines in fourfold space, 113, 203; by transformation from fivefold space, 208; number notation for lines and solids, 114; singular solids, 115, 124, 129; incidences proved algebraically, 115; six systems of planes, two of any system through a point, 123; locus for which two planes coincide, 123; planes found algebraically, 124; planes of a system meeting an arbitrary line, 136; reciprocating quadric for, 148
- Segre's cubic locus in fourfold space, regarded as a dual, 151; obtained from quadric surfaces through five points, 152; theorem for lines upon, 155; from fivefold space, 160
- Segre's quartic locus in fourfold space, irrational equation of, 125; is rational, 126; six conics through a point met by lines in related ranges, 127; tangent solid, 128; contains a symmetrical figure of sixteen points and planes, 130, 134; thirty-two related ranges in, 131; intersection with tangent solid, 138; particular form of equation, 142; tangential equation, 148; and Weddle surface, 159; representing an involution of threefold space, 210
- Severi, surface with no chords through arbitrary point, 55
- Similitude, circle of, obtained by projection, 15; centres of similitude for two circles, 66

- Sphere, determined by section of quadric in fourfold space, 36; centre of, by projection, 39; represented by a point of fourfold space, 39; angle of intersection of two spheres, 39; touching four given spheres, or plates, 57, 58; enveloping a Cyclide, 175
- Steiner: a theorem for two triads of points, 21; Steiner's quartic surface, 54; by projection from fourfold space, or from fivefold space, 191; circles cutting three given circles at given angles, 67; solution of Malfatti's problem, 68
- Stéphanos, figure of fifteen circles, in anticipation of Segre, 116, 203
- Study, mutual relations of four circles and four tangent circles, 86
- Taylor's theorem, for poles of a line in regard to six conics, 5
- Tetrads, Moebius, mutually inscribed, theorem for four, 141; fifteen associated with six conjugate linear complexes, 206; property of tetrad in regard to a line, considered in fivefold space, 239
- Tetrahedral complex, determined from planes in fourfold space, 32; in fivefold space, 239
- Tetrahedroid, or Wave surface, 217
- Tore, or Anchor Ring, and Dupin's Cyclide, 193, 199
- Transversals of two lines, and points of a quadric surface, 31
- Triangularly circumscribed conic, 3, 5
- Vaidyanathaswamy, property of a cubic curve, 147
- Veronese's surface in fivefold space, conics thereon, 52, 54; projecting to Steiner's quartic surface, 191; theorem of reciprocity of a general figure, 149
- Wakeford, on the theorem of a double-six of lines, 6; six conics triangularly circumscribed to another, 32; theorem for six lines with a common transversal, 60
- Wallace's theorem, and Moebius's figure of two tetrads, 18; additions to, 21; generalised, 10, 58
- Wave surface, or tetrahedroid, 217
- Weber's theorem, construction of six conjugate linear complexes from six points, 139
- Weddle's surface, origin of, 156; equation of, 158; from Segre's quartic locus, 159; and Kummer's surface, from four quadrics in threefold space, 160; correspondence of conjugate directions, 233
- White, equation of a conic triangularly circumscribed to six conics, 5; proof of Clifford's chain of theorems for circles, 64