

INDEX

(For authors' names not included in this index see *List of Books and Memoirs Cited*, pp. 212–229, where page references to citations in the text will be found. As a rule, botanical family names are indexed only when several of their genera are named in the text.)

- Abietineae, 129–131
 Abnormal structures, see Teratology
 Abú Sa'íd, 211
Acacia, phyllodes with pseudo-midrib, 102
Acacia alata, R.Br., leaf-like winged shoot, 87, 88 (Fig. 7, A), 118
Acer Pseudo-platanus L., effect of one-sided union of two leaves, 171, 172 (Fig. 36, A)
Acmopyle, Sahni on microphyllous shoots simulating compound leaves, 91; on reproductive shoot, 128
 Aconite, Winter, see *Eranthis hyemalis* Salisb.
Aconitum Napellus L., 178, 179 (Fig. 39, A)
Adoxa moschatellina L., terminal flower, 185
Aesculus Hippocastanum L., absciss layer of petiolules, 80; ternate stage in leaf development, 118
Agrimonia Eupatoria L., subsidiary leaflets, 85, 102, 103 (Fig. 13, D), 122, 123
Ailanthus Girdalii Dode, large-scale development of compound-pinnate leaf, 116
Akebia quinata Decne., approximation to peltation of leaf, 111, 113 (Fig. 17, C)
Albertus Magnus, *De vegetabilibus*, 24 *et seq.*; difference from Cesalpino, 28; on branch-like leaves, 73; on flowers, 27; on leaf-veins of *Plantago*, 26; on parts of plants, 25; on physical and chemical causes, 26, 27; on roots, 25, 26, 30, 134; on 'sap', 25; on styles and stamens, 27; on thorns and spines, 26, 27; relation to Theophrastus, 24, 25, 27
 lbrecht, S. A., editor of Jung's works, 33
 Aléchamp, J. d', on Theophrastus, 15
 Alexandrian laurel, see *Ruscus Hypophyllum* L.
 al-Ghazālī and the physicist, 201, 202
 Allen, G. S., criticism of the writer's theory of the root, 135 *n.*
 Allseed, see *Radiola Linoides* Roth
 Almond, see *Prunus Amygdalus* Stokes
 Alps, 28, 42
Amaranthus Blitum L., Theophrastus on, 16 *n.*
 Amaryllidaceae, *Galanthus*, 53 (Fig. 3, C), 55; *Narcissus*, 82, 103 (Fig. 13, E), 104, 143
Amherstia nobilis Wall., steles replacing bundles, in bracteole, 105, 106 (Fig. 14, A), in stamen, 56, 57 (Fig. 5, C)
Amicia Zygomeris DC., absence of terminal pinna, 98; stipules compared with cotyledons of *Fagus*, 81 (Fig. 6, A), 82
Ampelopsis Veitchii, see *Parthenocissus tricuspidata* Planch.
Anagallis, number of petals, 27
 Analogous variation, Darwin, C., and Huxley, T. H., on, 160
 Analogy between plants and animals, 11–14, 17, 26
 Anaxagoras, 11 *n.*
 Anderson, E. and Abbe, L. B., on 'compact' mutants of *Aquilegia vulgaris* L., 196
 Andrews, H. N., on shoot interpretation of 'leaves' in certain fossil genera, 74, 75
 Androecium, see Stamens
Angelica ampla Nelson, asymmetrical branching of pinnae, 177; opposite pinnae, 114, 115 (Fig. 18, B)
 ἀνομοιομερῆ, τὰ, 11
Anschaung, 209
Anthoxanthum, terminal flower, 94

INDEX

- Anthriscus sylvestris* Hoffm., reduced terminal umbel, 184, 185, 186 (Fig. 42); short fruit beak, 168, 169 (Fig. 35, A)
- Antirrhinum majus* L., peloria, 178, 187, 189 (Fig. 44, A); alternating sterile and fertile zones in raceme, 194 (Fig. 46, A), 195
- '*Apium hortense*', see *Carum Petroselinum* Benth. et Hook.
- Apple, see *Malus communis* DC.
- Aquilegia*, Albertus Magnus on, 27; effects of crowding in the flower-bud, 173; Jung on leaf form, 73
- Aquilegia glandulosa* Fisch., pressure effect on staminodes, 172 (Fig. 36, C) 173
- Aquilegia vulgaris* L., 'compacta' mutants, 196
- Arabic translation of *De plantis* (Nicolaus Damascenus), 24
- Araceae, reduction in flowers correlated with size of spathe, 170; *Arum*, 143, 193; *Cryptocoryne*, 145
- Archetype, first use of word, 65 *n.*
- Aristolochia*, Cesalpino on, 31; Troll on, 145
- Aristotelian school, 9–36, 76
- Aristotelianism, reaction to Platonic, and pre-Platonic views from, 37 *n.*
- Aristotle, in relation to Theophrastus, 11–13; on actual and potential, 78, 87; on analogy between plants and animals, 11, 12; on causation, 199, 200; on conditional necessity, 196; on form, 2; on morphological thought, 7, 8; on *voûs*, 209; on parts of the living body, 10, 11, 14; on psyche in living things, 9, 10; on rejuvenescence in plants, 19; on reproduction, 20, 78; studied by Cesalpino, 28
- Artemisia*, Jung on, 33
- Arum*, Engler on, 193; Malpighi on, 143
- Ash, see *Fraxinus excelsior* L.
- Asparagus*, correction of the author's former interpretation of the needle, 97; sterile distal region in inflorescence, 193, 194 (Fig. 46, B)
- Aspergillus*, Raistrick on biochemical classification, 195
- Asperula odorata* L., comparison of shoot with a lupin leaf, 111, 113 (Fig. 17, A, B)
- Aster, distribution of blue and yellow coloration, 147, 191
- Asymmetry and branching of laminae and pinnae, 120, 177
- Atraphaxis buxifolia* Jaub. et Spach, ovule, 174, 175 (Fig. 37, A)
- Augustine, St, 2
- Avicenna, 211
- Axes, competition between main and lateral, 93 *et seq.*
- Axillary bud, a new interpretation of bud morphology, 125–31 (summarised, 126); anatomical connexion with parent shoot, 88 (Fig. 7, B), 91, 92, 126, 127; inhibiting action of leaves on bud, 97
- Bacon, F., on abnormalities, 5; on compatibility of mechanism and teleology, 202; on man and the plant, 12; on numbering and measuring in natural history, 27; parallel with Goethe, 40
- Ballota nigra* L., 3-leaved whorls, 163 (Fig. 32, B), 164
- Banana, see *Musa sapientum* L.
- Banks, Sir J., 60
- Banyan, see *Ficus benghalensis* L.
- Barnes, B., on *Lathyrus Aphaca* L., 99
- Basil, see *Ocimum Basilicum* L.
- Bauhinia yunnanensis* Franch., reduction of terminal leaf-pinna, 98 (Fig. 10, A), 99; stamen-anatomy, 56, 57 (Fig. 5, A)
- Beech, see *Fagus sylvatica* L.
- Bellis perennis* L., effect of crowding of florets in ontogeny, 173; funnel-shaped leaf as abnormality, 106 (Fig. 14, D), 109; 'hen-and-chicken' variety, 153, 179 (Fig. 39, D); inflorescence called a 'flower', 144
- Berberidaceae, reduction of midrib in phyllomes, 102, 103 (Fig. 13, A–C); repetitive leaf-ternation, 138 (Fig. 25), 139; *Berberis*, 102, 103 (Fig. 13, C); *Diphylleia*, 102, 103 (Fig. 13, A); *Epimedium*, 120, 138 (Fig. 25), 139; *Podophyllum*, 102, 103 (Fig. 13, B)
- Berberis calliantha* Mull., reduction of midrib in honey-leaf, 102, 103 (Fig. 13, C)
- Bertalanffy, L. von, on physico-chemical description, 201
- Betula*, median flower, 178
- Bilateral and radial symmetry, 176
- Bildung*, 206, 207

Cambridge University Press

978-1-108-04505-6 - The Natural Philosophy of Plant Form

Agnes Arber

Index

[More information](#)

INDEX

- Bindweed, see *Calystegia sepium* R.Br.
 Blackberry, see *Rubus fruticosus* L.
 Blite, see *Amaranthus Blitum* L.
 Blue and yellow coloration, distribution in flowers and inflorescences, 147, 191
 Bock, J. (Tragus, H.), compared with Malpighi and Grew, 37
 Boethius, on self-maintenance in the plant world, 76, 77; on the root corresponding to the head, 12
 Borage, Bacon, F., on, 27
 Boraginaceae; *Borago*, 27; *Myosotis*, 147, 191
 Bower, F. O., on 'Size-Factor', 190
 Bracteoles, equivalent to prophylls, 48
Brassica oleracea L., ascidial peltate lamina, 109, 110 (Fig. 15, B1–B3); enations from lamina, 82; lamina-wings from midrib, 107, 108; shoot-like outgrowth from midrib, 110 (Fig. 15, B 4); tuft of laminae borne by leaf, 106 (Fig. 14, B)
Brassica sp., 115 (Fig. 18, A)
 Braun, A., short-shoot theory of the ovuliferous scale in conifers, 128–130
 Brieger, F. G., on chemical factors in the differentiation of the flower, 191 *n.*
 Brown, R., on connation, 171; on reticulate connexion of taxonomic groups, 66; on the relation of the leaf to the parts of the flower, 68; connexion with Corrêa da Serra, 60
 Browne, Sir T., on the rose calyx, 50
 Bud, axillary, a new interpretation of bud morphology, 125–31 (summarised, 126); anatomical connexion with parent shoot, 88 (Fig. 7, B), 91, 92, 126, 127; inhibiting action of leaves on axillary and terminal bud, 97
 'Buglosse', Bacon, F., on, 27
 Buitenzorg Garden, 149 *n.*
 Bulbous plants, Cesalpino on gynaecea of, 29
 Bundles, collateral, replaced by radial strands or steles, 104, 105, 106 (Fig. 14, A)
 'Bundles of insertion' for axillary bud, 88 (Fig. 7, B), 91, 92, 126, 127
Bupleurum rotundifolium L., pseudanthia 149, 151, 152 (Fig. 29, A)
 Butchers'-broom, see *Ruscus aculeatus* L.
 Buttercup, see *Ranunculus*; Water-buttercup, see *R. heterophyllus* Fries
 Cabbage, see *Brassica oleracea* L.
Caesalpinia japonica Sieb. et Zucc., 89, 90 (Fig. 8, A)
Caesalpinia sepiaria Roxb., see *C. japonica* Sieb. et Zucc.
Calendula officinalis L., 153, 154 (Fig. 30, C)
Calystegia sepium R.Br., leaf with second apex, 101 (Fig. 12, C), 104; Theophrastus on the corolla, 20
 Cambridge Platonists, 202
Camellia japonica L., transitions from stamens to petals, 54 (Fig. 4, C), 55
Campanula, Jung on heterophyly, 35; petaloid calyx, 51, 53 (Fig. 3, B)
Campanula rotundifolia L., palmately and pinnately veined leaves, 101 (Fig. 12, A), 102
 Candolle, A. Casimir P. de, on the nature of the leaf, 74 *et passim*; relationship to Candolle, A. P. de, 74
 Candolle, A. P. de, his ideas in relation to those of Goethe and Grew, 46; on abnormalities, 5; on corolla-androecium system, 55; on family archetypes, 63; on Goethe, 41; on parallelism of vegetative and floral members, 76; on sectorial analysis of shoot, 166; on symmetry, 174, 176, 177; on the description of marginal 'cutting' in leaves, 83, 84; on the flower, 46, 47; on the *noëd vital*, 30; on units of plant structure, 70
Canna, Goethe on, 44
Cannabis sativa L., Cesalpino on sterile and fertile flowers, 29
 Canterbury-bells, see *Campanula*
 Caper, see *Capparis spinosa* L.
 Capparidaceae, *Capparis*, 56, 57 (Fig. 5, D); *Gynandropsis*, 48, 49 (Fig. 1, D), 88 (Fig. 7, B), 92, 175 (Fig. 37, D); *Polanisia*, 51, 53 (Fig. 3, A)
Capparis spinosa L., stamen anatomy, 56, 57 (Fig. 5, D)
Caput radices of Cesalpino, 29, 30
 Carpels, corresponding to peltate leaves 58, 111; interpretation of, 44–6, 56, 58, 129–31; non-vascular, 58; terminal, 124; theory of Delpino, 129–31
 Carrot, see *Daucus Carota* L.
Carum Petroselinum Benth. et Hook., Jung on branch-like leaves, 73

INDEX

- Carum verticillatum* Koch, shoot-like compound-pinnate leaf, 116
- Cassia bicapsularis* L., stamen anatomy, 56, 57 (Fig. 5, B)
- Castor-oil-plant, see *Ricinus communis* L.
- Catananche caerulea* L., rachis-leaves, 85, 122 (Fig. 22, C)
- Caucalis nodosa* Scop., spiny fruit, 169 (Fig. 35, I), 170
- Causal morphology, 162, 196
- Causes, The four: material and efficient (mechanical or physico-chemical), formal and final (teleological), 199, 200 *et seq.*
- Čelakovský, L. J., on the carpel as a peltate leaf, 58
- Celandine, Lesser, see *Ranunculus Ficaria* L.
- Cell-wall, interpretation, 195, 196
- Cephalotus*, pitcher leaves, 157
- Ceratophyllum*, shoot resembling leaf of *Carum verticillatum* Koch, 116
- Cercis Siliquastrum* L., pseudo-simple leaf, 98 (Fig. 10, B), 99
- Cesalpino, A., as a precursor of Spinoza, 28; *De plantis*, 28–32; difference from Albertus, 28; New World flora, 29; on bud production, 28; on classificatory criteria, 31, 32; on climbing plants, 28, 29; on ‘coronary’ herbs, 31; on flowers, fruits, and seeds, 29–31; on ‘necessity’ and ‘purpose’, 202; on seed production as the plant’s goal, 30; on stem-like ‘leaves’, 29; on sterile (male) flowers, 29; on the anatomy of nodes, 28; on the *cor plantarum* (root-crown, shoot-initial) 29, 30; on the importance of the pith, 30, 31; on the psyche in plants, 29, 30; on the quadrate stem of *Marrubium*, 28; on the shoot, 28; on Theophrastus, 21; representing the Aristotelian school, 24
- Chamaerops humilis* L., Goethe’s palm at Padua, 42
- Cheiranthus Cheiri* L., pressure in ovule development, 175 (Fig. 37, C)
- Chelidonium majus* L., pinnate leaf and ‘stringing out’ of leaflets, 115 (Fig. 18, E), 122
- Chemical factors in the differentiation of the flower, 190, 191
- Chestnut, Horse-, see *Aesculus Hippocastanum* L.; Water, *Trapa natans* L.
- Chimaeras, periclinal, 134
- Chimonanthus fragrans* Lindl., bracts and perianth members, 51, 54 (Fig. 4, B)
- Choripetalae, comparison of petal coloration with that of pseudopetals of composites, 147
- Chrysanthemum Leucanthemum* L., 48, 49 (Fig. 1, A)
- Chrysogonum virginianum* L., 147
- Citrus* sp., Theophrastus on, 21
- Clematis heracleafolia* DC., 3-pinna leaf, 137 (Fig. 24, C), 139
- Clematis orientalis* L., leaf with ternation of second and third order, 137 (Fig. 24, E), 139
- Clematis Vitalba* L., leaf with ternate apical pinna, 137 (Fig. 24, D), 139
- Cleome graveolens* Rafin., see *Polanisia viscosa* DC.
- Clover, see *Trifolium repens* L.
- Codex Aniciae Julianae* (Dioscorides), 27, 123
- Coleridge, S. T., on conditions and causes, 196
- Collar, collet (*cor plantarum*), 29, 30
- Coloration, blue and yellow, distribution in flowers and inflorescences, 147, 191
- Columbine, see *Aquilegia*
- Components and parts, in relation to the whole plant, 157, 158
- Compositae, colour distribution in capitula, 147, 191; comparison of pseudopetals with petals of Choripetalae, 147, 148; effect of crowding in capitulum ontogeny, 173; florets replaced by inflorescences, 153; Grew on the capitulum, 143; inhibition of growth in length in inflorescence-shoot, 104; pseudanthia, 147 *et seq.*; *Artemisia*, 33; *Aster*, 147, 191; *Bellis*, 106 (Fig. 14, D), 109, 144, 153, 173, 179 (Fig. 39, D), 180; *Calendula*, 153, 154 (Fig. 30, C); *Chrysanthemum*, 48, 49 (Fig. 1, A); *Chrysogonum*, 147
- Echinops*, 149; *Filago*, 95; *Helipterum*, 148; *Lapsana*, 80, 114, 115 (Fig. 18, C); *Scorzonera*, 85; *Syncephalanthia*, 148; *Taraxacum*, 181 (Fig. 40, C), 182; *Tragopogon*, 153, 154 (Fig. 30, D)
- Compound inflorescences, 148, 149
- Compound-peltate leaves, comparison with shoots, 111, 113 (Fig. 17, A, B), 114

INDEX

- Compound-pinnate leaves, comparison with shoots, 114, 116, 118; interpretation, 119–23
- Conatus* of Spinoza, 77, 207
- Conditions *versus* causes, 196–8
- ‘Congenital fusion’, 171
- Conifers, female reproductive shoot, 127–31
- Connation, 171
- Convolvulus, see *Calystegia sepium* R.Br.
- Coppicing, effect on *Corylus Avellana* L., 177, 178
- Cor plantarum* of Cesalpino, 29, 30
- Corchorus olitorius* L., hair-like lobes and stipules, 140 (Fig. 26, D), 141
- Coriandrum sativum* L., pseudanthia, 151, 152 (Fig. 29, B)
- Coriaria myrtifolia* L., leaf-like branch system and stipular hairs, 89, 90 (Fig. 8, B), 142
- Corn, Theophrastus on the ear, 143
- ‘Coronary’ herbs, Cesalpino on, 31
- Corréa da Serra, J. F., life, 59, 60; on a plan underlying each plant group, 59–61; on parallelism, 160; on symmetry, 174; on vegetation of New Holland, 60
- Corylus Avellana* L., effect of coppicing, 177, 178; Theophrastus on the male catkin, 20, 21
- Cotyledons, and prophylls, 70, 71; and stipules, 80–2; Goethe on, 43; root-shoot members, 30
- Cow-parsley, see *Anthriscus sylvestris* Hoffm.
- Crambe maritima* L., inflorescences replacing flowers, 153, 154 (Fig. 30, A); narrowing of fruit base due to pressure, 197
- Crane’s-bill, Long-stalked, see *Gernium columbinum* L.
- Crassulaceae, *Pagella*, 148
- Crataegus oxyacantha* L., asymmetrical stipules, 52 (Fig. 2, A 3), 82; stipulate and exstipulate leaves, 51, 52 (Fig. 2, A 1, A 2)
- Crocus sativus* L., 153; C. sp., 203
- Cross, G. L., on comparison of scale-leaves and foliage-leaves, 86
- Cruciferae, ‘bractless’ raceme, 99, 127, 140 (Fig. 26, E); effect of pressure in flower development, 173, 175 (Fig. 37, C), 197; range of form in gynaeceum, 168, 170; *Brassica*, 80, 105, 106 (Fig. 14, B), 107–110 (Fig. 15, B), 115 (Fig. 18, A); *Brassica* (turnip), 31; *Cheiranthus*, 175 (Fig. 37, C); *Crambe*, 153, 154 (Fig. 30, A), 197; *Nasturtium*, 99, 140 (Fig. 26, E), 178, 179 (Fig. 39, B), 180; *Succowia*, 169 (Fig. 35, H), 170
- Cryptocoryne*, pseudanthia, 145
- Cryptogams, vascular, the plant body in, 74–6, 91, 116, 139
- Cryptomeria japonica* (L.-f.) Don, Hagerup on, 128
- Cudweed, see *Filago germanica* L.
- Cudworth, R., on mechanism and teleology, 202; on ‘Plastick Nature’, 202 n.; on the formal cause, 205
- Cupressus sempervirens* L., Theophrastus on fruit and seed, 22
- Cyclamen, Cesalpino on, 31
- Cydonia oblonga* Mill., Grew on leaf-like sepals, 39
- Cymose branching, 94
- Cypella Herberti* Herb., foliated lamina, 85
- Cyperus longus* L., Theophrastus on, 19
- Cypress, see *Cupressus sempervirens* L.
- Daisy, see *Bellis perennis* L.
- Daisy, Oxeye, see *Chrysanthemum Leucanthemum* L.
- D’Aléchamps (Dalechamps), J., emendation of Theophrastus, 15
- Danae*, interpretation of sterile phylloclade, 97
- Dante, 8, 39, 208
- Darlingtonia californica* Torr., ‘fishtail’ appendage, 195
- Darstellung*, 209
- Darwin, C., on analogous variation, 160
- Darwinian theory, see Pre- and Post-Darwinian morphology
- Date palm, see *Phoenix dactylifera* L.
- Daucus Carota* L., luxuriant umbel, 149, 150 (Fig. 28); pseudanthia, 151, 152 (Fig. 29, C); terminal umbellule, 182, 183 (Fig. 41), 184, 185, 190
- De causis plantarum* (Theophrastus) 13, 20, 22, 23
- De historia plantarum* (Theophrastus), 13–23
- Delpino, F., foreshadowing *Gestalt* concept, 155; interpretation of ovuliferous scale in conifers, and of the carpel in angiosperms, 129–31

INDEX

- Democritus, 12
De partibus animalium (Aristotle), 10, 11
De plantis (Nicolaus Damascenus), 24, 76
 Development, apical, and ontogeny, 162, 164, 166–70, 178, 205, 206
Dichophyllum, Andrews, H. N., on shoot interpretation of ‘leaves’, 74, 75
 Dicotyledony, Theophrastus on, 22
Dicraea, root-thalli, 135
 Digby, Sir K., on development from the seed, 206
Digitalis orientalis L., exaggeration of front petal, 169 (Fig. 35, E), 170
Digitalis purpurea L., peloria, 187, 188 (Fig. 43); proportions of corolla, 170
 Dioscorides, *Codex Aniciae Julianae*, 27, 123
Diphylleia cymosa Michx., reduction of leaf-midrib, 102, 103 (Fig. 13, A 1)
Dipsacus fullonum L., luxuriant central head, 182
Dipsacus sylvestris Mill., order of flower opening, 151
 ‘Dissimilar parts’ (τὰ ἀνομοιομερῆ), 10, 11, 158
 Dog’s-mercury, see *Mercurialis*
 Dog’s-tooth grass, Theophrastus on, 15
 Dorsiventrality and radiality in leaf and shoot, 34, 87–92, 107 *et seq.*
Echinops globifer Jacka (Compositae), comparison of inflorescence with *Spinifex* (Gramineae), 149
 Efficient cause, see Causes, The four ‘Egyptian bean’, see *Nelumbium speciosum* Willd.
Eichhornia speciosa Kth., ligular sheath, 99
 Eichler, A. W., on flower of *Plantago*, 171
 Elder, see *Sambucus nigra* L.
 Embryo-sac as a *Gestalt* type, 157
 Enations borne by foliage leaves, 82
 Epigyny, relation to axial abbreviation, 50; Theophrastus on, 20
Epimedium, garden hybrid, asymmetry of leaflets and repetitive ternation, 120, 138 (Fig. 25), 139
Eranthis hyemalis Salisb., comparison of leaf and reproductive shoot, 113 (Fig. 17, D), 114
Eryngium amethystinum L., transition from pinnae to hairs, 139, 140 (Fig. 26, A), 141
Eryngium Bourgati Gouan, lamina complexity arising out of ternation, 117 (Fig. 19, B), 118
Eryngium bromeliaefolium Delar., rachis-leaf, 85, 122 (Fig. 22)
Eryngium creticum Lam., transition from pinna-lobes to sclerotic tooth, 140 (Fig. 26, B), 141
Eryngium pandanifolium Cham. et Schlecht., elongated leaf, 168
 Euanthium and pseudanthium, 144 *et seq.*
Euphorbia, Cesalpino on ‘seeds’, 29; reduction of terminal cyathium, 182, 185; Troll on pseudanthia, 145
Euphorbia fulgens Karw., 145, 146 (Fig. 27, A)
Euphorbia helioscopia L., difference between terminal and lateral cyathia, 182
 Euphorbiaceae, *Euphorbia*, 29, 145, 146 (Fig. 27, A), 182; *Mercurialis*, 29; *Phyllanthus*, 88 (Fig. 7, D), 89
Fagus sylvatica L., cotyledons compared with stipules of *Amicia*, 81 (Fig. 6, B), 82
 Fern-frond, correspondence between whole-leaf, pinna, and pinnule, 139; prolonged apical growth, 116; Potonié, H., and Tansley, A. G., on leaf and stem, 74
 Ferns, coenopterid, Andrews, H. N., on shoot interpretation of ‘leaves’, 75
Ficus benghalensis L., Theophrastus on, 15
Ficus Krishnae C. de C. (hort. var. of *F. benghalensis* L.), hypoascial leaf, 110 (Fig. 15, C)
Filago germanica L., cymose inflorescence-complex, 95
 Filbert, see *Corylus Avellana* L.
 Final cause, see Causes, The four ‘Fine-leaved’ bay, Theophrastus on, 18
 Firs, Theophrastus on the cone compared with *Corylus* catkin, 21; on the roots, 15
 Five-faced-counsellor, see *Adoxa moschatellina* L.
 Flower-apices and inflorescence-apices, anatomical parallel, 151
 Flower, comparison with vegetative shoot, 39–58 *et passim*; *Gestalt*

Cambridge University Press

978-1-108-04505-6 - The Natural Philosophy of Plant Form

Agnes Arber

Index

[More information](#)

INDEX

- conception, 131, 143 *et seq.*; inhibition of growth in length and correlated features, 48, 50, 51
- 'Flower-de-Luces', Bacon, F., on, 27
- Form, 1–8, 40, 209 *et passim*
- Form and function, 3
- Formal cause, see Causes, The four
- Foster, A. S., and his school, on ontogeny of leaf and shoot, 86, 108, 118, 162
- Fraxinus excelsior* L., Theophrastus on the leaf, 18; transitions from simple to compound-pinnate leaves, 118, 119 (Fig. 20)
- Fruchtschuppe* of conifers, 127–31
- Fumarioideae, anatomy of inflorescence-apex, 94; effect of pressure on flower development, 173
- Function (Form and function), 3
- Galanthus nivalis* L., transitions between stamens and perianth members, 53 (Fig. 3, C), 55
- Galingale, see *Cyperus longus* L.
- Gall structures, 195
- Gaya Lyallii* E. G. Baker, palmately and pinnately veined leaves, 101 (Fig. 12, B), 102
- Genesis*, 64
- Geoffroy Saint-Hilaire, on abnormalities, 5
- Geranium*, comparison of leaf with reproductive shoot, 111
- Geranium columbinum* L., Jung on form of lamina, 34
- Germinis principium* of Cesalpino, 30
- Gestalt* in Goethe's sense, 206; *Gestalt* type, 144–61, 193, 206–8
- Geum* sp., proliferation and foliaceous sepals, 51, 54 (Fig. 4, A)
- Gladiolus tristis* L., cruciate lamina, 107
- Glaucium leiocarpum* Boiss., elongated ovary, 168
- Gleditsia caspica* Desf., abortive terminal pinna and pinnule, 98 and Frontispiece, B; simple or compound pinnae, 139 and Frontispiece, B; asymmetry in pinnae and pinnules, 120 *n.*
- Glisson, F., on plant anatomy, 16
- Glyptostrobis*, microphyllous shoots simulating compound leaves, 91
- Goat's-beard, see *Tragopogon pratensis* L.
- Godron, D. A., effect of pressure on flower development, 173
- Goebel, K. von, rejection of typology, 159
- Goethe, J. W. von, amateur approach to botany, 40, 41; as *Naturshauer*, 210; botanical autobiography, 41, 42; on abnormalities, 5; on *Anschaung* and *Darstellung*, 209; on *Bildung* and *Gestalt*, 206; on cotyledons, 43; on crowding in development, 44, 173; on juvenile leaves, 43; on parallelism of vegetative and floral members, 45–58, 76; on petals, stamens, staminodes, and nectaries, 44, 55; on shape as an index quality, 4; on shoot-like leaves, 73; on the 'flower' of composites, 45; on the gynaeceum, 44, 45; on the leaf as a type form, 42–4, 70; on the legume, 45; on the plant as a single individual only in the seed, 136; on the root, 132; on the transition from vegetative to flowering phase, 43; on the type concept in general, 59; on the urge of the leaf to become a branch, 93; on the vascular system as the 'cause' of shape, 195; on the visual sense, 210; *Versuch die Metamorphose der Pflanzen zu erklären*, 40–6, 73
- Goethean views from the standpoint of modern botany, 47–58
- Gola, Prof. G., on Goethe's palm, 42
- Gramineae, alternate distichous leaves, 114; correlation between elaborate inflorescences and reduced flowers, 170; effect of pressure in development, 173; parallelism within the family, 6, 160 *n.*; pseudospikelets and compound inflorescences, 148, 149; Theophrastus on, 15, 143; *Anthoxanthum*, 94; *Lygeum*, 94; *Ochlandra*, 56; *Paspalum*, 87, 88 (Fig. 7, C); *Schizostachyum*, 149; *Secale*, 55, 132; *Spinifex*, 149; *Streptochaeta*, 94; *Triticum*, 38
- Grapes, Theophrastus on, 143
- Gray, Asa, on the metamorphosis theory, 68
- Greek morphology, 7–23, 210 *et passim*
- Grete Herball, *The*, 153
- Grew, N., identification of foliage-leaves, sepals, and petals, 39, 40, 46; identification of leaf and branch, 72, 73; on buds and shoots, 38; on clas-

INDEX

- sification, 32; on *Crocus*, 203; on leaf veneration, 38; on physical and chemical causes, 27; on seedlings, 37, 38; on teleology and mechanism, 203; on the capitulum of the Compositae, 143; representing decline of classical tradition, 24, 37; *The Anatomy of Plants*, 37; *The Anatomy of Vegetables Begun*, 37–40
- Ground-ivy, see *Nepeta Glechoma* Benth.
- Growth, differential or relative, see Relative growth
- Grundorgan*, leaf as, 70
- Gymnosperms, ovuliferous scale, 127–31; Theophrastus on ‘firs’ and cypress, 15, 22; *Acropyle*, 91, 128; *Cryptomeria*, 128; *Glyptostrobus*, 91; *Pinus*, 75; *Sciadopitys*, 129–31; *Taxodium*, 91; *Thuja*, 91; *Thujopsis*, 91; *Voltzia*, 127, 128
- Gynaecium, Delpino’s view, 131; Goethe and de Candolle’s view, 44, 45, 46, 56, 58; pseudo-monomery (Eckardt, T.), 58 n.; Troll’s view, 58; Wilson, C. L., telome interpretation, 76
- Gynandropsis pentaphylla* DC., crowding in ovary, 175 (Fig. 37, D)
- Gynandropsis speciosa* DC., elongated internodes in flower, 48, 49 (Fig. 1, D); bundles of insertion, 88 (Fig. 7, B), 92
- Hacquetia Epipactis* DC., pseudanthium, 148
- Hagerup, O., on the ovuliferous scale of conifers, 128
- Hairs, see Trichomes
- Harebell, see *Campanula rotundifolia* L.
- Hare’s-ear, see *Bupleurum rotundifolium* L.
- Hawthorn, see *Crataegus Oxyacantha* L.
- Hazel, see *Corylus Avellana* L.
- Hedwig, G., denying importance of pith, 31
- Hegel, G. W. F., 71, 87, 158 n., 161
- Heliamphora nutans* Benth., ‘hood’ development, 195
- Helipterum Manglesii* F. Muell., pseudanthium, 148
- Hemp, see *Cannabis sativa* L.
- ‘Hen-and-chicken’ daisy, inflorescences replacing flowers, 153, 179 (Fig. 39, D), 180
- Heracleum Sphondylium* L., foliar origin of axillary buds, 125
- Herbaceous and tree habit, Takhtajan on, 108
- Herball, The Grete*, 153
- Historia animalium* (Aristotle), 10, 11, 61
- Historia plantarum* (Theophrastus), 13–24
- History of botany, 6, 7
- ἁμοιομερῆ, τὰ, 11
- Hop, see *Humulus Lupulus* L.
- Hop-hornbeam, see *Ostrya carpinifolia* Scop.
- Horehound, Black-, see *Ballota nigra* L.; White-, see *Marrubium*
- Horse-chestnut, see *Aesculus Hippocastanum* L.
- Houttuynia cordata* Thunb., pseudanthia, 145, 146 (Fig. 27, B)
- Humulus Lupulus* L., transition from simpler leaves to ternate lobing, 137 (Fig. 24, A), 139
- Hutchinson, J., comparison of pseudanthium of *Houttuynia cordata* Thunb. to a ranunculaceous flower, 145
- Huxley, J. S., on relative growth, 168, 170
- Huxley, T. H., on morphological relations between species, 64; on polyphyletic, 160; on the influence of the cell-wall, 196
- Hydrangea*, pseudanthia, 151, 152 (Fig. 29, D)
- Hypocotyl, 30, 134
- Identity-in-parallel, Frontispiece, 142, 143
- Impious herb, see *Filago germanica* L.
- Inflorescence-apices and flower-apices, anatomical parallel, 151
- Inflorescences, compound, 148; simulating flowers, see Pseudanthia
- Iridaceae, pseudo-midrib in leaves, 102; *Crocus*, 153, 203; *Cypella*, 85; *Gladiolus*, 107; *Iris*, 27 (“Flower-de-Luces”) 36, 45, 153, 163 (Fig. 32, A), 164
- Iris*, abnormal dimerous flower, 163 (Fig. 32, A), 164; Bacon on “Flower-de-Luces”, 27; Goethe on style, 45; Jung on style, 36, 45; Troll on ‘secondary flower’ 153
- ἴς, 12
- Italy, 28, 42

INDEX

- Jäger, G. F. von, on abnormalities, 5
Jasminum humile L., range of leaf form, and correspondence of shoot and leaf, Frontispiece, A, and 139
 Johnson, T., on Aristotle, 9
 Joshi, A. C., on the effect of pressure on ovule development, 174
 Judas tree, see *Cercis Siliquastrum* L.
 Juncaginaceae, *Lilaea*, 95
 Jung, J., *De Plantis Doxoscopiae*, 33; *Isagoge Phytoscopica*, 33; on botanical terminology, 34, 35; on branching, 33; on branch-like leaves, 73; on composite flowers and their 'doubling', 35, 36; on heterophylly, 35; on leaf form, 34, 35; on pappus of composites, 36; on physical and chemical causes, 27; on radial stem and dorsiventral leaf, 34; on subsidiary leaflets in *Spiraea Ulmaria* L., etc., 35; on the 'flower', 35; on the *fundus*, 33; on the root-crown, 33, 143; representing Aristotelian botany, 24
- Kalanchoë daigremontiana* R. Hamet et Perrier de la Bâthie, leaf-budding, 107, 108 (Fig. 14, C)
 Kant, I., on necessity and free will, 208; on parallelism of leaf and shoot, 73, 74; on teleology and mechanism, 203
 καρδία, 12
- Labiatae, *Ballota*, 163 (Fig. 32, B), 164; *Marrubium*, 28; *Nepeta*, 192 (Fig. 45, D), 193; *Ocimum*, 16 n.; *Salvia*, 169 (Fig. 35, F), 170; *Scutellaria*, 169 (Fig. 35, G), 170; *Thymus*, 22, 192 (Fig. 45, C), 193
Lapsana communis L., alternate pinnae and marginal relation of pinna to rachis, 80, 114, 115 (Fig. 18, C)
 Lateral structures taking up parental role, 93 *et seq.*; summarised, 105
Lathyrus Aphaca L., stipules dominating leaf, 99, 100 (Fig. 11)
 Laurel, Alexandrian, see *Ruscus Hypophyllum* L.
 Leaf, apex dominated by laterals, 97, 98; budding, 105 *et seq.*; Candolle, A. P. de, on leaf, 46, 83, 84; Candolle, Casimir de, on leaf, 74 *et passim*; Cesalpino on leaf, 28, 29; classical view of function of leaf, 17, 25, 28; compound-peltate and compound-pinnate, comparison with shoot, 111, 114, 116, 118; connexion of generations in leaf, 142, 143; cuttings, 107; dorsiventrality, 18, 34, 87 *et seq.*, 200; Goethe on leaf, 42 *et passim*; Grew on leaf, 38, 39; Jung on leaf, 34, 35; Malpighi on leaf, 38, 39; non-vascular, 58; ontogeny, 79, 107, 108, 178; partial-shoot theory of leaf, 70–123 *et passim*; peltate, Casimir de Candolle on, 109; rachis- or midrib-leaf, 85, 122 (Fig. 22), 123; radiality, 107 *et seq.*; relative growth and leaf anatomy, 170, 171; repetitive branching, 136–9; terminal leaf, 124; transition from foliage leaves to hairs, 139 *et seq.*; vernation, 38 (see also 'Petiolar phyllode'; references, *passim*, to foliage-leaves and other phyllomes)
 Leguminosae; bundles replaced by bundle-rings, 104, 105; flower compared with *Polygala*, 155; winged axis, 87; *Amherstia*, 56, 57 (Fig. 5, C), 104–6 (Fig. 14, A); *Amicia*, 81 (Fig. 6, A), 82; *Bauhinia*, 56, 57 (Fig. 5, A), 98 (Fig. 10, A), 99; *Caesalpinia*, 89, 90 (Fig. 8, A); *Cassia*, 56, 57 (Fig. 5, B); *Cercis*, 98 (Fig. 10, B), 99; *Gleditschia*, Frontispiece, B, 98, 120 n., 139; *Lathyrus*, 99, 100 (Fig. 11); *Lupinus*, 111, 113 (Fig. 17, A); *Medicago*, 121 (Fig. 21, D), 169 (Fig. 35, J), 170; *Scorpiurus*, 169 (Fig. 35, K), 170; *Trifolium*, 58, 155, 156 (Fig. 31)
 Lesser-celandine, see *Ranunculus Ficaria* L.
 Life-knot (*cor plantarum*), 30
 Lilac, see *Syringa vulgaris* L.
Lilaea subulata Humb. et Bonpl., anatomy of inflorescence, 94, 95
 Liliaceae, *Allium*, 15, 17, 18, 34; *Asparagus*, 97, 193, 194 (Fig. 46, B); *Danae*, 97; 'Lillies', 27; *Ruscus*, 18, 97; *Scilla*, 179 (Fig. 39, C), 180; *Semele*, 97; *Smilax*, 17; *Tulipa*, 44
 Lillie, R. S., on teleology and mechanism, 203
 'Lillies', Bacon, F., on, 27
 Lime tree, see *Tilia europaea* L. and *T. tomentosa* Moench
Linaria vulgaris Mill., peloria, 178, 189 (Fig. 44, C), 190
 Linnaean botany, 41

Cambridge University Press

978-1-108-04505-6 - The Natural Philosophy of Plant Form

Agnes Arber

Index

[More information](#)

INDEX

- Linnaeus, C., on the root as 'core' of the shoot, 134
- Loganberry, see *Rubus*? *Idaeus* L. $\times$ *R. fruticosus* L.
- Logical analysis and metaphysics, 208
- λόγος, 2
- Lotus, Sacred, see *Nelumbium speciosum* Willd.
- Lübeck, 33
- Lund, S., on relation of phyllomes and trichomes, 141
- Lupin, see *Lupinus*
- Lupinus* sp., comparison of compound-peltate leaf with a shoot, 111, 113 (Fig. 17, A)
- Luxuriance of basal and apical flowers of inflorescences, 178, 179 (Fig. 39, A–C), 180, 181 (Fig. 40), 182, 187 (Fig. 43)
- Luziola Spruceana* Benth., terminal flower, 94
- Lyceum garden, 21
- Lychnis diurna* Sibth., sectorial colour variation, 166, 167 (Fig. 34, A)
- Lycopersicum esculentum* Mill., shoots from leaves, 105
- Lygeum Spartum* Loebl., terminal pair of flowers, 94
- Lyte, H., 4
- McClure, F. A., on pseudospikelets of Gramineae, 149
- M'Cosh, J., on leaf-nervation and shoot-branching, 83 n.
- Macleod, J., on 'mechanical concordance', 155
- Madder, see *Rubia peregrina* L.
- Magnolia, elongated receptacle, 49 (Fig. 1, C), 50
- Majumdar, G. P., on the foliar origin of the buds of *Heracleum*, 125
- Malpighi, M., *Anatome Plantarum*, 37–9; on buds and shoots, 38, 41; on leaf development, 38, 39; on petaloid stamens, 55; on seedlings, 37, 38; on transitions from bud-scales to foliage-leaves, 39; on the inflorescence of *Arum*, 143; representing decline of classical tradition, 24, 37
- Malus communis* DC., comparison of leaf and branch, 72; Theophrastus on fruit, 143
- Malvaceae, *Gaya*, 101 (Fig. 12, B)
- Marchantia*, Sachs, J. von, on scale and structure, 190
- Marigold, Garden, see *Calendula officinalis* L.
- Marrubium*, Cesalpino on the quadrate stem, 28
- Material cause, see Causes, The four
- Mechanical interpretations *versus* teleology, 200 *et seq.*
- Mechanism of morphology, 162–98
- 'Median apple', Theophrastus on, 21
- Medicago arborea* L., coiled legume, 169 (Fig. 35, J), 170
- Medicago lupulina* L., branching of pinnae, 121 (Fig. 21, D)
- Meliaceae, leaf asymmetry, 120 n.
- Melianthus major* L., axillary stipules, 82, 115 (Fig. 18, D1, D3); opposite pinnae, 114, 115 (Fig. 18, D1); unifacial winged rachis, 80, 115 (Fig. 18, D1, D2), 118
- Mercurialis*, Cesalpino on sterile and fertile flowers, 29
- Merman from Suffolk, 67
- Metaphysics and logical analysis, 208
- Microcosm, 158, 159; branches of biology as microcosms, 1
- Microphyllous shoots simulating compound leaves, 88 (Fig. 7, D), 89, 90 (Fig. 8, B1), 91
- Midrib- or rachis-leaves, 85, 122 (Fig. 22), 123
- Mill, J. S., on parallelism, 160
- Mirabilis Jalapa* L., terminal carpel, 124
- Mirabilis longiflora* L., elongated flower-tube, 168
- Monkshood, see *Aconitum*
- Monocotyledonous leaves, cylindrical, correction of the writer's former interpretation as petiolar phyllodes, 108
- Monocotyledony, Theophrastus on, 22
- Montaigne, M. de, on abnormalities, 5, 6
- μορφή, 2
- Morphology, 1–4 *et passim*; 'causal', 162; interpretation, 199–211; mechanism, 162–198; relation to art, 210, 211
- Moschatel, see *Adoxa moschatellina* L.
- 'Motor act', 3
- Mousetail, see *Myosurus minimus* L.
- Müllerott, M., on *Zwischenfedern*, 123
- Musa sapientum* L., unusual lamina development, 85

Cambridge University Press

978-1-108-04505-6 - The Natural Philosophy of Plant Form

Agnes Arber

Index

[More information](#)

INDEX

- Myosotis*, distribution of blue and yellow coloration, 147
- Myosurus minimus* L., elongated receptacle, 49 (Fig. 1, B), 50
- Narcissus*, relation of corona and perianth, 82; Theophrastus on, 143
- Narcissus Bulbocodium* L., corona dominating parent phyllome, 103 (Fig. 13, E), 104
- Nasturtium officinale* R.Br., accessory flowers, 178, 179 (Fig. 39, B), 180; stipules, 99, 140 (Fig. 26, E)
- Negative conditions, 197, 198
- Nelumbium speciosum* Willd., Theophrastus on, 21, 22
- Neo-Platonism, 209
- Nepenthes*, pitcher leaves, 157
- Nepeta Glechoma* Benth., hermaphrodite and female flowers, 192 (Fig. 45, D), 193
- Nerium Oleander* L., Troll, W., on, 145
- Nettle, see *Urtica*
- New Holland, Corrêa da Serra on vegetation of, 60
- 'New Philosophy' of the seventeenth century, 37, 202
- Nicolaus Damascenus*, influence on Albertus Magnus, 24; on self-maintenance in plants, 76
- Nicotiana Tabacum* L., foliar development, 79; leaf anatomy depending on relative growth, 170, 171
- Noëud vital (*cor plantarum*), 29, 30
- Norman, J. M., on stipules in Cruciferae, 99 n.
- Numerical element in biology, 27
- "*Nymphaea micrantha*", see *N. stellata* Willd.
- Nymphaea stellata* Willd., leaf-budding, 107, 115 (Fig. 18, F)
- Nymphaeaceae, *Nelumbium*, 21; *Nymphaea*, 107, 115 (Fig. 18, F), 193
- Ochlandra setigera* Gamble, stamen anatomy, 56
- Ocimum Basilicum* L., Theophrastus on, 16 n.
- Oenone*, shoot-thalli, 135
- Oken, L., on dorsiventrality of leaf, 87; on the relation of leaf and whole plant, 158; on the relation of leaf nervation and shoot branching, 83
- Olea europaea* L., Theophrastus on the roots, 15
- Olive tree, see *Olea europaea* L.
- Onion, Jung on, 34; Theophrastus on, 15, 18, 34
- Ontogeny and apical development, 80, 162, 164, 166–8, 170, 171, 178
- Opuntia*, Cesalpino on, 29
- Origin of Species* (Darwin, C.), 63, 160
- Ostrya carpinifolia* Scop., Theophrastus on, 17
- Ovule, relation of atropous and anatropous forms, 174, 196, 197; shoot-leaf-, and trichome-interpretations, 131, 142
- Ovuliferous scale of conifers, 127–31; Braun's short-shoot theory, 128, 129; Sachs's theory and its variants, including Delpino's theory, 129, 130
- Oxalis enneaphylla* Cav., compound peltate leaf, 111, 112 (Fig. 16, B)
- Oxalis lasiandra* Zucc., 111 n.
- Oxalis lupinifolia* Jacq., compound-peltate leaf, 111
- Oxalis Ortgiesi* Regel, reduction of terminal leaflet-lobe, 98 (Fig. 10, D), 102
- Oxeye daisy, see *Chrysanthemum Leucanthemum* L.
- Padua, Botanical Garden and Goethe's palm, 42
- Paeonia*, Jung on, 73
- Paeonia Clusii* Stearn, foliaceous sepals, 49 (Fig. 1, F), 50
- Paeony, see *Paeonia*
- Pagella Archeri* Schonl., pseudanthium, 148
- Palm, Goethe's (*Chamaerops humilis* L.), 42
- Palm stem, Theophrastus on anatomy, 16
- Palmae, *Chamaerops*, 42; *Phoenix*, 38
- Papaver*, crowding of young ovules, 175 (Fig. 37, B); Goethe on double form, 44, 45; reduction of median region in carpel, 104
- Papaver Argemone* L., crowding in young ovary, 175 (Fig. 37, B)
- Papaveraceae, *Chelidonium*, 115 (Fig. 18, E), 122, 177; *Glaucium*, 168; *Papaver*, 44, 45, 104, 175 (Fig. 37, B)

Cambridge University Press

978-1-108-04505-6 - The Natural Philosophy of Plant Form

Agnes Arber

Index

[More information](#)

INDEX

- Parallelism in organisation, and in configuration or *Gestalt*, 86, 159–61, 207
- Parsley, see *Carum Petroselinum* Benth. et Hook.
- Parsley, Cow, see *Anthriscus sylvestris* Hoffm.
- Parthenocissus tricuspidata* Planch., transition from simple to 3-pinna lamina, 137 (Fig. 24, B), 139
- Partial-shoot theory, of the leaf, 70–123, 209 *et passim*; bearing upon other morphological problems, 124–35; of the root, 131–5; criticisms, 135 *n.*
- ‘Parts’ and ‘components’ in relation to the whole plant, 157, 158
- Paspalum* sp., leaf-like inflorescence-branches, 87, 88 (Fig. 7, C)
- Passiflora capsularis* L. var. *acutiloba*, domination of leaf-apex by lateral lobes, 98 (Fig. 10, C), 102
- Passifloraceae, torus, 193
- Pear, see *Pyrus communis* L.
- Pedicularis sylvatica* L., peloria, 189 (Fig. 44, B), 190
- Pelargonium*, comparison of leaf with reproductive shoot, 111
- Peloria, 178, 185, 187, 188 (Fig. 43), 189 (Fig. 44), 190, 197
- Peltate leaves, Casimir de Candolle on, 109, 111
- Penzig, O., on chemical factors in the differentiation of the flower, 191 *n.*, on Delpino’s view of the double needle of *Sciadopitys*, 129 *n.*
- περί φυτῶν αἰτιῶν (Theophrastus), 13, 20, 22–4
- περί φυτῶν ἱστορία (Theophrastus), 13–24
- Perigyny, 50
- Periploca graeca* L., stipulate sepals, 51, 52 (Fig. 2, B)
- Permian, *Voltzia* from, 127, 128
- ‘Persian apple’, Theophrastus on, 21
- Petals, chemical interpretation of their characteristics, 191
- ‘Petiolar phyllode’, correction of the author’s former theory of the monocotyledonous leaf, 85 *n.*, 108
- Petioles, unifacial and radial structure, 108, 109
- Philipson, W. R., criticism of partial-shoot theory of the leaf, 135 *n.*; on comparison of vegetative and reproductive ontogeny, 47; on the effect of pressure on floret development in Compositae, 173
- Philo Judaeus, and the word ‘archetype’, 65
- Phlox*, differentiation in leaf rudiment, 80
- Phoenix dactylifera* L., Malpighi on the seedling, 38
- Phyllanthus mimosoides* Sw., leaf-like shoot, 88 (Fig. 7, D), 89
- Phyllome, as a generalised term for foliar members, 42; considered in relation to the type concept and to parallelism, 86; effect of one-sided fusion, 171, 172 (Fig. 36, A); equivalence of phyllomes and trichomes, 139–142; floral phyllomes, ternate form, and subordination of median region, 104; terminal phyllomes, 124 (see also Leaf, and ‘Petiolar phyllode’)
- Phylogenetic trees, 65, 66
- Phylogenetics in relation to morphology, 63 *et seq.*
- Phyton theories, 166, 167
- Pimpernel, see *Anagallis*
- Pinus* needles, Dufrenoy, J., on shoot interpretation, 75
- Pitcher-leaves of *Nepenthes* and *Cephalotus* as representing one *Gestalt* type, 157
- Pith, 25, 26, 30, 31
- Placentae in angiosperm carpel, Delpino’s theory, 129–31
- Plagianthus Lyallii* A. Gray, see *Gaya Lyallii* E. G. Baker
- Plane-tree, see *Platanus orientalis* L. and *P. acerifolia* Willd.
- Plantago*, Albertus on the leaf-veins, 26; Eichler, A. W., on corolla and androecium, 171; inflorescences replacing flowers, 153, 154 (Fig. 30, B); ‘plume’ abnormality, 193, 194 (Fig. 46, C); rachis-leaf, 85, 122 (Fig. 22, A)
- Plantago Coronopus* L., inflorescences replacing flowers, 153, 154 (Fig. 30, B)
- Plantago major* L., rachis-leaf, 122 (Fig. 22, A); ‘plume’ abnormality, 193, 194 (Fig. 46, C)
- Plantain, see *Plantago*
- Platanus acerifolia* Willd., ascidial peltate lamina, 109, 110 (Fig. 15, A)

Cambridge University Press

978-1-108-04505-6 - The Natural Philosophy of Plant Form

Agnes Arber

Index

[More information](#)

INDEX

- Platanus orientalis* L., Theophrastus on plane-tree seed, 22, 23
- Plato, 'archetype', word not used by him, 65 *n.*; 'forms', 67, 159, 204, 205; myth of Creation, 64, 65; on human analogy for the root in plants, 12; on sensation, but lack of self-consciousness in plants, 9; on the marrow as the life-substance, 26, 30; on vision, 211; teacher of Aristotle and Theophrastus, 11
- Platonism, reaction to, in the sixteenth and seventeenth centuries, 37 *n.*
- Platonists, Cambridge, 202
- Pleuroplastic lamina development, 79, 84, 85; divergences from it, 85
- Pliny on cudweed, 95
- Plotinus, 158
- Podophyllum peltatum* L., absence of leaf midrib, 102, 103 (Fig. 13, B)
- Podostemaceae, root-thalli of *Dicraea* and shoot-thalli of *Oenone*, 134, 135
- Polanisia viscosa* DC., leaf-like petal, 51, 53 (Fig. 3, A)
- Polygala*, Delpino's comparison with flower of Papilionatae, 155
- Polygonaceae, *Atraphaxis*, 174, 175 (Fig. 37, A); *Polygonum*, 99
- Polygonum equisetiforme* Sibth. et Sm., reduced leaf, 99
- Polypetalae, Jung on corolla, 35
- Polyphyletic, 65, 66
- Poppy, see *Papaver*
- Post-Darwinian morphology, 7, 63–6, 114, 160
- Potamogeton densus* L., distichous, opposite-leaved shoot, 114
- Potato, see *Solanum tuberosum* L.
- Potentilla erecta* (L.) Hampe, see *P. Tormentilla* Neck.
- Potentilla reptans* L., pedate leaf, 121 (Fig. 21, A); variation in petal number, 164, 165 (Fig. 33, A)
- Potentilla Tormentilla* Neck., variation in petal number, 164, 165 (Fig. 33, B)
- Potonié, H., on the nature of the fern-leaf, 74
- Pre-Darwinian morphology, 5, 6, 66–8
- Priestley, J. H., on the unit of shoot growth, 166
- 'Primal patterns' of Owen, R., 67
- Primrose, see *Primula vulgaris* Huds.
- Primula sinensis* Sabine, atropous and anatropous ovules in abnormal form, 174
- Primula vulgaris* Huds., foliaceous calyx, 49 (Fig. 1, E), 51
- Primulaceae, *Anagallis*, 27; *Cyclamen*, 31; *Primula* 49 (Fig. 1, E), 51, 174
- Prophylls, equivalent to bracteoles, 48; compared with cotyledons, 70, 71
- Prunus Amygdalus* Stokes, Cesalpino on, 29; Malpighi on leaf transitions in the bud, 39; terminal carpel, 124
- Przibram, H., on polyphyletic, 65
- Pseudanthia, 144 *et seq.*
- 'Pseudo-lamina', correction of the author's former interpretation of certain monocotyledonous leaves, 85 *n.*
- Pseudo-monomery, 58
- Pseudospikelets of Gramineae, 149
- 'Psyche' in living things, 9, 10, 29, 30, 207
- Psymphyllum*, Andrews, H. N., on shoot interpretation of leaves, 74, 75
- Pterocarya fraxinifolia* K. Koch, bud-like apex of compound-pinnate leaf, 116, 117 (Fig. 19, A); marginal lamina, 80
- Pyrus communis* L., luxuriance of basal and apical flowers, 181 (Fig. 40, A), 182; Theophrastus on shape of pear leaf, 17
- Pyrus Sorbus* Gaertn., Theophrastus on, 18
- Queen-Anne's-lace, see *Anthriscus sylvestris* Hoffm.
- Quince, see *Cydonia oblonga* Mill.
- Rachis- or midrib-leaves, 85, 122 (Fig. 22), 123
- Radial and bilateral symmetry, 176
- Radiality and dorsiventrality in shoot and leaf, 34, 87–92, 107 *et seq.*
- Radiola Linoides* Roth, cymose branching 95, 96 (Fig. 9, C)
- Radiola Millegrana* Sm., see *R. Linoides* Roth
- Raistrick, H., on biochemical classification of *Aspergillus*, 195
- Ramus folialis* of Albertus Magnus, 73
- Ranunculaceae, comparison of flower with pseudanthium of *Houttuynia*, 145; *Aconitum*, 178, 179 (Fig. 39, A); *Aquilegia*, 27, 73, 172 (Fig. 36, C), 173, 196; *Clematis*, 137 (Fig. 24, C),

INDEX

- D, E), 139; *Eranthis*, 113 (Fig. 17, D), 114; *Myosurus*, 49 (Fig. 1, B), 50; *Paeonia*, 49 (Fig. 1, F), 50, 73; *Ranunculus*, 82, 96 (Fig. 9, A, B), 97, 106 (Fig. 14, E), 109, 110 (Fig. 15, D), 168, 169 (Fig. 35, C, D)
- Ranunculus*, bisexual and female flowers, 193; buttercup as illustrating the use of the term 'concept', 204; nectary scale, 82
- Ranunculus acris* L., female plant, 192 (Fig. 45, B)
- Ranunculus bulbosus* L., female plant, 192 (Fig. 45, A1); hypoascidial petal, 110 (Fig. 15, D) short fruit beak, 168, 169 (Fig. 35, C)
- Ranunculus falcatus* L., long fruit beak, 168, 169 (Fig. 35, D)
- Ranunculus Ficaria* L., peltate leaf as abnormality, 106 (Fig. 14, E), 109
- Ranunculus heterophyllus* Fries, reduced apical leaf-lobe, 96 (Fig. 9, A), 97
- Reason and consequent *versus* cause and effect, 205
- Reed, Theophrastus on, 16
- Relative growth, 167–71
- Repetitive branching, 136 *et seq.*
- ῥάχῃς, 12
- Rhyniaceae, 75
- Rhythm, and symmetry, 176, 178
- Ricinus communis* L., Malpighi on seedling, 38
- Robinet, J. B. R., on the type concept, 66, 67
- Root, 12, 15, 16, 25, 26, 29, 30; partial-shoot theory, 131–5
- Root-crown, 30, 33
- Rosaceae, *Agrimonia*, 35, 102, 103 (Fig. 13, D), 122, 123; *Crataegus*, 51, 52 (Fig. 2, A), 82; *Cydonia*, 39; *Geum*, 51, 54 (Fig. 4, A); *Malus*, 72, 143; *Potentilla*, 121 (Fig. 21, A), 164, 165 (Fig. 33); *Prunus*, 29, 39, 124; *Pyrus*, 17, 181 (Fig. 40, A), 182; *Rosa*, 29, 44, 50, 55, 143; *Rubus*, 51, 80, 81 (Fig. 6, C), 117 (Fig. 19, C), 118–21 (Fig. 21, B, C); *Spiraea*, 35, 166, 167 (Fig. 34, B)
- Rose, Cesalpino on, 29; Goethe on, 44; Browne, Sir T., on, 50; Malpighi on, 55; Theophrastus on, 143
- Rubia peregrina* L., 3-leaved nodes, 164
- Rubus fruticosus* L., leaf ternation followed by branching of lateral pinnae, 117 (Fig. 19, C), 118, 119
- Rubus?* *Idaeus* L. × *R. fruticosus* L., ternation followed by branching of apical pinna, 120, 121 (Fig. 21, C)
- Rubus laciniatus* Willd., foliaceous sepals, 51, 81 (Fig. 6, C3, C4); leaf sequence in seedling, 121 (Fig. 21, B1); petiole surviving leaflets, 80, 81 (Fig. 6, C1); ternation stage in development of pedate leaf, 118, 121 (Fig. 21, B)
- Rubus thyrsoides* Wimm., luxuriant terminal flower, 181 (Fig. 40, B) 182
- Rue, see *Ruta graveolens* L.
- Ruscus aculeatus* L. and *R. Hypophyllum* L.; interpretation of sterile phylloclade, 97; Theophrastus on, 18
- Rush, Theophrastus on, 16
- Russell, E. S., on 'parts-of-a-whole', 158
- Ruta graveolens* L., 181 (Fig. 40, D), 182
- Rye, see *Secale cereale* L.
- Sachs, J. von, artistic capacity, 210; on chemical factors in the differentiation of the flower, 191 *n.*; foreshadowing the *Gestalt* concept, 153, 155; on parallelism of leaf and shoot, 83; on scale and structure, 190; on the 'bush' analogy in phylogenetics, 66 *n.*; on the formal cause, 205; on the *noëud vital*, 30; on the ovule, 142; on the shoot, 71; view of Darwinism, 160
- Sacred lotus*, see *Nelumbium speciosum* Willd.
- Saffron, see *Crocus sativus* L.
- Sahni, B., on *Acmopyle* shoot, 91; on the ovuliferous scale of conifers, 128
- St Augustine, 2
- Salvia Sclarea* L., long upper lip of corolla, 169 (Fig. 35, F), 170
- Sambucus nigra* L., stipellar and leafy pinnules, 140 (Fig. 26, C), 141; Theophrastus on the inflorescence and flower, 143; three-leaved nodes, 164
- Sarracenia*, Goethe on the stigma, 45
- σάπξ, 12
- Saururaceae, *Houttuynia*, 145, 146 (Fig. 27, B)
- Saxifragaceae, *Hydrangea*, 151, 152 (Fig. 29, D)
- Scale and structure, 190

Cambridge University Press

978-1-108-04505-6 - The Natural Philosophy of Plant Form

Agnes Arber

Index

[More information](#)

INDEX

- Scandix Pecten-Veneris* L., elongated fruit-beak, 168, 169 (Fig. 35, B)
- Scheinblüte*, 145
- Schiller, J. C. F. von, discussion with Goethe on the archetypal plant, 61, 62
- Schizostachyum*, pseudospikelets, 149
- Schizostachyum longispiculatum* Kurz, 149 n.
- Schoute, J. C., on phyllotaxis of calyx, 51; of *Ulmus*, 89
- Schultz-Schultzenstein, K. H. von, on the nature of the leaf, 74; on the shoot-like leaf of lupin, 111
- Sciadopitys*, double needle, 129–31
- Scilla hispanica* Mill., 179 (Fig. 39, C), 180
- Scorpiurus subvillosa* L., coiled legume, 169 (Fig. 35, K), 170
- Scorzonera*, rachis-leaf, 85
- Scrophulariaceae, relation of flower type to that of Solanaceae, 177; *Antirrhinum*, 178, 187, 189 (Fig. 44, A), 194 (Fig. 46, A), 195; *Digitalis*, 169 (Fig. 35, E), 170, 187, 188 (Fig. 43); *Linaria*, 178, 189 (Fig. 44, C), 190, 197; *Pedicularis*, 189 (Fig. 44, B), 190
- Scutellaria orientalis* L., long tube of corolla, 169 (Fig. 35, G), 170
- Seakale, see *Crambe maritima* L.
- Secale cereale* L., comparative surface area of shoot and root, 132; vegetative versus spikelet development, 55
- Sedge, Theophrastus on, 16
- Self-maintenance in living things, 76–8, 207
- Semele*, interpretation of sterile phylloclade, 97
- Service-tree, see *Pyrus Sorbus* Gaertn.
- Shakespeare, W., 1
- Sharman, B. C., on *Dipsacus*, 151
- Sharrock, R., on abnormalities, 6; on 3-leaved whorls in *Rubia*, 164
- Shoot as a unit of the plant body, 18, 19, 70–3 *et passim*; comparison with leaf, 80 *et passim*; microphyllous, simulating compound leaf, 88–90; ontogeny compared with leaf ontogeny, 79, 107; units of growth, physiological and morphological, 166, 167
- ‘Shoot-initial’ of Cesalpino, 30
- Sicily, 42
- ‘Similar parts’ (τὰ ὁμοιομερῆ), 10, 11, 158
- ‘Size-Factor’, Bower, F. O., and Wardlaw, C. W., on, 190
- Smilax*, Theophrastus on, 17
- Snowdrop, see *Galanthus nivalis* L.
- ‘Social’ influences upon cells, 171
- Socrates, 204, 211 n.
- Solanaceae, *Lycopersicum*, 105; *Nicotiana*, 79, 170, 171; *Solanum*, 82, 95, 97, 123 n. 147
- Solanum Dulcamara* L., distribution of blue and yellow coloration, 147
- Solanum nigrum* L., shoot system, 95, 97
- Solanum tuberosum* L., prophylls, 82; *Zwischenfedern*, 123 n.
- Sorb, see *Pyrus Sorbus* Gaertn.
- Spencer, H., on phyton theory, 166 n.; on relative growth, 167, 168
- Spieghel, A., on plant form, 40; on the nature of the leaf, 72
- Spinifex* (Gramineae), inflorescence compared with *Echinops* (Compositae), 149
- Spinoza, B. de, Cesalpino as a precursor of, 28; on form as limitation, 197; on self-maintenance (*conatus*) in the living thing, 77, 78, 207; on the attributes, 208
- Spiraea Ulmaria* L., localised chlorosis in a leaf, 166, 167 (Fig. 34, B); subsidiary leaflets, 35
- Spurge, see *Euphorbia*
- Squill, Theophrastus on, 15
- Stamens, anatomical comparison with foliage-leaves, 55, 56, 57 (Fig. 5); Candolle, A. P. de, on, 46; foliaceous, 53 (Fig. 3, C), 54 (Fig. 4, C), 55; Goethe on, 44; Wilson, C. L., telome interpretation, 76
- Stefanoff, B., on the needle of *Asparagus*, 97
- Sterility and unisexuality in normally hermaphrodite species, 192 (Fig. 45), 193, 194 (Fig. 46, C)
- Stipules, as sole representatives of phyllome, 99; asymmetrical, 52 (Fig. 2, A3), 82; comparison with cotyledons, 80, 81 (Fig. 6, A, B), 82; non-vascular 99, 140 (Fig. 26, E), 141; relation to leaf-teeth, 141; thread-like, 140 (Fig. 26, D), 141
- Stocks, J. L., on the formal cause, 211 n.
- Stoics and self-maintenance, 76

INDEX

- Streptochaeta*, terminal flower, 94
 Subsidiary leaflets, see *Zwischenfiedern*
Succowia Balearica Medic., spiny fruit, 169 (Fig. 35, H), 170
 Swift, Jonathan., on the purpose of disputing, 71
 Sycamore, see *Acer Pseudo-platanus* L.
 Symmetry, 174, 176, 177; see also Radiality and dorsiventrality
 Sympetalae, Jung on flower of, 35
 Sympodial branching, 19, 94
Syncephalanthia decipiens Bartl., compound inflorescence simulating simple capitulum, 148
 Syriac translation of *De plantis* (Nicolaus Damascenus), 24
Syringa vulgaris L., replacement of main axis by laterals, 94
 Takhtajan, A., on herbaceous and tree habit, 108
 Tansley, A. G., on chemical factors in the differentiation of the flower, 191; on frond-like branch system of *Selaginella*, 91; on numerical variation in *Potentilla*, 164; on the nature of the fern-leaf, 74
Taraxacum officinale Web., fasciated central shoot, 181 (Fig. 40, C), 182
Taxodium, microphyllous shoots simulating compound leaves, 91
 Teasle, Fuller's, see *Dipsacus fullonum* L.; Wild, see *D. sylvestris* Mill.
 Telome theory (Zimmermann, W.), 75, 76; application to stamens and carpels (Wilson, C. L.), 76
 Telomes, sterile (phylloid) and fertile (sporangial), 75
 Teratology, 4–6, 44, 45, 51 *et seq.*, 109–11, 185, 187–90; illustrations, 49 (Fig. 1, A3, A4, E), 53 (Fig. 3, B2, C), 54 (Fig. 4, A, C2–C4), 81 (Fig. 6, C3, C4), 101 (Fig. 12, C2), 106 (Fig. 14, B, D, E2, E3), 110 (Fig. 15), 121 (Fig. 21, D2, D3), 150 (Fig. 28, A), 154 (Fig. 30), 163 (Fig. 32), 172 (Fig. 36, A), 179 (Fig. 39), 181 (Fig. 40, A, C), 188 (Fig. 43), 189 (Fig. 44), 194 (Fig. 46, C)
 Terminal phyllomes, 124; luxuriance or reduction in terminal shoots, 178 *et seq.*
 Theophrastus, botanical terminology, 11, 12; in relation to Albertus, 24, 25, 27; method of leaf description, 18; on analogy between plants and animals, 12–14; on bulbs, 20; on catkins, 20, 21; on classification of plant organs, 14, 143; on dicotyledony and monocotyledony, 22; on evergreen and deciduous trees, 17; on internodes, 16; on leaf-fall, 17; on mechanism and teleology, 202; on monopodial and sympodial growth, 19; on nodes, 16; on anatomy, 16, 17; on repetitive growth and the reproductive shoot, 143; on roots, 15, 16; on scope of botany, 13; on stem-like leaves, 18, 29; on stems, 16; on the compound leaf, 18, 143; on the flower, 20, 21, 143, 153; on the 'flowering tuft' of the pine, 20; on the leaf, 17, 18, 34; on the 'oak-moss', 20; on the seed, 22, 23; on tillering of cereals, 19; relation to Aristotle, 11 *n.*; use of φύλλον for petal, 39
 Thistle, Goethe on the flower, 45
 Thoday, D., on the chimaera nature of the rhizophore of *Selaginella*, 134
 Thompson, D'Arcy W., on mechanism and teleology, 201; on relative growth, 168
Thuja and *Thujopsis*, microphyllous shoots simulating compound leaves, 91
 Thyme, see *Thymus Serpyllum* L.
Thymus Serpyllum L., hermaphrodite and female flowers, 192 (Fig. 45), 193; Theophrastus on the seed, 22, 23
Tilia europaea L., sympodial branching, 94
Tilia tomentosa Moench, asymmetry in venation, 120 *n.*, 176 (Fig. 38), 177; bracteole and inflorescence, 109, 176 (Fig. 38)
 Tiliaceae, *Corchorus*, 140 (Fig. 26, D), 141; *Tilia*, 94, 109, 120 *n.*, 176 (Fig. 38) 177
 Tobacco, see *Nicotiana Tabacum* L.
 Tomato, see *Lycopersicum esculentum* Mill.
 'Torus' of Candolle, A. P. de, 191, 193
Tragopogon pratensis L., inflorescences replacing gynaecea, 153, 154 (Fig. 30, D)
 Tragus, H., see Bock, J.
Trapa natans L., Theophrastus on, 16

Cambridge University Press

978-1-108-04505-6 - The Natural Philosophy of Plant Form

Agnes Arber

Index

[More information](#)

INDEX

- Tree and herbaceous habit, Takhtajan, A., on, 108
- Trichomes, relation to phyllomes and shoots, 139–42
- Trifolium fragiferum* L., comparison of calyx with corolla of *T. procumbens* L., 155, 156 (Fig. 31)
- Trifolium repens* L., foliaceous carpels, 58
- Triglochin palustre* L., sterile carpels, 58
- Triticum*, Malpighi on seedling, 38
- Troll, W., on Goebel's rejection of typology, 159; on the carpel, 58; on his *Gestalt* theory of the flower, 144 *et seq.*; on the leaf as a *Grundorgan*, 70; on the metamorphosis theory, 68; on the type concept, 61; on *Zwischenfiedern*, 123
- Tropaeolum pentaphyllum* Lam., approximations to peltation, 111, 112 (Fig. 16, A)
- Truffle, Theophrastus on, 15
- Tulip, see *Tulipa*
- Tulipa*, Goethe on, 44
- Turnip, Cesalpino on, 31
- Turpin, P. J. F., on Goethe, 41, 62; on pen and pencil, 210
- Type concepts, 42 *et seq.*, 59–69, 144 *et seq.*; replaced by parallelism concepts, 86, 159–61
- Uittien, H., on relation of leaf nervation and shoot branching, 83, 102; on Umbelliferae, pinnate leaf, 119 *n.*, umbel, 182, 184
- Ulmus*, distichous lateral branches, 88 (Fig. 7, E), 89; Malpighi on leaf development, 38, 39; root-budding, 132, 133 (Fig. 23)
- Umbelliferae, *Angelica*, 114, 115 (Fig. 18, B) 116, 177; *Anthriscus*, 168, 169 (Fig. 35, A), 184–6 (Fig. 42); *Bupleurum*, 149, 151, 152 (Fig. 29, A) *Carum*, 73, 116; *Caucalis*, 169 (Fig. 35, I), 170; *Coriandrum*, 151, 152 (Fig. 29, B); *Daucus*, 149, 150 (Fig. 28), 151, 152 (Fig. 29, C), 182, 183 (Fig. 41), 184, 185; *Eryngium*, 85, 117 (Fig. 19, B), 118, 122 (Fig. 22, B), 139, 140 (Fig. 26, A, B), 141, 168; *Hacquetia*, 148; *Heracleum*, 125; *Scandix*, 168, 169 (Fig. 35, B)
- Unisexuality and sterility in normally hermaphrodite species, 192 (Fig. 45), 193, 194 (Fig. 46, C)
- Units of the plant body, 70–2, 166, 167 *et passim.*
- Urpflanze*, Goethe's, 59, 61–4; portrayals by Turpin and Schleiden, 62, 63
- Urtica*, Cesalpino on sterile and fertile flowers, 29
- Vascular supply, effect on form, 195
- Vegetative reproduction in phyllomes, 105, 106 (Fig. 14, C), 107, 115 (Fig. 18, F)
- Želenovský, J., on pseudo-simple leaves, 98, 99; on relation of leaf nervation and shoot branching, 83; on terminal flower of *Rubus thyrsoides* Wimm., 182
- Venus's-comb, see *Scandix Pecten-Veneris* L.
- Vialleton, L., on 'bush' analogy in phylogenetics, 66 *n.*
- Vienna manuscript of Dioscorides, 27, 123
- Virgula parvula* of Albertus Magnus, 73
- Visual thought in morphology, 210, 211
- Voigt, F. S., on distribution of blue and yellow coloration in flowers and inflorescences, 147
- Voltzia*, ovuliferous scale, 127, 128
- Wallflower, see *Cheiranthus Cheiri* L.
- Walton, J., on the cone-scale of a Permian *Voltzia*, 127, 128
- Wardlaw, C. W., on 'Size-Factor', 190 *n.*
- Warming, E., on bud and axillant leaf as a unit, 125
- Water-chestnut, see *Trapa natans* L.
- Watercress, see *Nasturtium officinale* R.Br.
- Water-hyacinth, see *Eichhornia speciosa* Kth.
- Wettstein, R. von, on the flower as a reduced inflorescence, 155, 157
- Wheat, see *Triticum*
- Whewell, W., on the type concept, 67, 68
- Whole plant in relation to its parts and components, 157, 158
- Whole-shoot-hood in leaf, urge to, 78, 79, 93–123
- Willis, J. C., on root- and shoot-thalli of Podostemaceae, 134, 135

INDEX

- Willow seed, Theophrastus on, 22
Wilson, C. L., telome interpretation of
 stamens and carpels, 76
Winged shoots compared with winged
 rachides, 118
Wing-nut, Caucasian, see *Pterocarya*
 fraxinifolia K. Koch
Wolff, C. F., on the foliar nature of the
 parts of the flower, 40, 42, 48
Woodger, J. H., on rhythm and sym-
 metry, 176
Woodruff, see *Asperula odorata* L.
Xyris sp., effect of crowding in in-
 florescence, 172 (Fig. 36, B), 173
Yellow and blue coloration, distribution
 in flowers and inflorescences, 147, 191
Zimmermann, W., on chemical factors
 in the differentiation of the flower,
 191 *n.*; telome theory, 75, 76
Zwischenfiedern (subsidiary leaflets), 35,
 102, 103 (Fig. 13, D), 122, 123, 167
 Fig. 34, B)