

TABLE OF CONTENTS.

VOL. II.

ON THE MULTIPLICATION OF PLANTS.

CONVERSATION XV.

ON THE MULTIPLICATION OF PLANTS BY SUBDIVISION.

Two Modes of multiplying Plants; by Seed, and by Subdivision. — Slips or Layers preserve the peculiar Qualities of the Fruit, tend to diminish the Quantity of Seed, and to improve the Fruit. — Germs. — Three Modes of subdividing Plants:

1. By Layers, various Modes of producing them. — Layers above Ground. — Multiplication of Plants by Slips.

CONVERSATION XVI.

ON GRAFTING.

The Graft or Scion. — Stock. — Grafting improves the Quality of the Fruit. — Plants of the same Family can alone be grafted on each other: other

A 3



νí

CONTENTS.

circumstances requisite. — Grafting accelerates the Period of Bearing of Seedling Trees. — Mechanical Part of the Process. — Three Classes, *Unitiges*, *Omnitiges*, *Multitiges*. — Various Modes of grafting: 1. By Approach; 2. By Scions; 3. By Bourgeons or Buds. — Page 17

CONVERSATION XVII.

ON THE MULTIPLICATION OF PLANTS BY SEED. —
THE FLOWER.

Organs of the Flower: Calyx, Corolla, Petals, Nectary, Ovary or Seed Vessel, Carpel, Style, Stigma, Pistil, Stamens, Filaments, Anthers. — Palm Trees. Pedunculus. — Torus, Pedicles, Receptacle, Inflorescence. — Calendar of Flora. — Period of Flowering. — Mode of advancing or retarding it. — Annular Section. — Fructification of the Seed prevented by Rain.

CONVERSATION XVIII.

ON COMPOUND FLOWERS.

A Head, or Capitate. — Receptacle. — Involucrum. — Compound Flowers. — Calyx. — Elongation of Tuft or Pappus. — Coma. — Akéne. — Description of the Florets of a Compound Flower. - 47



CONTENTS.

vii

CONVERSATION XIX.

ON FRUIT.

Conversion of the Flower into Fruit. — Pericarp. —
Pea Pod. — Legumen. — Epicarp. — Mesocarp. —
Endocarp. — Fruits composed of one Carpel. —
Drupa, or Stone Fruits. — Fruits composed of several Carpels resulting from a single Flower. —
Pœony. — Raspberry. — Fruits formed of several Carpels, with the Calyx adhering. — Pome: Apple, Pear, Quince, Medlar. — Bacca or Berry: Gooseberry, Grape, Currant, Strawberry, Mulberry. —
Fir Cone. — Akéne. — Magnolia Cone. — Spanish Chesnut. — Horse Chesnut. — Page 61

CONVERSATION XX.

ON THE SEED.

Structure of the Seed. — Spermoderm. — Testa. — Mesosperm. — Endopleura. — Amnios. — Embryo Plant. — Cotyledons. — Eye, or Hilum. — Cicatrix. — Albumen. — Nucleus, or Kernel. — Germination. — Embryo, consisting of the Radicle, Plumula, and Cotyledons. — Seeds in germinating require Moisture and Subtraction of Carbon. — Oxygen required for that Purpose. — Light injurious to Germination. — Temperature requisite. — Of sowing Seed. — Knight's Experiment on the Direction of the Roots and Stems. — Modes of sowing. — Advantage of sowing thin. — Sowing in Gardens. — Hot beds, Garden-pots, &c. — Transplantation. 83



viii

CONTENTS.

CONVERSATION XXI.

ON THE CLASSIFICATION OF PLANTS.

Necessity of some Mode of Classification. — Species. — Genera. — Nomenclature. — Character of Genera. — Page 107

CONVERSATION XXII.

ON ARTIFICIAL SYSTEMS OF CLASSIFICATION OF PLANTS.

Analytic Method. — Flore Francoise. — System of Linnæus. - 119

CONVERSATION XXIII.

ON THE NATURAL SYSTEMS OF CLASSIFICATION.

Method of Tatonnement. — Method of general Comparison of Mr. Adamson. — Method of Subordination of M. De Jussieu and M. De Candolle. — Comparative Importance of Organs. — Organs of Reproduction and of Nutrition.

CONVERSATION XXIV.

ON BOTANICAL GEOGRAPHY.

Geographical Distribution of Plants. — Difference of Habitation and Station of Plants. — Botanical Regions. — Their Boundaries. — Various Modes of conveying Seed from one Region to another. —



CONTENTS.

ix

Proportion of Monocotyledons in the Vegetable Kingdom.—Dycotyledons increase towards the Equator, Acotyledons towards the Pole.—Ligneous Plants increase towards the Equator, Herbaceous Plants towards the Pole.—Annuals most common in Temperate Climates.—Greater Variety of Plants in hot, than in cold Climates.—Social Plants.

Page 153

CONVERSATION XXV.

ON THE INFLUENCE OF CULTURE ON VEGETATION.

Races. — Varieties and Variations, derived from Species. — Influence of Culture in producing these Modifications. — Hebrides. — Dutch Tulips. — Hydrangia. — Effect of Soil on Variations. — Of Grafting. — Of Lopping. — Of Pruning Trees. — Resinous Trees. — Green-house Plants. — Fruit Trees.

CONVERSATION XXVI.

ON THE DEGENERATION AND THE DISEASES OF PLANTS.

Degeneration of Organs producing Monstrosities. —
Monopetals. — Failure of Seed. — Double-blossomed Flowers. — Conversion of Petioles into
Leaves. — Acacia of Arabia — Conversion of Petioles into Tendrils, of young Shoots into Thorns —
Difference of Thorns and Prickles. — Diseases of
Plants ranged under Six Heads: 1st Class, Constitutional Diseases — Variegated Leaves. — 2d



X CONTENTS.

Class, Diseases produced by Light, Heat, Water, Air, or Soil, improperly applied. — 3d Class, Diseases from Contusions and external Injury — Loss of Leaves and of Bark — Flagellations — Unskilful Pruning — Pollarding. — Page 177

CONVERSATION XXVII.

THE DISEASES OF PLANTS CONTINUED.

4th Class, Diseases arising from the Action of Animals on Plants — Devouring of Leaves by Quadrupeds or Insects, &c. — 5th Class, Action of Plants on each other — Parasitical Plants — True and False Parasites, External and Internal — Creeping Plants — Rhizomorpha — Mistletoe — Cuscuta — Orobanche — Fungi — Erishe — Rhizotonia — Smut — Rot — Ergot — Rust. — 6th Class, Diseases of resulting from Age, and causing the Death of Plants.

CONVERSATION XXVIII.

ON THE CULTIVATION OF TREES.



CONTENTS.

χį

CONVERSATION XXIX.

ON THE CULTIVATION OF PLANTS WHICH PRODUCE FERMENTED LIQUORS.

Fermentation. — Of Sap. — Of Fruits. — Of Grain, producing Wine, Beer, Cyder, Perry. — History and Culture of the Vine. — Page 239

CONVERSATION XXX.

ON THE CULTIVATION OF GRASSES, TUBEROUS ROOTS, AND GRAIN.

Meadows, perennial, renovated chiefly by the Spreading of the Roots, require a moist Climate.— Sustain the Soil on Declivities.— Artificial Grasses, not forming permanent Meadows, but entering into a Course of Cropping.— Clover.— Saintfoin.— Lucerne.— Leaves used as Forage for Cattle.— Tuberous Roots: Beet, Parsnips, Turnips, Carrots, Potatoes.— Of Corn: Straw, Grain, Flour, Husk, Beard.— Corn divided into Three Series: 1. Wheat, Fermentation of Bread, Gluten, Yest, Leven, Rye, Barley; 2. Series: Oats, Philares, Rice; 3. Series: Corn having the Pistils and Stamens in different Flowers.— 254

CONVERSATION XXXI.

ON OLEAGINOUS PLANTS AND CULINARY VEGETABLES.

Fixed Oils. — Volatile Oils. — Fixed Oils ranged under Three Heads: 1. Olive Oil, the Produce of Warm Climates; 2. Nut Oil, of Temperate Cli-



xii contents.

mates; 3. Oils obtained from Herbs. - Of the Olive Tree. - Of Nut Oil. - Oleaginous Herbs enter into a Course of Cropping. - Rape. - Poppy. - Flax. - Subterranean Arichis. - Culinary Vegetables. - Strong-flavoured: Artichoke, Asparagus, Rhubarb, Prussic Acid, Celery. - Insipid Plants. - From the Cruciform Family are obtained Cabbages, Brocoli, Cauliflower, Turnips, Radishes, Water Cresses, and Sea Cale. - From the Leguminous Family, Peas, Beans, Lentiles, and Kidneybeans. - From the Cucurbitaceæ Family, Cucumbers, Melons, and Pumpkins. - From the Umbelliferous Family, Carrots, Parsley, Lettuces, and Hemlock. - From the Solanum Family, Potatoes, Tomarta, Belladona. - From the Fungi Family. Mushrooms. - Page 280