

INDEX

- ACETYLALIC ACID, 117, 138, 157
 Acetylene dicarboxylic acid, on
 addition of bromine yields di-
 bromomaleic acid, 106
 α -Acrose, divided by means of
 yeast, 32
 Acrylic acid series, 101
 Active derivatives formed by the
 action of hydrochloric acid on
 albuminoids, 48
 — substances, list, 14
 Activity, disappearance of, in
 derivatives, 21
 — in which compounds it is re-
 tained, 21
 — optical, 9, 10
 — prediction of, confirmed, 19
 — refutation of supposed, 20
 — what is a sufficient cause for, 23,
 25
 Addition, mechanism of, in the
 formation and transformation of
 isomers, 105
 Adonite, 79, 86
 Alanine, no rotation as yet observed,
 163
 Aldehyde ammonia, 25
 — thio derivatives of acetyl-, &c.,
 117
 Aldoximes, stereomeric, 172
 Alkaloids exhibit the highest known
 optical rotation, 159
 Allyl alcohol, inactivity, 95
 Allylene type, second case of optical
 activity, 103
 Amido acids, slight activity, 163
 Ammonium bimalate, 11, 52
 — chlorides, substituted, inactive
 (except methylethylpropyliso-
 butyl), 181
 Ammonium chlorides, trimethyliso-
 butyl, two isomers, 180
 — — methyldiethylisoamyl, two
 isomers, 181
 — compounds, substituted, stereo-
 merism of, 180
 Ammoniums, substituted, stereo-
 merism of salts of, 180
 Amyl alcohol, 16, 31
 — — active rotation of derivatives,
 Guye's theory, 156
 — — chloride of secondary, active,
 24
 — — iodide of secondary, active, 24
 — — secondary, 20
 — alcohols, inactive, obtained from
 active carbohydrate by fermen-
 tation, 23
 — chloride, 16
 Amyl hydride, found to be inactive,
 21
 — iodide, 16
 Amylmalic acids, preparation of
 four, 57
 Amyl series, activity among mem-
 bers, 21
 Amylene, found to be inactive, 21
 Anethol, 101
 Angelic acid, 101
 Anil compounds, 173
 Apiol, 111
 Apocinchonine, specific rotation,
 137
 — hydrochlorate, specific rotation,
 101
 Arabinose, 85, 86
 — r- and l-, 64
 — multi-rotation, 149
 Arabite, its strikingly small rotation,
 162

- Arabonic acid, 64
 — — its rotation and that of its lactone, 148
 — — conversion into ribonic acid, 78
 Asparagine, poisoning power of *r*- and *l*-, 12
 — inactive, doubling by crystallisation, 39
 Aspartic acid, 15, 22, 32, 47
 — — inactive, obtained by mixing the dextro- and lævo-rotatory, 53
 — — rotation, 163
 Astracanine, formation from the simple salts, 39
 Asymmetric carbon, character of the isomerism due to the, 9
 — conditions of formation of compounds, 45
 Asymmetry, manifested in chemical, crystallographical, physical, and physiological properties, 10, 11
 Atropine, its activity, 73
 — contains at least two asymmetric carbon atoms, 61
 — its modifications, 61
 Atropines, dextro- and lævo-rotatory, 61

BENZALLEVULIC acid, 102
 Benzene, configuration, 1, 128
 — derivatives, 127
 — its acetylene nature, hence no prospect of stereomers, 127
 — the problem of the relative position of the six carbon atoms is not yet completely solved, 128
 — stability of, accounted for, 132
 Benzildioximes, configurations of the three isomers, 179
 Benzoin, hydro- and isohydro-, their constitution, 77
 Benzylmalimide, two isomers, 177
 Bi- and isobi-desyl, constitution, 77
 Bi-rotation, 148
 Boiling point, of racemic compounds, 43
 Boric acid, effect on rotation of polyatomic alcohols, 151
 Borneol as example of a transformation caused by changing one of the asymmetric atoms, 72, 74

 Borneol, rotation in benzene and in alcohol, 153
 Borneols, four active, and their derivatives prepared by reducing camphor, 58, 59
 Brassic acid, 101
 β -Bromacrylic acid, 101
 Bromethylenes, Paterno's explanation of their isomerism, 1
 Bromine, addition of, to cinnamic acid, giving an example of racemic compound containing two asymmetric atoms, 68
 — addition products (of crotonic and iso-crotonic acids, hypogæic and gaidic acids, oleic and elaidic acids, erucic and brassic acids, mesaconic and citraconic acids), as examples of racemic compounds as yet undivided, 69
 Bromnitroethane, 25
 Bromobenzene, inactive, converted in the animal organism into active bromophenylmercapturic acid, 45
 Bromocinnamic acid from cinnamic acid dibromide, 96
 — — inactivity of, 3, 96
 α - and β -Bromocinnamic acids, 101
 α - and β -Bromocrotonic, and bromisocrotonic acids, 101
 Bromoglycollic acid, 25
 Bromomalic acid, 24
 Bromomethacrylic acid, 101
 Bromophenylcystine, 18
 Bromophenyl-lactic acid, division of, as example of division of a compound containing two asymmetric carbon atoms, 69
 Bromopropionic acid, 24
 Bromopseudobutylene, 100
 Bromosuccinic acid, 24
 — — formation from malic acid, 24
 — — from malic acid, inactivity, 49
 Butyl alcohol, 14
 — — secondary, division, 31
 — alcohols, inactive, obtained from active carbohydrate by fermentation, 23
 Butylene glycol, division, 31

CAMPHOR, 19
 — modifications, two camphoric acids corresponding to each, 66

- Camphor series, synthesis of
 racemic bodies, 40
 — yields two isomeric borneols, 66
 Camphoric acid, 60, 102
 — — anhydride of the active, 49
 — — specific rotation, 139
 — acids, 60, 66
 Carbazides, 174
 Carbodi-imides, probably not stereo-
 meric, 175
 Carbohydrates, formed by plants
 from inactive material, 45
 — conspectus, 91
 Carbon atom, asymmetric, represen-
 tation by models, 8
 — atoms, graphic representation of
 several connected, 54
 — — asymmetric, formation of iso-
 mers containing several, 65
 — doubly linked, 99
 — — — graphic representation, 98
 — — — prediction of isomerism,
 99
 Carboxyls, proof of their neighbour-
 ing position, 113
 Carvol, 19
 Catalytic influence, bringing about
 inactivity, 48
 Catalysis, racemising by, 48
 Chitamic acid, 163
 Chloral alcoholate, 25
 — bornylates, prepared in four
 modifications, 63
 — hydrocyanide, 25
 — sulphhydrate, 25
 Chloroethylidene oxide, 26
 Chlorobromomethanesulphonic
 acid, 25
 α -Chlorocinnamic acid, 102
 α - and β -Chlorocrotonic and chlori-
 sacrotonic acids, 100
 Chlorofumaric acid, 20, 96
 — — obtained by treatment of tar-
 taric acid with PCl_5 , 22
 Chloromaleic acid, 20, 96
 Chloromalic acid, 24
 Chloroplatinates of substituted
 ammoniums, isomeric, 180
 Chloropropionic acid, 24
 Chlorosuccinic acid, 15, 22, 48, 96
 Cholalic acid, specific rotation, 139
 Cinnamic acid, 101
 — — addition of bromine to, 68
 — — dibromide, 68
 — — — division, 25
 Cinnamic acid, dichloride, 25
 — — series, 101
 Cinchonine, rotatory power and
 constitution, 71
 Cinchonidine, rotation in alcoh-
 ol and in aqueous solution, 141
 — rotatory power and constitution,
 71
 — specific rotation, 137
 Cinchonine, rotation in alcoholic
 and in aqueous solution, 141
 — rotatory power and constitution,
 71
 — equal rotation of the sulphate
 and selenate not due to their
 isomorphism, 136
 — specific rotation, 137
Cis and *trans* isomerism, 125
 Citraconic acid, 94, 102
 Citric acid, 23
 Cobalt, ammoniacal compounds of,
 186, 196
 Cocaine, 61
 Codeine, specific rotation, 137
 Conductivity, electric, of ammonia-
 cal compounds of cobalt, 186
 — — of ammoniacal compounds of
 platinum, 188
 — — of racemic compounds, 44
 Configuration, favoured, 55
 Configurations, determination of,
 in the sugar group, 81
 Coniferyl alcohol, from inactive
 coniferine, inactive, 96
 Conine, 17, 30
 — conversion into an isomer,
 177
 — *r*-malate of *r*- and *l*-, 66
 Conquinamine, rotation in alco-
 holic and in aqueous solution,
 141
 — specific rotation, 137
 Conversion, mutual, of active
 bodies, 47
 Copellidine, 18, 30
 Croceocobaltamine, salts of, 196
 Crotonchloral, gives with amides
 two isomers, 69
 Crotonic acid, 22, 95, 96
 Crotonylene bromide, 100
 Cumarine, inactive, 127
 Cumaric acid, 102
 Cystin, 165
 Cystine, 14
 — derivatives, rotation, 165

- DEXTROSE, multitrotation, 149, 165
 Diacetyltartaric acid and anhydride, 150
 — — methyl ester, rotation, 157
 α - γ -Diacipiperazines, 119
 Di-allyl, isomeric bromides, 78
 Diazotates, 175
 Dibromocinnamic acid, 48, 101
 Dibromoshikimic acid, 151
 Dibutyryltartaric acid, propylic ester, rotation, 157
 Dichlorosuccinic acid, as an example of isomerisation on bringing halogens into union with the asymmetric carbon atom, 24, 49
 Diethyl dipropionyl tartrate, rotation, 157
 Dihydrobenzene derivatives, 126
 Dihydroterephthalic acids, 127
 Dilatometer, 36
 Dimethyladipic acids, 78
 Dimethyldioxyadipic acids, two modifications, 77
 Dimethyldioxyglutaric acid, 78
 Dimethylene, ease with which it is saturated, 131
 Dimethyl-fumaric and -maleic acids, 102
 Diphenyl-fumaric and -maleic acids, 102
 Dimethylglutaric acid, 78
 Dimethyl diacetyl tartrate, rotation, 157
 Dimethylmaleic acid, 102
 Dimethyltricarballic acids, three inactive, 79
o-Dinitrostilbene, 101
 Di-oximes, isomeric, 173
 Dipentene, 47, 68
 Diphenyldiethylenediamine, 80
 Diphenylfumaric acid, 102
 Diphenylmaleic acid, 102
 Dipropionyltartaric acid, ethylic ester, rotation, 157
 Dipropylic dibutyryl tartrate, rotation, 157
 Dissociation by heat, theory of universal, 6
 — constants of fumaric and maleic acids, 113
 — electrolytic, 136, 138
 — — its bearing on the theory of rotation, 136, 145
 Ditoluyltartaric acid, rotation of esters, 155
 Divisibility of racemic compounds, by active compounds, 28
 — — — by organisms, 30
 — — — by synthesis, 40
 — — — spontaneous, 34
 Division into optical isomers, vain attempts, 21, 46, 77, 127, 170, 181
 Double linkage of carbon atoms, nature of, 97
 — molecules, liquids composed of, 134, 135
 Dulcitol, inactive indivisible, 89
 Dynamic hypothesis, 112
 ECGONINE, 61
 — l-, transformation into r-, 73
 — l- and r- yield the same active anhydroecgonine and the same ecgonic and tropic acids, 68
 Elaidic acid, 101
 Electrolytes, rotatory power, 136
 Electrolytic dissociation, 186
 — — and rotatory power, 136, 138
 Enantiomorphism of optical isomers, 11
 — causing opposite activity, 10
 Equilibrium between active isomers, conditions of, 49
 Erucic acid, 101
 Erythrene bromide, 76
 Erythrite, 74
 Ethoxysuccinic acid, 15, 29, 32
 — — rotation of salts, 138
 Ethyl alcohol, inactive, obtained by fermentation from active carbohydrate, 23
 — amyl, 17
 — cumaric acid, 102
 Ethylene derivatives, inactivity of, 96
 — — plane or three-dimensional configuration of, 82
 Ethylidene anilines, 173
 — chlorosulphinic acid, 26
 — iodobromide, 25
 — lactic acid, 14
 — methethylate, 26
 Ethylmalic acid, rotation, 157
 — — obtained by heating fumaric and maleic acids with sodium ethylate, 53
 α -Ethyl piperidine, 17, 30

- FAVoured configuration, 55
 Formylthymotic acid, 127
 Fumaric acid, 96
 — — inactivity of, 3
 — — indivisibility, 46
 — — series, 102
 Furfuracrylic acid, 101
 Furfurol, from active arabinose or xylose, is inactive, 22, 96
- GALDID acid, 101
 Galactonic acid, 29, 65, 89
 — — rotation compared with the lactone, 147, 149, 164
 — — from mucic acid, divisible, 53
 — — converted into talonic acid, 73
 Galactose, 89
 — multi-rotation, 149, 165
 α - and β -Glucoheptonic acids, formed from glucose, 67
 — — — rotation, 164
 — acid, rotation compared with that of the lactone, 148
 α -Gluco-heptose, rotation, 165
 Gluconic acid, 64
 — — its rotation compared with the lactone, 147, 164
 l-Gluconic acid, formed with l-mannonic acid from arabinose, 67
 Gluco-octonic acid, rotation, 164
 α - and β -Gluco-octonic acids, formed from heptose, 67
 — — — transformation of α - into β -, 73
 α -Gluco-octose, rotation, 165
 Glucose, 64
 — gives two isomeric glucoheptonic acids, 67
 — converted into levulose, 92
 — multi-rotation, 149
 α -Glucosectite, rotation, 161
 Glucoses, isomeric, 64, 82, 87, 165
 Glutamine, rotation, 163
 Glutamic acid, 17, 32, 163
 — — formed on heating albuminoids with baryta, 48
 Glutaric acid, type 78
 — acids, the two isomeric dimethyl- and dimethyldoxy-, 78
 — — methylethyl- and methylpropyl-, two modifications, 78
 Glyceric acid, 14, 31
 Glyceric acid, change in rotation, 146
 — — specific rotation, 138, 139
 — — esters rotation, as evidence for Guye's hypothesis, 158
 Glycols, substituted, two modifications, 69
 Gulonic acid, l- and r-, 65
 — — lactone, division, 39
 — — rotation, compared with that of the lactone, 148
 Gulose, configuration, 88
 — l- and r-, 64
- HALOGEN, derivatives containing asymmetric carbon, active, 24
 — — — inactive, 25
 Halogens, attached to the asymmetric carbon atom cause isomerisation, 24, 49
 Hexabromobenzenes, isomeric, 123
 Hexachlorbenzenes, isomeric, 123
 Hexahydrophthalic acids, two modifications, 121
 Hexaisophthalic acids, two modifications, 122
 Hexahydroterephthalic acids, two modifications, 121
 Hexamethylene derivatives, 120
 — — activity among, 123
 Hexyl alcohol, 17
 — — secondary, 17, 31
 — — — chloride, 24
 — — — iodide, 24
 — chloride, 17
 — iodide, 17
 Hexylic acid, 17
 Homoaspartic acid, division, 39
 Homometa-oxybenzoic acid, indivisible, 46
 β -o-Homomethoxybenzoic acid, visible, 127
 Homosalicylic acid, indivisible, 46
 Hydrazones, isomeric, 174
 Hydrobenzoin, 77
 Hydrochlorapocinchonine, specific rotation, 137
 Hydrogen, silver fulminate, 25
 Hydromellithic acid, and iso-, 122
 Hydronaphthalenediamine, 20
 Hydropiperic acid, α - and β -, 102
 Hydroshikimic acid, 123
 Hydrosorbic acid, 101
 Hydroterephthalic acids, 121

- Hyoscyamine, conversion into atropine, 48
 Hypogæic acid, isomers, 101
 Hyponitrous acid, 176
- IDOSACCHARIC acid**, rotation and configuration, 161
 Idose, 64
 Inactive indivisible type, 50, 74
 Inactivity among compounds containing the asymmetric carbon atom, 27
 — of a body arising from the incompatibility of its constitution with rotatory power, 23
 Indivisible inactive type, 50, 74
 Indivisibility, in absence of the asymmetric carbon atom, 46
 Influence, mutual, of the groups forming a molecule, 93, 110
 Inorganic compounds, stereomerism of, 185
 Inosite, 123
 — its strong rotatory power caused by ring formation, 161
 — indivisibility, 46
 Inosites, r- and l-, configuration, 124
 β -Iodacrylic acid, 101
 Iodohexyl, 49
 — from mannite, 24
 Ions, rotation of acid, 140
 Isapiol, 101
 Isocinnamic acid, 101
 Isocrotonic acid, 101
 Isodibromosuccinic acid, prepared from maleïc acid by addition of bromine, 108
 — from maleïc acid, converted into racemic acid, 109
 Isohydrobenzoïn, 77
 Isohydromellithic acid, 122
 Isomerism, due to the asymmetric carbon atom, character of, 9
 — due to doubly linked carbon, character of, 99
 Isomerisation, on attaching halogens to the asymmetric carbon, 24, 49
 Isomers, number due to several asymmetric carbons, 56
 — optical properties of, 10
 Isopropoxysuccinic acid, 15
 Isopropylphenylchloracetic acid, 24, 49
 Isopropylphenylglycollic acid, 18, 24, 29
 — — rotation, 159
 Isosaccharinic acid, rotation, 164
- KETOXIMES**, stereomeric, 173
 Kinetics of racemising, 49
- LACTIC acid**, 58
 — — division, 29, 31
 — — rotation, 159, 163
 — — — alteration of, on dilution, 144
 — — — decrease on standing, 146
 Lactid, 14
 — rotation, 147
 Lactone formation, 130
 — and rotatory power, 147
 Lactose, multi-rotation, 149
 Levulose, configuration, 88
 — yields on reduction the isomers mannite and sorbite, 67
 — multi-rotation, 149
 — breaks up on oxidation into glycollic acid and inactive tartaric acid, 90
 Leucic acid, rotation, 159
 Leucine, 17, 32, 47, 163
 — inactive, formed on heating albuminoids with baryta, 48
 Leucinephthaloylic acid, 19
 Limonene, 18, 40
 — nitrosochloride and its derivatives, 60, 67
 — — rotation, 166
 Linkage, treble, graphic representation, 104
 Lyxose, 86, 160
 Lyxonix acid, 87
- MALAMIDE**, 16
 Maleïc acid, 96
 — — dissociation constant, 113
 — — easy conversion into fumaric acid, 106
 — — proof of configuration, 111
 Malic acid, 15, 22
 — — acetyl, and Guye's theory, 117, 157
 — — — and its anhydride, rotation, 150

- Malic acid, acetyl, salts of, their equal rotatory power, 138
 — — alteration of rotation on diluting, 145
 — — rotation of salts, 153
 — — amylic esters, preparation of four, with two active malic acids, and two active amylic alcohols, 57
 — — behaviour in solution, with regard to rotation, 143
 — — conductivity of active and inactive the same, 44
 — — activity among derivatives of, 22
 — — increase of lævo-rotation in the diluted solution on warming, 144
 — — inactive, 51
 — — — its acid ammonium salt, 52
 — — prepared by heating fumaric or maleic acid with soda, identity of this with the divisible malic acid, 52, 53
 — — remarkable change in its rotation on diluting, 145
 — — salts of, their equal rotatory power, 139
 Maltose, multi-rotation, 149
 Mandelic acid 18, 29, 31
 — — rotation, 143, 159
 — — inactive from the *r*- and *l*-modifications, identical with synthetic, 40
 Mannite configuration, 88
 — hexachlorhydrin of, 25
 — isomers, 80
 — rotation, 161
 — yields on oxidation mannose and levulose, 92
 Mannoheptonic acid, rotation, 164
 Mannoheptose, rotation, 165
 Mannonic acid, 29, 64
 — — *l*-, formed from arabinose, 67
 — — *l*- and *r*-, 64
 — — rotation, 164
 — — compared with that of the lactone, 147
 Mannonose, rotation, 165
 Manno-octose, rotation, 165
 Mannosaccharic acid, configuration, 161
 — — rotation, compared with that of the lactone, 147, 164
 Mannose, configuration, 88
 — divided, 92
 Mechanism of addition in forming and transforming isomers, 105
 Mechanics of the atoms, 5
 Melting point, quantitative expression for lowering of, 43
 Menthol, 72
 Mesaconic acid, 95, 102
 β -Metahomosalicylic acid, indivisible, 127
 Methoxysuccinic acid, 15
 — — rotation of salts, 138
 Methoxytoluyllic acid, indivisible, 46, 127
 Methylamyl, 21
 Methylcumaric acid, 102
 Methylconine, 18
 Methylenehydropiperidine (copellidine), 18
 Methyl glucoside, high rotation of, 151
 — glucosides from glucose, 67
 Methyl benzoyltartrate, high rotation, 159
 Methylmalic acid, 16
 — — formed by the action of fungi on citraconic acid, 95
 — — formed by heating fumaric and maleic acids with sodium methylate, 53
 α -Methylpiperidine (pipercoline), 17
 — active, conversion into an isomer, 177
 β -Methylpiperidine, 20
 Mica, active combinations of, 11
 Models, 8, 55
 Molecular dimensions, equality of, in optical isomers, 10
 — rotation, 133
 Monochloropropylene, 100
 Morphine, specific rotation, 137
 Mucic acid derivatives, 89
 Multi-rotation, 148
 NICOTINE, 18
 — specific rotation, 137
 — acetate, rotation in alcoholic and in aqueous solution, 141
 Nitro-camphor and derivatives, rotation, 167
 Nitrogen, combined with carbon, configuration, 178
 — — — nitrogen, 179

- Nitrogen compounds, stereochemistry of, 169
 — trivalent, 169
 — — in rings, 176
 — pentavalent, 180
 — — graphic representation of isomers, 182
 Nitrolepipiperidine, 60
 o- m- and p-Nitrophenylcinnamic acids, 102
 Nitrosohexahydroquinolic acid, division, 30
 Nitrostilbene, constitution of bromides, 77
 Nitrostyrolene, 101
 Nitrotartaric acid, inactivity, 48
 Nitrothymotic acid, 127
- OBJECTIONS to the representation of benzene by tetrahedra, 128
 — to the theory of unsaturated isomers, 109
 Oleic acids, 101
 Optical isomerism and the asymmetric carbon atom, 13
 — isomers, properties of, 10
 Organism, formation of active compounds in the, 65
 Organisms, division of racemic compounds by, 30
 — action of modified, by transposition of groups, 32
 Oxalic acid formed from active sugar or active tartaric acid is inactive, 22
 — — indivisible, 46
 Oximes, isomerism of, 171
 Oxy-acids, small rotation of, 163
 — — change of rotation on diluting, 145
 — — high rotation on introducing a benzene residue, 165
 Oxybutyric acid, 15, 139, 159
 α -Oxybutyric acid, division, 29
 γ -Oxybutyric acid most readily forms a lactone, 130
 Oxyglutaric acid, 16
 — — rotation, 164
 Oxy-pyrotaric acid, 20
 Oxy-pyrotartaric acid, 20
- PAPAVERINE, 20
 Pentose group, 63, 82, 84
- Pentoses, rotation, 160
 Perseite, rotation remarkably small, 161
 Phenose, 125
 Phenoxacrylic acid, 150
 Phenylamidopropionic acid, 18, 163
 Phenylamyl, 19
 Phenylbromacetic acid from mandelic acid, 25, 49
 — — from phenylglycollic acid, 24
 Phenylbromolactic acid, 29, 150
 Phenylbromomercapturic acid, 18
 Phenylchloracetic acid, 24, 25, 49
 Phenylcystine, 18, 163
 Phenyl dibromobutyric acid, division, 29
 Phenylmercaptan, 22
 Phenyltrimethylenecarboxylic acids, three isomeric, 115
 Phenylurethane, rotation in various solvents, 154
 Phthalylamidocaproic acid, 19
 Physiological properties of optical isomers, 12
 — different, of the two optical isomers, 12
 β -Picoline, 20
 Picryl α - and β -naphthyl hydrazines, 179
 Picrylhydrazines, isomerism, 179
 Pinenedihydro-chlorides and -bromides, 122
 α -Pipicoline, 17, 30
 Piperidine monocarboxylic acids, five isomeric, 176
 Piperylene, isomeric bromides of, 78
 Platinum, bivalent, 187
 Platopyridine chloride, 193
 Platosammine salts, 188, 191
 Platosemidiammine salts, 188, 191
 Podocarpic acid, specific rotation, 139
 Polymethylene rings, Baeyer's hypothesis, 131
 — derivatives, 114
 Polymethylenes, relative stability, 131
 Praseocobaltammine, salts of, 196
 Product of asymmetry, 155
 Projections of models, 55
 Propionaldoxime, 171
 Propoxysuccinic acid, 15
 Propylene diamine, 14
 — glycol, 14, 150

- Propylene oxide, 14, 150
 Propyl alcohol, 20, 23
 Propylmandelic acid, rotation, 165
 α -Propylpiperidine (conine), 17
 Pyrotartaric acid, 29
 — — inactivity, 48
 Pyrrolylene bromide, example of inactive indivisible type, 76
- QUINAMINE salts, rotation of alcoholic and aqueous solutions, 141
 — specific rotation, 137
 Quinic acid, 123
 — — as an example of constancy of rotation on dilution, 144
 — — inactive, divisibility, 44
 — — salts, rotation in alcoholic and in aqueous solution, 140, 141
 — — specific rotation, 139
 Quinicine, rotatory power and constitution, 71
 Quinidine, rotation in alcoholic and in aqueous solution, 141
 — rotatory power and constitution, 71
 — specific rotation, 137, 160
 Quinine group, transformations in, 71
 — rotatory power and constitution, 71
 — specific rotation, 137
 — sulphate, rotation in alcoholic and in aqueous solution, 141
- RACEMATE of sodium and ammonium, 34
 — — — — potassium, 38
 Racemic acid, 74
 — — division, 29, 31, 74
 — — compounds, 28
 — — characteristics, 41
 — — properties of, 41
 Racemising by heat, by catalysis, 47, 48
 Rhammonic acid, rotation compared with that of the lactone, 147
 Rhamnose, multi-rotation, 149
 Ribonic acid, 64, 163, 164
 — — rotation compared with that of the lactone, 147
 Ribose, 160
- Ring formation, stability of, 129
 — — connection with unsaturated isomers of maleic type, 113
 — — influence on rotation, 146, 150, 164
 — — with boric acid and hydroxyl groups, 151
 — — stability, 129
 Rings, four carbon, 118
 — three carbon, 114
 — six carbon, 120
 Rotation, free, 55
 — — cessation of, in the case of doubly linked carbon, 97
 — — optical, causes which influence, 134
 — — influence of concentration, 142, 156
 — — influence of groups attached to the asymmetric carbon atom, 154
 — — influence of group weights, 155
 — — influence of ring formation, 146, 150, 164
 — — influence of solvent, 135
 — — influence of type, 162
 — — molecular, 133
 — — raised by addition of boric acid, 151
 — — relative, of singly linked carbon atoms, 54
 Rotatory power and electrolytic dissociation, 136, 138
 — — and lactone formation, 147
 — — of non-electrolytes, 153
 — — and ring formation, 146
 — — Guye's hypothesis, 154
 — — of electrolytes, 136
 — — of imperfect electrolytes 141
 — — remarkable cases, 165
 — — numerical value of, 133
 — — — — — in relation to constitution, 134
 — — — — — tables, 137, 139
 — — — — — variation with varying conditions, 134
 — — varying, of tartaric acid solutions, 136
- SACCHARIC acid, configuration, 161
 — — group, rotation, 160, 164

- Saccharic acid, rotation compared with that of the lactone, 147, 164
 — — type, 70, 79
 Saccharic acids, 80
 Saccharin, multi-rotation, 149
 Saccharinic acid, rotation compared with that of the lactones, 147, 164
 Salicylic aldehyde, inactive, 127
 Salts of active bases and acids, rotatory power, 136, 138
 — — multivalent metals with glyceric acid, rotation, 138
 — — — — polyatomic acids, rotation, 152
 Santonine derivatives, rotation, 160, 167
 — — — the highest known, 159
 Scacchi's salt, 37
 Scyllite, 125
 Serine, no rotation as yet observed, 163
 Shikimic acid, rotation, constancy on dilution, 145
 — — — of derivatives, 166
 — — — of salts, 108, 138
 Sodium ammonium racemate, 34
 — nitro-ethane, 25
 — potassium racemate, 38
 Sorbite, configuration, 88
 — rotation, remarkably small, 162
 Spontaneous division of racemic compounds, 34
 Stability, absolute criterion of, 111
 — of unsaturated stereomers, unequal, 100
 — equal, of active isomers, 47
 Strychnine, specific rotation, 137
 — isomorphous sulphate and selenate, equal rotation of, 136
 Styrolene, 20, 95
 Succinic acid, inactive, 22
 — — formed from asparagine, is inactive, 22, 23
 — — obtained by reduction of malic acid, 22
 — acids, bisubstituted, obtained in two modifications, 69, 77
 Sugars, configuration, 82
 — conspectus, 91
- TALOMUCIC acid, rotation, 161, 164
 — — — compared with that of the lactone, 147, 164
- Talomucic acid, configuration, 161
 Talonic acid, 65, 89
 Talose, 89
 Tartar emetic, abnormally large rotation of, 140, 152
 Tartaric acid, 81
 — — activity among derivatives, 21
 — — conversion into racemic acid by oxide of iron or of aluminium, 48
 — — derivatives, 22
 — — esters, rotation in different solvents, 154
 — — formation, from racemic acid, by Pasteur, 28
 — — imides, isomeric, 178
 — — inactive, indivisible, 51, 74, 81
 — — rotation and Guye's hypothesis, 158
 — — rotation specific, 139, 142, 164
 — — — — many circumstances affecting, 135, 143
 — — — — in tartar emetic, 140
 — — type, 74
 Temperature, effect on equilibrium, 50
 — of conversion of active substances, 35
 Tensimeter, 37
 Terephthalic acid, di- and tetrahydrides of, 125
 Terpenes, 122
 Terpines, structure, 122
 Tetrabromobutane, 76
 Tetrahedron theory, 129
 Tetrahydrobenzene derivatives, 125
 Tetrahydrobenzoic acid, bromides of, 122
 Tetrahydronaphthylene diamine, 19, 30
 Tetrahydroterephthalic acids, 126
 Tetrahydroterpenes, α - and β -structure, 122
 Tetramethylene derivatives, 118
 Tetroses, 82
 Thermodynamics of racemising, 49, 50
 Thiacetones, polymeric, configuration, 116
 Thialdehydes, polymeric, configuration, 116
 Thiobenzylcrotonic acid, 101
 Thiodilactic acids, two modifications, 77

INDEX

211

- β -, Thio-ethylcrotonic acid, 101
 β -Thio-phenylcrotonic acid, 101
 Tiglic acid, 95, 101
 Tolane bromides, 100
 — chlorides, 100
 o -Toluidine, indivisible, 46, 127
 Treble linkage of carbon atoms,
 representation of, 104
 Trimethylene, 114
 Trimethylene dicarboxylic acids,
 isomeric, 115, 117
 Trimethylethylstibine iodide, 20
 Tri-oxyglutaric acid, a second in-
 active indivisible type, 78
 — — rotation, 164
 — acids, 87
 Trithiodimethylmethylene, 116
 Trithiomethylene, 114
 Tropaic acid, 18, 159
 — — rotation, 159
 Tropine, 62
 Truxillic acids, 118
 Turpentine yields two borneols,
 66
 Tyrosine, 18, 163
 — inactive, obtained on heating
 albuminoids with baryta, 48
 URAMIDOSUCCINAMIDE, 16
 Uramidosuccinic acid, active, 22
 VALERIC acid, 16
 — — esters, rotation and Guye's
 theory, 158, 159
 Vanilline, inactive, 127
 Violeocobaltammies, salts of, 196
 XYLITE, 79, 86
 Xylonic acid, 64, 87, 164
 — — rotation compared with that
 of the lactone, 147, 164
 Xylose, 86, 160
 — multi-rotation, 149, 165
 YEAST, action on various sugars, 32

Errata.

- Page 52, line 11 from top, for *ammonia* read *ammonium*.
 Page 87, the formula of lyxose should be as on page 88.
 Page 95, line 12 from bottom, for *styrol* read *styrolene*.
 Page 101, line 4 from top, for *nitrostyrol* read *nitrostyrolene*.
 Page 166, line 3 from bottom, for *limonenenitroso chloride* read
limonene nitrosochloride.

PRINTED BY
 SPOTTISWOODE AND GO., NEW-STREET SQUARE
 LONDON