

Index

- abdomen
 examination, 49–51, 59–61
 visceral trauma, 247–8
- Aberdeen Area Infant Mortality Study (AAIMS), 198–9
- abscesses, 102
- accident and emergency (A&E) departments
- accidental death, 41
- acid-base disorders, 213
- acute tubular necrosis, 219–20
- acylcarnitine (AC) profiles
 bile, 121
 blood, 117–20
 cultured fibroblasts, 122
- adenoviruses, 224
- adrenal glands, 51
- adrenal haemorrhage, 221, 223, 248
- advocacy groups, 41
- agency policies and practices, improving, 78–9
- alcohol consumption, maternal, 134
 arousability and, 192–3
 bed-sharing and, 150–1
 brainstem 5-HT deficits and, 199–202
- alveolarcapillary dysplasia, 231–2
- ambulance service, 29
- American Academy of Pediatrics (AAP), 149, 151
- amino acid testing, 119–21
- anion gap, 108, 112
- anogenital injuries, 242–3
- anterior chest wall, 58–9
- anterior torso, musclocutaneous dissection, 58–9
- aortic stenosis/atresia, 226
- arcuate nucleus, 96, 162, 203
- arm chairs, cushioned, 137
- arousal, 189–94
 brainstem 5-HT mechanisms, 190–191
 breastfed infants, 143
 definition, 189
 failure, 178, 180, 189, 193–4
 inner-ear defects and, 164, 207–9
 intermittent hypoxia and, 189–90
 pontine Kölliker-Fuse nucleus, 162
 response to hypoxia and hypercapnia, 178–80, 189
- risk factors and, 191–3
 secondary depression phase, 180
- arrest, of suspects, 106
- arrhythmogenic right ventricular cardiomyopathy (ARVC), 160
- asphyxia, 236–7
 arousal response, 189
 cause of death classification, 41
 causes of sleep-related, 236–7
 parents' perspective, 6
 positional, 237
 sleep-related, 236–7
- asplenia, 219, 222
- autopsy, 48–53
 ancillary investigations, 51, 53
 brain examination
 cerebral white matter, 98
 brainstem, 96–7
 hippocampus, 97
 removal and fixation, 52, 95–6
 clinico-pathological summary, 53
 external examination, 48–9, 57–8
 forensic (see joint pediatric/forensic autopsy),
 histology samples, 49, 52
 imaging findings, 89–93
 initial incision, 49–50
 instruments, 49
 internal examination, 49–52
 joint forensic/paediatric. *See* joint forensic/paediatric autopsy
 microscopic examination, 53
 minimally invasive, 71–3
 molecular. *See* genetic testing
 NAME recommendations, 39–40
 parental consent, 12, 16, 95
 parents' perspective, 15, 46
 pathologists performing, 21–22, 24, 48
 release of body after, 22, 24
 religious objection laws, 41
 second post-mortem, 24
 suspicious signs, 30
 tissue retention, 22, 249–50
 UK guidelines, 55
 UK legal framework, 21–22
 US legal framework, 40–1
- autoresuscitation, 180, 189, 196
- axonal injury, 96–7, 257–8
- baby boxes, 138
- Baby Friendly Hospital Initiative (BFHI), 142–3, 146
- baby nests (baby pods), 138
- back-to-sleep campaigns
 bed-sharing SIDS after, 149–50
 recurrent SIDS after, 130
 SIDS rate after, 136–7
- bacteraemia, 100–1
- bacterial infections, 163, 219–23
- bacterial translocation, post-mortem, 100
- bacteriology
 interpretation, 100–4, 222–3
 samples, 53, 101–2
- Barth syndrome, 160
- battered-child syndrome, 237–9
- batteries, swallowed, 232–3
- BCGitis/BCGosis, 225
- bedding, 138–9
- beds, child and adult, 137–8
- bed-sharing, 149–53. *See also* overlaying
 autopsy findings, 53
 Bowden's studies, 2
 breastfed infants, 143, 151
 fall in SIDS rates and, 149–50
 hazardous interactions, 132, 134, 149–51
 risk in nonhazardous situations, 150–1
 risk-reduction strategies, 151–3
 substance misuse and, 134, 149–50
- bedside sleepers (side cars), 138
- bereavement, 12–15
 risk impeding
- β-amyloid precursor protein (βAPP)
 immunohistochemistry, 257–9
- beta-blockers, 175
- bile
 analysis, 119, 121
 samples, 109, 117
- biochemical analysis
 inherited metabolic disease, 119
 poisoning, 106–9
 tissue samples, 117–18
- biological risk factors, 127–8
- biopsy, targeted, 92
- birth-related injuries
 fractures, 245, 250

- retinal haemorrhages, 254
- subdural bleeding, 254
- bite marks, 240
- blankets, 139
- bleeding disorders, 241, 253
- blood cultures, 100–1
- blood samples, post-mortem, 51, 108
 - biochemistry and toxicology, 107–9
 - collection, 50
 - drugs detected, 112
 - inherited metabolic disease, 117–21
- blunt trauma, 239–41, 247
- body
 - external examination, 48–9, 57–8
 - measurements, 48
 - parents' perspective, 12
 - police actions, 29
 - release after autopsy, 22, 24
- bone scan, 243
- bony lesions, 56
- Bordetella pertussis*, 219
- botulism, 102
- Bowden, K. M., 1–3
- brachycephaly, 251
- brain
 - forensic pathology, 255–8
 - gross examination, 95–6, 255–6
 - microscopic examination, 257–8
 - removal, 52, 61–3, 95
 - sectioning and sampling, 52, 95–6, 249–50
 - size abnormalities, 252
 - swelling, 250, 255–8
- brainstem
 - arousal response, 190–1
 - cardiorespiratory control, 180–3
 - effects of inflammation, 183
 - gliosis, 96
 - mechanisms of SIDS, 95, 178–83
 - pathology, 96–7
 - serotonin system, 195–204. *See* serotonin (5-HT) system, brainstem
 - upstream influences on, 96–8, 182
- breastfeeding, 142–3
 - bed-sharing and, 143, 151
 - mechanisms of protection, 143
 - methadone-using mothers, 135
 - pacifier/dummy use and, 143, 146–7
 - protective effects, 142–3
- breathing control
 - brainstem circuitry, 180–3, 196
 - effects of inflammation, 183
 - neuromodulation, 182–3
- brief resolved unexplained events (BRUE), 237
- Brugada syndrome, 160–1, 175
- bruises, 238–41
 - ageing, 240–1
 - mimicking abuse, 241
- signs of abuse, 239–41
- bucket-handle fractures, 245
- bumper pads, 81–2, 139
- burns, 241–2
- C1-neurons, noradrenergic, 179–80
- caffeine, 134
- calcium (Ca²⁺)-related gene mutations, 173–4
- calcium, blood, 108
- car seats, 138
- carbohydrate disorders, inherited, 212, 215–16
- cardiac arrhythmias
 - channelopathies, 160–1, 172–6
 - fatty acid oxidation defects, 214
- cardiac disease
 - inherited, 160–1, 213
 - sudden deaths in infancy, 225–31
- cardiac tumours, 229
- cardiomyopathies, 160, 176, 214, 226–9
- cardiopulmonary resuscitation (CPR), 245, 247–8
- cardiorespiratory control, 180–3
- care of next infant (CONI)
 - programme, 3, 15–16
- carers (caregivers). *See also* parents
 - initial accounts, 31
 - interview by police, 32
 - risk factors for abuse, 30, 238–9
 - taking statements from, 33–4
- carnitine palmitoyltransferase type 2 deficiency, 120
- carotid bodies, 182, 190
- case-control studies, 136
- catecholaminergic polymorphic ventricular tachycardia (CPVT), 161, 175–6, 229
- cause of death
 - determination in US, 40–1
 - diagnostic shift, 6, 78
 - discussion with parents, 41, 87
 - final case discussion (UK), 86
 - inquest conclusions, 26–7, 35
 - parents' perspective, 6, 9
 - trends in US, 127–8
 - undetermined. *See* undetermined
- Centers for Disease Control and Prevention (CDC)
 - investigation of SUID, 37–9
 - SUID Case Registry, 82–3
- central nervous system
 - forensic pathology, 249–59
 - histology samples, 52
 - inherited defects, 162
 - pathways, 161–2
- cerebellum, delayed maturation, 96
- cerebral venous and dural sinus thrombosis (CVDST), 254–5
- cerebral white matter, 98
- cerebrospinal fluid (CSF) samples
 - collection, 51–2, 61–2
 - inherited metabolic disease, 117, 121
 - poisoning, 109
- cervical spine, excision, 66–7
- channelopathies, cardiac, 160–1, 172–6
- chemoreceptors, central, 190
- chest. *See also* rib fractures
 - examination, 48–50, 58–61
 - internal injuries, 248
 - skeletal trauma, 244–5
- child abuse, 237–59. *See also* suspicious deaths
 - autopsy findings, 239–59
 - mimics, 241, 246–7
 - pathologists' role, 238
 - risk factors and signs, 30, 238–9
- child-care settings, 156
- Child Death Overview Panel (CDOP), 23, 87–8
- child death review (CDR), 11, 75–83
 - home visit, 45–6
 - improving reporting of SUID, 80–3
 - information shared, 77
 - objectives, 45, 77
 - participants, 76–7
 - role in improving outcomes, 78–80
 - steps in case discussion, 77
 - steps in process, 76
- Child Death Review Case Reporting System (CDR-CRS), US National, 80–2
- child fatality review board, 40
- Child Practice Review (Wales), 23
- chloride, blood, 108
- choanal atresia, 231
- chorioamnionitis, 102
- chromosomal abnormalities, 164
- chromosomal microarray (CMA), 164–5
- classification, sudden infant deaths, 41, 236
- clavicular fractures, 245
- clinical exome sequencing, 166
- clinico-pathological summary, 53
- Clostridium difficile*, 222
- Clostridium perfringens*, 221
- coagulation disorders, 241, 253
- coliforms, 104
- commensal flora, respiratory tract, 104
- computed tomography, post-mortem (PMCT), 72–3, 89–91
 - intraosseous needle tracts, 90–1
 - normal variants, 89–90
 - post-mortem change, 89
- conclusion, inquest, 26–7, 35
- congenital heart disease, 225–6
- congenital infections, 102

Index

- contamination, microbiological, 100–4
- coronary arteries
anomalous origin, 226
fibromuscular dysplasia, 226
- Coroners
autopsy decision, 22
further enquiries, 22–3
history, 19–20
independence, 20
initial investigation, 21–2
inquest conclusion, 26–7
inquest decision, 23–4
inquest procedure, 25
inquest remit, 24–5
legislation, 20
notifying child deaths to, 20–1
paediatric post mortems, 22
suspicious deaths, 24
tissue retention, 22
UK legal framework, 19–27
unnatural deaths, 21
US, 19, 36–9
working with police, 21–2, 24, 29, 34–5
- Coroners (Inquests) Rules 2013, 20
- Coroners (Investigations) Regulations 2013, 20
- Coroners and Justice Act 2009, 20
- co-sleepers, bedside and in-bed, 138
- co-sleeping, 149–53. *See also* bed-sharing; sofa-sharing
hazardous interactions, 132, 134, 149–50
Māori practices, 152–3
- costochondral junctions, 59–61, 246
- cot death, 9, 53
- Cot Death Foundation (Netherlands), 11
- Cot Death Parent Association (Netherlands), 9, 11, 16
- cots (cribs), 138
- couches, sharing. *See* sofa-sharing
- coxsackieviruses, 223–4
- craniosynostosis, 251–2
- craniotabes, 52
- crib death, 9
- cribs. *See* cots
- criminal cases, 34–5
- Crown Prosecution Service (CPS), 34
- cyst-like brain lesions, 256
- cytomegalovirus (CMV), 102, 224
- Davidson, W. H., 1
- day-care environment, 156
- Dbx1 neurons, 182
- death certificate, 20
- death scene
doll re-enactment, 39, 43, 56
examination, 43, 56
first responders, 43
home visit, 32, 45–6
initial police actions, 29–30
physical evidence, 31
residues, toxicology, 109
unsafe sleep environment, 236
- dehydration, 111–12, 248–9
- delayed death in SIDS/SUID, 98, 255, 257
- dentate fascia, hippocampal, 97
- Department of Health and Human Services, US, 76
- dilated cardiomyopathy (DCM), 160, 227, 229
- disseminated intravascular coagulation (DIC), 219–20
- diving reflex, 97–8, 208–9
- DNA analysis. *See* genetic testing
- doctors, death certification, 20
- doll re-enactment, 39, 43, 56
- Dravet syndrome, 233
- drug misuse. *See also* substance abuse
bed-sharing and, 150
child deaths, 109–11
post-mortem screening, 111–12
pregnancy, 134–5
- drugs. *See also* toxicological analysis
numbers of deaths, 109–10
post-mortem screening, 111–12
- dual process model, grief, 13
- dummies. *See* pacifiers
- duodenal injuries, 248
- dura, 98, 252–5
acute intradural bleeding, 98, 252
incision, 62
sinuses, 254–5
subdural bleeding. *See* subdural bleeding
- dysmorphic features, 116, 213
- ears, examination, 48
- e-cigarettes, 132
- Ehlers-Danlos syndrome, 241
- electron microscopy, 121–2
- electrospray ionisation-tandem mass spectrometry, 118–21
- emergency departments, 29, 31–2, 45–6. *See* Accident and Emergency departments
- emergency services, 28–9, 43
- Emery, J. L., 1–3
- encephalitis, viral, 224–5
- encephalomalacia-like lesions, 256
- encephalopathy, 257–8
- endocardial fibroelastosis, 229
- endocarditis, 102
- enteroviruses, 223–4
- environmental risk factors, 136–40. *See also* sleep environment
- day-care environment, 156
- death scene investigation, 43
- epidemiology of SIDS
Netherlands, 10–11
UK, 153, 159
US, 127–8, 159
- epigenetic changes, 203
- epiglottitis, 219
- epilepsy, 233, 256
- Escherichia coli*, 163, 219
- ethnic/racial differences
bed-sharing, 151
infant sleep position, 136
SIDS rates, 127–8, 159
- ethylene glycol poisoning, 112–13
- eupnoea, fictive, 179–80
- European Convention on Human Rights (ECHR), Article 2, 19–20, 25–7
- evidence
expert, 34
forensic trace, 56
securing and collecting, 29–30, 32–33
valuable physical, 31
- external post-mortem examination, 48–9, 57–8
- eye movement desensitisation and reprocessing (EMDR), 14
- eyes
external examination, 48, 57–8
removal, 65–6
- face
dissection, 64–5
external examination, 48, 57
trauma, 247
- falls, 239, 251
- families. *See also* carers; parents
explaining cause of death to, 87
improving services for, 78–9
initial care, 11–12
perspective on SIDS, 5–8
recurrent SIDS, 3, 129–30
tissue retention and, 22
- family doctors, 11, 16
- family history, 46
- family studies, recurrent infant mortality, 129
- fat deposition, 121–2
- fatty acid oxidation disorders, 212–14
- fetal and infant mortality review (FIMR), 75
- fibroblast cultures, 117–18, 122
- fibromuscular dysplasia, 226
- filicide, 3
- final case discussion, UK, 86–7
- finger-bone fractures, 245
- first responders, 29, 43
- FISH/QF-PCR, 164

- fluorescent in situ hybridisation (FISH) QF-PCR
FMO3 gene polymorphisms, 163
 fontanelles, 48, 52, 57, 252
 forensic autopsy, 24, 48, 55–69. *See also*
 joint forensic/paediatric autopsy
 forensic pathologist, 24, 55
 forensic pathology, 235–59
 fractures, 243–7. *See also* rib fractures;
 skull fractures
 abuse mimics, 246–7
 ageing, 244
 detection, 243–4
 facial, 247
 long bones, 245
 post-mortem radiology, 89–90
 frozen tissue samples, 53, 101
 fructose intolerance, hereditary, 215–16
 fructose-1,6-bisphosphatase deficiency, 216
 funerals, 12

 GABAergic neurons, 192–3
 galactosaemia, 215–16
 Garrow, I., 1
 gas embolism, 90
 gas chromatography mass spectrometry (GC-MS), 111–12
 gasping
 arousal response, 179–80
 medullary 5-HT neurons, 196
 neuromodulation, 182–3
 SIDS pathogenesis, 180
 gastric contents, sampling, 109
 gastroenteritis, 225
 gene sequencing, specific, 165
 general practitioners, 11, 16
 genetic testing
 cardiac arrhythmias, 176
 metabolic disease, 117, 119, 122
 sample collection, 49, 53, 249
 technologies, 164–6
 genetics, 159–66
 brainstem 5-HT changes, 162, 202
 cardiac arrhythmias, 173–4
 recurrent SIDS, 129–30
 SIDS susceptibility, 159–64
 genital injuries, 242–3
 genitourinary system, examination, 51
 genotyping, 165
 gentamicin, intratympanic, 207–9
 gliosis, brainstem, 96
 glucose testing, 108, 121
 glutaryl-CoA dehydrogenase deficiency, 215
 glycogen deposition, 121–2
 glycogen storage diseases, 216
 gram-negative bacilli, 104

 grief, 13
 complicated, 13
 constructive, 13–14
 factors impeding, 15
 individual variations, 13
 models of mourning, 13
 group A streptococcus, 104, 219
 group B streptococcus (GBS), 104, 219, 221
 guilt, 13
 gut microbiome, 203
 Guthrie cards
 ante-mortem newborn screening, 118–19, 123
 blood acylcarnitine profiles, 118–19
 blood amino acid profiles, 119–21
 post-mortem samples, 108, 117, 119

 haematological malignancies, 232–3
 haemophagocytosis, 219, 222
Haemophilus influenzae, 219, 221
 haemosiderosis, 237, 254–5
 hair analysis, 109
 head
 external examination, 48, 57
 internal examination, 51–2
 shape and size abnormalities, 251–2
 head injuries
 cerebral venous and dural sinus thrombosis, 255
 scalp lesions, 250
 skull fractures, 250–1
 subcortical contusional tears, 256–7
 subdural bleeding, 252–5
 healthcare workers
 home visit, 32, 45
 support for parents, 12
 Healthy People (HP) 2020 objective, 76
 hearing defects, 164, 207
 heart
 examination, 50
 microbiology, 100–101
 heart rate
 arousal response, 180
 brainstem circuitry, 180–3
 Haemophilus influenzae,
 hematological malignancies,
 hepatitis, fulminant, 102
 hepatoblastoma, 231–2
 hippocampus, 97
 histiocytoid cardiomyopathy, 228–9
 histochemistry, 121–2
 histology samples
 abdomen and thorax, 49
 central nervous system, 52
 inherited metabolic disease, 118
 staining, 53
 history of SIDS, 1–3
 history, clinical
 home visit, 46

 importance, 2
 neuropathological findings and, 95
 home monitors, 15–16
 home visit, 32, 45–6
 who and when, 45–6
 obtaining history, 46
 homicide, 24
 hookah, 132
 hospital records, 32
 5-HT (5-hydroxytryptamine). *See*
 serotonin
 5-HT receptors
 changes in SIDS, 197–9
 prenatal adverse exposures and,
 199–202
 5-HT transporter (5-HTT)
 changes in SIDS, 197–8, 200
 gene polymorphisms, 162, 202
 5-HT_{1A} receptors
 changes in SIDS, 198–9
 nicotine interactions, 164
 risk factor interactions, 199
 5-HT_{2A} receptors, 164, 197, 199
 5-HT₃ receptors, 197
 Human Rights Act 1998, 19–20, 25
 3-hydroxy-3-methylglutaryl (HMG)
 CoA lyase deficiency, 214
 hypercapnia
 arcuate nucleus function, 162
 arousal response, 179, 189
 brainstem 5-HT mechanisms, 190–1
 inner-ear defects and, 164, 207–8
 hyperflexion/extension injuries, neck,
 66, 258
 hypernatraemia, 111–12
 hypertrophic cardiomyopathy (HCM),
 160, 227, 229
 hypoglycaemia, 121, 213–14
 hypoplastic left heart ventricle, 226
 hypostasis, 57–8
 hypoxia
 arousal response, 178–80, 189
 brainstem 5-HT deficits and, 203
 brainstem 5-HT mechanisms,
 190–1
 diving reflex, 208–9
 habituation to intermittent, 189–90
 inner-ear defects and, 164, 207–8
 nicotine exposure and, 191–2
 post-mortem MRI, 93
 prenatal alcohol exposure and,
 192–3
 hypoxic-ischaemic brain injury, 97,
 180, 259

 identification, formal legal, 34–5, 40
IL-10 gene polymorphisms, 163
 imaging, post-mortem, 89–93
 inherited metabolic disease, 116
 minimally invasive autopsy, 71–3

Index

- imaging, post-mortem (cont.)
 pre-autopsy, 56
 immune dysfunction, 163
 immunodeficiency syndromes, 225
 inborn errors of metabolism. *See*
 inherited metabolic disease
 infant and child death review. *See* child
 death review
 infanticide, 3
 infections, 219–25
 microbiological diagnosis, 100–4
 pathogenic role, 163
 post-mortem skin samples, 117–18
 inferior olivary nucleus, gliosis, 96
 inflammation, breathing control and,
 183
 inflicted injuries. *See* child abuse
 influenza virus, 224
 inherited metabolic disease
 causes of sudden death, 212
 clinical presentation, 212–16
 investigations and analyses, 118–23
 pathophysiology, 163, 211–16
 post-mortem investigation, 116–23
 samples, 116–19
 inner-ear abnormalities, 164, 207–9
 arousal failure, 164, 207–9
 diving reflex and, 208–9
 inquest, 24–7, 35
 adjournment, 24
 Article 2, 25–7
 conclusions, 26–7, 35
 deciding on, 21, 23–4
 history in US, 36–7
 opening the, 25
 inspiratory control
 brainstem circuitry, 180–3
 neuromodulation, 182–3
 interested persons, 25
 interleukin 6 (IL-6), 163
 international normalised ratio (INR),
 108
 interviews
 home visits, 46
 police, 32–4
 intestines, examination, 50
 intra-abdominal injuries, 247–8
 intradural bleeding, 98, 252
 intraosseous needle tracts, 90–2
 investigation of sudden infant death
 advocacy groups in US, 40
 Child fatality review board, 40
 Coroners, 19–27
 determining cause and manner of
 death, 40–1
 final case discussion (UK), 86–7
 first responders, 43
 home visit, 32, 45–6
 Medical examiner, 38–9
 NAME guidelines, 39–40
 Netherlands, 11
 police, 28–35
 role of child death reviews, 78
 UK legal framework, 19–35
 US legal framework, 36–41
 *Investigation Questionnaire for Sudden
 Unexpected Deaths in Infants*,
 56
 joint forensic/paediatric autopsy,
 55–69
 brain and spinal cord removal,
 61–4
 cervical spine excision, 66–7
 decision to perform, 24, 48
 evisceration and special dissections,
 58–69
 external examination, 57–8
 eyes removal, 65–66
 facial dissection, 64–5
 information on circumstances of
 death, 55–6
 information on death scene, 56
 long bone removal, 69
 pre-autopsy radiography, 56
 procuring trace evidence, 56
 rib cage removal, 66–7
 jury, inquest, 25
 juvenile myelomonocytic leukaemia,
 232
 karyotyping, 164
 KCNH2 gene mutations, 160, 172
 KCNQ1 gene mutations, 160, 172,
 176
 Kennedy Report, 55
 ketone body production/utilisation
 defects, 214–15
 3-ketothiolase deficiency, 214–15
 kidneys, 51, 100–1, 118
 Kölliker-Fuse nucleus (KFN), 162
 labial frenulum, lacerations, 57–8,
 247
 lactate, blood, 108
 lactic acidosis, congenital, 212
 laryngeal chemoreflex, 97–8, 196
 laryngeal immune system, 163
 laryngomalacia, 231
 leptomeninges, 255
 lips, abrasions, 57, 247
 liquid chromatography-mass
 spectrometry (LC-MS),
 111–12
 liver
 external examination, 50–1
 samples, 100–1, 107, 109,
 118–23
 liver disease, 213–14
 Local Safeguarding Children Board
 (LSCB), 22–3, 31, 88
 long bones
 fractures, 245
 removal, 69
 long QT syndrome (LQTS), 160–1,
 172–5, 229
 Lullaby Trust, 3
 lungs
 examination, 50
 microbiology, 100–1, 104
 lysergic acid diethylamide, ³H-labelled
 (³H-LSD), 197–8, 200
 macrocephaly/macroencephaly, 252
 magnesium, blood, 108
 magnetic resonance imaging (MRI),
 post-mortem
 findings, 91–3
 inherited metabolic disease, 116
 minimally invasive autopsy, 72–3
 post-mortem changes, 92
 Make a Memory, 12
 malaria, 102
 manner of death, determination, 24,
 26–7, 40–1
 Māori, New Zealand, 138, 152–3
 mattresses, 137–8
 measurements, body, 48
 Medical Certificate as to Cause of
 Death (MCCD), 20
 Medical examiners
 UK, 20
 US, 36–9
 medical records, 31
 medical treatment, deaths related to,
 21
 medium-chain acyl-CoA
 dehydrogenase (MCAD)
 deficiency
 clinical presentation, 163, 214
 post-mortem diagnosis, 118,
 120–1
 medullary 5-HT system., 198–9. *See*
 serotonin system, brainstem
 melanosis, congenital dermal, 241
 meninges, spinal, 258
 meningitis, bacterial, 219, 221–3
 meningococcus (*Neisseria*
 meningitidis), 104, 219, 221–3
 meningoencephalitis, 102
 metabolic bone disease, 246–7
 metabolic disorders, inherited. *See*
 inherited metabolic disease
 metabolites, plasma, 121
 metabolomics, 118
 metaphyseal fractures, 245–7
 metatarsal fractures, 245

- methadone
poisoning, child deaths, 109–10
-using mothers, SIDS risk, 134–5
- methanol poisoning, 112
- microbiology, post-mortem, 100–5
additional samples, 100, 102
interpretation, 100–4
minimal sampling protocol, 53, 101
- microcephaly/microencephaly, 252
- Middleton (R (Middleton) v West Somerset Coroner [2004])*, 20, 26–7
- mini-arousals, 207, 209
- minimally invasive autopsy, 71–3
- minor illness, prior, 2
- mitochondria, abnormal, 122
- mitochondrial 2-methylacetoacetyl-CoA thiolase deficiency, 214–15
- mitochondrial respiratory chain disorders, 122, 212
- mitogenic cardiomyopathy, 228–9
- molecular analysis
inherited metabolic disease, 117, 119, 122
pathogens, 101–2, 104–5
- molecular autopsy. *See* genetic testing
- Mongolian spot, 241
- monoamine oxidase A (MAOA), 162, 197, 202
- mortuary staff, 46
- mourning. *See* grief
- mouth
examination, 48, 57–8
injuries within, 247
- multi-agency approach
Child Death Overview Panels, 87
child death reviews, 76–7
final case discussion, 86–7
infant death investigation, 31
- multiple carboxylase deficiency, 215
- multiple system atrophy, 182
- muscle biopsy, 96, 108, 118–19, 122
- Mycobacterium tuberculosis*, 104
- myelitis, 102
- myocarditis
eosinophilic, 225–6
tissue samples, 102
viral, 223–4
- myositis, diaphragmatic, 230, 232
- National Association of Medical Examiners (NAME), 1, 37, 39–40
- National Center for Fatality Review and Prevention (NCFRP), 77–8, 80, 83
- National Cot Death Study Group (LWW) (Netherlands), 11
- natural deaths, 41
- neck
examination, 48–50
hyperflexion/extension injuries, 66, 258
layered dissection, 58–60
negative feelings/perceptions, 6
neglect, 237–9. *See also* child abuse
contributory, 26–7
pathologists' role, 238
starvation, 248–9
- Neisseria meningitidis* (meningococcus), 104, 219, 221–3
- Netherlands, 9–16
- neurological deterioration, metabolic disease, 213
- neuromodulation, 182–3
- neuromuscular disease, 232
- neuropathology, 95–8, 249–59
- New Zealand, 138, 152–3
- newborn screening
hearing tests, 207
metabolic disorders, 211
stored Guthrie cards, 118–19, 123
- next-generation sequencing (NGS), 165–6, 176
gene panels, 165–6
- nicotine. *See also* smoking
alternative delivery systems, 132
arousal response and, 191–2
metabolism, genetic variations, 163–4
prenatal and postnatal exposure, 131–3
replacement therapy, 133
- NODOK procedure, 11–12
- non-accidental injury. *See* child abuse
- Noonan syndrome, 227, 229
- noradrenergic brainstem nuclei, 182
- Norris, Charles, 37
- NOS1AP* gene variants, 161
- nose, examination, 48
- nucleus ambiguus* (NA), 180
- nucleus tractus solitarius* (nTS), 181–2, 196
- nurses, home visit, 45
- obstetric risk factors, 127
- oesophagus, examination, 50
- oncocyctic cardiomyopathy, 228–9
- opiate misuse, 134–5
- optic nerve sheath haemorrhages, 253–4
old, 254
- orexin, 162
- organ retention, 22, 249–50
- organic acids, urinary, 121
- organic acidurias, 212, 215
- osmolar gap, 108
- osteogenesis imperfecta, 241, 246
- out-of-home settings, 82
- overheating, 137, 139, 162
- overlying
autopsy findings, 236–7
Bowden's studies, 2
substance misuse and, 150
- oxygen-conserving reflexes (OCR), 97–8, 208–9
- pacifiers (dummies), 146–8
breastfeeding and, 143, 146–7
protective effects, 146–7
- paediatric pathologists, specialist, 21–22, 55
- paediatricians
emergency department, 32
home visit, 32, 45–6
- pancreas, examination, 50–1
- parabrachial nucleus, 190–1
- paracetamol, 106
- parafacial respiratory group. *See* retrotrapezoid nucleus
- parasitosis, systemic, 102
- parents. *See also* carers; families
consent to autopsy, 12, 16, 95
cooperation with researchers, 9–11
expectations, 12
explaining cause of death to, 41, 87
follow-up support, 14–15
home visit, 46
initial accounts, 31
initial care, 11–12
interview by police, 32
perspective on SIDS, 5–8
relations with police, 30–1
support and care, 9–16, 46
support groups, 7–8, 12, 40–1
taking statements from, 33–4
- parvovirus B 19, 224
- pathologists
briefing prior to autopsy, 21–2, 55–6
performing the autopsy, 21–2, 24, 48
responsibility in US, 40–1
role in child abuse/neglect, 238
tissue retention, 22
- patterned injuries, 240
- pelvis, examination, 50–1
- Pépi-Pod™, 138, 152–3
- pericardial fluid sampling, 109
- pericardium, 50
- perinatal infections, 102
- peripherally located cortical defects, 256
- peritonitis, bacterial, 102, 222
- periventricular leucomalacia, 256
- pertussis (whooping cough), 219, 221
- petechiae
face, neck and chest, 57–8, 237
intrathoracic, 209, 236
- photographs, 12, 48

Index

- Phox2B* gene polymorphism, 162
 Pierre Robin syndrome, 231
 pillows, 138
 pituitary adenylate-cyclase-activating polypeptide (PACAP), 162
 placenta, 116
 plagiocephaly, 251
 pleura, 49
 pneumococcus. *See Streptococcus pneumoniae*
 pneumonia, 102, 219–21
 pneumonitis, viral, 224
 pneumothorax, 49
 poisoning, 106–14
 agents involved, 109–11
 drug screening tests, 111–12
 iatrogenic, 106
 malicious, 107, 111–13
 numbers of deaths, 109–10
 sampling methods, 107–9
 police, 28–35
 assessing suspicious deaths, 30
 decision on criminal process, 34
 emergency department actions, 31–2
 first actions at death scene, 29–30
 guidance documents, 28
 home visit, 32, 45–6
 initial notification, 28–9
 level of follow-up response, 30
 major enquiry, 32–3
 parents' perspective, 15
 records of carers' accounts, 31
 statements from witnesses, 33–4
 valuable physical evidence, 31
 working with coroner, 21–2, 24, 29, 34–5
 posterior torso, layered dissection, 61
 post-inspiratory complex (PiCo), 180
 post-mortem examination. *See* autopsy
 post mortem computed tomography, 180
 post mortem MRI, 180
 post mortem radiography, 180
 post-traumatic stress disorder/reactions, 14
 potassium (K⁺) channel gene mutations, 172–5
 potassium, blood, 108
 pre-Bötzinger complex (preBötC)
 arousal response, 179–80
 cardiorespiratory coupling, 180
 control of inspiration, 180, 182–3
 pregnancy
 brainstem 5-HT deficits and, 199–202
 drug misuse, 134–5
 maternal smoking, 131–3
 second-hand smoking, 132
 smoking cessation, 131–3
 pre-inquest review (PIR), 25, 35
 premature babies, 95, 189
 prone sleeping position, 136–7
 asphyxia related to, 236–7
 child-care settings, 156
 interaction with smoking, 132
 pathophysiology, 136–7
 soft surfaces, 137
 protective factors. *See also* risk factors
 breastfeeding, 142–3
 pacifiers/dummies, 146
 prothrombin time, 108
 pseudomembranous colitis, 221–2
 pulmonary haemorrhage, 237
 pulmonary haemosiderosis, 237
 pulmonary vascular disease, 230–2
 QT interval, prolongation, 134, 172–5
 quantitative fluorescence-polymerase chain reaction (QF-PCR), 164
 ‘question mark’ incision, 51, 61
 racial differences. *See* ethnic/racial differences
 radiography, post-mortem skeletal, 89
 ageing of fractures, 244
 diagnostic accuracy, 89
 excised ribs, 66–7, 243
 findings, 89–90
 minimally invasive autopsy, 71–2
 pre-autopsy, 56
 ‘Rainbow’ children, 7
 raphe nuclei, 161, 181–3, 190–1
 Red Book, 46
 Registrar of Births and Deaths, 20
 religious objection laws, 41
 research, parental involvement, 9–11
 respiratory chain disorders, 212
 mitochondrial, 122, 212
 respiratory failure, neuromuscular disease, 231–2
 respiratory syncytial virus (RSV), 163, 224
 respiratory tract
 commensal flora, 104
 disorders, 231–2
 respiratory tract infections
 bacterial, 219–21
 Bowden's studies, 1–2
 microbiology, 102, 104
 viral, 163, 224
 retinal haemorrhages, 65, 253
 old, 254
 retroperitoneal injuries, 247–8
 retropharyngeal abscess, 221
 retrotrapezoid nucleus (RTN), 180–1, 191
 rhabdomyomas, cardiac, 229
 rib cage
 examination, 50, 59–61
 removal, 66–8
 rib fractures, 244–5
 abuse mimics, 246
 birth-related, 245
 cardiopulmonary resuscitation, 245
 detection at autopsy, 59–61, 243–4
 inflicted, 244–5
 post-mortem imaging, 89–90
 removal of rib cage, 66–8
 spinal cord removal and, 67
 ribs, histological examination, 67–8
 rickets, 227, 246
 risk factors. *See also* protective factors
 arousal and, 191–3
 bed-sharing, 149–53
 biological, 127–8
 brainstem 5-HT deficits and, 199–203
 child death review-based studies, 81–2
 day-care environment, 156
 drug misuse in pregnancy, 134–5
 Emery's studies, 3
 environmental, 136–40
 information gathering, 46
 interactions between, 131–2, 149–50
 nicotine exposure, 131–3
 role of child death reviews in reducing, 79–80
 triple-risk model. *See* triple-risk hypothesis
 room sharing, 149, 151–2
 RYR2 gene mutations, 161, 175–6
 safe-sleep campaigns. *See also* back-to-sleep campaigns
 Māori initiative, 152–3
 recurrent SIDS after, 130
 Salmonella, 104
 salt poisoning, 111–12
 samples
 ante-mortem, 107, 116
 general post-mortem, 51, 53, 108
 inherited metabolic disease, 116–18
 microbiology. *See* microbiology, post-mortem
 toxicology, 107–9
 scalds, 242
 scalp
 incision, 51, 61
 injuries, 250
 scapular fractures, 245
 scene of infant death. *See* death scene
 SCN5A gene mutations
 Brugada syndrome, 161, 175
 long QT syndrome (LQT3), 160, 172, 175
 secondary depression, 180
 Senior Investigating Officer (SIO), 24, 29–30
 septicaemia/sepsis, 102, 219–20

- Serious Case Review (SCR), 23, 31
 serotonin (5-HT)
 brainstem content, 197–200
 neurochemical mediators in brain, 196–7
 post-mortem serum levels, 204
 serotonin (5-HT) neurons
 immaturity, 198, 203
 numbers and density, 197–8, 200
 serotonin (5-HT) system, brainstem
 abnormalities in SIDS, 182–3, 197–204
 anatomy and function, 195–6
 arousal response, 190–1
 caudal, 96, 195–6
 developmental origin of deficits, 203
 genetic studies, 161–2, 202
 medullary abnormalities, 198–200
 nicotine exposure and, 163–4, 191–2
 postnatal origin of deficits, 203
 risk factor interactions, 199–202
 prenatal adverse exposure, 202
 genetic susceptibility, 199–202
 rostral, 195–6
 serotonin receptors. *See* 5-HT receptors
 serotonin transporter. *See* 5-HT transporter
 sex-ratio statistics, 2, 127
 sexual abuse, 242–3
 shaken baby syndrome, 235, 253–4, *See also* triad
 short QT syndrome (SQTS), 176
 siblings
 impact of SIDS, 7
 risk of SIDS in subsequent, 129–30
 support and care, 12, 14
 witness statements, 33
 side cars (bedside sleepers), 138
 side sleeping position, 136
 siderophages, 237, 241
 siderosis, 254–5
 SIDS. *See* sudden infant death syndrome
 sighs
 arousal response, 178–80
 brainstem circuitry, 179, 182
 neuromodulation, 182–3
 skeletal survey, post-mortem. *See* radiography, post-mortem skeletal
 skeletal trauma, 56, 243–7, *See also* fractures
 skin injuries, 57, 239–43
 skin samples, 51, 108
 collection, 49, 53
 for fibroblast culture, 117–18
 inherited metabolic disease, 119
 skull, 250–2
 deformation, 251–2
 developmental abnormalities, 250
 examination, 52, 61, 63
 opening, 52, 62–3
 skull fractures, 250–1
 birth related, 250
 inspection for, 61, 63
 mimics, 250
 short falls, 251
 sleep devices, 137–8
 sleep enablers (in-bed sleepers), 138
 sleep environment, 136–40 *See also* bed-sharing
 child-care settings, 156
 safe, 140
 unsafe, 41, 81–2, 236
 sleep position, 136–7, 156
 sleep surfaces, 81, 137–8
 firmness, 137
 sleeping sacks, infant, 139
 small-bowel injuries, 248
 smoking. *See also* nicotine
 arousal response and, 191–2
 bans in public places, 132
 bed-sharing and, 132, 149–51
 brainstem 5-HT deficits and, 199–202
 cessation in pregnancy, 131–3
 maternal, 131–3
 outdoors vs indoors, 132
 paternal, 131
 smothering, 236–7
 social care records, 31
 sodium
 blood, 108
 fractional excretion, 112–13
 sodium (Na⁺) channel gene mutations, 173–4 *See also* SCN5A gene mutations
 sofa-sharing, 82, 137, 150–1
 soft-tissue infections, 102
 soft-tissue injuries, 57, 239–43
 spinal cord
 forensic pathology, 259
 injuries, 258
 inspection, 52, 63–4
 removal, 63–4, 67
 spinal cord injury without radiological abnormality (SCIWORA), 258
 spine
 examination, 51–2
 forensic pathology, 258–9
 spleen, 50–1, 100–1
 stains, histological, 53
Staphylococcus aureus, 104, 163, 219
Staphylococcus, coagulase negative, 221–2
 starvation, 248–9
 statements, witness, 33–4
 sternal fractures, 245
 sterols, plasma, 121
 stillbirths, 21
 stomach, 50–1
Streptococcus agalactiae. *See* group B streptococcus
Streptococcus pneumoniae (pneumococcus), 104, 219, 221
Streptococcus, beta-haemolytic, 104, 219
 stressors, exogenous, 180, 183
 subarachnoid bleeding, spinal, 258–9
 subarachnoid spaces, benign enlargement, 256
 subcortical contusional tears, 256–7
 subdural bleeding
 acute, 252–3
 detection, 62–3, 98
 non-accidental head injury and, 253–4
 old, 254–5
 spinal, 258–9
 substance abuse. *See also* alcohol consumption; drug misuse; smoking
 bed-sharing and, 134, 149–50
 child deaths, 109–11
 post-mortem screening, 111–12
 pregnancy, 134–5
 substance P, 98, 182–3
 Sudden Death in the Young Case Registry (SDY-CR), 83
 sudden infant death syndrome (SIDS)
 CDC guidelines, 39
 clinico-pathological diagnosis, 53
 definition, 9, 236
 neuropathology, 95–8
 parents' perspective, 6, 9
 post-infancy, 40
 recent trends, 151–3
 recurrent, 3, 129–30
 use of term, 1, 41
 Sudden Infant Death Syndrome Act 1974, 40
 sudden unexpected death in infancy (SUDI/SUID)
 causes, 219–33
 child death review process, 75–83
 definition, 236
 forensic pathology, 235–59
 improving reporting, 80–3
 investigation. *See* investigation of sudden infant death
 microbiology, 100–1
 minimally invasive autopsy, 71–3
 police involvement, 28–30
 prevention, 79–80
 recent trends, 151–3
 standard questionnaires, 39, 56
 trends by cause, 127–8
 use of term, 9, 41, 236

Index

- Sudden Unexpected Infant Death Case Registry (SUID-CR), 82–3
- sudden unexplained death in childhood (SUDC), 40
- Sudden Unexplained Infant Death Investigation form, 39
- SUDI. *See* sudden unexpected death in infancy
- suffocation, accidental, 2, 78, 81–2
- SUID. *See* sudden unexpected death in infancy
- supine sleeping position, 136
- support, 9–16, 46
- support groups, parent, 7–8, 12, 40–1
- suspicious deaths
- autopsy findings, 30, 235–59
 - compassion for parents, 30–1
 - coroner's perspective, 24
 - development of a major enquiry, 32–3
 - differential diagnosis, 238–9
 - forensic autopsy, 24, 48, 55
 - parents' responses, 15
 - police response, 29–31
 - referral to police, 29
 - risk factors and signs, 30, 238–9
- suture lines, cranial, 250
- swabs, microbiological, 101–2
- swaddling, 139–40
- Swinscow, D., 1–3
- synovial fluid samples, 109
- syphilis, congenital, 102
- temperature
- ambient room, 139, 180
 - recordings, death scene, 30
- thermoregulation, 162, 196
- thoracic injuries, 244–5, 248
- thorax. *See* chest
- thymus, 49–50
- tissue retention, 22, 249–50
- tissue samples
- general, 53, 108
 - inherited metabolic disease, 116, 118, 121–3
 - timing of collection, 116–17
 - toxicology, 107, 109
- tobacco use. *See also* nicotine; smoking
- alternative products, 132
 - prenatal and postnatal exposure, 131–3
- tonsillitis, clostridial, 221
- total anomalous pulmonary venous return, 226
- Touche (R v Inner N. London Coroner ex p Touche [2001])*, 21
- toxicological analysis, 106–14
- chain-of-custody procedures, 106–7
 - sampling, 107–9
 - screening tests, 111–12
 - service provision (UK), 110–11
 - suspicious deaths, 249
- toxoplasmosis, 102
- traceless impact, 253–4
- tracheomalacia, 231
- transposition of the great arteries, 226
- triad (encephalopathy, subdural and retinal haemorrhages), 30, 235, 253–4
- cerebral venous and dural sinus thrombosis, 255
 - delayed death and, 255
 - subdural bleeding, 253
- trigemino-cardiac reflex, 97–8
- triple-risk hypothesis, 159, 178
- as basis for investigation, 43
 - Bowden's research, 1
 - brainstem 5-HT deficits, 202–3
 - prone sleeping position and, 136–7
 - role of inflammation, 183
- TRPM4* (transient potential melastatin 4) gene mutations, 161
- truncus arteriosus, 226
- tryptophan, 191–2, 197
- tryptophan hydroxylase 2 (TPH2), 197
- brainstem levels, 190, 197–200
 - risk factor interactions, 199
- tuberculosis, 102
- tuberous sclerosis, 229
- tumour necrosis factor- α (TNF- α), 163
- tumours, 229, 232–3
- ultrastructure, 121–2
- unascertained (cause of death), 53
- undetermined (cause of death)
- child death review-based studies, 81–2
 - diagnostic shift to, 6, 78
 - parents' perspective, 6
 - recommended use, 1, 41, 53
- unexpected deaths, 20, 28
- United Kingdom (UK)
- bed-sharing, 152
 - Child Death Overview Panel. *See* Child Death Overview Panel
 - final case discussion, 86–7
 - home visit, 32, 45–6
 - legal investigation framework, 19–35
 - recent trends in SIDS/SUDI, 152
- United States (US)
- child death reviews, 75–83
 - coroners, 19, 36–9
 - legal investigation framework, 36–41
 - recent trends in SIDS/SUDI, 151–2
 - SUID Case Registry, 82–3
- unnatural deaths, 20–1, 28, 41
- urea cycle disorders, 119, 121, 212, 215
- urine samples, 51, 108–9
- collection, 51
 - drug screening, 111–12
 - inherited metabolic disease, 117, 119, 121
- ventilation, room, 139
- verdict, inquest, 26
- vertebrae, 258
- very-long-chain fatty acids (VLCFAs), 121
- virology
- molecular analyses, 104–5
 - samples, 53, 101–2
- viral infections, 163, 223–5
- visceral trauma, 247–8
- vitamin C deficiency, 246
- vitamin D deficiency, 227, 229, 246–7
- vitreous humour samples, 107, 109, 117, 121
- wahakura, 138, 152–3
- water, fractional excretion, 112–13
- Waterhouse-Friedrichson syndrome, 221, 223
- Waterlow classification, 249
- wedding, 237
- Werne, J., 1
- whole-exome sequencing (WES), 166
- whole-genome sequencing (WGS), 166
- whooping cough (pertussis), 219, 221
- Williams syndrome, 232
- witnesses, 25, 33–4
- Worcestershire Coroner* case (2013), 23
- wormian bones, 250