

Index

- ACE test, 15
 ADAS-Cog test, 15
 adjustment disorder with depressed mood (DSM-5), 62
 age-based geriatric services, 118–119
 agoraphobias, 90
 AIDS (Acquired Immune Deficiency Syndrome), 3, 31
 alcohol abuse
 alcohol metabolism, 100
 health effects of, 100–101
 late onset alcoholism, 101
 management, 101–102
 outcomes, 102
 prevalence of, 99–100
 alcohol abuse dementia, 31
 Alzheimer's Association (USA), 2
 Alzheimer's disease (AD)
 clinical research in, 115
 mild cognitive impairment (MCI) and, 32
 mixed vascular and Alzheimer's dementia, 30
 pathophysiology of, 29
 Alzheimer's Disease Assessment Scale Cognitive subscale (ADAS-Cog), 10, 15, 35
 Alzheimer's Disease International (ADI), 2
 Alzheimer's Society, 2
 American Association of Geriatric Psychiatry (AAGP), 115
 American Psychiatric Association, 26
 Angiotensin-converting enzyme (ACE), 73
 anticholinergic drugs, 53, 81
 as cause of psychotic symptoms, 81
 anticonvulsants, 47
 antidepressants, 47
 antipsychotic drugs
 for BPSD, 46–47
 for schizophrenia, 79–80
 anxiety
 drug treatment, 93–94
 generalised anxiety disorder (GAD), 90
 apolipoprotein E, 29
 Arie, Tom, 115
 Arie model, 105
 aripiprazole, 46, 48, 65
 assessment
 additional tests, 17
 carer/informant involvement, 6–7
 cognitive assessment, 9–16
 cognitive assessment and capacity, 16
 history taking, 7–8
 location, 5–6
 mental state examination, 8–9
 neuroimaging, 17–18
 physical examination, 16
 specialist referral, 18
 Automatic Thought Questionnaire, 92
 barbiturate dependency, 103
 Beck Depression Inventory, 92
 behavioural and psychological symptoms of dementia (BPSD). *See also* dementia
 assessment of, 44
 natural history of, 43
 origin of, 42
 summary of, 42–43
 behavioural and psychological symptoms of dementia (BPSD) management
 non-pharmacological, 45–46
 pharmacological, 46–48
 principles, 44
 benzodiazepine dependency, 103
 biomarkers, 117
 bipolar disorders
 age of onset, 70
 and cerebrovascular disorders, 71
 classification (DSM-5), 69
 classification (ICD-11), 67–68
 classification difficulties, 67
 clinical course, 71–72
 determinants in later life, 72
 epidemiology of, 68–69
 and mania, 67–68
 neuroimaging, 71
 neurological comorbidity, 71
 in older populations, 67
 outcomes, 72
 pathophysiology of, 71
 bipolar disorders treatment
 carer/informant involvement, 73
 lithium, 73
 bipolar type 1 disorder (DSM-5), 62
 bipolar type I disorder (ICD-11), 68
 bipolar type II disorder (ICD-11), 68
 Cambridge Cognitive Examination (CAMCOG), 10, 15, 35
 Canadian Association of Geriatric Psychiatry, 115
 carbamazepine, 47, 73
 carer support, dementia, 40
 carer/informant involvement
 accuracy of statements, 7
 in assessment of old age psychiatric disorders, 6–7
 bipolar disorders, 73
 for depression, 66
 residential care, 108–110
 respite care, 110
 CARITAS principles, 106–107
 cerebrovascular adverse events (CVAEs), 47
 cholinesterase inhibitors
 for BPSD, 47
 for dementia, 38–39
 chronic fatigue, 95
 cimetidine, 55

- clinical neuroscience and the future of psychiatry of old age (POA)
 biomarkers, 117
 division between neurology and psychiatry, 115–116
 environmental and biological factor
 understanding, 116–117
 future training in, 117–118
 research milestones, 115
- Clock Drawing Test (CDT), 11
- clomipramine, 94
- CLOX test, 11
- cognitive assessment
 head turning sign, 10
 importance of, 9
- cognitive assessment tests
 ACE, 15
 ADAS-Cog, 15
 CAMCOG, 15
 Clock Drawing Test (CDT), 11
 CLOX test, 11
 executive function, 11–15
 mini-cog, 11, 13
 Mini-Mental State Examination (MMSE), 10–11, 51
 Montreal Cognitive Assessment (MoCA) test, 15
 summary of, 10
- cognitive assessment tests and capacity, 16
- cognitive behaviour therapy (CBT), 91–92, 94
- cognitive screening during assessment, 8–9
- Cohen Mansfield Agitation Inventory (CMAI), 44
- comorbid medical conditions management, 37
- Confusion Assessment Method (CAM), 51
- Consortium to Establish a Registry for Alzheimer's Disease (CERAD), 11
- Creutzfeldt Jakob disease, 18
- Cummings, Jeffrey, 117
- cytokines, 53
- delayed recall screening instrument, 13
- delirium
 clinical features, 50–51
 differential diagnosis, 51–52
 epidemiology of, 54
 history taking (patient), 20–25
 hyperactive, 51, 52
 hypoactive, 51, 52
 and late onset psychosis, 80–81
 medical comorbidity, 21–22
 neuropathophysiology, 52–53
 as one of 3 Ds, 20
 prevention, 58
 prognosis, 55–56
 risk factors, 54–55
- delirium management
 behaviour control, 56–57
 complication prevention, 57
 effective diagnostic assessment, 56
 rehabilitation, 57–58
- Delirium Rating Scale (DRS), 51
- dementia. *See also* behavioural and psychological symptoms of dementia
 defined, 26
 and depression, 60–61
 epidemiology of, 32–33
 history taking (patient), 20–25
 increase in prevalence in future, 2–3
 and late onset psychosis, 81
 medical comorbidity, 21–22
 as one of 3 Ds, 20
- dementia assessment
 ECG, 36
 full blood examination, 36
 medical, 35
 neuroimaging, 36
 neuropsychological testing, 35
 patient/informant history, 34–35
 specialised tests, 36
- dementia causes
 Alzheimer's disease (AD), 29
 associated with alcohol abuse, 31
 dementia with Lewy bodies (DLB), 30
 mixed vascular and Alzheimer's, 30
 summary of, 28
 vascular, 29
- dementia diagnostic criteria
 DSM-5, 27
 ICD-11, 27
- dementia management drug treatments
 cholinesterase inhibitors, 38–39
 memantine, 39
 vascular dementia, 39
- dementia medical management
 comorbid medical conditions, 37
 impacting diagnosis, 36–37
 outcomes, 40
- dementia praecox, 76
- Dementia Praecox and Paraphrenia, 76
- dementia social management
 carer support, 40
 community support, 39–40
- dementia symptoms
 mild, 33
 moderate, 33–34
 severe, 34
- dementia with Lewy bodies (DLB), 30
- depression. *See also* neurotic disorders
 classification (DSM-5), 61–62
 classification (ICD-11), 60–61
 and cognitive impairment, 61
 and dementia, 60–61
 due to a general medical condition (DSM-5), 61–62
 epidemiology of, 64–65
 history taking (patient), 20–25
 and late onset psychosis, 81
 medical comorbidity, 21–22
 as one of 3 Ds, 20
 pathophysiology of, 64
 and suicide, 63–64
 vascular depression hypothesis, 63
 vascular disease and, 62
- depression treatment
 carer/informant involvement, 66
 efficacy-effectiveness gap, 66
 electroconvulsive therapy (ECT), 66
 for mild depression, 66

- monoamine oxidase inhibitors (MAOIs), 66
 outcomes, 66–67
 second line, 65
 SSRIs and older adults, 65
 TCA, 65
 transcranial magnetic stimulation (TMS), 66
 depressive episode (ICD-11), 61
 detoxification for alcohol abuse, 101–102
 Detre, Thomas, 117
 differential diagnosis
 common diagnostic pitfalls, 24
 depression, delirium and dementia, 20
 family history, 22–24
 medical comorbidity, 21–22
 mental state examination, 24–25
 mild cognitive impairment (MCI), 32
 psychotic symptoms, 25
 digoxin, 55, 81
 disinhibition syndrome, 68
 donepezil, 38–39
 double jeopardy of cognitive impairment, 77
 Dysfunctional Attitude Scale, 92
 dysthymic disorder (DSM-5), 62
 dysthymic disorder (ICD-11), 61
 early onset schizophrenia, 77
 efficacy-effectiveness gap, 66
 electroconvulsive therapy (ECT), 66
 executive function tests
 fluency tests, 14–15
 frontal assessment battery (FAB), 11
 extra-pyramidal side effects, 30, 46
 family. *See* carer/informant involvement
 family history, 22–24
 fibromyalgia, 95
 fluency tests, 15
 4AT, 51
 frontal assessment battery (FAB), 11
 frontotemporal dementia (FTD), 29, 30–31
 future of psychiatry of old age (POA)
 clinical neuroscience, 115–118
 future of geriatric services, 118–119
 historical influences, 114–115
 two major issues in, 114
 galantamine, 38–39
 generalised anxiety disorder (GAD), 90, 94
 geriatric medicine, 1–2
 Gilles de la Tourette syndrome, 115
 glucocorticoid cascade hypothesis, 30–31
 governmental health policy towards old age, 2
 GPCOG screening instrument, 10
 haloperidol, 46, 57
 head turning sign, 10
 history taking (patient)
 3 Ds, 20–25
 age of psychiatric onset, 7
 importance of, 7
 interaction of common medical conditions and psychiatric disorders, 7
 hyperactive delirium, 51, 52
 hypoactive delirium, 51, 52
 hypomanic episode (DSM-5), 69
 Insel, T.R., 116–117
 insomnia. *See* sleep disorders
 International Consensus Model on Organisations of Care
 best practices, 106
 CARITAS principles, 106–107
 International Psychogeriatric Association (IPA), 42, 115
 Kandel, Eric, 116
 lamotrigine, 47, 73
 late onset alcoholism, 101
 late onset psychosis
 history of, 76–77
 pharmacological causes of, 81–82
 late onset psychosis, differential diagnosis
 delirium and, 80–81
 dementia, 81
 depression, 81
 neurological comorbidity, 81
 late onset schizophrenia (LOS)
 clinical features, 77–78
 cognitive changes during, 79
 defined, 76
 epidemiology of, 78
 neuroimaging, 78
 late onset schizophrenia (LOS) management
 pharmacological, 79–80
 physical, 79
 psychosocial, 80
 residential care, 80
 Lewy, Friedrich, 30
 lithium, 73
 lorazepam, 56
 major depressive disorder (DSM-5), 62
 mania
 and bipolar disorders, 67–68
 and cerebrovascular disorders, 70
 manic episode (DSM-5), 69
 medical comorbidity (3 Ds), 21–22
 memantine, 39
 mental state examination
 3 Ds, 24–25
 cognitive screening, 8–9
 form of, 8
 white roots sign, 9
 mild cognitive impairment (MCI) differential diagnosis, 32
 mini-cog test, 11, 13
 Mini-Mental State Examination (MMSE), 10–11, 51
 mirtazapine, 47
 monoamine oxidase inhibitors (MAOIs), 66
 Montreal Cognitive Assessment (MoCA) test, 15

- mood disorders
 bipolar disorders, 67–74
 depression, 60–67
 Motor Neuron Disease (MND), 31
- National Health Service (NHS), 1, 115
- needs-based geriatric services, 118–119
- neuroimaging assessment tests, 17–18
- Neuropsychiatric Inventory (NPI), 44
- neuropsychological testing, dementia assessment, 35
- neurotic disorders. *See also* anxiety; depression, characterisation of, 83
- neurotic disorders aetiology
 care experience of, 83–85
 destabilisation factors, 85–86
 restitution factors, 86
 vulnerability factors, 85
- neurotic disorders
 classification
 DSM-5, 83
 ICD-11, 83
- neurotic disorders differential
 diagnosis
 delirium, 89
 dementia, 88–89
 depression, 88
 personality disorders, 97
 physical illness, 89
 schizophrenia, 89
 sleep disorders, 89
- neurotic disorders
 epidemiology
 behavioural problems, 88
 clinical features, 86
 psychological symptoms, 86–87
 risk factors, 86
 somatic symptoms, 87–88
- neurotic disorders
 management
 drug treatment, 93
 psychological approaches, 91–93
- neurotic disorders, types of
 generalised anxiety disorder (GAD), 90
 obsessive compulsive disorder (OCD), 90
 panic disorder, 90
 phobias, 90
 post-traumatic stress disorder (PTSD), 90–91
 somatic disorders, 91
 non-steroidal anti-inflammatory drugs (NSAID), 73
- obesity, 2, 30
- O'Brien, John, 117–118
- obsessive compulsive disorder (OCD)
 clinical features, 87
 drug treatment, 94
 general overview of, 90
- oxazepam, 56
- oxcarbazepine, 73
- panic disorder, 90, 94
- paraphrenia, 76
- Parkinson's disease dementia (PDD), 30, 39
- paroxetine, 94, 95
- persistent depressive disorder (DSM-5), 62
- personality disorders
 characterisation of, 95–96
 lack of studies in, 97
 management, 98
 and neurosis, 97
 in old age, 95
 and organic brain disorders, 97
- personality disorders
 classification
 DSM-5, 96
 ICD-11, 96
- phenelzine, 65, 66
- phobias, 90, 94
- physical examination, 16
- Pitt, Brice, 67, 115
- Post, Felix, 9
- post-traumatic stress disorder (PTSD), 90–91, 95
- prednisolone, 55
- professional services
 advocacy, 111
 community-based
 assessment, 110
 creative innovations, 112
 history of, 105
- International Consensus Model on Organisations of Care, 105–107
- leisure and spiritual needs, 111
- medical care, 110
- multidisciplinary team, 107–108
- public education, 111
- residential care, 108–110
- respite care, 110
- surround them with care principle, 111–112
- psychiatry of old age (POA)
 defined, 1
 future of, 2–3
 geriatric medicine, 1–2
 governmental health policy, 2
 history of, 1
- psychiatry of old age (POA)
 assessment
 additional tests, 17
 carer/informant involvement, 6–7
 cognitive assessment, 9–16
 cognitive assessment and capacity, 16
 history taking, 7–8
 location, 5–6
 mental state
 examination, 8–9
 neuroimaging, 17–18
 physical examination, 16
 specialist referral, 18
- psychiatry of old age (POA)
 differential diagnosis
 common diagnostic pitfalls, 24
 depression, delirium and dementia, 20
 family history, 22–24
 medical comorbidity, 21–22
 mental state examination, 24–25
 mild cognitive impairment (MCI), 32
 psychotic symptoms, 25
- psychiatry of old age (POA)
 future
 clinical neuroscience, 115–118
 future of geriatric services, 118–119
 historical influences, 114–115
 two major issues in, 114
- psychiatry of old age (POA)
 mood disorders
 bipolar disorders, 67–74
 depression, 60–67

psychiatry of old age (POA) <ul style="list-style-type: none">professional servicesadvocacy, 111community-based<ul style="list-style-type: none">assessment, 110creative innovations, 112history of, 105International Consensus Model on Organisations of Care, 105–107leisure and spiritual needs, 111medical care, 110multidisciplinary team, 107–108public education, 111residential care, 108–110respite care, 110surround them with care<ul style="list-style-type: none">principle, 111–112	residential care, 108–110 <ul style="list-style-type: none">risperidone, 46, 73rivastigmine, 38–39Roth, Martin, 1, 76, 114Rowland Universal Dementia Assessment Scale (RUDAS), 10Royal College of Psychiatrists, 115selective serotonin reuptake inhibitors (SSRIs), 65serotonin and noradrenergic reuptake inhibitors (SNRIs), 65sertraline, 47, 95Shulman, Ken, 114sleep disorders, 89somatic disorders<ul style="list-style-type: none">drug treatment, 95symptoms of, 87–88as type of neurotic disorder, 91substance abuse<ul style="list-style-type: none">alcohol abuse, 99–102illicit substances, 102–103prescribed medicine iatrogenesis, 103suicide<ul style="list-style-type: none">alcohol abuse and, 101depression and, 63–64	surround them with care <ul style="list-style-type: none">principle, 111–112 syphilis, 17, 31, 36
psychiatry of old age (POA) <ul style="list-style-type: none">substance abuse<ul style="list-style-type: none">alcohol abuse, 99–102illicit substances, 102–103prescribed medicine iatrogenesis, 103psychodynamic therapy, 92–93		3 Ds <ul style="list-style-type: none">common diagnostic pitfalls, 24defined, 20history taking (patient), 20–25medical comorbidity, 21–22 Tourette’s syndrome, 115
		Trail Making Test, 10, 15
		tricyclic antidepressants (TCA), 47, 81
		vascular dementia <ul style="list-style-type: none">causes of, 29drug treatment, 39mixed with Alzheimers, 30
		vascular depression <ul style="list-style-type: none">hypothesis, 63
		venlafaxine, 94
		very-late-onset schizophrenia (VLOSP), 77
		Wertheimer, Jean, 106
		white roots sign, 9
		Whitehead, Tony, 1
		World Psychiatric Association, 2, 105