

Index

Numbers: **bold**=table, *italics*=figure.

- Abbey pain scale, 130
- abbreviated mental test score (AMTS), 114
- abuse and neglect, 82
 - barriers to disclosure, 89
 - carers, 93–4
 - crime scenes, 88
 - disclosure (what to do), **91**
 - indicators and examples, **84**
 - information-sharing, 93
 - perpetrators, 88
 - professional barriers, 90
 - referral outcomes, 92–3
 - referrals (what to expect), 91–2
 - reporting, 92
 - risk assessment decision tree, 92
 - types, 83–8
 - victimhood (factors), 88–9
- acceptance and commitment therapy (ACT), 194
- accident and emergency, 76–7, 98, **192**, 248
- acetylcholinesterase (AChE)
 - inhibitors, 116, 132, 133, 272, 280
 - side effects, 116
- activities of daily living, 15, 283, 286, 287
 - types ('basic' versus 'instrumental'), 305
- activities of daily living (assessment), 305–7
 - Bayer-ADLS, 306
 - Bristol-ADLS, 306
 - CSADL, 306
 - DAD, 307
 - DS, 306
- acute care settings, 25, 29, 39, 114, 150, 200–17
 - anxiety, 212
- acute elderly care teams, 287
- acute hospital admission, 171, 173, 174
 - psychological sequels, 12–13
- acute hospital settings, 164, 166, 169, 170, 298
- acute hospitals, 38, 82, 125, 127, 128, 134, 144, 150, 226, 294
- beds, 26
- delirium (incidence), 140
- delirium management (liaison old-age psychiatry), 140–52
- delirium-screening, 145
- health care professionals, 110
- mental health disorders (influence of COVID-19), 32–4
- mental illness in older people (epidemiology), 24–34
- Parkinson's disease and related disorders, 266–81
- privacy and dignity, 59–68
- psychiatric disorders (prevalence), 32–4
- acute kidney injury, 174, 247, 248, 251, 256, 260
 - postoperative (hip patients), 257
- acute myocardial infarction, 256, 258
 - BPSD, 124–39
 - fear and anxiety, 200–17
- acute stress reaction
 - diagnostic criteria, 204–17
 - exclusion criteria, 204
- acute trusts, 292, 293, 294, 296
- adaptive information processing (AIP) system, 213
- Addenbrooke's cognitive examination III (ACE-III), 115
- cognitive screening tool, 301–2
- languages, 302
- adjustment disorders, 30, 158, 161, 162
- adjustment reaction:
 - diagnostic criteria, 204
- Admiral Nurses, 106
- Admission or Assessment Unit, 98
- adult multi-agency safeguarding referral form, 92
- Adult Support and Protection (Scotland) Act (2007), 82
- adult-safeguarding, 82–95, *See also* abuse and neglect
 - abuse and neglect (types), 83–8
 - definition, 82
 - principles and proactive interventions, 83
 - statutory responsibilities, 82
- advance care planning, 54, 98, 284
- advance decisions, 75
- advocates, 75–6
- affective disorder, 223
- ageing, 9, 186, 214, 239
 - barriers to communication, 49–50
 - negative stereotypes (internalisation), 186
 - normal biological and social process, 24
 - successful (Rowe-Kahn model), 10
- ageing population, 2–3, 5, 59, 246, 289
 - statistics, 26
- ageing process, 285–6
- ageism, 48, 50, 186, 194, 225, 235, 235, 236, 239, 286
 - 'age discrimination', 59
- aggression, 125, 126, 127, 132, 146
- agitation, 126, 127, 128, 132, 143, 146, 178, 190, 201, 276
- akinesia, **269**
- alcohol, 172, **184**
 - COVID-19 pandemic, 32
 - dementia risk factor, 113

- alcohol misuse, 28, 30, 187,
 208, *See also* drug and
 alcohol misuse
 acute presentations, 236–8
 CAGE questionnaire, 235
 community and hospital
 settings, 29
 definition (UK), 233
 prevalence, 31
 secondary psychosis in later
 life, 223
 alcohol withdrawal, 215, 236–8
 treatment, 238
 alcohol-dependence
 syndrome: diagnostic
 criteria, 238
 alcohol-related brain damage,
 223, 234, 238, 240–1
 alcohol-related dementia,
 241
 amnesic syndromes, 240–1
 frontal lobe damage, 241
 alcohol-related dementia, 241
 diagnostic criteria, 241
 alcohol-related seizures, 243
 alien limb, 278, 278
 Alzheimer's dementia, 67, 111,
 115, 116, 126, 127, 133,
 159, 280, 301
 BEHAVE-AD, 307
 CERAD-BRSD, 308
 DAD, 307
 dependence scale (DS), 306
 drugs, 116
 early onset, 112
 late onset, 113
 AMHPs. *See* approved mental
 health professionals
 amisulpride, 227, 229
 amitriptyline, 167, 234
 amnesic syndromes, 240–1
 amphetamines, 234, 240, 242
 anaemia, 248, 256
 older hip-fracture patients,
 250
 postoperative (hip patients),
 257
 predictor of increased
 mortality, 257
 analgesia, 259
 analgesic ladder, 249
 analgesics, 127, 249, 259
 analytical psychotherapy, 210
 angiotensin converting
 enzyme (ACE) inhibitors,
 252, 257
 anosmia: definition, 268
 antecedent behaviour
 consequence (ABC)
 charts, 129, 130
 anterograde amnesia, 215, 241
 antibiotics, 235, 250
 anticholinergic side effects,
 114, 125, 133, 146, 216,
 259, 276
 anticoagulants, 248, 284
 hip patients, 253
 anticonvulsants, 133, 176, 217,
 221, 223
 antidepressant therapy, 127,
 132, 133, 160, 162, 166,
 176, 193, 217, 221, 223,
 259, 277
 absorption, 167
 anxiety treatment, 216–17
 choice, 167
 commonly-prescribed
 (alternative forms), 166
 prescribing considerations,
 216
 pros and cons, 168
 suicide-prevention role, 188
 tricyclic, 133
 antihistamines, 178, 208, 259
 antiplatelet agents, 248, 252
 antiplatelet treatments: hip
 patients, 252–3
 antipsychotic drugs, 125, 132,
 134, 146, 148, 149, 169,
 178, 225, 227, 229, 277,
 280
 atypical, 217, 229, 281
 choice (late-onset
 psychosis), 229–30
 as treatment adjuvants, 168
 use in delirium, 149
 anxiety disorders, 12, 30, 113,
 125, 126, 127, 129, 133,
 172–3, 276
 acute settings, 150, 200–17
 comorbidity, 208
 degree of severity (basis for
 decision-making), 209
 features, 202
 HADS, 304–5
 management, 208–9
 negative beliefs (types), 210
 Parkinson's disease patients,
 277
 pharmacological treatment,
 215–17
 prevalence, 29, 31
 prevalence in hospital
 setting, 208
 psychosocial treatments,
 209–15
 reactions (types), 202–8
 resistance to treatment
 (factors), 201
 symptoms, 172
 anxiety rating scales, 209
 anxiolytics, 193
 apathy, 126, 127, 133
 apixaban, 253
 apomorphine, 255, 272
 side-effects, 256
 approved mental health
 professions: LPOP,
 292–3
 ARBD. *See* alcohol-related
 brain damage
 ARD. *See* alcohol-related
 dementia
 aripiprazole, 168, 178, 227
 aromatherapy, 48, 117, 131
 arteriosclerotic Parkinsonism,
 280
 arthritis, 105, 127, 234
 artificial feeding, 169
 aspirin, 168, 252, 288
Assessment of Mental Capacity
 (Law Society and BMA,
 2015 edition), 81
 assessment tools for older
 adults: practical
 guidelines, 300
 assisted living facilities, 117
 Association of Anaesthetists
 of Great Britain and
 Ireland (AAGBI), 248,
 250, 253
 atherosclerosis, 113
 atherosclerotic vascular
 disease, 113
 ATLAS study, 229, 232
 atrial fibrillation, 253, 257
 hip patients, 252
 attachment, 67
 attention-deficit hyperactivity
 disorder (ADHD), 172
 audiology, 221
 AUDIT (alcohol use disorders
 identification test), 236,
 237
 autonomic nervous system,
 202, 267
 autonomy, 60, 61, 62, 64, 65,
 67, 189, 195

- basal ganglia, 266, 267, 281
 basal insulin, 254, 255
 Bayer activities of daily living scale (Bayer-ADLS), 306
 behavioural and psychological symptoms of dementia, 30, 113, 124–39
 assessment, 128–30
 challenges presented to acute settings, 125
 classification, 125–6
 coinage of term (1996), 124
 definition, 124
 behavioural and psychological symptoms of dementia (assessment), 307–8
 BEHAVE-AD, 307–8
 CERAD-BRSD, 308
 NPI, 307
 behavioural and psychological symptoms of dementia (causes and contributors), 126–8
 non-patient (external) factors, 128
 patient (internal) factors, 126–8
 behavioural and psychological symptoms of dementia (management), 130–4
 guiding principle (prevention better than cure), 130
 non-pharmacological strategies, 130–1
 behavioural and psychological symptoms of dementia (pharmacological strategies), 131–4
 aggression and agitation, 132
 mood disorders and anxiety symptoms, 133
 psychotic symptoms, 133–4
 sleep disturbance, 133
 toolkit, 134
 behavioural pathology in Alzheimer's disease (BEHAVE-AD), 129, 307–8
 behavioural syndromes: prevalence rates (ED patients, UK), 28
 being well in oneself: biomedical definition, 10
 benzodiazepines, 30, 31, 127, 133, 146, 148, 160, 178, 203, 208, 216, 233, 234, 238, 239, 259, 276
 alcohol withdrawal, 215
 anxiety treatment, 215
 key to appropriate use, 215
 side-effects, 215
 withdrawal symptoms, 240
 bereavement, 183, 229
 aftermath of suicide, 196
 best interests, 71–3, 75, 76, 77, 78, 80, 100, 105, 106, 149, 169, 229, 242
 beta-blockers, 217, 252
 biographical details, 67
 biopsychosocial care plans, 229
 bipolar affective disorder, 30, 127, 174, 176, 178
 prevalence, 29, 31
 bipolar disorder, 170, 174
 Black and minority ethnic people, 44, 50
 use of interpreters, 51
 blood brain barrier, 160, 271
 blood glucose, 254, 255, 256
 blood tests, 115, 284
 blood transfusion, 257
 body language, 48, 56
 Bolam test, 73
 Bournemouth case (2004), 77
 BPSD. *See* behavioural and psychological symptoms of dementia
 bradyarrhythmias, 256
 bradykinesia, 268
 definition, 266, 269
 brain, 247, 268, 303
 deafferentiation process, 127
 brain damage, 233, *See also* alcohol-related brain damage
 breaking bad news, 49
 brief assessment schedule
 assessment cards (BASAC), 163
 brief assessment schedule
 depression cards (BASDEC), 304
 brief confusion assessment method, 144
 Bristol activities of daily living scale (Bristol-ADLS), 306
 test-retest reliability, 306
 buspirone, 133, 169, 216
 side-effects, 216
 Butterfly Scheme, 55, 101, 102
 CAGE questionnaire, 235
 cancer, 13, 49, 76, 105, 162, 184
 cancer care, 62
 cannabis, 223, 233, 234, 239–40, 242
 changes in strength, 239
 cannabis psychosis: presentations to liaison service, 228–9
 capacity to consent, 77, 78, 79, 80, 178, 228
 capacity: common law definition, 70
 carbidopa, 271, 272
 cardiovascular disease, 113, 201
 care
 holistic models, 25
 versus cure, 3
 Care Act (England, 2014), 82, 86, 90
 people in positions of trust, 90
 Section 42 enquiries, 92, 95
 Section 45 (supply of information), 93
 care and welfare, 73, 74, 76
 care culture
 developing positive, 18–20
 individualised approach, 18
 value base, 18
 care homes, 5, 76, 77, 78, 96, 97, 103, 104, 134, 287
 care plans, 55, 105
 Care Quality Commission, 27, 71, 80, 101, 104
 care triangle, 97
 caregivers
 psycho-education, 130
 stress, 307
 CAREING acronym, 18–19
 carer groups, 102
 carers (informal), 55, 61, 64, 93–4, 118, 286, 290
 abuse (either as agent or victim), 106
 definition (Carers Act 2014), 97
 essential requirements, 5
 impact of COVID-19 pandemic, 106–7

- carers (informal) (cont.)
 needs, 96–109
 needs before patient's admission to hospital, 98
 needs during and after patient's transition from hospital, 104–5
 needs during patient's time in hospital, 98–104
 own health, 105–6
 patient's transition into hospital, 98
 rights, 97
 stress and loss of personal freedom, 105
 stress mitigation, 106
 time devoted and equivalent economic value, 96
- Carers Act (2014)
 carers' rights, 97
- cathcol-o-methyltransferase (COMT) inhibitors, 255, 271, 272
- central neuropathic pain, 127
- cerebellar dysfunction, 236, 241
- cerebrovascular disease, 12, 112, 201, 223, 240, 280
- CGA. *See* comprehensive geriatric assessment
- Charles Bonnet Syndrome (CBS), 221, 277
- Cheshire West* case (2014), 78
- childhood: adverse experiences, 188
- children, 32, 79, 212
 carers for parents or grandparents, 96
 safeguarding, 90
- chlordiazepoxide, 215, 238
- chronic kidney disease, 246, 250, 251, 257
- chronic obstructive pulmonary disease (COPD), 242, 246
- citalopram, 133, 167–8, 216
- Cleveland scale for activities of daily living (CSADL), 306
- clinical institute withdrawal assessment of alcohol scale, revised (CIWA-Ar), 238
- clinical opiate withdrawal scale (COWS), 240
- clonazepam, 178, 277
- clopidogrel, 252, 253
- clozapine, 134, 277, 281
- cocaine, 234, 240, 242
- codeine, 234, 235, 249
- cognitive ageing, 9, 214
- cognitive assessment method (CAM), 258
- cognitive behavioural therapy (CBT), 13, 166, 194, 259, 278
 anxiety, 210
- cognitive impairment, 30, 47, 54, 65, 90, 110, 112, 119, 129, 150, 208, 215, 217, 233, 234, 235, 239, 243, 249, 274, 281, 301, 307, 308
 causes, 111
 exacerbation, 133
 medications, 116–17
 suicide risk, 195
- cognitive screening tools, 300–3
 Addenbrooke's cognitive examination III (ACE-III), 301–2
 frontal assessment battery (FAB), 298–303
 mini-Addenbrooke's cognitive examination (MACE), 302
 mini-cog, 303
 Montreal cognitive assessment (MoCA), 302
 RUDAS, 303
- cognitive stimulation therapy, 130
- collateral history, 126, 129, 142, 143
- common law, 70, 73, 93
- communication, 34
 definition (*OED*), 44
 general points, 46
 important part of healing process, 44
 liaison psychiatry services, 296
- communication difficulties, 17, 49, 161
- communication skills, 2, 13, 41, 44, 45, 65, 224, 287
- communication with older people, 44–57
 age-related barriers, 49–50
 breaking bad news, 49
- environmental factors, 46, 47
- factors affecting healthcare professionals, 45–6, 46
- factors affecting patients, 46, 47
- non-verbal communication, 46, 48–9
- social context, 47–8
- community mental health teams, 192, 228
- comorbidities, 24, 32, 55, 114, 132, 157, 161, 195
- comprehensive geriatric assessment, 4, 61, 98, 146, 222, 248, 283–91
 advantage (important), 286
 barriers to implementation, 284
 benefits, 283
 clinical practice, 287–8
 components, assessment, intervention, 288
 concept, 284–5
 criticism, 287
 critics as well as supporters, 284
 definition, 283
 delivery (practical tips), 289
 delivery (skills required), 289
 effectiveness, 285
 further research, 285, 290
 history, 283–4
 insufficient evidence (some areas), 286–7
 key aspects, 286
 old age versus ageing process, 285–6
 polypharmacy, 288–9
 tips (top ten), 290
 toolkit domains, 283
- confidentiality, 39, 49, 51, 54, 91, 93, 103
- confusion, 28, 103, 114
- confusion assessment method (CAM), 141, 144, 145, 151
- confusion assessment method: intensive care unit (CAM-ICU), 144, 151
- confusional states, 12, 48, 239
- congestive cardiac failure, 252, 256, 257
- conivaptan, 252
- consent, 66, 91, 92

- consent form, 73
 constipation, 248, 268, 275
 contextual safeguarding:
 definition, 90
 coping skills, 31, 183, 185, 213
 coping strategies, 13, 15, 41, 212
 coping styles, 9, 13, 202, 213
 Cornell scale for depression in dementia, 164
 coroners, 78, 183, 186, 196
 corticobasilar degeneration (CBD), 278, 278
 corticosteroids, 160, 176, 208
 Court of Appeal, 78
 Court of Protection, 74, 75
 covert medication, 80
 COVID-19 pandemic, 27, 49, 55
 alcohol, 32
 delirium patients, 142
 disruption to mental health services, 32
 impact on people with dementia and their family carers, 106–7
 influence on prevalence of mental disorders in acute hospitals, 32–4
 liaison psychiatry, 32
 vaccination, 32
 creatinine, 251, 251, 258
 Crime and Disorder Act (1998), 93, 95
 Cronbach's alpha, 301, 305
 cuckooing, 84
 definition, 95
 cumulative deficit model, 4
 cycle of discontent (Jurgens et al.), 17, 101
 cytochrome P450, 216, 217
 cytokines, 160, 259

 dabigatran, 253
 Data Protection Act (2018), 93
 DDS. *See* dopamine dysregulation syndrome
 death, 33, 76, 78, 100, 103–4, 151
 aftermath practicalities, 103
 alcohol-specific, 234
 dying process (alien behaviour compared with usual persona), 104
 in hospital, 67
 idealised scenario, 67
 proximity, 103
 decision-making, 49, 52, 55, 66
 capacity, 70
 deep vein thrombosis (DVT), 253
 hip-fracture patients, 258
 dehydration, 79, 186, 191, 248, 257, 258, 294
 delirium, 5, 12, 29, 30, 32, 33, 78, 80, 100, 101, 103, 113, 114, 125, 172, 220, 259, 285
 4AT screening tool, 141, 144, 145, 151, 258
 additional resources, 141, 142
 antipsychotic medication, 149
 awareness-raising, 140
 complex-case management, 141–2
 definition, 140, 258
 detection and screening, 141
 diagnostic features, 141
 distress caused, 140
 hospital patients, 39
 management (liaison old age psychiatry), 140–52
 medical emergency, 163
 mortality, 258
 overlap with dementia, 30
 Parkinson's disease patients, 276
 pathways and aftercare, 150
 patient outcomes, 140
 patients and carers (support), 142
 postoperative, 227
 postoperative (hip patients), 258–9, 260
 postoperative complication, 285
 prevalence in community and hospital settings, 29
 prevalence rates, 30
 prodromal phase, 144
 risk factors, 140
 secondary psychosis in late life, 222–3
 types, 140, 144
 delirium (clinical management), 143–50
 1. take good history, 143
 2. use screening tool, 143–5
 3. identify and treat all possible causes, 142–6
 4. prioritise non-pharmacological management, 146
 5. use pharmacological management as last resort, 146–9
 6. have clear follow-up plan on discharge, 149–50
 delirium care
 additional information, 143
 awareness-raising, 151
 education and peer supervision, 142
 influencing local services, 142
 key principles, 143
 knowledge, 152
 psychiatric liaison services (role), 140–3
 skills, 151
 training, 144
 Delirium Dementia Outreach Team (DDOT, Sunderland), 151
 delirium education, 151
 delirium rating scale, 145
 delirium room model (Saint Louis University Hospital), 40
 delirium superimposed on dementia, 150
 delirium tremens, 215, 236–8
 mortality, 238
 delirium triage screen (DTS), 144
 delirium wheel, 147
 delusional disorder, 221, 226–7, 229
 key differences versus EOS and VLOSP, 226
 presentations to liaison service, 228
 delusional parasitosis, 226
 delusions, 113, 126, 222, 225
 dementia, 1, 3, 5, 12, 29, 30, 32, 48, 51, 64, 79, 96, 100, 101, 102, 103, 106, 110–23, 163, 200, 217, 226, 248, 301, 305
 alcohol-related ~, 241
 atypical cause (indicators), 115
 atypical cause (investigations suggested), 115

- dementia (cont.)
 behavioural and
 psychological symptoms
 [*qv*], 124–39
 blood tests, 115
 Bristol-ADLS, 306
 causes, 111
 clinical management, 114
 cognitive assessment tools,
 114–15
 community support,
 117–18
 complex interplay
 (Kitwood), 66
 COVID-19 pandemic, 106–7
 CSADL, 306
 definition, 110
 degenerative, 111
 and delirium, 172
 diagnosis (cognitive
 assessment ‘essential
 part’), 300
 discharge to place of
 residence, 117–18
 disempowerment, 48
 epidemiology, 110
 funding and care challenges,
 30
 hospital admission
 (negative outcomes), 17
 incidence (UK), 110
 investigations, 115–16
 medications for cognitive
 impairment, 116–17
 neuroimaging, 115–16
 non-pharmacological
 interventions, 117
 overlap with delirium, 30
 patient report and
 informant history, 114
 pharmacological
 interventions, 116–17
 presentation (hospital
 setting), 113–14
 raising awareness (global
 priority), 110
 results, 110
 risk factor for postoperative
 delirium, 248
 secondary psychosis in late
 life, 222–3
 suicidal behaviour, 195
 UK incidence (2040
 projection), 5
 dementia (risk factors),
 112–13
 age, 112
 alcohol, 113
 depression, 113
 ethnicity, 113
 gender, 113
 genetics, 112
 head injury, 113
 smoking, 113
 dementia (sub-types), 111–12
 Alzheimer’s disease, 111
 dementia with Lewy bodies,
 112
 mixed cases, 112
 vascular dementia, 111
 Dementia Action Alliance, 134
 dementia care, 1
 staff satisfaction, 118–19
 vital element, 195
 dementia knowledge, 118–19
 dementia nurses, 104
 dementia patients
 admission to general
 hospital, 96
 restrictive practices, 128
 dementia with Lewy bodies
 (DLB). *See* Lewy-body
 dementia
 dementia-friendly acute care
 (D-FAC) settings, 39
 dementia risk factor, 112
 Department of Health, 52, 134
 dependence scale (DS):
 activities of daily living
 (assessment), 306
 dependence: definition, 233
 dependency, 283, 286
 depot medication, 229, 230
 depression, 11, 12, 12, 14, 29,
 30, 79, 113, 125, 127, 129,
 133, 157–82, 186, 201, 287
 aetiology, 158
 artificial feeding and
 hydration, 169
 associated medical
 conditions, 160
 association with alcohol
 misuse, 234
 clinical management and
 therapeutic issues,
 164–70
 comorbidity with
 depression, 158
 dementia risk factor, 113
 diagnostic criteria, 162
 driver of suicide (but
 treatable), 188–91
 epidemiology, 158
 hip fractures, 246
 as medication side-effect,
 160
 Parkinson’s disease patients,
 277
 postoperative (hip patients),
 259–60
 prevalence (hospital
 settings, UK), 158
 prevalence rates, 29, 30
 psychotic symptoms, 294
 with psychotic symptoms,
 158
 recurrent episodes, 158
 risk factors, 158–60
 standardised diagnostic
 tools, 163–4
 treatment-resistant, 294
 treatment-resistant (use of
 ECT), 169–70
 depression (common
 presentations in hospital
 settings)
 adjustment disorders, 161
 appropriate mood reactions,
 161
 communication difficulties,
 163
 dementia and delirium, 163
 depressive disorders, 161–2
 depressive disorders with
 psychotic symptoms, 163
 depression scales, 303–5
 BASDEC, 304
 GDS, 304–5
 HADS, 304–5
 PHQ, 305
 depressive disorders, 161–2
 psychotic symptoms, 163
 depressive psychosis, 221, 223
 depressive symptoms (non-
 pharmacological
 management), 164–6
 active monitoring, 164
 psychological therapies, 166
 removing triggers, 165
 sleep hygiene, 165
 social interventions, 165–6
 somatic symptom control,
 165
 depressive symptoms
 (pharmacological
 interventions), 166–9
 antidepressant therapy
 (choice), 167

- antidepressants
 (absorption), 166, 167
 drug administration route, 166
 treatment adjuvants, 167–9
 deprivation of liberty, 77–9
 six qualifying requirements, 77
 deprivation-of-liberty
 safeguards (DOLS), 77,
 78, 79, 80, 100, 142, 242
 despair, 10, 13, 204
 dextrose, 254, 255
 diabetes, 3, 113, 114, 234, 238,
 288
 elderly patient and surgery
 (general rules), 255
 hip patients, 254–5
 type one, 254
 type two, 254, 255
 diagnostic overshadowing:
 definition, 90
 dialectical behaviour therapy
 (DBT), 194
 diazepam, 178, 215
 dignity
 acute hospitals, 59–68
 concepts, 61–2
 definition (*OED*), 61
 end-of-life care, 67–8
 etymology, 61
 Nordenfelt's categorisation,
 61–2
 nutrition, 65
 personal hygiene, 65
 person-centred care, 65–7
 staff interaction, 64–5
 dignity and environment, 62–4
 clocks, 64
 colours, 63
 communal eating areas, 64
 day rooms, 64
 floors, 63
 lighting, 63
 noise, 64
 photographs, 64
 signage, 63
Dignity in Practice (Tadd et
 al., 2011), 60
 dignity of merit (Nordenfelt),
 61, 62
 dignity of moral stature
 (Nordenfelt), 62
 dignity of personal identity
 (Nordenfelt), 61, 62, 65,
 66, 68
 digoxin, 127, 252
 direct oral anticoagulants
 (DOACs), 253
 disability, 10, 15, 16, 50, 212
 adaptive coping strategies,
 13–16
 disability assessment for
 dementia (DAD), 307
 disability-adjusted life years
 (DALYs)
 definition, 27
 discharge planning, 55, 243–4
 discrimination: indicators and
 examples, 87
 disinhibition, 126, 127, 146,
 170, 184, 215, 242, 308
 dissociative anxiety: definition,
 206
 distraction, 14, 16, 56, 146, 300
 distress, 185
 doctrine of necessity, 77
 DOLS. *See* deprivation-of-
 liberty safeguards
 domestic abuse, 14, 92
 indicators and examples, 85
 domestic violence: telephone
 helpline numbers, 94
 domperidone, 256, 272, 273
 donees: definition, 74
 donepezil, 116, 132
 dopamine, 255, 266, 270, 281
 dopamine agonists, 271, 272,
 274, 278
 withdrawal syndrome, 270
 dopamine antagonists, 271,
 273, 280
 dopamine dysregulation
 syndrome, 270, 276
 Parkinson's disease patients,
 266–78
 dopamine receptor blockade,
 281
 dopamine transporter (DaT),
 112, 115
 dopaminergic cell loss
 (progressive), 267
 dopaminergic levels, 268
 dopaminergic medication,
 255, 271, 272, 276, 278,
 280, 281
 dopaminergic pathways, 267,
 278, 281
 drawing, painting, writing, 48
 driving
 alcohol-related accidents
 (UK), 243
 legal entitlement (alcohol
 use), 243
 drug and alcohol misuse,
 233–44
 acute presentations of
 alcohol misuse and
 dependence, 236–8
 adverse reactions
 (pharmacodynamic,
 pharmacokinetic), 234
 alcohol withdrawal, 236–8
 alcohol-related brain
 damage, 240–1
 assessment, 235–8
 comorbidity, 241–2
 definitions, 233
 discharge planning, 243–4
 dual diagnosis (with mental
 disorders), 234
 early versus late onset, 233
 epidemiology, 233–4
 identification (barriers),
 235
 integrated care, 242
 interactions, 234–5
 medico-legal aspects, 242–3
 substance misuse and
 driving, 243
 suicide and self-harm, 238
 drug misuse
 acute presentation, 239–40
 intoxication and overdose,
 239–40
 opioids, cannabis,
 stimulants, 239–40
 psychosis, 223, 240
 withdrawal, 240
 drug-induced Parkinsonism,
 267, 281
 DSM-IV, 305
 delirium, 258
 DSM-V, 161, 171, 174, 224
 depressive episode
 (diagnostic criteria), 162
 manic episode (symptoms),
 171
 DT. *See* delirium tremens
 duloxetine, 167, 217
 dynamic hip screw (DHS),
 257
 dysarthria, 13, 47, 112, 270
 dyskinesia, 272, 274
 definition, 269
 dysphagia, 269, 270
 dysphasia, 47, 163
 dystonia, 269, 278

- early-onset schizophrenia (EOS), 224, 225–6, 229
 key differences versus DD and VLOSP, 226
 relapse, 227
 ECHR. *See* European Convention on Human Rights
 Edinburgh: Royal Victoria Hospital, 39
 EDs. *See* emergency departments
 education, 11, 13, 18, 65, 102, 141, 142, 151, 152, 164, 210, 301, 303
 elated mood, 170, 171
 active monitoring, 176–7
 acute stage, 176
 diseases causing, 175
 neurological conditions, 175
 non-pharmacological interventions, 176–7
 pharmacological interventions, 177–9
 relapse, 176
 removal of triggers, 177
 risk factors, 174–5
 sleep hygiene, 177
 somatic symptom control, 177
 treatment, 176–9
 elated mood (differential diagnoses), 172–4
 anxiety disorders, 172–3
 bipolar affective disorder, 174
 delirium, 172
 intoxication with psychoactive substances, 172
 personality disorders, 173–4
 schizophrenia spectrum disorders, 173
 stimulants, hallucinogens, alcohol, 172
 elder abuse, 41, 242
 elderly-friendly care settings, 38–42
 elderly-friendly clinician: top qualities, 41–2
 elderly-friendly hospitals, 38–9
 delirium room model, 40
 environmental strategies to enhance safety, 40
 environmental strategies to maintain orientation, 40
 orientation strategies, 40
 ward features, 41
 elderspeak, 52, 65
 definition, 50
 useful aspects, 65
 electrocardiograms, 131, 258
 electroconvulsive therapy, 71, 76, 193, 194
 definition, 294
 liaison psychiatry services for older people, 294
 use in treatment-resistant depression, 169–70
 electrolyte abnormalities, 248, 259
 electrolyte imbalance, 248, 250, 251, 258
 emergency departments, 5, 26, 33, 38, 40, 142, 183, 228, 287
 delirium (screening tools), 144
 delirium cases, 141
 older people (prevalence of mental health disorders), 26–8
 patient satisfaction (low level among elderly), 27
 patients with psychiatric conditions (increase 2016–2020), 28
 emergency services, 26, 93
 emotionally unstable personality disorder, 173
 empathy, 18, 41, 45, 49, 52, 59, 62, 64, 209
 empowerment, 18, 19
 encephalitis, 116, 175, 222
 endocrine diseases, 160, 175
 end-of-life care, 39, 42, 54, 67–8, 287
 gold standards framework, 103
 enduring power of attorney (EPA), 74
 enoxaparin, 258
 epidemiology
 definition, 24
 dementia, 110
 depression, 158
 drug and alcohol misuse, 233–4
 mental illness in older people (acute hospitals), 24–34
 self-harm and suicide, 186–7
 escitalopram, 167–8, 216, 217
 estimated glomerular filtration rate (eGFR), 117, 257
 ethnicity: dementia risk factor, 113
 European Convention on Human Rights, 77
 European Court of Human Rights, 77
 Evans Liverpool depression rating, 163
 executive function, 114, 226, 260, 280, 301, 302, 303
 exercise, 11, 14, 34, 166, 214, 283
 experience, 62, 118
 expert generalist role, 3
 extrinsic motivation, 118
 eye desensitisation and reprocessing (EMDR), 210, 213
 falls, 39, 41, 60, 113, 217, 236, 239, 255, 278, 281, 287
 Parkinson's disease, 274–5
 prevention plans, 260
 familial prion dementia, 115
 family, 13, 34, 51, 52, 76, 105, 117, 165, 172, 194, 211
 characteristics, 212
 closed versus open, 212
 congruent communication, 212
 enmeshment versus differentiation, 212
 goodness of fit, 212
 negative outcomes (highest risk), 213
 problems, 212
 role flexibility, 212
 themes, beliefs, patterns, scripts, 212
 use as interpreters, 51
 family carers, 44, 54–5, 96, 97, 100, 101, 107, 118, 195,
 See also carers (informal)
 family hierarchies, 212
 family therapy, 14, 209
 fasting, 254, 255
 fear, 13, 17, 33, 103, 161, 186, 196, *See also* anxiety
 female genital mutilation, 84
 Ferreira case (2017), 78
 financial abuse: indicators and examples, 84
 fire safety, 244

- fludrocortisone, 272, 275
 fluid balance, 250–1, 260
 fluoxetine, 133, 167–8, 216, 235, 252
 folic acid, 250
 frailty, 4–6, 12, 13, 33, 61, 62, 101, 287, 292
 characteristics, 4
 definition, 4
 delirium risk factor, 140
 impact of minor triggers on functional ability, 4
 frailty index, 4
 frailty phenotype model, 4
 frailty services, 143, 146
 frailty syndromes, 4, 287
 Francis report (2013), 59, 60, 65, 86, 95
 frontal assessment battery (FAB)
 cognitive screening tools, 298–303
 languages, 303
 frontotemporal dementia, 111, 127, 278, 301, 303, 308
 functional ability, 5, 16, 18, 236, 307
 risk factor (modifiable) for suicide, 188
 functional impairment, 186, 191, 307
 further research
 comprehensive geriatric assessment, 285, 290
 mental health conditions in older people (acute settings), 34
 Parkinson's disease, 268
 future directions: late-onset psychosis, 230

 GABA (γ -aminobutyric acid), 215
 gabapentinoids, 233, 234
 galantamine, 116
 gas lighting, 84
 definition (figurative), 95
 gastroenterology, 242, 275
 Gélinas I, 307
 gender
 dementia risk factor, 113
 differences (healthcare professional versus patient), 45
 gene mutations, 112, 268
 general anaesthetic, 78, 80, 169, 294
 General Data Protection Regulation (GDPR), 93
 general hospitals, 80, 125, 140, 141, 160, 169, 178, 220, 222, 223, 229, 292
 admissions (dementia, delirium, psychosis), 222
 beds, 26
 covert medication, 80
 delirium (screening tools), 144–5
 General Medical Council, 81, 82, 91
 general practitioners, 51, 55, 75, 80, 92, 116, 117, 149, 166, 189, 190, 192, 193, 228, 292
 services outside of hospital, 294–5
 generalised anxiety disorder (GAD), 208, 215, 216, 217
 autonomic arousal
 symptoms, 203
 diagnostic criteria, 203–19
 exclusion criteria, 203
 general symptoms, 207
 non-specific symptoms, 203
 symptoms concerning brain and mind, 207
 symptoms concerning chest and abdomen, 203
 symptoms of tension, 203
 genetics: dementia risk factor, 112
 geriatric anxiety inventory (GAI), 209
 geriatric depression scale (GDS), 163, 303, 304–5
 five item subset, 190
 test-retest reliability, 304
 geriatric giants (listed), 1, 289
 geriatric medicine, 1–6
 ageing population, 2–3
 definition, 1
 evolution of specialities, 1–2
 frailty and assessment, 4–6
 largest hospital medical specialty, 61
 pioneers (1940s), 1
 scale of problem, 3–4
 Gillick competence, 79
 glucose, 115, 254, 255, *See also* blood glucose
 grandiosity, 170, 171
 haematoma, 111, 222, 253, 258
 haemoglobin, 250, 256, 257
 hallucinations, 113, 126, 221, 222
 definition, 223
 Parkinson's disease patients, 276–7
 hallucinoses, 221, 223, 225
 haloperidol, 148, 178, 276
 Hamilton depression scale, 304
 hate crime, 89, 95
 definition (Metropolitan Police), 95
 head injury, 258
 dementia risk factor, 113
 Health and Social Care Act (2008), 65
 health and welfare, 72, 73
 health records, 66, 164
 health status, 10, 20, 160, 189
 healthcare professionals, 82
 communication with dementia patients (improvement tips), 56
 communication with family and family carers, 54–5
 communication with patients, 45–6
 information-sharing, 55–6
 motivation (extrinsic versus intrinsic), 118
 hearing aids, 47, 56, 90, 127, 146, 300
 hearing impairment, 48, 221
 hearing loss, 225, 300
 heart disease, 3, 62, 105, 190, 234
 ischaemic, 250, 252, 257
 hebephrenic schizophrenia, 173
 heroin, 228, 242
 comorbidities, 234
 high dependency unit, 250, 252
 hip-fracture patients
 mortality (factors), 247
 perioperative medical management (lessons), 246–60
 hip-fracture surgery
 reasons to delay, 256
 hippocampal formations, 111
 history-taking, 223, 227, 228, 240

- histrionic personality disorder, 173
 HIV-associated dementia, 115
 holistic approach, 41, 67, 90, 151, 165
 home treatment teams, 192, 195, 230, 293
 hopelessness, 13, 185, 195, 204
 hospice, 103
 hospital admissions, 125
 approaches to calmness, 99
 dissonance, 101
 first impressions, 98–9
 intensive investigation of physical ill-health, 99–100
 mental capacity, 100
 mental disorders, 100
 relocation with hospital, 100–1
 stressful and upsetting, 172
 hospital admissions (good practice), 101–4
 altered mental function (specialist services), 104
 carer initiatives, 102
 death, 103–4
 diagnosis and prognosis, 102
 making carers welcome, 101–2
 persisting symptoms, 103
 progress within ward, 102
 resolution, 103
 hospital admitted patient care activity, 2
 hospital anxiety and depression scale (HADS), 164, 304–5
 hospital design: decision-making (involvement of older people), 39
 hospital environment: effects upon recovery, 17
 hospital settings, 134, 142
 BPSD, 124–39
 hospitalisation
 adaptive coping strategies, 13–16
 COVID-19 pandemic (impact on people with dementia and their carers), 106–7
 older people, 38
 hospitals, 78
 elderly-friendly, 38–42
 night-health hazards (older people), 39–40
 physical environment, 17
 human beings: rational nature (Kant), 61
 human dignity (Nordenfelt), 61
 human rights, 60, 82
 Human Rights Act (1998), 93
 humour, 13
 Huntington's chorea, 111, 115, 175, 195
 hydration, 59, 63, 169
 hyperactive delirium, 143
 confused with mania or psychosis, 141
 hyperglycaemia, 248, 254, 255
 hypertension, 234, 287
 hypnotics, 178, 193
 hypoactive delirium, 143, 144, 157, 162, 275
 misdiagnosed as depression or dementia, 141
 hypoglycaemia, 222, 248, 254, 255
 hypomania, 173, 174, 177, 178
 definition, 174
 hypomimia, 269
 hyponatraemia, 160, 216, 250, 251
 older hip-fracture patient, 251–2
 severity scale, 252
 underlying cause (identification and treatment), 252
 hypotension, 160, 256, 257
 hypovolaemia, 248, 250
 ICD-10, 144, 161, 171, 174, 224
 acute stress reaction, 204–17
 adjustment reaction, 204
 delirium, 258
 depressive episode (diagnostic criteria), 162
 generalised anxiety disorder, 202–3
 manic episode (symptoms), 171
 obsessive compulsive disorder, 207–8
 panic disorder, 206–7
 phobias, 206
 psychiatric disorders (UK prevalence), 28
 PTSD, 204–5
 ICDs. *See* impulse control disorders
 ill-being (characteristics), 11–12
 bitterness, 11
 difficult life, 11
 negative self, 11
 negative world, 11
 purposelessness, 11
 illness (idiosyncratic and contextual variables), 12–13
 ability to communicate, 13
 compliance to treatment regimes, 13
 compounding difficulties, 12
 coping styles, 13
 level of awareness, 12
 illness scenarios (adaptive coping), 13–15
 appraisal-focused strategies, 13
 emotion-focused strategies, 14–15
 solution-focused strategies, 14
 illness scenarios (negative coping), 15–16
 avoidance, 16
 overcompensation, 15
 surrender, 16
 illness-specific conditions: causes, 12
 IMCA. *See* Independent Mental Capacity Advocate
 IMPACT intervention (depression), 191
 implantable cardiac defibrillators, 254
 improving access to psychological therapy (IAPT), 294
 impulse control disorders, 270, 276, 277
 Parkinson's disease patients, 277–8
 inappropriate sexual behaviour, 128, 131
 incidence: definition, 27
 independence, 11, 61, 64, 65, 67

- independent domestic violence advisors (IDVAs), 92
- independent mental capacity advocate, 75–6, 93
- inequality, 16, 87, 188
- infectious diseases, 175
- with depressive symptoms, 175
- informant questionnaire on cognitive decline in elderly (IQCODE), 150
- informant-based response, 306
- inpatient care, 17, 293
- inpatient services: LPOP, 292–3
- inpatients, 3, 110, 114, 184, 192, 273
- insulin, 184, 254, 255
- integrated care, 82, 143, 149, 242
- intensive care units, 78, 250
- interleukin 1 (IL-1), interleukin 6 (IL-6), 259
- intermittent pneumatic compression (IPC) devices, 258
- International Classification of Diseases* (tenth edition). See ICD-10
- International Consensus on VLOSP, 229
- international normalized ratio (INR), 253
- interpersonal skills, 118–19
- interpersonal therapy (IPT), 194
- interpreters, 50, 51, 201, 235
- use of family, 51
- interRAI PC scale, 286
- intrinsic motivation, 118
- irritability, 170, 173, 201
- John's Campaign, 64, 101
- ketoacidosis, 254
- Korsakoff's Syndrome (KS), 240
- language, 45, 50, 102
- lasting power of attorney, 66, 72, 73–5, 76, 81, 84, 98
- two types, 73
- late paraphrenia (Kay and Roth), 224
- late-onset psychosis, 220–30
- antipsychotic medication (choice), 229–30
- assessment, 228–9
- biopsychosocial framework, 222
- common medical causes, 222
- future directions, 230
- incidence, 221
- management, 229
- primary, 224
- symptoms, 220–1
- late-onset psychosis (presentations to liaison service)
- cannabis psychosis, 228–9
- delusional disorder, 228
- personality problems merging into VLOSP, 227
- postoperative delirium, 227
- relapse of EOS, 227
- VLOSP, 228–9
- late-onset schizophrenia (LOS), 224, 225
- Law Commission, 100
- law: liaison psychiatry, 70–81
- learning disabilities, 46, 88, 90
- levodopa, 160, 255, 271, 272, 274, 278, 280, 281
- Lewy bodies, 267
- definition, 112
- Lewy-body dementia, 112, 115, 116, 132, 134, 148, 223, 266, 278, 281, 301
- Parkinson's plus syndromes, 280
- liaison old age psychiatry
- alcohol misuse, 233
- delirium management (acute hospitals), 140–52
- education and peer supervision, 142
- influencing local services, 142
- liaison psychiatrists, 157, 169, 179, 266
- mental illness in older people (growing challenge), 24–34
- Parkinson's plus syndromes (need to know), 278–80
- Parkinson's disease, 273, 276
- providing reports, 80
- role, 3
- liaison psychiatry
- COVID-19 pandemic, 32
- definition, 1
- drug and alcohol misuse, 233
- goal, 25
- impact 'limited and sometimes variable', 25
- importance (acute hospitals), 25
- increasing demand, 25
- law, 70–81
- models, 25
- referral rate (2021 report), 28
- liaison psychiatry services for older people
- AMHPs, 292–3
- crisis teams, 293
- electroconvulsive therapy, 294
- general community services, 293–4
- inpatient services, 292–3
- interface with wider community services, 292–7
- role, 292
- services outside of hospital, 292–4
- transfer of patients from acute trusts to psychiatric wards, 292–3
- transfer of patients from psychiatric wards to acute trusts, 293
- liaison psychiatry team, 1, 15, 27, 29, 160, 164, 172, 174, 176, 178, 277
- liberty protection safeguards, 78, 142
- changes from DOLS, 78
- legal basis, 79
- scheduled to be implemented on 1 April 2022, 78
- life expectancy, 2, 26, 38, 226
- life-saving treatment, 76, 78
- lifestyle counselling, 214
- building self-confidence, 214
- life-sustaining treatment, 75
- lifetime risk: definition, 27
- lip-reading, 48, 49, 300
- transparent masks 'helpful', 90

- listening, 44, 48, 56, **91**, 96,
 106, 201, 209, 284
 lithium, 178, 188, 277, 281
 lithium adjuvant therapy, 168
 lithium toxicity, 160
 liver disease, 32, 233, 236, 238,
 242
 living meaningful life, 10
 local authorities, 77, 78, 82, 97
 local authority designate
 officer (LADO), 90
 loneliness, 10, 31, 61, 161, 188
 long COVID, 33
 long-term care, 50, 283, 284
 lorazepam, 148, 178, 238
 loss, 161, 208
 low mood, 28, 126, 194
 low-income countries, 32
 low-molecular-weight heparin
 (LMWH), 253
 LPA. *See* lasting power of
 attorney
 LPOP. *See* liaison psychiatry
 services for older people
 LPS, 79. *See* liberty protection
 safeguards
 LTC. *See* long-term care
 lung function, 247
- making safeguarding personal,
 82, 90
 malignant social psychology
 (Kitwood), 48
 Mallory-Weiss tears, 241
 malnutrition, 17, 38, **41**, 79,
 158
 Malnutrition Task Force, 65
 mania, 170–2, 173, 174, 175,
 178, 223
 aetiology, 171–2
 as secondary syndrome, 174
 epidemiology, 170
 presentation, 170–1
 symptoms, 171
 material abuse: indicators and
 examples, **84**
 MCA. *See* Mental Capacity
 Act (2005)
 McMaster quality of life scale,
 286
 MDT. *See* multi-disciplinary
 teams
 meaning and purpose to life,
 10
 meaninglessness, 11
 causal factors, 11
 strong predictor of mental
 health problems, 11
 medical surgical treatment:
 mental capacity and
 consent, 71
 medication-related harm, 288,
 289
 medications, 105, 208, 258
 causing elated mood, **176**
 interactive effects, 12
 medicinal cannabis: side-
 effects, 239
 melatonin, 133, 277
 memantine, 116–17, 132, 277
 memorial delirium assessment
 scale, 145
 memory, 260, 300
 memory aids, 56
 memory assessment services,
 293–4
 memory clinics, 150
 memory loss, 1, 111, 114
 memory problems, 52, 114
Menschenwurde, 61, 62, 67
 mental capacity, 31, 52, 54,
 100
 Mental Capacity
 (Amendment) Act
 (2019), 78, 79
 Mental Capacity Act (2005),
 52, 70, 100, 131, 142, 148,
 149, 164, 178, **192**, 242
 accident and emergency,
 76–7
 accompanied by Code of
 Practice, 71, 72, 76, 77
 advance decisions, 75
 advocates, 75–6
 capacity and consent,
 70–3
 children and young people,
 79
 consideration of best
 interests, 71–3
 court-appointed deputy, 75
 definition of ‘nearest
 relative’, 73
 definition of capacity, 70, 71
 deprivation of liberty, 77–9
 five principles, 70–1, 75
 LPA, 73–5
 no definition of ‘next of
 kin’, 73
 restraint, 73
 mental capacity and consent,
 70–3
 proportionality (major
 versus minor decision-
 making), 71
 provision of information to
 patient, 73
 Mental Health Act (1983), 71,
 77, 78, 79–80, 142, 158,
 169, 178, **192**, 227, 228,
 230, 242, 293, 294
 amended (2007), 70
 reform consultation
 completed, 81
 Mental Health Crisis
 Assessment Service, 33
 mental health disorders, 3, 164
 influence of COVID-19
 (acute hospitals), 32–4
 older people (annual cost,
 UK), 24
 prevalence in community
 and hospital settings, **29**
 prevalence in older people
 (EDs), 26–8
 mental health law, 70–81
 accident and emergency,
 76–7
 advance decisions, 75
 advocates, 75–6
 capacity and consent, 70–3
 children and young people,
 79
 covert medication, 80
 deprivation of liberty, 77–9
 LPA, 73–5
 restraint, 73
 mental health services, 5, 25
 delayed contact, 26
 liaison service, 104
 perceived advantage, 1
 mental state examination, 144,
 172, 175, 228
 methadone, 234, 235
 metoclopramide, 235, 273,
 276, 280, 281
 MHA. *See* Mental Health Act
 (1983)
 midodrine, 272, 275
 migration theory, 225
 mild cognitive impairment
 (MCI), 114, 115, 302
 mindfulness, 214–15
 MINE mnemonic, 222, 222
 mini-Addenbrooke’s cognitive
 examination (MACE)
 cognitive screening tool,
 302

- mini-cog (cognitive screening tool), 303
 minimal clinically important difference (MCID), 299
 mini-mental-state examination (MMSE), 114, 115, 296, 303, 306
 test-retest reliability, 301
 minimum alcohol pricing, 187
 mirtazapine, 133, 160, 167–8, 167, 168, 216
 mitochondria DNA, 9
 MMSE. *See* mini-mental-state examination
 MOBID-2 pain scale, 129
 mobility, 11, 13, 15, 49
 model of human occupation screening tool (MOHOST), 15
 modern slavery: indicators and examples, 85
 modified confusion assessment method for emergency departments (mCAM-ED), 144
 Mokken scaling, 302
 monoamine oxidase (MAO) inhibitors, 216, 256, 271, 272, 277
Montgomery case, 73
 Montreal cognitive assessment (MoCA), 114, 296
 cognitive screening tool, 302
 test-retest reliability, 302
 translations, 302
 mood disorders, 30, 133, 157, 164, 165, 170, 171, 223, 281
 aetiology, 165
 development or exacerbation (risk factors), 158–60
 prevalence rates (ED patients, UK), 28
 mood disturbance, 28, 116, 173
 mood stabilisers, 174, 178
 morbidity: definition, 27
 morphine, 184, 234, 249
 mortality, 9, 33, 40, 119, 140, 260, 285, 286
 hip fractures, 246, 247
 mortality rate: definition, 27
 motor symptoms, 267, 268, 270, 271, 277
 Parkinson's disease, 269
 multi-agency risk assessment conference (MARAC), 92
 multidisciplinary teams, 41, 148, 189, 243, 244, 247, 283, 287, 289, 295
 multimorbidity, 4, 9, 10, 38, 186, 220
 multiple comorbidities, 3, 44, 50, 143, 287
 multi-system atrophy (MSA), 278, 278, 280
 musculoskeletal system, 127, 130
 music therapy, 130
 myocardial infarction, 168, 257
 narcissistic wounds, 186
 nasogastric tube, 169
 National Confidential Inquiry into Suicide (2018), 193
 National Dementia Strategy, 110
 national early warning score (NEWS), 251
 National Helpline (domestic violence), 94
 national hip fracture database (NHFD), 248
 National Suicide Prevention Alliance, 196
 Neelon and Champagne (NEECHAM) confusion scale, 144
 neuroimaging, 115–16, 126, 175
 neuroleptic malignant syndrome, 255, 273
 neurological diseases, 175, 184
 with depressive symptoms, 175
 neuropsychiatric inventory (NPI), 126, 129, 307
 test-retest reliability, 307
 neuropsychiatric symptoms, 130, 131, *See also* BSPD
 examples, 124
 neuropsychological tests, 301
 neuropsychometric assessments, 115
 newer oral anticoagulants (NOACs), 253
 next of kin, 73, 132, 169
 N-methyl-D-aspartic acid receptor antagonists, 116–17
 noise, 64, 130, 146, 160, 200
 non-verbal communication, 46, 48–9
 Nordenfelt L: categorisation of dignity, 61–2
 normal pressure hydrocephalus (NPH), 267
 NPI-Q (Kaufer et al.), 129
 nurses, 40, 51, 55, 119, 191, 222, 242, 256, 259, 287, 288
 nurses' delirium screening checklist, 145
 nursing, 1, 40, 67, 72, 125, 145
 communication skills, 44
 nursing homes, 96, 117, 286, 307, 308
 covert medication, 80
 nursing shifts, 65
 nutrition, 59, 60, 63, 64, 65, 102
 obsessive compulsive disorder, 31, 213, 216
 components, 207
 diagnostic criteria, 207–8
 occupational therapy, 1, 14, 15, 117, 194, 214, 222, 243, 247, 295
 olanzapine, 148, 168, 178, 229, 276
 old age psychiatry, 1–6
 evolution of specialities, 1–2
 older patients, older people
 attendance at EDs (unexpected scale of increase, 2007–2014), 28
 factors affecting communication, 47
 hospitalisation, 38
 lack of physiological reserve, 247
 meaningful activities tailored to hospital settings, 39
 mental health disorders (ED prevalence), 26–8
 night-health hazards in hospitals, 39–40
 perioperative medical management (lessons from hip-fracture patients), 246–60
 perspectives (incorporation in care delivery), 18

- older patients, older people (cont.)
 preference for single rooms, 39
 self-harm and suicide (prevention), 183–96
- older person assessment and liaison (OPAL) teams, 285
- oncology, 29, 145, 166, 284, 287
- opiates, 239, 249, 276
 withdrawal, 240
- opioids, 127, 233, 234, 239, 249, 259
- organisational abuse: indicators and examples, 86
- osteoporosis, 236, 242, 255, 260
- overdose, 186, 187
- oxazepam, 215, 238
- Oxford English Dictionary*, 44, 61
- oxycodone, 235, 249
- pacemakers, 248, 254, 263
- pain, 11, 47, 67, 125, 127, 129, 162, 165, 177, 184, 194, 246, 248, 249, 259
- pain control, 260
 preoperative management (hip patients), 249
- palliative care, 286
 suicide risk, 195–6
- panic, 203, 216
- panic attacks, 206, 213
- panic disorder
 autonomic arousal symptoms, 207
 characteristics, 206
 diagnostic criteria, 200–7
 exclusion criteria, 203
 general symptoms, 207
 symptoms concerning brain and mind, 207
 symptoms concerning chest and abdomen, 207
- paracetamol, 188, 235, 249
- paramedics, 98, 295
- paranoia, 223, 239
- paraphrenia (Kraepelin), 224
- Parkinson's disease, 13, 47, 49, 112, 148, 159, 248, 295
 acute hospitals, 266–81
 dopaminergic drugs, 272
 hip patients (surgery), 255–6
 idiopathic, 278, 281
 medications, 270–1
 medications (workings), 271
 motor symptoms, 268
 non-motor symptoms, 268–70
 and related disorders, 266–81
 staging, 270
- Parkinson's disease (acute mental health issues), 276–8
 anxiety and depression, 277
 delirium, 276
 dopamine dysregulation syndrome, 266–78
 hallucinations, 276–7
 impulse control disorders, 277–8
- Parkinson's disease
 dementia, 277
 REM sleep behaviour disorder, 277
- Parkinson's disease (common problems in hospital), 273–8
 1. missed medication, 273
 2. swallowing difficulties and nil-by-mouth patients, 273–4
 3. falls, 274–5
 4. postural hypotension, 275
 5. gastroenterological problems, 275–6
 6. acute mental health issues, 276–8
- Parkinson's disease (emergency presentations), 273–8
- Parkinson's disease dementia, 132, 134, 270, 272, 276, 280
 acute mental health issues, 277
 hallucinations, 276
- Parkinson's mimics, 266, 267
- Parkinson's plus syndromes, 266, 276, 278–80
 Lewy-body dementia, 280
- liaison psychiatrists (need to know), 278–80
- Parkinsonism, 112, 267
 causes, 267
 commonest cause, 266
 definition, 266
 other causes, 280–1
- Parkinsonism-hyperpyrexia syndrome, 273
- paroxetine, 167, 216
- patient activated analgesia (PCA), 249
- patient health questionnaire (PHQ), 189, 305
- patient reported outcome measures (PROMS), 287
- PDs. *See* personality disorders
- people in positions of trust, 90
- performance measures, 306
- perioperative medical management (lessons from hip-fracture patients), 246–60
 postoperative management, 256–60
 preoperative management, 247–56
- peripheral neuropathy, 236, 242
- persecution, 163, 227, 229
- personal care, 62, 63, 64, 65, 67, 88, 228
- personal hygiene, 39, 59, 60, 65, 110
- personal identity, 61, 62
- personal protective equipment (PPE), 49
- personality disorders, 173–4
 prevalence, 31
 prevalence rates (ED patients, UK), 28
- person-centred care, 18, 20, 44, 50, 55, 56, 59, 65–7, 229
- CAREING acronym, 18–19
 principle, 66
- pharmacological treatment (anxiety), 215–17
 antidepressants, 216–17
 benzodiazepines, 215
 other medications, 217
- phobias, 213
 definition, 205
 diagnostic criteria, 206
 most common fears, 205
- physical abuse: indicators and examples, 84
- physical environment, 62, 64, 85
- physical health, 12, 16, 20, 24, 25, 26, 29, 31, 34, 100, 117, 127, 128, 129, 131,

- 171, 174, 177, 188, 189,
 190, 192, 194, 226, 242,
 266, 304
 physiologic reserve, 247, 260
 physiotherapy, 166, 194, 243,
 250, 295
 PINCH ME mnemonic, 143
 pneumonia, 248, 255, 259, 273
 incidence in pre-operative
 hip fractures, 250
 postoperative complication,
 285
 poisoning, 28, 234
 polypharmacy, 10, 38, 143,
 157, 186, 233, 234, 246,
 289
 comprehensive geriatric
 assessment, 288–9
 poor recovery from illness, 10,
 11, 20, 202
 Popoff index of depression,
 164
 positron emission tomography
 (PET), 112, 115
 postoperative cognitive
 dysfunction (POCD)
 hip patients, 260
 postoperative delirium, 246,
 248, 249, 259
 risk factor, 249
 postoperative management
 (hip patients), 256–60
 acute kidney injury, 257
 anaemia, 257
 cognitive dysfunction, 260
 delirium, 258–9
 depression, 259–60
 diabetic elderly patient and
 surgery, 254–5
 prevention of venous
 thromboembolism, 258
 vascular complications,
 257–8
 posttraumatic stress disorder,
 200, 216, 234
 diagnostic criteria, 204–5
 postural hypotension, 168,
 217, 268, 272, 274, 275,
 278, 288
 postural instability, 278, 281
 potassium, 251
 pregabalin, 133, 217
 pregnancy, 76
 preoperative management
 (hip patients), 247–56
 anticoagulants, 253
 antiplatelet treatments,
 252–3
 atrial fibrillation, 252
 general considerations,
 247–8
 hyponatraemia, 251–2
 optimizing anaemia prior to
 surgery, 250
 optimizing electrolyte
 imbalance, 251
 optimizing fluid balance,
 250–1
 pacemakers and
 implantable cardiac
 defibrillators, 254
 pain control, 249
 Parkinson's disease
 comorbidity, 255–6
 pneumonia and other
 infections, 250
 reasons to delay surgery,
 256
 prescription drugs, 223, 234
 pressure sores, 86, 246, 247,
 248, 260
 prevalence: definition, 27
 primary care, 2, 105, 188
 primary prevention:
 definition, 187
 primary psychosis, 223
 definition, 220
 late onset, 224
 PRIME study: key to
 abbreviation, 289
 prion disease, 115
 privacy, 20, 47, 51, 56, 99, 103,
 235
 acute hospitals, 59–68
 problem adaptation therapy
 (PATH), 195
 problem solving therapy
 (PST), 195
 prochlorperazine, 273, 276,
 280, 281
 progressive supranuclear palsy
 (PSP), 278, 278
 property and finances, 73, 74
 PROSPECT intervention
 (depression), 191
 protected characteristics, 87
 prothrombin complex
 concentrate (PCC), 253
 proxy response, 306
 psychiatric disorders, 170
 prevalence in acute general
 hospitals, 32–4
 prevalence rates (ED
 patients, UK), 28
 psychiatric inpatients: suicide
 risk, 191–5
 psychiatric patients: transfer
 to general hospital, 79
 psychiatric wards, 292, 293
 psychological abuse:
 indicators and examples,
 84
 psychological defence
 mechanisms, 211
 psychological sequels of
 illness, 12–13
 idiosyncratic and contextual
 variables, 12–13
 illness-specific factors, 12
 psychological therapies
 (anxiety), 209–13
 analytical psychotherapy,
 210
 cognitive behavioural
 therapies, 210
 EMDR, 213
 purpose, 209
 systemic therapies, 211–13
 psychological therapies
 (depressive symptoms),
 166
 psychological well-being in
 later life, 9–20
 psychology of ageing, 10–12
 psychometric measures (old
 age psychiatry), 298–308
 psychometric robustness, 298,
 299
 psychometric tests
 false-positive values, 300
 precision, 299
 quality, 298–300
 responsiveness, 299
 validity, 299
 psychometric tools
 clinical utility, 298
 definition, 298
 psychomotor retardation, 162,
 163, 267
 psychosis, 28, 33, 126, 129
 drug-induced, 240
 late-onset (*qv*), 220–30
 psychosocial treatments
 (types), 209–15
 lifestyle counselling and
 coaching, 214
 mindfulness-based
 techniques, 214–15

- psychosocial treatments:
 definition, 209
 psychotherapy, 14, 191, 194,
 209, 210
 psychotic symptoms, 133–4,
 158, 173
 public health
 backbone, 24
 suicide reduction, 187–8

 QTc, 132, 133, 146, **168**, 216
 qualitative studies, 30, 118,
 283, 285, 299
 quality of life, 9, 10, 11, 13, 15,
 17, 18, 124, 194, 201, 246
 aspects, 11
 quantitative analysis, 283, 299
 questionnaires, 129, 150, 163,
 189, 189, 236, 298, 304,
 305
 quetiapine, 132, 168, 178, 217,
 229, 273, 277, 281

 racial discrimination, 16
 radicalisation, 88
 rapid access clinics for older
 people (RACOPs), 287
 rational suicide, 195
 readmissions, 4, 38, 286
 refusal of treatment, 76
 rehabilitation, 1, 6, 39, 214,
 259, 260, 266, 283, 296
 renal function, 169, 239, 247
 repression, 210, 211
 resilience, 11, 20, 183, **185**
 resources, 9, 11, 20, 24, 26, 97,
 99, 140, 164, 166, 213,
 220, 293
 respect, 47, 50, 56, 60, 61, 65,
 67, 68, 96, 99
 responsible clinician, 80, 293
 restraint, 73
 definition (§6 of MCA
 2005), 73
 retirement, 9
 Richmond agitation sedating
 scale (RASS), 144
 risk assessment, 92, 94, **192**,
 194, 238, 259
 risk enablement, 18, 19
 risperidone, 132, 134, 148,
 168, 178, 217, 228, 229,
 276
 rivaroxaban, 253
 rivastigmine, 116, 148, 272,
 277

 rotigotine, 255, 272, 274
 Rowland universal dementia
 assessment scale
 (RUDAS)
 cognitive screening tool, 303
 languages, 303
 test-retest reliability, 303

 sadness, 161, 190
 safeguarding, 82–95, **193**, 194,
 233, 236, 242, 244, 295
 safeguarding adult boards, 94
 safeguarding legislation, 82
 safeguarding support, 94
 SafeLives (domestic violence
 helpline), 94
 schizoaffective disorder
 (manic type), 173
 schizophrenia, 30, 127, 220
 age-of-onset criterion, 224
 prevalence, **29**, 31
 prevalence), **28**
 Scottish Intercollegiate
 Guidelines Network
 (SIGN), 253
 delirium, 141, 142
 risk reduction poster, *147*
 secondary care, 2, 26, 150, 283,
 288
 secondary psychosis in late
 life, 220, 221–3
 affective disorder, 223
 alcohol misuse, 223
 definition, 221
 dementia and delirium,
 222–3
 illicit drug use in youth, 223
 integrated assessment, 222
 physical disease states, 221
 security, 11, 202, **211**
 selective noradrenaline
 reuptake inhibitors
 (SNRIs), 167, **168**, 217
 selective optimisation and
 compensation (Baltes
 and Baltes), 10, 214
 selective serotonin reuptake
 inhibitors (SSRIs), 133,
 160, 167, **168**, 193, 216
 side-effects, 216
 self-confidence, 18, 20, 214
 self-esteem, 15, 18, 61, 62, 214
 self-harm, 28, 76
 drug and alcohol misuse, ,
 238
 epidemiology, 187

 prevention (older people),
 183–96
 repetition and subsequent
 suicide, 187
 self-neglect, 12, 51, 82, **84**,
 157–8, 169, 202, 228, 243
 indicators and examples, **85**
 self-respect, 61, 62, 65, 67
 self-starvation, 186, 191
 senior friendly hospital (SFH),
 39
 sensory impairment, 46, 47,
 48, 88, 114, 143, 200, 225,
 226, 235, 300
 sepsis, 228, 273, 275
 serotonin auto receptors, 216
 serotonin syndrome, 160, 168,
 216, 277
 sertraline, 133, 216, 252
 serum ketones, 254
 services outside of hospital,
 292–6
 general practice, 294–5
 third sector (charity and
 voluntary services), 296
 sexual abuse: indicators and
 examples, **84**
 sexual disinhibition, 31, 126
 sexually inappropriate
 behaviour, 170
 Short Anxiety Screening Test
 (SAST), 209
 siblings, 211
 signage, 63
 single photon emission
 computed tomography
 (SPECT), 112, 115
 single question to identify
 delirium (SQID), 141, 145
 single rooms, 62
 sleep, 31, 39, 40, 60, 63, 64,
 106, 125, 126, 131, 133,
 170, 173, 177, 277, 305
 REM behaviour disorder,
 269, 272, 277, 280
 sleep disturbance, 133, 162
 sleep hygiene, 165, 177
 smoking, 242
 dementia risk factor, 113
 snoezelen: definition, 130
 social isolation, 10, 11, 16, 88,
 165, 202, 224, 225, 227,
 228, 238
 social networks, 34, **185**, 195,
 202, 236, 284
 social phobia, 216, 217

- social services, 55, 103
 Social Services and Well-Being Act (Wales, 2014), 82
 social support, 22, 38
 social workers, 191, 192, 194, 222, 295
 socioeconomic status, 16
 link with multimorbidity, 3
 sodium valproate, 174, 178, 277, 281
 somatic symptoms, 157, 161, 162, 165, 173, 177, 200
 somatoform disorders, 26, 28, 204, 207
 spice (synthetic cannabinoids), 240
 stem cells, 9, 250
 stigma, 89, 102, 157, 235
 stimulants, 172, 240
 stress, 117, 119, 161, 174, 200, 201
 stroke, 33, 47, 49, 53, 111, 113, 132, 159, 201, 240, 242, 256, 257, 258
 stroke risk, 146
 substance history, 208
 substance misuse, 170
 definition, 233
 prevalence rates (ED patients, UK), 28
 substantia nigra, 267
 successful ageing, 10
 suicidal ideation, 28, 188, 194
 risk factors, 183
 suicide, 11, 17, 32
 drug and alcohol misuse, 238
 epidemiology, 186–7
 indicated prevention (four high-risk groups), 191–6
 multidimensional, integrated approaches, 183
 postvention, 196
 precipitants (recent events), 183
 pre-existing factors, 183
 prevention (older people), 183–96
 primary prevention efforts, 187, 191
 public health measures, 187–8
 safer care in mental health services (key elements), 194
 secondary prevention (early intervention), 187
 social factors, 186
 suicide (high-risk groups), 191–6
 1. presenting with new suicidal ideas or actions, 191
 2. psychiatric inpatients, 191–5
 3. older adults with cognitive impairment, 195
 4. palliative care older patients, 195–6
 suicide methods, 186
 suicide notes, 191
 suicide risk, 216, 234
 five Ps approach, 183, 184
 questionnaire, 189
 syndrome of inappropriate antidiuretic hormone secretion (SIADH), 252
 TADA (tolerate, anticipate, don't agitate) approach, 40
 Ten Point Dignity Challenge, 61
 tetrahydrocannabinol (THC) potency, 239
 therapeutic lying, 52
 perspectives of people with dementia, 53
 reflective case, 53
 therapist and client relationship, 210
 thiamine (vitamin B1), 241
 This Is Me, 56, 67, 102, 146
 thromboembolic disease, 247, 260
 tinnitus, 221
 titration, 216
 Together in Dementia
 Everyday (TIDE), 106, 107
 tramadol, 235, 249
 transient ischaemic attacks (TIAs), 111, 246, 256
 translators, 102
 gender preference, 91
 trauma, 119, 166, 239, 247, 254
 trauma-focused CBT, 210
 treatment adjuvants, 167–9
 treatment regimens: ability to comply, 13
 triangle of care, 96
 tricyclics, 168, 217
 tumour necrosis factor alpha (TNF alpha), 259
 unified Parkinson's disease rating scale (UPDRS), 270
 unmet needs model of agitated behaviour, 127
 urinary tract infections, 113, 255
 variable rate intravenous insulin infusion (VRIII), 254, 255
 vascular dementia, 111, 113, 115, 117
 vascular Parkinsonism, 267, 280–1
 vasopressin receptor antagonists, 252
 venlafaxine, 133, 167, 168, 217
 venous thromboembolism (VTE), 248, 250, 253, 256, 273
 postoperative prevention (hip patients), 258
 ventricular failure, 256, 260
 verbal rating scale, 129
 very late onset schizophrenia-like psychosis (VLOSP), 221, 224–5, 229, 230
 ascertaining problem, 224
 ascertiveness of care, 224
 distinctive features, 225
 key differences versus DD and EOS, 226
 nosological uncertainty, 224
 personality problems merging into, 227
 presentations to liaison service, 228–9
 Victim Support (domestic violence helpline), 94
 visiting hours, 39, 41, 102
 visual analogue scale, 129
 visual impairment, 225, 275, 277
 voluntary services, 165, 296
 vortioxetine, 217
 waiting times, 27, 60
 wards
 changes, 47
 delirium cases, 141
 dementia-friendly design, 63

334	Index	
wards (cont.)		
environment, 20, 125,		
170		
factors promoting recovery,		
19		
managers, 77		
warfarin, 168, 248, 253, 253,		
284, 290		
well-being, 10, 82		
faith, 17		
hospital environment, 17		
racial discrimination, 16		
social support, 22		
	socio-economic status, 16	
	themes, 21	
	well-being (properties)	
	agency, 11	
	belonging, 11	
	enrichment, 11	
	integrity of other, 11	
	integrity of self, 11	
	security, 11	
	Wernicke’s Encephalopathy	
	(WE), 240	
	Wernicke-Korsakoff	
	syndrome, 238	
		Wessely review, 78, 81
		Whooley questions, 305
		wider community services:
		interface with liaison
		psychiatry services,
		292–7
		wish to hasten death
		(WTHD), 195
		Z drugs, 233, 234
		zolpidem, 178, 234
		zopiclone, 178, 234
		Zung depression scale, 304