

## LONDON MATHEMATICAL SOCIETY LECTURE NOTE SERIES

Managing Editor: Professor M. Reid, Mathematics Institute,  
 University of Warwick, Coventry CV4 7AL, United Kingdom

The titles below are available from booksellers, or from Cambridge University Press at  
<http://www.cambridge.org/mathematics>

- 325 Lectures on the Ricci flow, P. TOPPING
- 326 Modular representations of finite groups of Lie type, J.E. HUMPHREYS
- 327 Surveys in combinatorics 2005, B.S. WEBB (ed)
- 328 Fundamentals of hyperbolic manifolds, R. CANARY, D. EPSTEIN & A. MARDEN (eds)
- 329 Spaces of Kleinian groups, Y. MINSKY, M. SAKUMA & C. SERIES (eds)
- 330 Noncommutative localization in algebra and topology, A. RANICKI (ed)
- 331 Foundations of computational mathematics, Santander 2005, L.M. PARDO, A. PINKUS, E. SÜLI & M.J. TODD (eds)
- 332 Handbook of tilting theory, L. ANGELERI HÜGEL, D. HAPPEL & H. KRAUSE (eds)
- 333 Synthetic differential geometry (2nd Edition), A. KOCK
- 334 The Navier-Stokes equations, N. RILEY & P. DRAZIN
- 335 Lectures on the combinatorics of free probability, A. NICA & R. SPEICHER
- 336 Integral closure of ideals, rings, and modules, I. SWANSON & C. HUNEKE
- 337 Methods in Banach space theory, J.M.F. CASTILLO & W.B. JOHNSON (eds)
- 338 Surveys in geometry and number theory, N. YOUNG (ed)
- 339 Groups St Andrews 2005 & I, C.M. CAMPBELL, M.R. QUICK, E.F. ROBERTSON & G.C. SMITH (eds)
- 340 Groups St Andrews 2005 & II, C.M. CAMPBELL, M.R. QUICK, E.F. ROBERTSON & G.C. SMITH (eds)
- 341 Ranks of elliptic curves and random matrix theory, J.B. CONREY, D.W. FARMER, F. MEZZADRI & N.C. SNAITH (eds)
- 342 Elliptic cohomology, H.R. MILLER & D.C. RAVENEL (eds)
- 343 Algebraic cycles and motives I, J. NAGEL & C. PETERS (eds)
- 344 Algebraic cycles and motives II, J. NAGEL & C. PETERS (eds)
- 345 Algebraic and analytic geometry, A. NEEMAN
- 346 Surveys in combinatorics 2007, A. HILTON & J. TALBOT (eds)
- 347 Surveys in contemporary mathematics, N. YOUNG & Y. CHOI (eds)
- 348 Transcendental dynamics and complex analysis, P.J. RIPPON & G.M. STALLARD (eds)
- 349 Model theory with applications to algebra and analysis I, Z. CHATZIDAKIS, D. MACPHERSON, A. PILLAY & A. WILKIE (eds)
- 350 Model theory with applications to algebra and analysis II, Z. CHATZIDAKIS, D. MACPHERSON, A. PILLAY & A. WILKIE (eds)
- 351 Finite von Neumann algebras and masas, A.M. SINCLAIR & R.R. SMITH
- 352 Number theory and polynomials, J. MCKEE & C. SMYTH (eds)
- 353 Trends in stochastic analysis, J. BLATH, P. MÖRTERS & M. SCHEUTZOW (eds)
- 354 Groups and analysis, K. TENT (ed)
- 355 Non-equilibrium statistical mechanics and turbulence, J. CARDY, G. FALKOVICH & K. GAWEDZKI
- 356 Elliptic curves and big Galois representations, D. DELBOURGO
- 357 Algebraic theory of differential equations, M.A.H. MACCALLUM & A.V. MIKHAILOV (eds)
- 358 Geometric and cohomological methods in group theory, M.R. BRIDSON, PH. KROPHOLLER & I.J. LEARY (eds)
- 359 Moduli spaces and vector bundles, L. BRAMBILA-PAZ, S.B. BRADLOW, O. GARCÍA-PRADA & S. RAMANAN (eds)
- 360 Zariski geometries, B. ZILBER
- 361 Words: Notes on verbal width in groups, D. SEGAL
- 362 Differential tensor algebras and their module categories, R. BAUTISTA, L. SALMERÓN & R. ZUAZUA
- 363 Foundations of computational mathematics, Hong Kong 2008, F. CUCKER, A. PINKUS & M.J. TODD (eds)
- 364 Partial differential equations and fluid mechanics, J.C. ROBINSON & J.L. RODRIGO (eds)
- 365 Surveys in combinatorics 2009, S. HUCZYNSKA, J.D. MITCHELL & C.M. RONEY-DOUGAL (eds)
- 366 Highly oscillatory problems, B. ENGQUIST, A. FOKAS, E. HAIRER & A. ISERLES (eds)
- 367 Random matrices: High dimensional phenomena, G. BLOWER
- 368 Geometry of Riemann surfaces, F.P. GARDINER, G. GONZÁLEZ-DIEZ & C. KOUROUNIOTIS (eds)
- 369 Epidemics and rumours in complex networks, M. DRAIEF & L. MASSOULIÉ
- 370 Theory of  $p$ -adic distributions, S. ALBEVERIO, A.YU. KHRENNIKOV & V.M. SHELKOVICH
- 371 Conformal fractals, F. PRZYTYCKI & M. URBAŃSKI
- 372 Moonshine: The first quarter century and beyond, J. LEPOWSKY, J. MCKAY & M.P. TUIE (eds)
- 373 Smoothness, regularity and complete intersection, J. MAJADAS & A. G. RODICIO
- 374 Geometric analysis of hyperbolic differential equations: An introduction, S. ALINHAC
- 375 Triangulated categories, T. HOLM, P. JØRGENSEN & R. ROUQUIER (eds)
- 376 Permutation patterns, S. LINTON, N. RUŠKUC & V. VATTER (eds)
- 377 An introduction to Galois cohomology and its applications, G. BERTHUY
- 378 Probability and mathematical genetics, N. H. BINGHAM & C. M. GOLDIE (eds)
- 379 Finite and algorithmic model theory, J. ESPARZA, C. MICHAUX & C. STEINHORN (eds)
- 380 Real and complex singularities, M. MANOEL, M.C. ROMERO FUSTER & C.T.C. WALL (eds)
- 381 Symmetries and integrability of difference equations, D. LEVI, P. OLVER, Z. THOMOVA & P. WINTERNITZ (eds)

- 382 Forcing with random variables and proof complexity, J. KRAJÍČEK
- 383 Motivic integration and its interactions with model theory and non-Archimedean geometry I, R. CLUCKERS, J. NICAISE & J. SEBAG (eds)
- 384 Motivic integration and its interactions with model theory and non-Archimedean geometry II, R. CLUCKERS, J. NICAISE & J. SEBAG (eds)
- 385 Entropy of hidden Markov processes and connections to dynamical systems, B. MARCUS, K. PETERSEN & T. WEISSMAN (eds)
- 386 Independence-friendly logic, A.L. MANN, G. SANDU & M. SEVENSTER
- 387 Groups St Andrews 2009 & in Bath I, C.M. CAMPBELL *et al* (eds)
- 388 Groups St Andrews 2009 & in Bath II, C.M. CAMPBELL *et al* (eds)
- 389 Random fields on the sphere, D. MARINUCCI & G. PECCATI
- 390 Localization in periodic potentials, D.E. PELINOVSKY
- 391 Fusion systems in algebra and topology, M. ASCHBACHER, R. KESSAR & B. OLIVER
- 392 Surveys in combinatorics 2011, R. CHAPMAN (ed)
- 393 Non-abelian fundamental groups and Iwasawa theory, J. COATES *et al* (eds)
- 394 Variational problems in differential geometry, R. BIELAWSKI, K. HOUSTON & M. SPEIGHT (eds)
- 395 How groups grow, A. MANN
- 396 Arithmetic differential operators over the  $p$ -adic integers, C.C. RALPH & S.R. SIMANCA
- 397 Hyperbolic geometry and applications in quantum chaos and cosmology, J. BOLTE & F. STEINER (eds)
- 398 Mathematical models in contact mechanics, M. SOFONEA & A. MATEI
- 399 Circuit double cover of graphs, C.-Q. ZHANG
- 400 Dense sphere packings: a blueprint for formal proofs, T. HALES
- 401 A double Hall algebra approach to affine quantum Schur-Weyl theory, B. DENG, J. DU & Q. FU
- 402 Mathematical aspects of fluid mechanics, J.C. ROBINSON, J.L. RODRIGO & W. SADOWSKI (eds)
- 403 Foundations of computational mathematics, Budapest 2011, F. CUCKER, T. KRICK, A. PINKUS & A. SZANTO (eds)
- 404 Operator methods for boundary value problems, S. HASSI, H.S.V. DE SNOO & F.H. SZAFRANIEC (eds)
- 405 Torsors, étale homotopy and applications to rational points, A.N. SKOROBOGATOV (ed)
- 406 Appalachian set theory, J. CUMMINGS & E. SCHIMMERLING (eds)
- 407 The maximal subgroups of the low-dimensional finite classical groups, J.N. BRAY, D.F. HOLT & C.M. RONEY-DOUGAL
- 408 Complexity science: the Warwick master's course, R. BALL, V. KOLOKOLTSOV & R.S. MACKAY (eds)
- 409 Surveys in combinatorics 2013, S.R. BLACKBURN, S. GERKE & M. WILDON (eds)
- 410 Representation theory and harmonic analysis of wreath products of finite groups, T. CECCHERINI-SILBERSTEIN, F. SCARABOTTI & F. TOLLI
- 411 Moduli spaces, L. BRAMBILA-PAZ, O. GARCÍA-PRADA, P. NEWSTEAD & R.P. THOMAS (eds)
- 412 Automorphisms and equivalence relations in topological dynamics, D.B. ELLIS & R. ELLIS
- 413 Optimal transportation, Y. OLLIVIER, H. PAJOT & C. VILLANI (eds)
- 414 Automorphic forms and Galois representations I, F. DIAMOND, P.L. KASSAEI & M. KIM (eds)
- 415 Automorphic forms and Galois representations II, F. DIAMOND, P.L. KASSAEI & M. KIM (eds)
- 416 Reversibility in dynamics and group theory, A.G. O'FARRELL & I. SHORT
- 417 Recent advances in algebraic geometry, C.D. HACON, M. MUSTAŢĂ & M. POPA (eds)
- 418 The Bloch-Kato conjecture for the Riemann zeta function, J. COATES, A. RAGHURAM, A. SAIKIA & R. SUJATHA (eds)
- 419 The Cauchy problem for non-Lipschitz semi-linear parabolic partial differential equations, J.C. MEYER & D.J. NEEDHAM
- 420 Arithmetic and geometry, L. DIEULEFAIT *et al* (eds)
- 421 O-minimality and Diophantine geometry, G.O. JONES & A.J. WILKIE (eds)
- 422 Groups St Andrews 2013, C.M. CAMPBELL *et al* (eds)
- 423 Inequalities for graph eigenvalues, Z. STANIĆ
- 424 Surveys in combinatorics 2015, A. CZUMAJ *et al* (eds)
- 425 Geometry, topology and dynamics in negative curvature, C.S. ARAVINDA, F.T. FARRELL & J.-F. LAFONT (eds)
- 426 Lectures on the theory of water waves, T. BRIDGES, M. GROVES & D. NICHOLLS (eds)
- 427 Recent advances in Hodge theory, M. KERR & G. PEARLSTEIN (eds)
- 428 Geometry in a Fréchet context, C. T. J. DODSON, G. GALANIS & E. VASSILIOU
- 429 Sheaves and functions modulo  $p$ , L. TAEMLAN
- 430 Recent progress in the theory of the Euler and Navier-Stokes equations, J.C. ROBINSON, J.L. RODRIGO, W. SADOWSKI & A. VIDAL-LÓPEZ (eds)
- 431 Harmonic and subharmonic function theory on the real hyperbolic ball, M. STOLL
- 432 Topics in graph automorphisms and reconstruction (2nd Edition), J. LAURI & R. SCAPELLATO
- 433 Regular and irregular holonomic D-modules, M. KASHIWARA & P. SCHAPIRA
- 434 Analytic semigroups and semilinear initial boundary value problems (2nd Edition), K. TAIRA
- 435 Graded rings and graded Grothendieck groups, R. HAZRAT
- 436 Groups, graphs and random walks, T. CECCHERINI-SILBERSTEIN, M. SALVATORI & E. SAVA-HUSS (eds)
- 437 Dynamics and analytic number theory, D. BADZIAHIN, A. GORODNIK & N. PEYERIMHOFF (eds)
- 438 Random walks and heat kernels on graphs, M.T. BARLOW
- 439 Evolution equations, K. AMMARI & S. GERBI (eds)
- 440 Surveys in combinatorics 2017, A. CLAESON *et al* (eds)
- 441 Polynomials and the mod 2 Steenrod algebra I, G. WALKER & R.M.W. WOOD
- 442 Polynomials and the mod 2 Steenrod algebra II, G. WALKER & R.M.W. WOOD
- 443 Asymptotic analysis in general relativity, T. DAUDÉ, D. HÄFNER & J.-P. NICOLAS (eds)
- 444 Geometric and cohomological group theory, P.H. KROPHOLLER, I.J. LEARY, C. MARTÍNEZ-PÉREZ & B.E.A. NUCINKIS (eds)

Cambridge University Press  
978-1-108-41448-7 — Polynomials and the mod 2 Steenrod Algebra  
Grant Walker, Reginald M. W. Wood  
Frontmatter  
[More Information](#)

London Mathematical Society Lecture Note Series: 441

# Polynomials and the mod 2 Steenrod Algebra

Volume 1: The Peterson Hit Problem

GRANT WALKER  
*University of Manchester*

REGINALD M. W. WOOD  
*University of Manchester*



Cambridge University Press  
978-1-108-41448-7 – Polynomials and the mod 2 Steenrod Algebra  
Grant Walker, Reginald M. W. Wood  
Frontmatter  
[More Information](#)

## CAMBRIDGE UNIVERSITY PRESS

University Printing House, Cambridge CB2 8BS, United Kingdom  
One Liberty Plaza, 20th Floor, New York, NY 10006, USA  
477 Williamstown Road, Port Melbourne, VIC 3207, Australia  
4843/24, 2nd Floor, Ansari Road, Daryaganj, Delhi – 110002, India  
79 Anson Road, #06-04/06, Singapore 079906

Cambridge University Press is part of the University of Cambridge.

It furthers the University's mission by disseminating knowledge in the pursuit of education, learning, and research at the highest international levels of excellence.

[www.cambridge.org](http://www.cambridge.org)  
Information on this title: [www.cambridge.org/9781108414487](http://www.cambridge.org/9781108414487)  
DOI: 10.1017/9781108333368

© Grant Walker and Reginald M. W. Wood 2018

This publication is in copyright. Subject to statutory exception and to the provisions of relevant collective licensing agreements, no reproduction of any part may take place without the written permission of Cambridge University Press.

First published 2018

Printed in the United Kingdom by Clays, St Ives plc

*A catalogue record for this publication is available from the British Library.*

ISBN – 2 Volume Set 978-1-108-41406-7 Paperback  
ISBN – Volume 1 978-1-108-41448-7 Paperback  
ISBN – Volume 2 978-1-108-41445-6 Paperback

Cambridge University Press has no responsibility for the persistence or accuracy of URLs for external or third-party internet Web sites referred to in this publication, and does not guarantee that any content on such websites is, or will remain, accurate or appropriate.

## Contents of Volume 1

---

<i>Contents of Volume 2</i>	<i>page xi</i>
<i>Preface</i>	xv
<b>1 Steenrod squares and the hit problem</b>	<b>1</b>
1.0 Introduction	1
1.1 The total square Sq	2
1.2 The action of matrices on $P(n)$	4
1.3 Some properties of $Sq^k$	7
1.4 The hit problem	8
1.5 Spikes	12
1.6 The Kameko maps for $P(2)$	12
1.7 The duplication map for $P(2)$	14
1.8 The hit problem for $P(2)$	15
1.9 Remarks	17
<b>2 Conjugate Steenrod squares</b>	<b>19</b>
2.0 Introduction	19
2.1 Power series	19
2.2 The total conjugate square $Xq$	22
2.3 Some properties of $Xq^k$	23
2.4 The numerical function $\mu$	26
2.5 The $\chi$ -trick and the Peterson conjecture	28
2.6 Catalan numbers and $Xq^k$	30
2.7 Remarks	31
<b>3 The Steenrod algebra <math>A_2</math></b>	<b>33</b>
3.0 Introduction	33
3.1 The Adem relations	34

3.2	The action of $A_2$ on $P(n)$	37
3.3	Symmetric functions and blocks	40
3.4	The admissible basis of $A_2$	44
3.5	The Milnor basis of $A_2$	46
3.6	Remarks	52
<b>4</b>	<b>Products and conjugation in <math>A_2</math></b>	<b>54</b>
4.0	Introduction	54
4.1	The Milnor product formula	54
4.2	The Bullett–Macdonald identity	59
4.3	Conjugation in $A_2$	61
4.4	Conjugation and the Milnor basis	64
4.5	Remarks	66
<b>5</b>	<b>Combinatorial structures</b>	<b>67</b>
5.0	Introduction	67
5.1	Sequences, blocks and partitions	69
5.2	Dominance	74
5.3	Binary partitions	76
5.4	Minimal spikes	79
5.5	Spike partitions	82
5.6	Maximal spikes	84
5.7	Dominance in $A_2$	86
5.8	The excess function on $A_2$	91
5.9	Remarks	93
<b>6</b>	<b>The cohit module <math>Q(n)</math></b>	<b>96</b>
6.0	Introduction	96
6.1	Steenrod and matrix actions on blocks	97
6.2	Filtrations of $Q^d(n)$	100
6.3	Concatenation of blocks	103
6.4	Splicing	107
6.5	The Kameko maps	109
6.6	The duplication map	111
6.7	Tail sequences	114
6.8	Head sequences	116
6.9	Tail and head modules	118
6.10	Remarks	122
<b>7</b>	<b>Bounds for <math>\dim Q^d(n)</math></b>	<b>123</b>
7.0	Introduction	123
7.1	The boundedness theorem	124

*Contents of Volume 1*

vii

7.2	Combinatorial splicing	127
7.3	Semi-standard blocks	132
7.4	Young tableaux	135
7.5	Remarks	138
<b>8</b>	<b>Special blocks and a basis for <math>Q(3)</math></b>	139
8.0	Introduction	139
8.1	Blocks with large head	139
8.2	Blocks with long tail	146
8.3	A monomial basis of $Q^d(3)$	151
8.4	Blocks of head length $n - 1$	159
8.5	Remarks	162
<b>9</b>	<b>The dual of the hit problem</b>	164
9.0	Introduction	164
9.1	The divided power algebra $DP(n)$	165
9.2	The action of matrices on $DP(n)$	168
9.3	Down squares $Sq_k$	170
9.4	The Steenrod kernel $K(n)$	174
9.5	Dual Kameko and duplication maps	177
9.6	Tail and head sequences for $DP(n)$	180
9.7	Dual tail modules	182
9.8	Dual head modules	184
9.9	$K^d(3)$ for $d = 2^s - 1$	186
9.10	Remarks	187
<b>10</b>	<b><math>K(3)</math> and <math>Q(3)</math> as <math>\mathbb{F}_2GL(3)</math>-modules</b>	189
10.0	Introduction	189
10.1	Some irreducible $\mathbb{F}_2GL(n)$ -modules	191
10.2	Transpose duality	193
10.3	$K^d(3)$ for $d = 2^{s+1}$	195
10.4	$K^d(3)$ for $d = (2^{s+t} - 1) + (2^t - 1)$	196
10.5	The flag module $FL(3)$	198
10.6	$K^d(3)$ and $Q^d(3)$ as $\mathbb{F}_2GL(3)$ -modules	203
10.7	Submodules of $Q^d(3)$	206
10.8	Remarks	210
<b>11</b>	<b>The dual of the Steenrod algebra</b>	211
11.0	Introduction	211
11.1	Hopf algebras	212
11.2	The coproduct $A_2 \rightarrow A_2 \otimes A_2$	218
11.3	The Hopf algebra structure of $A_2$	221

11.4	Duality for Hopf algebras	223
11.5	The dual algebra $A_2^*$	226
11.6	Conjugation in the Milnor basis	229
11.7	Remarks	233
<b>12</b>	<b>Further structure of <math>A_2</math></b>	234
12.0	Introduction	234
12.1	Some conjugation formulae	235
12.2	The Milnor elements $Sq_t^k$	237
12.3	The subalgebras $A_q$ , $q = 2^f$	240
12.4	The subalgebras $A_2(\lambda)$	243
12.5	2-atomic numbers and bases	246
12.6	The Arnon C basis	251
12.7	Remarks	252
<b>13</b>	<b>Stripping and nilpotence in <math>A_2</math></b>	254
13.0	Introduction	254
13.1	Cap products for Hopf algebras	255
13.2	The action of $A_2^*$ on $A_2$	257
13.3	The halving map of $A_2$	260
13.4	The odd subalgebra $Od$	262
13.5	Nilpotence of $Sq^{2^s}$	266
13.6	Relations in $A_q$	269
13.7	Nilpotence of $P_t^s$	275
13.8	Remarks	278
<b>14</b>	<b>The 2-dominance theorem</b>	279
14.0	Introduction	279
14.1	The excess theorem	280
14.2	Stripping and conjugation for $\mu(d) = 1$	282
14.3	The numerical function $\gamma$	285
14.4	Stripping and conjugation for general $d$	287
14.5	Proof of the excess theorem	292
14.6	$Xq^{[k;d]}$ in the Milnor basis	294
14.7	Remarks	296
<b>15</b>	<b>Invariants and the hit problem</b>	297
15.0	Introduction	297
15.1	Dickson invariants	298
15.2	The hit problem for $D(2)$	302
15.3	The action of $A_2$ on Dickson invariants	303
15.4	The hit problem for $P(2)^{H_2}$	308



<i>Contents of Volume 1</i>	ix
15.5 The hit problem for $P(3)^{H_3}$	311
15.6 The relative hit problem for $D(3)$	316
15.7 Elementary Dickson monomials	318
15.8 The relative hit problem for $D(n)$	320
15.9 Remarks	322
<i>Bibliography</i>	325
<i>Index of Notation for Volume 1</i>	337
<i>Index for Volume 1</i>	339
<i>Index of Notation for Volume 2</i>	343
<i>Index for Volume 2</i>	345

## Contents of Volume 2

---

<i>Contents of Volume 1</i>	<i>page ix</i>
<i>Preface</i>	xv
<b>16 The action of <math>GL(n)</math> on flags</b>	1
16.0 Introduction	1
16.1 Permutation matrices	2
16.2 The Bruhat decomposition of $GL(n)$	8
16.3 The flag module $FL(n)$	12
16.4 Partial flags and parabolic subgroups	14
16.5 The direct sum decomposition of $FL(n)$	20
16.6 The structure of $FL(3)$	23
16.7 Remarks	27
<b>17 Irreducible <math>\mathbb{F}_2GL(n)</math>-modules</b>	29
17.0 Introduction	29
17.1 The Hecke algebra $H_0(n)$	31
17.2 The Schubert monoid $Sch(n)$	33
17.3 The permutation matrices $W_0^I$	36
17.4 Generators of the summands of $FL(n)$	38
17.5 The submodule $s_I(FL(n))$ of $FL_I(n)$	43
17.6 Irreducible submodules of $FL(n)$	45
17.7 Remarks	49
<b>18 Idempotents and characters</b>	53
18.0 Introduction	53
18.1 Idempotents and direct sums	55
18.2 Splitting fields and conjugacy classes	58
18.3 The Steinberg idempotents $e(n)$ and $e'(n)$	60

18.4	Embedding $H_0(n)$ in $\mathbb{F}_2\text{GL}(n)$	63
18.5	Brauer characters	67
18.6	The representation ring $R_2(\text{GL}(n))$	70
18.7	Remarks	72
<b>19</b>	<b>Splitting <math>P(n)</math> as an <math>A_2</math>-module</b>	<b>75</b>
19.0	Introduction	75
19.1	The $A_2$ -modules $P(n, \lambda)$	76
19.2	Poincaré series of $P(n, \lambda)$	80
19.3	The coinvariant algebra $C(n)$	84
19.4	$B(n)$ -invariants and irreducible modules	87
19.5	Irreducible submodules of $P(n)$	91
19.6	Remarks	93
<b>20</b>	<b>The algebraic group <math>\overline{G}(n)</math></b>	<b>95</b>
20.0	Introduction	95
20.1	Polynomial representations of $\overline{G}(n)$	96
20.2	The $\overline{G}(n)$ -module $\overline{P}(n)$	100
20.3	Weyl modules	104
20.4	Weyl modules and flag modules	106
20.5	Weyl modules and the hit problem	109
20.6	First occurrences of irreducibles in $P(n)$	111
20.7	Remarks	113
<b>21</b>	<b>Endomorphisms of <math>P(n)</math> over <math>A_2</math></b>	<b>114</b>
21.0	Introduction	114
21.1	Embedding $\mathbb{F}_2M(m, n)$ in $\text{Hom}_{A_2}(P(m), P(n))$	116
21.2	Detecting squares and Dickson invariants	116
21.3	Kernels of squaring operations	119
21.4	Root generation	121
21.5	Key monomials	124
21.6	Values on key monomials	127
21.7	Examples	132
21.8	Remarks	137
<b>22</b>	<b>The Steinberg summands of <math>P(n)</math></b>	<b>138</b>
22.0	Introduction	138
22.1	Steenrod operations on rational functions	140
22.2	The cyclic $A_2$ -module $T(n)$	143
22.3	The Mitchell–Priddy module $MP(n)$	146
22.4	The hit problem for $MP(n)$	150
22.5	The symmetric Steinberg summand	152

<i>Contents of Volume 2</i>		xiii
22.6	The $B(n)$ -invariant Steinberg summand	153
22.7	Remarks	156
<b>23</b>	<b>The <math>d</math>-spike module <math>J(n)</math></b>	<b>158</b>
23.0	Introduction	158
23.1	Flag $d$ -polynomials	160
23.2	Zip $d$ -monomials	163
23.3	Permutation tableaux	167
23.4	$J^d(n)$ for 1-dominant degrees	171
23.5	Zip monomials and $Q^d(n)$	175
23.6	Remarks	179
<b>24</b>	<b>Partial flags and <math>J(n)</math></b>	<b>180</b>
24.0	Introduction	180
24.1	$d$ -spikes and $d$ -duplication	182
24.2	The Grassmannian case	184
24.3	Zip $d$ -monomials for $FL_J(n)$	187
24.4	Lower bounds for $J^\omega(4)$	194
24.5	Counterexamples to Kameko's conjecture	202
24.6	Remarks	203
<b>25</b>	<b>The symmetric hit problem</b>	<b>204</b>
25.0	Introduction	204
25.1	The hit problem for permutation groups	205
25.2	The hit problem for $S(2)$	209
25.3	The hit problem for $S(3)$	211
25.4	The case $n = 3$ , $\mu(d) = 1$	214
25.5	The Wu formula	217
25.6	Remarks	218
<b>26</b>	<b>The dual of the symmetric hit problem</b>	<b>220</b>
26.0	Introduction	220
26.1	The dual algebra $DS$	221
26.2	The symmetric Steenrod kernel $K(DS(3))$	227
26.3	The case $n = 3$ , $\mu(d) = 1$	231
26.4	The bigraded Steenrod algebra $\tilde{A}_2$	235
26.5	Remarks	239
<b>27</b>	<b>The cyclic splitting of <math>P(n)</math></b>	<b>240</b>
27.0	Introduction	240
27.1	The twisted $A_2$ -module $\tilde{P}(n)$	242
27.2	The cyclic splitting and blocks	246

27.3	The Kameko map for $\tilde{P}(n)$	249
27.4	Twisted tails and heads	252
27.5	The twisted cohits $\tilde{Q}^\omega(3)$	255
27.6	Remarks	263
<b>28</b>	<b>The cyclic splitting of <math>DP(n)</math></b>	264
28.0	Introduction	264
28.1	The twisted $A_2$ -module $\tilde{DP}(n)$	264
28.2	Relations in $\tilde{J}(n)$	269
28.3	The Kameko map for $\tilde{DP}(n)$	275
28.4	Calculation of $\tilde{K}(2)$	280
28.5	Calculation of $\tilde{K}(3)$	281
28.6	Remarks	285
<b>29</b>	<b>The 4-variable hit problem, I</b>	286
29.0	Introduction	286
29.1	$Q^d(4)$ and $K^d(4)$ for $d = 7, 8$ and $16$	286
29.2	Non-decreasing $\omega$ -sequences	293
29.3	A basis for $Q^\omega(4)$ , $\omega = (2, \dots, 2)$	297
29.4	Remarks	306
<b>30</b>	<b>The 4-variable hit problem, II</b>	307
30.0	Introduction	307
30.1	The large body case $t \geq 4$	308
30.2	Strongly spike-free polynomials	315
30.3	The spike-free modules $SF^\omega(4)$	316
30.4	$Q^\omega(4)$ and $K^\omega(4)$ for decreasing $\omega$	329
30.5	Remarks	332
	<i>Bibliography</i>	335
	<i>Index of Notation for Volume 2</i>	347
	<i>Index for Volume 2</i>	349
	<i>Index of Notation for Volume 1</i>	351
	<i>Index for Volume 1</i>	353

## Preface

---

This book is about the mod 2 Steenrod algebra  $A_2$  and its action on the polynomial algebra  $P(n) = \mathbb{F}_2[x_1, \dots, x_n]$  in  $n$  variables, where  $\mathbb{F}_2$  is the field of two elements. Polynomials are graded by degree, so that  $P^d(n)$  is the set of homogeneous polynomials of degree  $d$ . Although our subject has its origin in the work of Norman E. Steenrod in algebraic topology, we have taken an algebraic point of view. We have tried as far as possible to provide a self-contained treatment based on linear algebra and representations of finite matrix groups. In other words, the reader does not require knowledge of algebraic topology, although the subject has been developed by topologists and is motivated by problems in topology.

There are many bonuses for working with the prime  $p = 2$ . There are no coefficients to worry about, so that every polynomial can be written simply as a sum of monomials. We use a matrix-like array of 0s and 1s, which we call a ‘block’, to represent a monomial in  $P(n)$ , where the rows of the block are formed by the reverse binary expansions of its exponents. Thus a polynomial is a set of blocks, and the sum of two polynomials is the symmetric difference of the corresponding sets. Using block notation, the action of  $A_2$  on  $P(n)$  can be encoded in computer algebra programs using standard routines on sets, lists and arrays. In addition, much of the literature on the Steenrod algebra and its applications in topology concentrates on the case  $p = 2$ . Often a result for  $p = 2$  has later been extended to all primes, but there are some results where no odd prime analogue is known.

We begin in Chapter 1 with the algebra map  $Sq : P(n) \rightarrow P(n)$  defined on the generators by  $Sq(x_i) = x_i + x_i^2$ . The map  $Sq$  is the total Steenrod squaring operation, and the Steenrod squares  $Sq^k : P^d(n) \rightarrow P^{d+k}(n)$  are its graded parts. The linear operations  $Sq^k$  can be calculated using induction on degree and the Cartan formula  $Sq^k(fg) = \sum_{i+j=k} Sq^i(f)Sq^j(g)$ , which is equivalent to

the multiplicative property of Sq. A general Steenrod operation is a sum of compositions of Steenrod squares.

The multiplicative monoid  $M(n)$  of  $n \times n$  matrices over  $\mathbb{F}_2$  acts on the right of  $P(n)$  by linear substitution of the variables. Thus  $P^d(n)$  gives a representation over  $\mathbb{F}_2$  of  $M(n)$  and of  $GL(n)$ , the general linear group of invertible matrices. This matrix action commutes with the action of the Steenrod squares, and the interplay between the two gives rise to a host of interesting algebraic problems.

One of these, the ‘hit’ problem, is a constant theme here. A polynomial  $f$  is ‘hit’ if there are polynomials  $f_k$  such that  $f = \sum_{k>0} Sq^k(f_k)$ . The hit polynomials form a graded subspace  $H(n)$  of  $P(n)$ , and the basic problem is to find the dimension of the quotient space  $Q^d(n) = P^d(n)/H^d(n)$ . We call  $Q(n)$  the space of ‘cohits’. Since  $P(n)$  is spanned by monomials,  $Q(n)$  is spanned by their equivalence classes, which we refer to simply as ‘monomials in  $Q(n)$ ’. A monomial whose exponents are integers of the form  $2^j - 1$  is called a ‘spike’, and cannot appear as a term in a hit polynomial. It follows that a monomial basis for  $Q(n)$  must include all the spikes.

At a deeper level, the hit problem concerns the structure of  $Q(n)$  as a representation of  $GL(n)$  or  $M(n)$ . We develop the tools needed to answer this in the 1- and 2-variable cases in Chapter 1. These include the 2-variable version of the maps introduced by Masaki Kameko to solve the 3-variable case, and a map which we call the duplication map. We hope that this opening chapter is accessible to graduate students and mathematicians with little or no background in algebraic topology, and that it will serve as an appetizer for the rest of the book.

Chapter 2 introduces a second family of Steenrod operations, the conjugate Steenrod squares  $Xq^k : P^d(n) \rightarrow P^{d+k}(n)$ . These are useful in the hit problem because of a device known as the ‘ $\chi$ -trick’. This states that the product of  $f$  and  $Xq^k(g)$  is hit if and only if the product of  $Xq^k(f)$  and  $g$  is hit. We use the  $\chi$ -trick to prove that  $Q^d(n) = 0$  if and only if  $\mu(d) > n$ , where  $\mu(d)$  is the smallest number of integers of the form  $2^j - 1$  (with repetitions allowed) whose sum is  $d$ . This establishes the 1986 conjecture of Franklin P. Peterson which first stimulated interest in the hit problem.

Here is a rough guide to the structure of the rest of the book, in terms of three main themes: the Steenrod algebra  $A_2$ , the Peterson hit problem, and matrix representations. Volume 1 contains Chapters 1 to 15, and Volume 2 contains Chapters 16 to 30.

Chapters 3 to 5 develop  $A_2$  from an algebraic viewpoint.

Chapters 6 to 10 provide general results on the hit problem, together with a detailed solution for the 3-variable case.

Chapters 11 to 14 introduce the Hopf algebra structure of  $A_2$  and study its structure in greater depth.

Chapter 15 introduces the theme of modular representations by relating the hit problem to invariants and the Dickson algebra.

Chapters 16 to 20 develop the representation theory of  $GL(n, \mathbb{F}_2)$  via its action on ‘flags’, or increasing sequences of subspaces, in an  $n$ -dimensional vector space  $V(n)$  over  $\mathbb{F}_2$ .

Chapter 21 explores a fundamental relation between linear maps  $V(m) \rightarrow V(n)$  and Steenrod operations, leading to a maximal splitting of  $P(n)$  as a direct sum of  $A_2$ -modules.

Chapter 22 studies the  $A_2$ -summands of  $P(n)$  corresponding to the Steinberg representation of  $GL(n, \mathbb{F}_2)$ .

Chapters 23 and 24 develop the relation between flag modules and the dual hit problem.

Chapters 25 and 26 study the hit problem for symmetric polynomials over  $\mathbb{F}_2$ .

Chapters 27 and 28 study the splitting of  $P(n)$  as an  $A_2$ -module obtained using a cyclic subgroup of order  $2^n - 1$  in  $GL(n, \mathbb{F}_2)$ .

Chapters 29 and 30 return to Peterson’s original problem, with a partial solution of the 4-variable case.

The contents of Chapters 3 to 30 are summarized below in more detail.

In Chapter 3 we interpret the operations  $Sq^k$  as generators of a graded algebra  $A_2$ , subject to a set of relations called the Adem relations. The algebra  $A_2$  is the mod 2 Steenrod algebra, and the operations  $Sq^k$  of Chapter 1 provide  $P(n)$  with the structure of a left  $A_2$ -module. If  $f = \sum_{k>0} Sq^k(f_k)$ , then  $f$  can be reduced to a set of polynomials of lower degree modulo the action of the positively graded part  $A_2^+$  of  $A_2$ . A monomial basis for  $Q(n)$  gives a minimal generating set for  $P(n)$  as an  $A_2$ -module. Thus the hit problem is an example of the general question of finding a minimal generating set for a module over a ring. The structure of  $A_2$  itself is completely determined by its action on polynomials, in the sense that two expressions in the generators  $Sq^k$  are equal in  $A_2$  if and only if the corresponding operations on  $P(n)$  are equal for all  $n$ . For example, the results  $Sq^1 Sq^{2k}(f) = Sq^{2k+1}(f)$  and  $Sq^1 Sq^{2k+1}(f) = 0$  of Chapter 1 imply the Adem relations  $Sq^1 Sq^{2k} = Sq^{2k+1}$  and  $Sq^1 Sq^{2k+1} = 0$ .

In Chapter 3 we also establish the two most important bases for  $A_2$  as a vector space over  $\mathbb{F}_2$ . These are the admissible monomials in the generators  $Sq^k$ , due to Henri Cartan and Jean-Pierre Serre, and the basis introduced by John W. Milnor by treating  $A_2$  as a Hopf algebra. As mentioned above,



we represent a monomial by a ‘block’ whose rows are the reversed binary expansions of its exponents, and whose entries are integers 0 or 1. We use blocks to keep track of Steenrod operations on monomials. This ‘block technology’ and ‘digital engineering’ works well for the prime 2, and greatly facilitates our understanding of techniques which can appear opaque when expressed in more standard notation.

Chapter 4 begins with the multiplication formula for elements of the Milnor basis. This combinatorial formula helps to explain the ubiquity of the Milnor basis in the literature, as a product formula is not available for other bases of  $A_2$ . We also discuss the compact formulation of the Adem relations due to Shaun R. Bullett and Ian G. Macdonald. We use this to construct the conjugation  $\chi$  of  $A_2$ , which interchanges  $Sq^k$  and  $Xq^k$ .

Chapter 5 provides combinatorial background for the algebra  $A_2$ , the hit problem and the representation theory of  $GL(n)$  over  $\mathbb{F}_2$ . Sequences of non-negative integers appear in various forms, and we distinguish ‘finite sequences’ from ‘sequences’. A ‘finite sequence’ has a fixed number of entries, called its ‘size’, while a ‘sequence’ is an infinite sequence  $R = (r_1, r_2, \dots)$  with only a finite number of nonzero terms, whose ‘length’ is the largest  $\ell$  for which  $r_\ell > 0$ . However, a sequence  $R$  is usually written as a finite sequence  $(r_1, \dots, r_n)$ , where  $n \geq \ell$ , by suppressing some or all of the trailing 0s. The modulus of a sequence or a finite sequence is the sum of its terms. For example, the degree of a monomial is the modulus of its sequence of exponents. The set of all sequences indexes the Milnor basis of  $A_2$ .

For brevity, we call a sequence  $R$  ‘decreasing’ if  $r_i \geq r_{i+1}$  for all  $i$ , i.e. if it is non-increasing or weakly decreasing. Thus a decreasing sequence of modulus  $d$  is a partition of  $d$ . Such a partition can alternatively be regarded as a multiset of positive integers with sum  $d$ . We discuss two special types of partition; ‘binary’ partitions, whose parts are integers of the form  $2^j$ , and ‘spike’ partitions, whose parts are integers of the form  $2^j - 1$ .

We introduce two total order relations on sequences, the left (lexicographic) order and the right (reversed lexicographic) order, and two partial order relations, dominance and 2-dominance. The  $\omega$ -sequence  $\omega(f) = (\omega_1, \omega_2, \dots, \omega_k)$  of a monomial  $f$  is defined by writing  $f$  as a product  $f_1 f_2^2 \cdots f_k^{2^{k-1}}$ , where  $f_i$  is a product of  $\omega_i$  distinct variables. In terms of blocks,  $\omega(f)$  is the sequence of column sums of the block representing  $f$ , and the degree of  $f$  is  $\omega_1 + 2\omega_2 + 4\omega_3 + \cdots + 2^{k-1}\omega_k$ , the ‘2-degree’ of  $\omega(f)$ . The set of decreasing sequences of 2-degree  $d$  has a minimum element  $\omega^{\min}(d)$ , which plays an important part in the hit problem, and is the same for the left, right and 2-dominance orders. We end Chapter 5 by relating this combinatorial material to the admissible and Milnor bases of  $A_2$ .

In Chapter 6 we return to the hit problem and introduce ‘local’ cohit spaces  $Q^\omega(n)$ . A total order relation on  $\omega$ -sequences of monomials gives a filtration on  $P^d(n)$  with quotients  $P^\omega(n)$ . For the left and right orders, this passes to a filtration on  $Q^d(n)$  with quotients  $Q^\omega(n)$ . A polynomial  $f$  in  $P^\omega(n)$  is ‘left reducible’ if it is the sum of a hit polynomial and monomials with  $\omega$ -sequences  $< \omega$  in the left order, and similarly for the right order. We prove the theorem of William M. Singer that  $Q^\omega(n) = 0$  if  $\omega < \omega^{\min}(d)$  in the left order. We introduce the ‘splicing’ technique for manufacturing hit equations, extend the Kameko and duplication maps of Chapter 1 to the  $n$ -variable case, and determine  $Q^\omega(n)$  for ‘head’ sequences  $\omega = (n-1, \dots, n-1)$  and ‘tail’ sequences  $\omega = (1, \dots, 1)$ .

We begin Chapter 7 by proving that  $\dim Q^d(n)$  is bounded by a function of  $n$  independent of  $d$ . Thus only finitely many isomorphism classes of  $\mathbb{F}_2\text{GL}(n)$ -modules can be realized as cohit modules  $Q^d(n)$ . We extend splicing techniques and show that  $Q^\omega(n) = 0$  if  $\omega$  is greater than every decreasing  $\omega$ -sequence in the left order. A correspondence between blocks with decreasing  $\omega$ -sequences and Young tableaux is used to define ‘semi-standard’ blocks (or monomials), and we show that  $Q^\omega(n)$  is spanned by such blocks when  $\omega = \omega^{\min}(d)$ .

In Chapter 8, we obtain reduction theorems for  $Q^\omega(n)$  when the sequence  $\omega$  has a ‘head’ of length  $\geq n-1$  or a ‘tail’ of length  $\geq n$ . It follows that  $\dim Q^d(n) = \prod_{i=1}^n (2^i - 1)$  for degrees  $d = \sum_{i=1}^n (2^{a_i} - 1)$ , when  $a_i - a_{i+1} \geq i+1$  for  $i < n$  and when  $a_i - a_{i+1} \geq n - i + 1$  for  $i < n$ . We complete a solution of the 3-variable hit problem by giving bases for  $Q^\omega(3)$  in the remaining cases.

The techniques introduced so far are useful for obtaining upper bounds for  $\dim Q^d(n)$ , but are less efficient for obtaining lower bounds, where we may wish to prove that no linear combination of a certain set of monomials is hit. Chapter 9 introduces the dual problem of finding  $K^d(n)$ , the simultaneous kernel of the linear operations  $Sq_k : \text{DP}^d(n) \rightarrow \text{DP}^{d-k}(n)$  dual to  $Sq^k$  for  $k > 0$ . Here  $\text{DP}(n)$  is a ‘divided power algebra’ over  $\mathbb{F}_2$ , whose elements are sums of dual or ‘d-monomials’  $v_1^{(d_1)} \dots v_n^{(d_n)}$ . As a  $\mathbb{F}_2\text{GL}(n)$ -module,  $K^d(n)$  is the dual of  $Q^d(n)$  defined by matrix transposition, and so  $\dim K^d(n) = \dim Q^d(n)$ . Thus we aim to find upper bounds for  $\dim Q^d(n)$  by using spanning sets in  $Q^d(n)$ , and lower bounds by using linearly independent elements in  $K^d(n)$ .

An advantage of working in the dual situation is that  $K(n)$  is a subalgebra of  $\text{DP}(n)$ . Since the dual spikes are in  $K(n)$ , they generate a subalgebra  $J(n)$  of  $K(n)$  which is amenable to calculation. In the cases  $n = 1$  and  $2$ ,  $J(n) = K(n)$ , and when  $n = 3$ ,  $K^d(n)/J^d(n)$  has dimension 0 or 1. We explain how to construct the dual  $K^\omega(n)$  of  $Q^\omega(n)$  with respect to an order relation. We study the duals of the Kameko and duplication maps, and solve the dual hit problem for  $n \leq 3$ . In Chapter 10 we extend these results by determining  $K^d(3)$  and

$Q^d(3)$  as modules over  $\mathbb{F}_2\text{GL}(3)$ . Here the flag module  $\text{FL}(3)$ , given by the permutation action of  $\text{GL}(3)$  on subspaces of the defining module  $V(3)$ , plays an important part. We describe tail and head modules in terms of the exterior powers of  $V(3)$ .

Hopf algebras are introduced in Chapter 11. A Hopf algebra  $A$  has a ‘coproduct’  $A \rightarrow A \otimes A$  compatible with the product  $A \otimes A \rightarrow A$ , and an ‘antipode’  $A \rightarrow A$ . We show that the coproduct  $Sq^k \mapsto \sum_{i+j=k} Sq^i \otimes Sq^j$  and the conjugation  $\chi$  provide the mod 2 Steenrod algebra  $A_2$  with the structure of a Hopf algebra. For a graded Hopf algebra  $A$  of finite dimension in each degree, the graded dual  $A^*$  is also a Hopf algebra. In this sense, the divided power algebra  $\text{DP}(n)$  is dual to the polynomial algebra  $P(n)$ . We show that the graded dual  $A_2^*$  of  $A_2$  is a polynomial algebra on generators  $\xi_j$  of degree  $2^j - 1$  for  $j \geq 1$ , and determine its structure maps. We conclude this chapter with the formula of Zaiqing Li for conjugation in  $A_2$ , which complements Milnor’s product formula of Chapter 4.

Chapters 12 and 13 give more detail on the internal structure of  $A_2$ . In Chapter 12 we focus on two important families of Hopf subalgebras of  $A_2$ , namely the subalgebras of ‘Steenrod  $q$ th powers’  $A_q$ , where  $q$  is a power of 2, and the finite subalgebras  $A_2(n)$  generated by  $Sq^k$  for  $k < 2^{n+1}$ . We also introduce some more additive bases of  $A_2$ . We continue in Chapter 13 by introducing a ‘cap product’ action of the dual algebra  $A_2^*$  on  $A_2$ , which can be used to obtain relations in  $A_2$  by a process which we call ‘stripping’. We use the ‘halving’ map (or *Verschiebung*) of  $A_2$  to explain why its action on  $P(n)$  reproduces itself by doubling exponents of monomials  $Sq^A$  and squaring polynomials. This map sends  $Sq^k$  to 0 if  $k$  is odd and to  $Sq^{k/2}$  if  $k$  is even. Since it is the union of the finite subalgebras  $A_2(n)$ , the algebra  $A_2$  is nilpotent. We apply the stripping technique to obtain the nilpotence order of certain elements of  $A_2$ .

Chapter 14 is devoted to a proof of the 2-dominance theorem of Judith H. Silverman and Dagmar M. Meyer. This deep result states that a monomial  $f$  in  $P^d(n)$  is hit if  $\omega(f)$  is not greater than  $\omega^{\min}(d)$  in the 2-dominance order. This strengthens the Peterson conjecture of Chapter 2 and the theorem of Singer from Chapter 6. One consequence is the Silverman–Singer criterion, which states that if  $g$  and  $h$  are homogeneous polynomials such that  $\deg g < (2^k - 1)\mu(\deg h)$ , where  $\mu$  is the numerical function of Chapter 2, then  $f = gh^{2^k}$  is hit.

In Chapter 15, we consider the Dickson algebra  $D(n)$  of  $\text{GL}(n)$ -invariants in  $P(n)$ . Following Nguyen H. V. Hung and Tran Ngoc Nam, we show that all Dickson invariants of positive degree are hit in  $P(n)$  when  $n \geq 3$ . There is a large class of similar problems: given a subgroup  $G$  of  $\text{GL}(n)$ ,

the subalgebra of  $G$ -invariant polynomials  $P(n)^G$  is an  $A_2$ -module, and the ‘relative’ hit problem asks for the elements of  $P(n)^G$  which are hit in  $P(n)$ . The corresponding ‘absolute’ hit problem asks for a minimal generating set for  $P(n)^G$ . We consider the absolute problem for the Weyl subgroup  $G = W(n)$  of permutation matrices in  $GL(n)$  in Chapter 25.

In the chapters which follow, we shift attention to the representation theory of  $GL(n)$  over  $\mathbb{F}_2$ . We begin in Chapter 16 by studying the flag module  $FL(n)$ , which is defined by the permutation action of  $GL(n)$  on the right cosets of the Borel subgroup  $B(n)$  of lower triangular matrices. This module is isomorphic to  $Q^d(n)$  when the degree  $d$  is ‘generic’ in the sense of Chapter 8. The Bruhat decomposition  $A = BWB'$  of a matrix  $A$  in  $GL(n)$  is used to define certain subspaces of  $FL(n)$  which we call ‘Schubert cells’. Here  $B, B' \in B(n)$  are lower triangular matrices and  $W \in W(n)$  is a permutation matrix. We show that  $FL(n)$  is the direct sum of  $2^{n-1}$  submodules  $FL_I(n)$ , where  $I \subseteq \{1, 2, \dots, n-1\}$  is the set of dimensions of the subspaces in the ‘partial’ flags given by right cosets of parabolic subgroups of  $GL(n)$ .

The main aim of Chapter 17 is to construct a full set of  $2^{n-1}$  irreducible  $\mathbb{F}_2GL(n)$ -modules  $L(\lambda)$ . Following C. W. Curtis,  $L(\lambda)$  is defined as the head of the summand  $FL_I(n)$  of  $FL(n)$ , where  $\lambda$  is the ‘column 2-regular’ partition corresponding to  $I$ , i.e.  $\lambda_i - \lambda_{i+1} = 1$  if  $i \in I$ , 0 if  $i \notin I$ . The summand of  $FL(n)$  corresponding to complete flags is the Steinberg module  $St(n)$ . We use the Hecke algebra  $H_0(n)$  of endomorphisms of  $FL(n)$  which commute with the action of  $GL(n)$ , and follow the methods of R. W. Carter and G. Lusztig.

In Chapter 18, we review the background from modular representation theory that we use to study  $P(n)$  and  $DP(n)$  as  $\mathbb{F}_2GL(n)$ -modules. We explain the role of idempotents in obtaining direct sum decompositions, and introduce the Steinberg idempotent  $e(n) = \overline{B(n)}\overline{W(n)} \in \mathbb{F}_2GL(n)$ , the sum of all products  $BW$ , where  $B \in B(n)$  and  $W \in W(n)$ . We study  $e(n)$  and the conjugate idempotent  $e'(n) = \overline{W(n)}\overline{B(n)}$  by means of an embedding of  $H_0(n)$  in the group algebra  $\mathbb{F}_2GL(n)$  due to N. J. Kuhn. We also discuss Brauer characters and the representation ring  $R_2(GL(n))$ .

In Chapter 19 we use idempotents in  $\mathbb{F}_2GL(n)$  to split  $P(n)$  as a direct sum of  $A_2$ -submodules  $P(n, \lambda)$ , each occurring  $\dim L(\lambda)$  times. We discuss the problem of determining the number of factors isomorphic to  $L(\lambda)$  in a composition series for  $P^d(n)$ . Following Ton That Tri, we use the Mui algebra of  $B(n)$ -invariants in  $P(n)$  to determine the minimum degree  $d$  in which  $P^d(n)$  has a submodule isomorphic to  $L(\lambda)$ .

As Weyl modules and their duals are central topics of modular representation theory, it is no surprise that they appear here also. As these modules are defined over infinite coefficient fields, we begin Chapter 20 by reviewing some

results on modular representations of the algebraic group  $\overline{G}(n)$  of nonsingular  $n \times n$  matrices over  $\overline{\mathbb{F}}_2$ , the algebraic closure of  $\mathbb{F}_2$ . We then introduce the ‘restricted’ Weyl module  $\Delta(\lambda, n)$  over  $\mathbb{F}_2$  and its transpose dual  $\nabla(\lambda, n)$ , and show that if  $\lambda$  is column 2-regular and if the ordering on  $\omega$ -sequences is suitably chosen, then  $\Delta(\lambda, n) \cong K^\omega(n)$  and  $\nabla(\lambda, n) \cong Q^\omega(n)$ , where  $\omega$  is the partition conjugate to  $\lambda$ . We use the theory of polynomial  $\overline{G}(n)$ -modules to determine the minimum degree  $d$  in which  $L(\lambda)$  occurs as a composition factor of  $P^d(n)$ .

Chapter 21 gives a self-contained proof of an important result of J. F. Adams, J. Gunawardena and H. R. Miller. This states that every degree-preserving  $A_2$ -module map  $P(m) \rightarrow P(n)$  is given by a sum of linear substitutions given by the action of  $m \times n$  matrices over  $\mathbb{F}_2$ . It follows that the  $A_2$ -summands in a maximal splitting of  $P(n)$  obtained using idempotents in  $\mathbb{F}_2 M(n)$  rather than  $\mathbb{F}_2 GL(n)$  are indecomposable. Hence such a splitting is a maximal direct sum decomposition of  $P(n)$  as an  $A_2$ -module.

Chapter 22 is concerned with the  $A_2$ -summands of  $P(n)$  corresponding to the Steinberg representation  $St(n)$  of  $GL(n)$ . We discuss the ‘internal’ model  $MP(n)$  of the  $A_2$ -module  $P(St(n))$  defined by Stephen A. Mitchell and Stewart B. Priddy using admissible monomials of length  $n$  in  $A_2$  itself. Although the hit problem for  $P(n)$  can be split into a corresponding problem for  $P(n, \lambda)$  for each  $\lambda$ , the Steinberg summand is the only case where this problem has been solved for all  $n$ , and we give minimal generating sets for the summands given by the idempotents  $e(n)$  and  $e'(n)$  of Chapter 18.

In Chapter 23, we identify the module  $J^d(n)$  of  $K^d(n)$  generated by the dual spikes in degrees  $d = \sum_{i=1}^n (2^{a_i} - 1)$ , where  $a_1 > a_2 > \dots > a_n$ , in terms of the flag module  $FL(n)$ . We show that  $Q^d(n) \cong FL(n)$  in ‘generic’ degrees  $d$ , and extend the method in Chapter 24 to obtain results of Tran Ngoc Nam on  $J(n)$  relating cohit modules to partial flag modules. Following Nguyen Sum, we give counterexamples for  $n \geq 5$  to Kameko’s conjecture that  $\dim Q^d(n) \leq \dim FL(n)$  for all  $d$ .

In Chapters 25 and 26 we discuss the hit problem for the action of  $A_2$  on the algebra of symmetric polynomials  $S(n)$ , the invariants in  $P(n)$  of the group  $W(n)$  of permutations of the variables. More generally, we discuss the ‘absolute’ hit problem for any subgroup  $G$  of  $W(n)$ , and show that the Peterson conjecture, the Kameko map and Singer’s minimal spike theorem have analogues for  $P(n)^G$ . We solve the symmetric hit problem for  $n \leq 3$ , using the dual problem to obtain the lower bound in the case  $n = 3$ . Following Singer, we introduce the ‘bigraded Steenrod algebra’  $\tilde{A}_2$ , which is obtained by omitting the relation  $Sq^0 = 1$  in the definition of  $A_2$ , and apply  $\tilde{A}_2$  to the dual problem.

In Chapters 27 and 28 we consider the cyclic subgroup  $C(n)$  of order  $2^n - 1$  in  $GL(n)$ . This is obtained by regarding  $P^1(n)$  as the underlying vector space of the Galois field  $\mathbb{F}_{2^n}$ . The action of  $C(n)$  on  $P(n)$  is diagonalized over  $\mathbb{F}_{2^n}$  by a change of variables which ‘twists’ the action of  $A_2$ , in the sense that  $Sq^1(t_i) = t_{i-1}$  in the new variables, which are indexed mod  $n$ . Following H. E. A. Campbell and P. S. Selick, we show that the polynomial algebra  $\tilde{P}(n) = \mathbb{F}_2[t_1, \dots, t_n]$  splits as the direct sum of  $2^n - 1$   $A_2$ -modules  $\tilde{P}(n, j)$  corresponding to the 1-dimensional representations of  $C(n)$ . In particular,  $\tilde{P}(n, 0)$  can be identified with the ring of  $C(n)$ -invariants of  $P(n)$ . In Chapter 28, we solve the dual cyclic hit problem for  $n = 3$  by using the twisted analogue  $\tilde{J}(n)$  of the d-spike module  $J(n)$ .

In Chapters 29 and 30, we collect some results on the hit problem in the 4-variable case as further illustration of our methods. Nguyen Sum has extended the method introduced by Kameko to find a monomial basis for  $Q^d(4)$  for all  $d$ . We include without proof some of the results of Sum, and also some results which we have verified only by computer using MAPLE. Thus there remain some challenging aspects of the hit problem even in the case  $n = 4$ .

The Steenrod algebra was originally defined for all primes  $p$ , but we have restricted attention to the case  $p = 2$ . All the problems have analogues for odd primes, but in general much less is known, and a number of difficulties arise in trying to extend our techniques to the odd prime case. The 2-variable hit problem for the action of  $A_p$  on the polynomial algebra  $\mathbb{F}_p[x, y]$  has been solved by Martin D. Crossley, but little appears to be known even for the 3-variable case. In common with many authors on the Steenrod algebra, we have therefore confined ourselves to the prime 2.

There are several good textbooks on topology which include material on the Steenrod algebra and its applications, such as those by Brayton I. Gray [70] and by Robert E. Mosher and Martin C. Tangora [147], in addition to the classic *Annals of Mathematics Study* [196], based on lectures by Steenrod himself, and the Cartan seminars [33]. A treatment of the Steenrod algebra from an algebraic viewpoint, including Steenrod operations over an arbitrary finite field, is given in Larry Smith’s book [190] on invariant theory. The book of Harvey R. Margolis [129] treats the general theory of modules over the Steenrod algebra. Still other approaches to the Steenrod algebra are possible. The survey article [233] treats Steenrod operations as linear differential operators with polynomial coefficients, and is a precursor for this book.

We sometimes introduce definitions and constructions for a small number of variables and extend them to the general case in later chapters. Although this can involve a certain amount of repetition, it has the advantage of leading to

interesting results at an early stage by elementary methods. We hope that our approach will appeal to readers whose main interests are in algebra, especially in the modular representation theory of linear groups, or in the combinatorics related to symmetric polynomials and to the invariant theory of finite groups.

In order to avoid interruptions to the text, citations and background material are collected in the ‘Remarks’ sections at the end of each chapter. The occasional reference to topology may occur in these, but we have not tried to explain the topology. We have also omitted important topics such as the Singer transfer map and its applications to the homotopy groups of spheres through the Adams spectral sequence. These would require another volume, which we are not qualified to write. For similar reasons, we do not treat the theory of analytic functors and the category  $\mathcal{U}/\mathcal{N}il$  of unstable  $A_2$ -modules modulo nilpotent objects due to Hans-Werner Henn, Jean Lannes and Lionel Schwartz. Finally, in a subject which crosses several disciplines, notation presents a problem because the traditional symbols of one area may be in conflict with those of another. A list of symbols for the main ingredients of our subject appears at the end of the book, together with an index of the main terms defined in the text.

We should like to offer our sincere thanks to the School of Mathematics of the University of Manchester for providing us with office space and computing facilities during work on this text. We should also like to thank several colleagues at Manchester for mathematical help and support, and in particular Peter Eccles, Nige Ray, Peter Rowley and Bob Sandling. Of colleagues farther afield, we should like to thank Nguyen H. V. Hung, Ali S. Janfada, Bill Singer and Larry Smith for their interest in our project. The first author would also like to thank Stephen R. Doty for teaching him something about modular representation theory. Finally we should like to thank Roger Astley and his colleagues at Cambridge University Press for their encouragement, support and patience. We set out with the modest aim of providing a beginning graduate student in topology and algebra with a basic primer on the Steenrod algebra, illustrated by our favourite application to a problem proposed by Frank Peterson, but the project has expanded substantially in scope over the past eight years.