

Index

abciximab, 262
 abdominal examination, 8
 abdominal surgery, 215
 Abiomed Impella, 120
 ablation procedures, cardiac, 136
 absent pulmonary valve syndrome, 163
 acid-base management, 42, 46
 cerebral protection, 196
 pharmacological effects, 200
 activated clotting time (ACT), 40, 43
 acute kidney injury (AKI), 61–4
 management, 64
 paediatric patients, 184
 prevention, 62
 risk factors, 63
 acute lung injury (ALI), 59
 acute renal failure (ARF), 61–4
 acute respiratory distress syndrome (ARDS), 59
 Adamkiewicz artery, 101
 adrenaline. *See* epinephrine
 adult congenital heart disease (ACHD), 186–90
 anaesthetic management, 188–9
 antibiotic prophylaxis, 189
 blood management, 189
 classification, 186
 postoperative care, 189–90
 preoperative assessment, 187–8
 after-drop hypothermia, 195
 afterload, 55, 57
 paediatric patients, 182
 air embolism (AE), 46, 210–12
 management, 211–12
 physical principles, 210–11
 air entrainment
 CPB, 210
 ECMO, 220
 air warmers, forced, 192
 alfentanil, 26
 α-stat blood-gas management, 42, 196
 paediatric patients, 150–1
 pH-stat management vs, 206–7
 ε-aminocaproic acid, 152, 264
 anaemia. *See also* haemotocrit
 congenital haemolytic, 262
 dilutional, during CPB, 41, 207
 failure to wean from CPB, 47
 postoperative, 267

preoperative, 261
 anaesthesia, 24–8
 checklist prior to surgery, 28
 during CPB, 42
 induction, 26
 maintenance, 28
 pregnancy, 272
 preparation for, 25
 transfer to operating room, 27–8
 vascular access, 25–6
 anaesthetic agents
 ideal cardiac, 26
 pregnancy, 272
 preparation for use, 26
 analgesia, postoperative. *See*
 postoperative pain management
 anasarca, 7
 andexanet alfa, 263
 angina, 6
 aortic stenosis, 69
 chronic refractory, 283
 antibiotics. *See* antimicrobials
 anticoagulation
 AF, 58
 CPB, 40, 264
 ECMO, 219–20
 IABP, 48
 MV valve repair, 80–1
 off-pump coronary surgery, 93
 perioperative management, 262–3
 pregnancy, 272
 preoperative cessation, 24, 114–15
 regional anaesthesia and, 278
 antifibrinolytics, 264
 antimicrobials
 Clostridioides difficile infection, 288
 endocarditis prophylaxis, 189, 286
 endocarditis therapy, 285–6
 heart transplant recipients, 128
 resistance and stewardship, 288
 surgical prophylaxis, 27, 286–7
 antiplatelet agents, 262
 neuraxial blockade, 277
 preoperative cessation, 25, 262
 antithrombin deficiency, 40
 aorta
 atheroma, 65, 204
 cannulation, 2–3, 40–1

coarctation, 161–2, 171
 thoracic. *See* thoracic aortic surgery
 aortic aneurysms, 99
 complications of surgery, 105–6
 descending aorta, 102–3
 indications for surgery, 99
 preoperative assessment, 100–1
 aortic arch surgery, 101–2
 aortic clamping/cross-clamping, 3, 41
 clamp-off, 4
 left heart bypass, 104
 minimally invasive surgery, 97
 off-pump surgery, 92
 aortic disease
 imaging, 17–19, 101
 pregnancy, 272
 thoracic, 99–100
 TOE assessment, 258–9
 aortic dissection, 99
 AR, 72
 classification, 99–100
 during CPB, 213
 indications for surgery, 1
 pregnancy, 272
 preoperative assessment, 100–1
 thrombolytic therapy, 263
 TOE assessment, 258–9
 aortic interruption, 161
 aortic pseudoaneurysms, 99
 aortic regurgitation (AR), 71–3
 anaesthesia, 73
 grading of severity, 248
 investigations, 72–3
 mechanisms, 249
 pathophysiology, 72
 TOE assessment, 248
 aortic root surgery, 101
 aortic stenosis (AS), 69–71
 anaesthesia, 70–1
 grading of severity, 248–9
 investigations, 70
 pathophysiology, 69–70
 percutaneous interventions, 138, 140
 pregnancy, 272
 TOE assessment, 248–9
 aortic valve (AV), 69
 balloon dilation, 140
 bicuspid, 69, 248
 TOE evaluation, 248

- aortic valve replacement (AVR), 3
 minimally invasive, 5
 post-CPB hypertension, 71
 Ross procedure, 172–4
 transcatheter implantation, 5, 137–40
- aortic valve surgery, 3, 69–73
 anaesthesia, 70–1, 73
 key points, 73
 surgical approach, 73
- aortography, 21–2
- apex beat, 8
- apixaban, 263, 278
- aprotinin, 152, 264
- argatroban, 264
- arrhythmias
 adult congenital heart disease, 188
 after MV repair, 81
 CPB termination, 42–3, 46
 CVP waveforms, 225
 device-based therapy, 131
 electrophysiological procedures, 134–6
 mapping, 136
 paediatric patients, 183
 postoperative, 55, 58
- arrhythmogenic right ventricular cardiomyopathy (ARVC), 111–12
- arterial BP monitoring, 223–4
 artefacts, 47
 underdamping and overdamping, 223
- arterial cannulae, 33
 cerebral perfusion, 101
 removal, 53–4
- arterial cannulation, 25
 BP monitoring, 223
 CPB, 40–1, 43
 arterial oxygen content (CaO_2), 226
- arterial shunts, 154–5
- arterial switch procedure, 177–8
- arteriovenous fistula, haemodialysis patients, 64
- arteriovenous oxygen content difference, 20, 211
- aspirin, 25, 262
- atelectasis, 53, 59
- atherosclerosis, aortic, 65, 204
- atracurium, 26
- atrial fibrillation (AF)
 ablation procedures, 136
 after MV repair, 81
 DC cardioversion, 135
 MV disease, 76–8, 81
 postoperative, 52, 58
 TOE assessment, 255
- atrial septal defect (ASD), 156
 percutaneous closure, 141
 surgical closure, 156–7
- TOE assessment, 254–5
- atrial septostomy, balloon
 PHT, 113
 transposition of the great arteries, 147, 177
- atrioventricular (AV) block, 12
 indications for pacing, 132
 VSDs, 180
- atrioventricular septal defects (AVSDs), 156, 158–9
- auditory evoked potentials (AEPs), 233
- auscultation, 8–10
- balloon aortic valve dilation, 140
- balloon atrial septostomy. *See* atrial septostomy, balloon
- balloon pulmonary angioplasty, 116
- base excess, 42
- benzodiazepines, 201
- Bernoulli equation, 239
- bidirectional cavopulmonary shunt (BCPS), 160
- bivalirudin, 264
- biventricular support devices, 118
- Blalock–Taussig (B-T) shunt, modified, 147, 154–5
 pulmonary atresia, 172
- bleeding. *See* haemorrhage/bleeding
- blood donation, preoperative autologous, 261
- blood flow
 effects of CPB, 199
 myocardial oxygen supply, 30
- blood pressure (BP)
 invasive monitoring, 223
 paediatric patients, 145
- blood transfusion
 adult congenital heart disease, 189
 after aortic surgery, 105
 coagulopathies, 264
 during CPB, 207
 paediatric patients, 182
- blood-gas management, 42, 206–7
- bloody tap, 278
- Boyle's law, 210–11
- bronchospasm, CPB termination, 46
- bubble oxygenators, 210
- bubble traps, 38
- bupivacaine, hyperbaric, 275–6
- Caesarean section, 270–1
- calcium
 coronary, scoring, 16
 pharmacological use, 52
- CaO_2 (arterial oxygen content), 226
- carboprost, 271
- carcinoid heart disease, 87–9
- carcinoid syndrome, 87
- cardiac ablation procedures, 136
- cardiac allograft vasculopathy, 129
- cardiac arrest. *See also* resuscitation
 AV surgery, 71
 CPB for resuscitation, 216–17
 MS, 77
 postoperative, 55, 67
 therapeutic hypothermia, 191–2
- cardiac catheter laboratory procedures
 DC cardioversion, 135–6
 electrophysiological, 134–6
 mapping and ablation, 136
 rhythm management, 131–4
 structural heart disease, 137–42
- cardiac catheterization, 20
- cardiac disease
 maternal deaths, 269–70
 physical signs, 7–10
 pregnancy and delivery, 270–1
 symptoms, 6–7
- cardiac index (CI), 20, 45
 neonates, 145
- cardiac magnetic resonance imaging (CMR), 18–20
 adult congenital heart disease, 188
 coronary artery disease, 17, 19
 valvular disease, 19
- cardiac masses, 19–20, 255–6
- cardiac output (CO). *See also* low cardiac output state
 CPB, 41
 measurement, 20, 226
 postoperative optimization, 52, 55
 shock, 228
- cardiac resynchronization therapy (CRT), 131–2
- cardiac rhythm. *See also* arrhythmias during CPB, 42–3
 management devices, 131–4
- cardiac surgery, 1–5, *See also*
 cardiopulmonary bypass
 early postoperative care, 50–4
 minimally invasive (MICS), 5, 94–7
 myocardial protection, 3, 30–3
 off-pump, 4–5, 91–4
 operative principles, 2–4
 patient selection, 1–2
 pregnancy, 271–3
 risk assessment, 2
 setting up, 2–3
- cardiac tamponade
 haemodynamic parameters, 228
 postoperative, 57–8
 TOE assessment, 256
- cardiogenic shock
 critical, 119–20
 haemodynamic parameters, 228
- cardiomyopathies, 107–12
- cardioplegia, 3, 32–3, *See also* deep hypothermic circulatory arrest

Index

- cardioplegia (cont.)
 - AV surgery, 3
 - delivery, 33–4
 - minimally invasive surgery, 95
 - myocardial protection, 30
 - normothermic versus hypothermic, 205
- cardioprotection. *See* myocardial protection
- cardiopulmonary bypass (CPB), 30–44
 - air embolism, 45–6, 210–12
 - anaesthesia, 42
 - anticoagulation, 40, 264
 - blood transfusion, 207
 - blood-gas management, 42, 206–7
 - cannulation, 40–1
 - cardiac rhythm, 42–3
 - circuit. *See* cardiopulmonary bypass circuit
 - coming off, 4
 - communication, 39–40
 - conduct, 39–43
 - controversies, 203–8
 - de-airing, 4
 - decannulation, 43
 - deep hypothermic circulatory arrest, 192–7
 - descending aorta surgery, 102–3
 - drug pharmacokinetics, 199–200
 - emergencies, 208
 - femoro-femoral, 215
 - haematocrit, 41, 207
 - haemodynamic management, 41–2
 - haemostasis and closure, 43
 - hypothermia, 42, 192–7
 - inadequate flow, 212–13
 - inadequate oxygenation, 212
 - initiation, 41
 - metabolic management, 42
 - mini, 39, 207–8
 - minimally invasive surgery, 95–7
 - myocardial protection, 30–3
 - non-cardiac applications, 215–17
 - normothermia versus hypothermia, 205
 - off-pump surgery conversion to, 93
 - paediatric patients, 150–2
 - pregnancy, 272–3
 - preparation for, 40
 - pressure versus flow, 203–5
 - prothrombotic risk, 263
 - pulsatile flow, 205–6
 - resuscitation using, 216–17
 - setting up, 2–3
 - temperature management, 191–8
 - ventilation and oxygenation, 208
 - weaning. *See* weaning from CPB
 - cardiopulmonary bypass circuit, 33–9
 - air entrainment, 210
 - components, 33–8
 - filters and bubble traps, 38–9
 - mini, 39, 207–8
 - safety standards, 39
 - sequestration of drugs, 200
 - cardiotomy suction, 39
 - cardiovascular complications, 55–8
 - cardiovascular monitoring, 223–8
 - cardioversion
 - CPB weaning, 46
 - DC, 135–6
 - postoperative arrhythmias, 52, 55, 58
 - carotid sinus hypersensitivity, 132
 - Carpentier classification, 248
 - catheter-based procedures. *See* cardiac catheter laboratory procedures
 - cavopulmonary shunt, bidirectional, 160
 - cell salvage, 39
 - central shunt, 154
 - central venous access
 - aortic surgery, 103
 - cardiac surgery, 25–7
 - CVP monitoring, 224
 - heart transplantation, 127
 - paediatric cardiac anaesthesia, 150
 - rhythm-management-device implantation, 134
 - central venous oxygen saturation (ScvO_2), 228
 - central venous pressure (CVP). *See also* right atrial pressure
 - CPB, 41, 46–7
 - monitoring, 224–5
 - shock, 228
 - waveforms, 224–5
 - centrifugal pumps, 35, 194
 - CentriMag® ventricular assist device, 119–20
 - cerebral artery blood flow velocity (CBFV), 234
 - cerebral autoregulation, 203–5
 - cerebral ischaemia, EEG changes, 230–1
 - cerebral oxygenation monitoring, 102, 233–5
 - cerebral perfusion, 101–2
 - CPB perfusion pressure and, 203–4
 - intermittent, 197
 - monitoring, 102, 233–5
 - retrograde, 101–2, 197, 211–12
 - selective antegrade, 101–2, 197
 - cerebral protection
 - aortic arch surgery, 101–2
 - deep hypothermic circulatory arrest, 196–7
 - paediatric patients, 150–1
 - cerebrospinal fluid (CSF) drainage, 104
 - cerebrovascular disease, pre-existing, 204
 - CHARGE association, 163
 - Charles' law, 210–11
 - CHD. *See* congenital heart disease
 - chest drains, removal, 53
 - chest pain, 6
 - chest radiography (CXR), 14
 - cholecystitis, postoperative, 60
 - chordae tendinae, 75
 - chronic refractory angina, 283
 - chronic thromboembolic pulmonary hypertension (CTEPH), 113–16
 - chylothorax, postoperative, 184–5
 - clonidine, 25
 - clopidogrel, 262, 278
 - Clostridioides difficile* infection, 288
 - coagulation testing, 264
 - coagulopathy
 - blood product replacement, 264
 - heart transplant recipients, 127–8
 - paediatric patients, 152
 - postoperative, 53, 267
 - coarctation of the aorta (CoA), 161–2, 171
 - cognitive dysfunction, postoperative, 64, 66–7
 - cold agglutinins, 262
 - colour flow Doppler (CFD), 239
 - communication, 39–40, 282
 - complement activation, during CPB, 263
 - computed tomography (CT), 15–18
 - computed tomography angiography (CTA)
 - aortic disease, 18, 101
 - coronary artery disease, 15–18
 - congenital heart disease (CHD), 145, 154–80, *See also* paediatric cardiac anaesthesia
 - adult, 186–90
 - anaesthesia, 148–50
 - closed cardiac surgery, 148
 - cyanotic, 148
 - failure to wean from CPB, 47
 - outcome of surgery, 185
 - pathophysiology, 149
 - postoperative care, 181–5
 - postoperative problems, 182–5
 - surgical strategy, 147–8
 - constrictive pericarditis, 109, 111
 - restrictive cardiomyopathy versus, 111
 - TOE assessment, 256
 - continuous haemodiafiltration, 64
 - continuous veno-venous haemofiltration (CVVHF), 64
 - continuous wave Doppler (CWD), 239
 - contractility, paediatric patients, 182

- contrast echocardiography, 14
 COPS protocol, 105
 cor triatriatum, 176
 coronary angiography, 21–2
 aortic disease, 101
 intravascular ultrasound versus, 22–3
 non-invasive, 17
 coronary artery bypass graft (CABG)
 surgery
 conduit harvesting, 2
 indications, 1
 off-pump (OPCAB), 4–5, 91–4
 operative principles, 3
 coronary artery calcium scoring, 16
 coronary artery disease (CAD)
 angiography. *See* coronary angiography
 cardiac magnetic resonance, 17, 19
 CT, 15–16, 18
 exercise ECG, 11–13
 indications for surgery, 1
 intravascular ultrasound, 22–3
 non-invasive imaging, 17
 coronary blood flow (CBF), 30
 coronary computed tomography
 angiography (CTA), 15–18
 coronary fistulae/sinusoids, 172
 coronary perfusion, PHT, 114
 Corrigan's sign, 8
 CPB. *See* cardiopulmonary bypass
 creatinine kinase MB (CK-MB), 57
 creatinine, serum, 63
 cryoablation, 136
 cryohydrocytosis, 262
 cryoprecipitate, 264
 CvO_2 (mixed venous oxygen content), 226
 cyanosis, 8
 adult congenital heart disease, 187–8
 CHD, 148
 differential, ECMO, 219
 Fallot's tetralogy, 163
 total anomalous pulmonary venous drainage, 176
 dabigatran, 263, 278
 DC cardioversion, 135–6
 de Musset's sign, 7
 de-airing, 4, 211–12
 DeBakey classification, 99–100
 deep hypothermic circulatory arrest (DHCA), 192–7. *See also* cardioplegia; hypothermia
 anaesthesia, 193–4
 aortic surgery, 101
 circulatory arrest, 195
 cooling, 195
 CPB modifications, 194–5
 failure to wean from CPB, 46
 indications, 193
 neurological monitoring, 197, 231–2
 neurosurgery, 216
 paediatric cardiac surgery, 150–1
 postoperative care, 197
 rewarming, 195–6
 surgical aspects, 194
 delirium, postoperative, 64, 66–7, 276
 depth-of-anaesthesia (DOA)
 monitoring, 26–7, 233
 descending thoracic aorta surgery, 102–5
 desflurane, 201
 Di George syndrome (22q11 deletion), 148, 163, 179
 diabetes mellitus, 65, 204
 diagnostic tests, 11–23
 diastolic dysfunction, LV, 243
 diathermy, pacemaker/ICD recipients, 134
 dilated cardiomyopathy (DCM), 108–9
 dimensionless index (DI), 248
 direct antiglobulin test, 262
 direct-acting oral anticoagulants (DOACs), 24–5, 263, 278
 distal limb ischaemia, 219
 dobutamine, 151
 dopamine, 63, 151
 Doppler ultrasound, 239
 Down syndrome, 158
 drugs
 pharmacological effects of CPB, 199–201
 preoperative cessation, 24–5
 dual source computed tomography (DSCT), 15–16
 ductus arteriosus (DA), 170
 closure, 145–6, 170
 dependent circulation, 146, 154
 patent (PDA), 146, 170
 Duke Activity Status Index (DASI), 7
 Duke criteria, endocarditis, 285
 Duroziez's sign, 9
 dyspnoea, 6
 dysrhythmias. *See arrhythmias*
 echocardiography, 13–14
 contrast, 14
 stress, 13
 transoesophageal. *See* transoesophageal echocardiography
 transthoracic (TTE), 13
 ECMO. *See* extracorporeal membrane oxygenation
 Eisenmenger syndrome, 187
 ejection fraction (EF)
 3D right ventricular, 245
 left ventricular, 242
 elderly, CPB perfusion pressure, 204
 electrical power failure, 213
 electrocardiography (ECG), 11–13
 abnormalities, 12
 ambulatory, 13
 atrial, 183
 exercise, 11–13
 implantable recorders, 13
 intraoperative, 28
 limitations to resting, 11
 perioperative myocardial infarction, 57
 electroencephalography (EEG), 231–2
 deep hypothermic circulatory arrest, 230–3
 depth-of-anaesthesia monitoring, 26, 233
 electrolyte management
 CPB, 42, 46
 paediatric patients, 181–2
 postoperative, 52
 electromagnetic interference (EMI), 134
 electron beam computed tomography (EBCT), 15–16
 electrophysiological disorders, 135
 electrophysiological procedures, 134–6
 embolus filters, 263
 emergencies, CPB, 208
 emphysema, 216
 end-diastolic volume index, 20
 endo-aortic balloon clamp, 97
 endocarditis, infective, 285–6
 aetiology, 286
 after MV surgery, 81
 AR, 72
 Duke criteria, 285
 MR, 78
 prophylaxis, 189, 286
 TOE assessment, 256
 endotracheal tubes, paediatric, 150
 end-systolic pressure–volume relationship (ESPVR), 55
 enflurane, 201
 epidural anaesthesia
 neuraxial bleeding, 276
 paediatric cardiac surgery, 150
 thoracic, 26, 274–5, 282
 epidural haematoma, 276–8
 epinephrine
 paediatric patients, 151

Index

- epinephrine (cont.)
 - postoperative cardiac arrest, 55
 - ventricular assist device insertion, 121–3
- ϵ -aminocaproic acid, 152, 264
- eptifibatide, 262
- ergometrine, 271
- etomidate, 26
- EuroSCORE (European System for Cardiac Operative Risk Evaluation), 2
- evoked potentials, 104–5, 233
- exercise electrocardiography (exECG), 11–13
- exertional syncope, 7
- extracorporeal life support, after paediatric cardiac surgery, 182–3
- extracorporeal membrane oxygenation (ECMO), 218–20
 - cannulation, 219
 - cardiogenic shock, 119
 - equipment, 218–19
 - management, 219–20
 - outcomes, 220
 - paediatric patients, 182–3
 - patient selection, 218
 - weaning, 220
- factor VIIa, recombinant, 264
- factor VIII deficiency, 263
- factor IX deficiency, 263
- Fallot's tetralogy, 163–4
 - anatomy, 163
 - physiology, 147, 163
 - surgery, 163–4
- family, 54, 148
- fast-tracking, 50
- fatigue, 7
- femoral arterial cannulation, 219
- femoral vein cannulation, 224
- femoro-femoral cardiopulmonary bypass, 215
- fentanyl, 26, 150, 274
- fibrinogen concentrate, 264
- fibrinolysis, 264
- filters, CPB circuit, 38
- flow index, effect of hypothermia, 46
- flow rate, CPB pump, 41–2
 - inadequate, 212
 - perfusion pressure and, 203–5
- fluid management
 - aortic surgery, 105
 - off-pump surgery, 93
 - paediatric patients, 181–2
 - postoperative, 52–3
- fluid warming, 192
- Fontan procedure, 165–6
 - postoperative management, 165
- surgery in adulthood, 186, 190
- foramen ovale
 - closure, 145
 - patent. *See* patent foramen ovale
- fractional area change, right ventricular (RVFAC), 245
- fractional flow reserve (FFR), 22
- fractional shortening (FS), 243
- fresh frozen plasma (FFP), 264
- functional status, 6
- fundal examination, 8
- furosemide, 52–3
- gadolinium diethylene triamine pentaacetic acid (Gd-DTPA), 19
- gastric tube, 27
- gastrointestinal (GI) complications, 59–61
- gastrointestinal (GI) haemorrhage, 60–1
- gastrointestinal (GI) perforation, 60
- Glenn procedure, 160
- global longitudinal strain (GLS), 243
- glucose management, 42, 197
- glyceryl trinitrate (GTN), 52, 151
- greater radicular artery (of Adamkiewicz), 101
- haematocrit, 41
 - correction prior to weaning, 47
 - deep hypothermic circulatory arrest, 196
 - minimum acceptable, 207
- haematology, 261–7
 - intraoperative care, 263–7
 - postoperative problems, 267
 - preoperative care, 261–3
- haemodialysis patients, 64
- haemodilution
 - cerebral protection, 196
 - during CPB, 41
 - pharmacokinetic effects, 199
- haemodynamics
 - CPB, 41–2
 - off-pump coronary surgery, 93
 - postoperative, 51–2, 55
 - shock, 228
- haemofilters, 194
- haemofiltration, intraoperative, 64
- haemoglobinopathies, 262
- haemolytic anaemia, congenital, 262
- haemophilia, 263
- haemoptysis, 6
- haemorrhage/bleeding
 - blood product replacement, 264
 - failure to wean from CPB, 47
 - GI, 60–1
- neuraxial, 276–8
- postoperative, 53, 267
- haemostasis
 - aortic surgery, 105
 - paediatric cardiac surgery, 152
 - post-bypass, 4, 43, 196
 - preoperative considerations, 261
 - handover (to ICU), 50–1
 - paediatric patients, 181
- Harlequin syndrome, 219
- heart block, 12, 132
 - after paediatric cardiac surgery, 183
 - VSDs, 180
- heart failure
 - acute, mechanical circulatory support, 119–20
 - carcinoid heart disease, 87–8
 - cardiac resynchronization therapy, 131–2
 - diastolic dysfunction, 243
 - dilated cardiomyopathy, 108–9
 - heart transplantation, 125–6
 - paediatric patients, 146, 148
 - VSDs, 118
- heart rate (HR)
 - after paediatric cardiac surgery, 182
 - paediatric patients, 145
- heart sounds, 9
- heart transplantation, 125–9
 - anaesthesia, 126–7
 - antimicrobial therapy, 128
 - bridge to, 118–21
 - early complications, 128
 - immunosuppression, 128
 - late complications and morbidity, 128–9
 - organ donor harvest, 276
 - patient selection, 125–6
 - reperfusion and weaning from CPB, 127–8
- HeartMate 3, 121
- HeartMate II LVAD, 121–2
- HeartMate PHP, 120
- HeartMate VE/XVE LVAD, 121
- HeartWare HVAD, 121–2
- heat conservation, 92–3, 191
- heat exchangers, 38
- heat loss, 191
- Henry's law, 210–11
- heparin (unfractionated heparin; UFH)
 - alternatives, 264
 - CPB, 40
 - ECMO, 219–20
 - IABP, 48
 - off-pump coronary surgery, 93
 - pharmacokinetic interactions, 199
 - preoperative management, 25, 263
 - resistance, 40
 - reversal, 43

heparin-induced thrombotic thrombocytopenic syndrome (HITT), 40, 267
 hepatic dysfunction, 60
 history, clinical, 6–7
 humidification, 192
 5-hydroxyindoleacetic acid (5-HIAA), 88–9
 hyoscine hydrobromide, 25
 hyperglycaemia, 52
 hyperkalaemia, 52
 hypertension
 CPB perfusion pressure, 204
 postoperative, 52, 71
 hyperthermia, postoperative, 181
 hypertrophic obstructive cardiomyopathy (HOCM), 107–8
 pathophysiology, 107
 surgical complications, 47
 hypokalaemia, 52
 hypomagnesaemia, 52
 hypoplastic left heart syndrome (HLHS), 167–8
 Fontan procedure, 165
 Norwood procedure, 167–8
 hypotension
 AV surgery, 71, 73
 artefactual, during CPB, 47
 carcinoid heart disease, 89
 MV surgery, 77, 79
 pharmacokinetic effects, 199
 postoperative, 51
 postural, 7
 hypothermia, 191–7. *See also* deep hypothermic circulatory arrest
 after-drop, 195
 aortic surgery, 101
 cerebral protection, 196
 during CPB, 42, 192–7
 failure to wean from CPB, 46
 inadvertent, 181
 myocardial protection, 30, 33
 normothermia versus, 205
 paediatric cardiac surgery, 150–1
 pathophysiology, 51, 193
 pharmacological effects, 199–201
 postoperative rewarming, 51
 therapeutic, cardiac arrest, 191–2
 hypovolaemic shock, 228

IABP. *See* intra-aortic balloon pump

ICU. *See* intensive care unit

idarucizumab, 263

ileus, postoperative, 60

imaging, cardiovascular, 14–20

immunosuppression, 128

implantable-cardioverter defibrillators (ICDs), 131

DC cardioversion, 135

explantation, 132–3
 implantation, 131–3
 perioperative management, 133

implantable device infection, 287

implantable ECG recorders, 13

induction of anaesthesia, 26
 paediatric surgery, 149–50
 procedures following, 27–8

infection, 285–8
 after MV repair, 81
 postoperative, 286–8

infection prevention and control interventions, 288

inferior vena cava (IVC)
 interruption, 160
 renal tumours involving, 216

inhalational anaesthetics. *See* volatile anaesthetics

inotropic agents
 paediatric patients, 151
 weaning from CPB, 47

insulin, 52

intensive care unit (ICU)
 early postoperative care, 51–2
 family involvement, 54
 fast-tracking, 50
 first postoperative day, 54
 paediatric patient transfer, 153, 181
 postoperative complications, 53, 55–67
 spinal cord perfusion, 105
 transfer, admission and handover, 50–1
 VADs, 123–4

INTERMACS patient profiles, 118

internal jugular vein (IJV) cannulation, 25–6, 224

intra-aortic balloon pump (IABP).
See also ventricular assist devices
 critical cardiogenic shock, 119
 failure to wean from CPB, 48–9
 indications, 49

intrathecal pressure (ITP), 104–5

intravascular ultrasound (IVUS), 22–3

intravenous anaesthesia, 42

invasive diagnostic tests, 20–3

iron deficiency, 261

ischaemia reperfusion injury (IRI), 31

ischaemic conditioning, 31

ischaemic heart disease (IHD). *See* coronary artery disease

ischaemic postconditioning, 31

ischaemic preconditioning (IPC), 31–2

isoflurane, 201

isoproterenol (isoprenaline), 151

jugular bulb oxygen monitoring, 234

jugular vein pressure levels and waveforms, 8

jugular venous oxygen saturation ($SjvO_2$), 234

junctional ectopic tachycardia, 183

Kawashima operation, 160

ketamine, 149

Kussmaul's sign, 8

labour, 270–1

Laplace's law, 55

laryngeal nerve injury, 184

left anterior hemiblock, 12

left atrial appendage (LAA)
 occlusion, 142
 thrombus, 255

left atrial hypertrophy, 12

left atrial myxoma, 255–6

left atrial pressure (LAP)
 elevated, 75–6, 78
 estimation, 226

left bundle branch block (LBBB), 12

left coronary artery, anomalous, 163

left heart bypass (LHB), 102–4

left heart catheterization, 20–2

left internal mammary artery (LIMA), 2–3

left posterior hemiblock, 12

left ventricular aneurysms, 110

left ventricular assist devices (LVADs), 118
 long-term, 119
 postoperative care, 123–4
 preoperative assessment, 120–1
 short-term, 118–19

left ventricular diastolic function, 243

left ventricular dysfunction
 postoperative, 55
 TOE assessment, 243

left ventricular ejection fraction (LVEF), 242

left ventricular end-diastolic pressure (LVEDP)
 AR, 72
 AS, 69–70
 estimation, 226
 mitral regurgitation, 78

left ventricular end-diastolic volume (LVEDV), 243

left ventricular end-systolic volume (LVESV), 243

left ventricular hypertrophy (LVH), 12, 69, 72

left ventricular internal dimension (LVID), 243

left ventricular outflow tract (LVOT)
 obstruction
 hypertrophic cardiomyopathy, 107
 management, 108

MV repair, 80–1

Index

left ventricular pressure, 20
 left ventricular pressure-volume loops, 57
 AR, 72
 AS, 70
 MR, 78
 MS, 76
 left ventricular systolic function, 243
 regional, 243–4
 strain imaging, 243
 left-to-right shunts, 20, 147
 leucocyte depleting filters, 38
 Levine's sign, 6
 levobupivacaine, 274
 lidocaine, 274
 liver transplantation, 216
 long saphenous vein (LSV), 2
 loop diuretics, 63
 lorazepam, 25
 low cardiac output state (LCOS)
 after Fallot's tetralogy surgery, 163–4
 after paediatric cardiac surgery, 181–3
 post-bypass, 45
 low-molecular-weight heparin (LMWH), 263
 lung
 auscultation, 9
 isolation, pharmacokinetic effects, 200
 lung fields, examination, 8
 lung transplantation, 216
 lupus anticoagulant, 263
 lymphoma, 262
 magnesium sulphate, 52
 magnetic resonance imaging (MRI)
 cardiac. *See* cardiac magnetic resonance
 pacemaker/ICD interference, 134
 magnets, application of, 134
 maintenance of anaesthesia, 28
 major aorto-pulmonary collateral arteries (MAPCAs), pulmonary atresia with, 172–3
 mannitol, 63
 mapping, electrophysiological, 136
 masses, cardiac, 19–20, 255–6
 maternal deaths, cardiac disease, 269–70
 mattress, heated, 192
 mean arterial pressure (MAP)
 acute kidney injury, 62–3
 aortic surgery, 104
 CPB, 41–2, 46–7, 203–4
 invasive monitoring, 223
 postoperative management, 55

mechanical circulatory support (MCS), 118–24, *See also* ventricular assist devices
 failure to wean from CPB, 47–9, 120
 long-term, 119–21
 short-term, 119–20
 median sternotomy, 2
 mediastinal surgery, 216
 mediastinitis, post sternotomy, 287
 membrane oxygenators, 37–8, 210
 mesenteric ischaemia, 60–1
 metaraminol, 93
 methadone, 25
 midazolam, 26
 milrinone, 151
 mini cardiopulmonary bypass (mini CPB), 39, 207–8
 minimally invasive cardiac surgery (MICS), 5, 94–7
 anaesthesia, 95–7
 MV, 75
 postoperative care, 97
 preoperative assessment, 94–5
 types, 94
 minimally invasive extracorporeal circulation (MiECC) techniques, 39
 MitraClip® system, 140–1
 mitral regurgitation (MR), 78–81
 anaesthesia, 78
 Carpentier classification, 248
 catheter-based therapies, 140–1
 grading of severity, 250–1
 pathophysiology, 78
 persistent, 80
 surgery, 79
 TOE assessment, 79, 248–50
 mitral stenosis (MS), 75–8
 after MV repair, 80
 anaesthesia, 77
 catheter-based therapies, 140–1
 investigations, 76–7
 pathophysiology, 75–6
 pregnancy, 272
 surgical options, 77–8
 TOE assessment, 250
 mitral valve (MV), 75
 myxomatous degeneration, 78
 orifice area (MVA), 75–6
 pressure gradient, 75–6
 systolic anterior motion. *See* systolic anterior motion of anterior MV leaflet
 TOE evaluation, 248
 mitral valve clipping, 140–1
 mitral valve commissurotomy, balloon, 141
 mitral valve surgery, 3–4, 75–81
 anaesthesia, 77–8
 catheter-based procedures, 140–1
 failure to wean from CPB, 47
 minimally invasive, 5
 MV repair, 79–80
 MV replacement, 80
 pregnancy, 272
 mitral valvotomy, percutaneous transeptal, 77–8
 mixed venous oxygen content (CvO₂), 226
 mixed venous oxygen saturation (SvO₂), 226–8
 during CPB, 42
 paediatric cardiac anaesthesia, 151
 shock, 228
 variables affecting, 228
 modified ultrafiltration, 152
 monitoring
 cardiovascular, 223–8
 deep hypothermic circulatory arrest, 193–4
 heart transplantation, 127
 neurological. *See* neurological monitoring
 off-pump surgery, 92
 paediatric cardiac anaesthesia, 150
 postoperative transfer to ICU, 50–1
 pregnancy, 272
 vascular access, 25–6
 morphine, 25, 275–6
 motor evoked potentials (MEPs), 104–5, 233
 mouth examination, 8
 Müller's sign, 8
 multidetector row computed tomography (MDCT), 15–18
 murmurs, cardiac, 8–9
 muscle relaxants. *See* neuromuscular blocking drugs
Mycobacterium chimaera, 38, 286
 myocardial hibernation, 30–1
 myocardial infarction (MI), acute
 cardiogenic shock, 228
 ECG changes, 12
 perioperative, 47, 55–7
 myocardial ischaemia, 12
 exercise ECG, 12
 intraoperative, 28, 30–2
 postoperative, 51–2
 reperfusion injury, 31
 myocardial oxygen delivery, 30
 myocardial oxygen demand, 30
 myocardial perfusion imaging, 15
 myocardial protection, 3, 30–3
 cardioplegia, 30, 32–3

- ischaemic conditioning, 31
- myocardial stunning, 30–1, 45
- myocardial viability assessment, 15, 19
- myxomas, 255–6
- nail beds, examination, 8
- near infrared spectroscopy (NIRS), 235–6
- aortic surgery, 102
- paediatric cardiac anaesthesia, 151–2
- neck examination, 8
- necrotizing enterocolitis, 184
- neonates
 - cardiopulmonary interactions, 147
 - coarctation of the aorta, 161
 - duct-dependent circulation, 146, 154
 - normal physiology, 145
 - systemic and pulmonary circulations, 146–7
 - transitional circulation, 145–6
- neuraxial blockade, central, 274–8, 282
- neurological complications, 64–7
 - aortic surgery, 104–6
 - cardiac ablation procedures, 136
 - CPB perfusion pressure and, 204
 - deep hypothermic circulatory arrest, 195
 - diagnosis, 65–6
 - management, 66–7
 - neuraxial blockade, 276–8
 - normothermia versus hypothermia, 205
 - paediatric cardiac surgery, 184
 - prevention, 65
 - risk factors, 65
- neurological monitoring, 230–7
 - aortic surgery, 102, 104–5
 - cerebral perfusion and oxygenation, 233–5
 - deep hypothermic circulatory arrest, 197, 231–2
 - neuronal function, 230–3
- neuromuscular blocking drugs
 - effects of CPB, 201
 - induction, 26–7
- neuropathic pain, 280–1
- neuroprotection. *See also* cerebral protection
 - air embolism, 212
 - aortic surgery, 101–2, 104–5, 197
 - paediatric patients, 150–1
- neurosurgery, 215–16
- New York Heart Association (NYHA)
 - functional capacity classification, 6
- nitric oxide (NO), inhaled
 - paediatric patients, 152, 155, 184
- ventricular assist device insertion, 123–4
- nitrous oxide (N_2O), 149
- nociception, 280
- non-compaction cardiomyopathy (NCC), 110
- non-invasive diagnostic tests, 11
- non-steroidal anti-inflammatory drugs (NSAIDs), 51, 282
- norepinephrine, paediatric patients, 151
- Norwood procedure, 167–8
- novel oral anticoagulants. *See direct-acting oral anticoagulants*
- nuclear cardiology, 14–15
- nutrition, postoperative, 182
- obesity, DC cardioversion, 135
- observation, clinical, 7–8
- octreotide, 89
- oedema, 7, 70
- off-pump cardiac surgery, 4–5, 91–4
 - off-pump coronary artery bypass (OPCAB), 4–5, 91–4
 - anaesthetic management, 92–3
 - conversion to on-pump, 93
 - history, 91
 - outcome, 94
 - postoperative care, 93
 - rationale, 91–2
 - surgical approach, 91–2
- Ohm's law, 30
- oliguria, 53
- operating room
 - arrival of patient, 25
 - transfer to, 27–8
 - transfer to ICU from, 50–1
- opioids
 - effects of CPB, 201
 - induction of anaesthesia, 26–7
 - postoperative analgesia, 51, 282
- optical coherence tomography, 23
- oral anticoagulants, direct/novel, 24–5, 263, 278
- osteomyelitis, sternal, 287
- oxygen administration
 - air embolism, 211
 - preanaesthesia, 25
- oxygen consumption (VO_2), 226
- oxygen delivery (DO_2), 226
- oxygen demand, 226
- oxygen extraction ratio (O_2 ER), 226
- oxygen, myocardial supply and demand, 30
- oxygenation
 - during CPB, 208
 - inadequate, during CPB, 212
- oxygenators
 - changeout during CPB, 212
- CPB, 37–8
- ECMO, 218
- failure during CPB, 212
- oxytocin, 271
- pacemakers, permanent (PPMs), 131
 - classification, 131
 - DC cardioversion, 135
 - explantation, 132–3
 - implantation, 131–3
 - perioperative management, 133
- pacing, temporary epicardial, 131
 - AVR, 71, 73
 - heart transplantation, 127
 - intraoperative, 43
 - off-pump surgery, 93
 - postoperative, 51, 54
- paediatric cardiac anaesthesia, 145–53, *See also* congenital heart disease
 - conduct of, 148–50
 - CPB, 150–2
 - drug doses, 151
 - induction, 149–50
 - outcome of surgery, 185
 - post-bypass management, 152
 - postoperative care, 181–5
 - premedication, 148–9
 - preoperative assessment, 148
 - principles, 145–8
 - transfer to intensive care, 153
 - vascular access and monitoring, 150
- pain
 - assessment, 281
 - physiology, 280
 - postoperative. *See* postoperative pain
- palpitation, 8
- palpitations, 7
- pancreatitis, acute, 60
- pancuronium, 26–7
- papillary fibroelastoma, 256
- papillary muscle dysfunction, 78
- rupture, 80
- paracetamol, 51, 282
- paraplegia, 104–6, 278
- parasternal blockade, 278–9
- paravalvular leaks (PVLs)
 - grading of severity, 253
 - percutaneous closure, 141
 - transcatheter aortic valve implantation, 138
- paravertebral blockade, 279
- patent ductus arteriosus (PDA), 146, 170
- patent foramen ovale (PFO), 254
- percutaneous closure, 141
- total anomalous pulmonary venous drainage, 176

Index

- percussion, 8
 percutaneous balloon mitral valvotomy (PBMV), 141
 percutaneous septal closure, 141
 percutaneous therapies. *See* cardiac catheter laboratory procedures
 perfusion pressure, CPB, 203–5
 pericardial disease
 anaesthesia, 110
 imaging, 18–20
 TOE assessment, 256–8
 pericardial tamponade. *See* cardiac tamponade
 pericarditis, constrictive. *See* constrictive pericarditis
 peritoneal dialysis, 64
 paediatric patients, 181–2, 184
 persistent PHT of the newborn, 145, 147
 pharmacodynamics, effects of CPB, 200
 pharmacokinetics, effects of CPB, 199–200
 phenylephrine, 151
 phosphodiesterase (PDE) inhibitors, paediatric patients, 151
 phrenic nerve injury, 136, 184
 pH-stat acid–base management, 42
 α-stat management vs, 206–7
 cerebral protection, 196
 paediatric patients, 150–1
 pharmacokinetic effects, 200
 PHT. *See* pulmonary hypertension
 physical examination, 7–10
 plasma protein binding, during CPB, 199
 platelet counts, 261
 platelet transfusion, 53, 261
 pneumonia, ventilator-associated (VAP), 287–8
 polycythaemia, 188
 positive end-expiratory pressure (PEEP), 51
 positron emission tomography (PET), 14–15, 17
 postoperative care. *See also* intensive care unit
 fast-tracking, 50
 first day, 53–4
 minimally invasive surgery, 97
 off-pump coronary surgery, 93
 paediatric, 181–5
 routine early, 50–4
 postoperative complications, 53, 55–67
 haematological, 267
 paediatric patients, 182–5
 postoperative pain, 280–3
 acute, 281–2
 adverse effects, 281
 causes, 280–1
 chronic, 282–3
 minimally invasive cardiac surgery, 97
 prevention, 281–2
 postoperative pain management, 51, 281–2
 aortic surgery, 105
 chronic, 282–3
 minimally invasive cardiac surgery, 97
 multimodal approach, 282
 non-pharmacological, 282
 postural hypotension, 7
 prasugrel, 262
 praecordium, examination, 8
 prednisolone, for thrombocytopenia, 261
 pregnancy, 269–73
 cardiac disease, 270–1
 cardiac surgery, 271–3
 physiological changes, 270
 preload, 46
 paediatric patients, 182
 postoperative optimization, 55, 57
 premedication, 25
 preoperative assessment, 24–5
 pressure half-time (PH-T), 76
 pressure support ventilation (PSV), 51
 pre-syncope, 7
 primary graft dysfunction, 128
 propofol
 anaesthesia, 26, 28
 pharmacology during CPB, 200
 postoperative sedation, 51
 prostacyclin I₂ (PGI₂), 264
 prostaglandin E₂ (PGE₂), 146
 prosthetic heart valves
 carcinoid heart disease, 89
 malfunction, failure to wean from CPB, 47
 paravalvular leaks. *See* paravalvular leaks
 pregnancy, 272
 TOE evaluation, 253
 transcatheter aortic implantation, 137
 protamine, 43
 Protek Duo catheter, 120
 prothrombin complex concentrate (PCC), 264
 proximal isovelocity surface area (PISA), 77
 pulmonary angioplasty, balloon, 116
 pulmonary artery (PA)
 banding, 171, 177
 catheterization, 26, 225, 227
 iatrogenic injury, 116, 225
 sarcomas, 256
 pulmonary artery floatation catheter (PAFC), 26, 225, 227
 pulmonary artery pressure (PAP), 20, 113
 measurement, 225
 waveform, 227
 pulmonary artery wedge pressure (PAWP), 55
 measurement, 225
 PHT, 113
 pulmonary atresia, 172–3
 pulmonary blood flow (PBF)
 balancing. *See* pulmonary:systemic blood flow balance
 CHD, 147–8
 drug pharmacokinetics, 200
 optimization, univentricular heart, 175
 truncus arteriosus, 179
 pulmonary circulation
 balancing. *See* pulmonary:systemic blood flow balance
 duct-dependent, 146, 154
 perinatal changes, 146
 pulmonary embolism, massive, 228
 pulmonary hypertension (PHT), 113–16
 adult congenital heart disease, 187
 chronic thromboembolic (CTEPH), 113–16
 classification, 113
 failure to wean from CPB, 47
 investigations, 20–1
 management, 113–16
 mitral stenosis, 76
 paediatric cardiac surgery, 183–4
 persistent, of newborn, 145, 147
 postoperative, 152
 RV pathophysiology, 113–15
 truncus arteriosus surgery, 179
 VSDs, 180
 pulmonary oedema, reperfusion, 116
 pulmonary regurgitation (PR), 86–7, 253
 pulmonary stenosis (PS), 85–6, 251
 pulmonary thromboendarterectomy (PTE), 114–16
 pulmonary valve (PV), 85
 absent, 163
 carcinoid heart disease, 87–8
 Ross procedure, 172–4
 TOE evaluation, 251–3
 pulmonary vascular disease, 113–16
 pulmonary vascular resistance (PVR), 20, 113
 CPB, 46–7
 neonates, 145, 147
 partitioning, 20
 right-sided valve surgery, 85–7

- pulmonary vasodilators, 114
- pulmonary:systemic blood flow balance (Q_p:Q_s)
- adult congenital heart disease, 187
 - arterial shunt procedures, 154–5
 - measurement, 20
 - neonates, 146–7
 - Norwood circulation, 167–8
 - postoperative care, 181–2
 - PA banding, 171
 - pulse, 8
 - pulsed-wave Doppler (PWD), 239
 - pulsus alternans, 8
 - pulsus bifidens, 8
 - pumps, blood, 35
 - failure, 213
- Qp:Qs. *See* pulmonary:systemic blood flow balance
- radial artery, cannulation, 25
- radiant heaters, 192
- radiofrequency ablation, 134, 136
- recurrent laryngeal nerve injury, 184
- regional anaesthesia, 274–9
- Caesarean section, 271
 - paediatric cardiac surgery, 150
- regional cerebral tissue oxygen saturation (rSO₂), 102, 235
- regional wall motion abnormalities (RWMA), 243–4
- ReliantHeart aVAD, 121
- REMATCH trial, 118
- remifentanil, 26
- remote ischaemic preconditioning (RIPC), 32
- renal failure. *See also* acute kidney injury
- heart transplant recipients, 129
 - postoperative, 61–4
 - pre-existing, 63–4
- renal replacement therapy (RRT), 64
- paediatric patients, 181–2, 184
- renal tumours, IVC, 216
- reperfusion injury, 31, 116
- respiratory complications
- CPB termination, 46
 - postoperative, 53, 58–9
- respiratory failure
- ECMO, 218
 - postoperative, 53, 58–9
- respiratory rate (RR), 8, 145
- restrictive cardiomyopathy, 109, 111
- resuscitation. *See also* cardiac arrest
- CPB for, 215–17
 - postoperative cardiac arrest, 55, 67
- retrograde cerebral perfusion (RCP), 197
- aortic surgery, 101–2
- massive air embolism, 211–12
- Reveal® implantable cardiac monitoring (ICM) systems, 13
- rewarming, 192
- deep hypothermic circulatory arrest, 195–6
 - postoperative, 51
 - rates, 42
- rheumatic heart disease, 78, 82, 141
- RIFLE classification, 61
- right atrial cannulation, for CPB, 2–3
- right atrial hypertrophy, 12
- right atrial pressure, 20, 225
- measurement, 225–6
 - tricuspid disease, 82, 84
 - waveforms, 224–5
- right bundle branch block (RBBB), 12
- right heart catheterization, 20–1
- right ventricle (RV)
- ejection fraction (RVEF), 245
 - fractional area change (RVFAC), 245
 - free wall strain, 245
 - function, TOE assessment, 244–5
 - hypertrophy, 12
 - PHT, 113–15
 - pressure, 20, 227
- right ventricular (RV) dysfunction/failure
- after pulmonary thromboendarterectomy, 116
 - carcinoid heart disease, 87–8
 - grading, 245
 - paediatric patients, 152
 - postoperative, 56–7
 - TOE assessment, 244–5
 - VADs, 121, 123
- right ventricular assist devices (RVADs), 118
- indications, 119, 123
- right ventricular outflow tract (RVOT) obstruction
- Fallot's tetralogy, 163–4
 - pulmonary atresia, 172
- risk assessment, cardiac surgery, 2
- rivaroxaban, 263, 278
- rocuronium, 26
- roller pumps, 35
- Ross procedure, 172–4
- Ross-Konno procedure, 174
- rSO₂ (regional cerebral tissue oxygen saturation), 102, 235
- S' velocity, tricuspid annulus, 245
- saphenous neuralgia, 282–3
- sarcomas, cardiac, 256
- SAVE (Survival After Veno-arterial ECMO) score, 218
- scars, surgical, 7
- scintigraphy, 14–15
- screen filters, CPB circuit, 38
- ScvO₂ (central venous oxygen saturation), 228
- sedation, postoperative, 51, 105
- seizures, 7
- selective antegrade cerebral perfusion, 101–2, 197
- sensory evoked potentials (SEPs), 233
- septal closure, percutaneous, 141
- serotonin, 88–9
- shock, haemodynamic parameters, 228
- shunt fraction. *See* pulmonary:systemic blood flow balance
- shunts
- bidirectional cavopulmonary (BCPS), 160
 - Blalock-Taussig. *See* Blalock-Taussig (B-T) shunt, modified
 - diagnostic evaluation, 20
 - left-to-right, 20, 147
 - systemic arterial to pulmonary, 147, 187
 - unrestricted, adult congenital heart disease, 187
- sickle cell disease, 262
- single photon emission computed tomography (SPECT), 14, 17
- single ventricle circulation. *See* univentricular circulation
- sinus node dysfunction, 132
- SjvO₂ (jugular venous oxygen saturation), 234
- skin, examination, 7–8
- sodium nitroprusside, 151
- somatic pain, 280
- somatosensory evoked potentials (SSEPs), 105, 233
- spherocytosis, hereditary, 262
- spinal anaesthesia, 275–6
- neuraxial bleeding, 276–8
 - organ donor harvest, 276
- spinal cord
- function monitoring, 104–5, 233
 - injury/ischaemia, 104
 - protection, 104–5, 197
- spinal cord perfusion pressure (SCPP), 104–5
- spinal haematoma, 276
- spongiform cardiomyopathy, 110
- stabilizer devices, off-pump coronary artery surgery, 91
- Stanford classification, aortic dissection, 100
- sternal closure, delayed, 152
- sternal osteomyelitis, 287
- sternotomy
- chronic pain after, 282
- J-shaped upper partial (mini), 73

Index

- sternotomy (cont.)
 - median, 2
 - postoperative mediastinitis, 287
- Stokes–Adams attacks, 7
- stomatocytosis, hereditary, 262
- strain imaging
 - left ventricle, 243
 - right ventricle, 245
- stress echocardiography, 13
- stress imaging, cardiac, 15
- stress response, surgery, 274
- stroke
 - CPB perfusion pressure and, 204
 - diagnosis, 65–6
 - LA appendage occlusion, 142
 - perioperative, 64–7
 - previous history, 65
 - thrombolytic therapy, 263
- subclavian flap angioplasty, 161
- subclavian vein (SCV) cannulation, 25–6, 224
- sufentanil, 26
- surgical site infection, 286–7
- SvO₂. *See* mixed venous oxygen saturation
- Syncardia Total Artificial Heart, 118, 121, 123
- synchronized intermittent mandatory ventilation (SIMV), 51
- syncope, 7
 - AS, 69
 - exertional, 7
- Syntometrine®, 271
- systemic arterial to pulmonary shunts, 147, 187
- systemic circulation
 - balancing. *See* pulmonary:systemic blood flow balance
 - duct-dependent, 146, 154
- systemic inflammatory response syndrome (SIRS), 62
- measures to reduce, 264–7
- paediatric patients, 182
- systemic vascular resistance (SVR), 20
 - CPB, 41, 45–7
 - measurement, 226
 - neonates, 147
 - shock, 228
- systolic anterior motion of anterior MV leaflet (SAM), 80–1, 107
- tachyarrhythmias, paediatric patients, 183
- tachycardia, postoperative, 52
- Takotsubo cardiomyopathy, 110–11
- tamponade. *See* cardiac tamponade
- temazepam, 25
- temperature. *See also* hypothermia
 - after-drop, 195
- cerebral metabolism and, 196
- measures for maintaining, 191
- monitoring, 27, 42, 102
- regulation, 191
- temperature management, 191–8
 - EEG monitoring, 231–2
 - paediatric patients, 150
- tetralogy of Fallot. *See* Fallot's tetralogy
- thermodilution method, 226
- thienopyridines, 24
- thiopental, 26
- thoracic aortic surgery, 99–106
 - anaesthesia, 101–5
 - aortic arch, 101–2
 - aortic root and ascending aorta, 101
 - complications, 105–6
 - descending aorta, 102–5
 - pathology, 99–100
 - postoperative care, 105–6
 - pregnancy, 272
 - preoperative preparation, 100–1
 - spinal cord perfusion, 104–5
- thoracic epidural anaesthesia, 26, 274–5, 282
- neuraxial bleeding risk, 276
- pros and cons, 275
- thoracic surgery, non-cardiac, 215–16
- Thoratec pVAD, 120
- thrombocytopenia, 261
- thromboelastography (TEG), 265
- thrombolytic agents, 263
- thrombophilia, 263
- thrombosis, postoperative, 185
- thrombus, intracardiac, 255
- ticagrelor, 262, 278
- ticlopidine, 278
- tirofiban, 25, 262
- TOE. *See* transoesophageal echocardiography
- total anomalous pulmonary venous drainage (TAPVD), 147, 176
- Total Artificial Heart, Syncardia, 118, 121, 123
- total cavopulmonary connection (TCPC). *See* Fontan procedure
- tracheal intubation, 27
- tranexamic acid (TXA), 152, 264
- transcatheter aortic valve implantation (TAVI), 5, 137–40
 - anaesthesia, 139–40
 - devices and techniques, 137
 - echocardiography, 140
 - patient selection, 138
 - procedural risks, 137
- transcranial Doppler (TCD), 233
- transfusion. *See* blood transfusion
- transitional circulation, 145–6
- transoesophageal echocardiography (TOE), 239–41
- adult congenital heart disease, 189
- aortic disease, 258–9
- ASDs, 254–5
- basic views, 240
- cardiac masses, 255–6
- complications, 239
- indications, 239
- LA appendage occlusion, 142
- MV clipping, 140–1
- off-pump surgery, 92
- paediatric patients, 152
- percutaneous septal closure, 141
- pericardial disease, 256–8
- physical principles, 239
- pregnancy, 272
- probe insertion, 27
- valvular heart disease, 248–53
- VAD insertion, 121, 123
- ventricular function, 242–7
- transposition of the great arteries (TGA), 177–8
- arterial switch procedure, 177–8
- balloon atrial septostomy, 147, 177
- PA banding, 171, 177
- transpulmonary gradient (TPG), 160
- transthoracic echocardiography (TTE), 13
- Traube's sign, 9
- tricuspid annular plane systolic excursion (TAPSE), 244
- tricuspid annulus, S' velocity, 245
- tricuspid regurgitation (TR), 83–5
 - carcinoid heart disease, 87
 - grading of severity, 251
 - TOE assessment, 251
- tricuspid stenosis (TS), 82–3, 251
- tricuspid valve (TV), 82
 - carcinoid heart disease, 87
 - surgery, 83–5
 - TOE evaluation, 251
- troponin I, 57
- truncus arteriosus (TA), 179
- tubing, CPB circuit, 33
- tumours, cardiac, 255–6
- 22q11 deletion (Di George syndrome), 148, 163, 179
- ultrasound, physics, 239
- univentricular circulation
 - arterial shunts, 154–5
 - bidirectional cavopulmonary shunt, 160
- Fontan procedure, 165–6
- postoperative care, 181–2
- pulmonary artery banding, 171
- staged palliation, 175
- urinary catheterization, 27, 54
- urine output, 41, 45, 63

urological surgery, 216
 uterotronics, peripartum, 271

VACTERL syndrome, 163
 VADs. *See* ventricular assist devices
 valvular heart disease
 imaging, 18–19
 indications for surgery, 2
 percutaneous procedures, 137–41
 pregnancy, 272
 prosthetic valves. *See* prosthetic heart valves
 TOE assessment, 248–53
 vascular access, 25–7
 adult congenital heart disease, 189
 aortic surgery, 101–3
 CPB, 40–1
 heart transplantation, 127
 minimally invasive surgery, 95–6
 paediatric cardiac anaesthesia, 150
 vasoconstrictors, paediatric patients, 151
 vasopressin, 151
 vecuronium, 26
 vegetations, 256
 venous cannulae, 33, 95
 venous cannulation
 CPB, 41, 43
 minimally invasive surgery, 95–6
 preoperative, 25
 venous oximetry, 226–8
 venous reservoir, CPB, 34
 ventilation, mechanical
 after induction, 27

 during CPB, 208
 ECMO, 220
 neonates with CHD, 147
 paediatric patients, 181
 post-bypass failure, 46
 postoperative continuation, 51
 weaning from, 51
 ventilator-associated pneumonia (VAP), 287–8
 ventricular aneurysms, 110
 ventricular assist devices (VADs), 118–24, *See also* intra-aortic balloon pump (IABP)
 bridge to recovery, 118–19
 bridge to transplantation, 118–21
 design, 118
 destination therapy, 118–21
 implantation, 121–3
 indications and patient selection, 118
 long-term, 119
 paediatric patients, 183
 postoperative care, 123–4
 preoperative assessment, 120–1
 short-term, 119–20
 ventricular dysfunction
 postoperative, 52, 55–7
 TOE assessment, 242–7
 weaning from CPB, 45
 ventricular dysrhythmias, postoperative, 183
 ventricular fibrillation (VF), 30, 42–3
 ventricular function, TOE assessment, 242–7

ventricular pseudoaneurysms, 110
 ventricular septal defects (VSDs), 180
 percutaneous closure, 141
 PA banding, 171
 pulmonary atresia with, 172–3
 surgical closure, 180
 transposition of the great arteries, 177
 ventricular septum, 180
 ventricular tachycardia (VT), postoperative, 42–3
 ventricular thrombus, 255
 ventriculography, 21–2
 vents, CPB, 39, 95–7
 visceral pain, 280–1
 vocal cord palsy, 184
 volatile anaesthetics, 28, 201
 cardioprotection, 32
 during CPB, 42
 paediatrics, 149
 pharmacology during CPB, 201
 VSDs. *See* ventricular septal defects

warfarin, 114–15, 262–3
 ‘waterhammer’ pulse, 8
 weaning from CPB, 43, 45–9
 causes of failure, 45
 correctable problems, 45–7
 ECMO for failure, 219
 heart transplantation, 127–8
 mechanical support, 47–9, 120
 non-correctable problems, 47
 paediatric patients, 151–2
 pharmacological support, 47