

# Index

- acidosis, 70, 94, 131
  - shoulder dystocia and, 210
- Adverse Outcome Index (AOI), 293
- air embolism, 27
- all-fours position, 199–200
- amniotic fluid embolism (AFE), 26–7
- anaesthetic emergencies, 47–8
  - extubation and recovery, 55
  - failed tracheal intubation, 48–55
  - high regional block, 56–9
  - local anaesthetic toxicity, 59–63
- anaphylactic reactions, 25
- aneurysm rupture, 21
- angina, 24
- anteartum haemorrhage, 136,
  - 146–51, *See also* maternal haemorrhage
- causes, 146
- clinical presentation, 146
- major haemorrhage management,
  - 146–51
  - actions, 149
  - assessment, 150
  - call for help, 149
  - expediting the birth, 150
  - initial management, 146
  - ongoing care, 167
  - situational awareness, 170
- antibiotics, maternal sepsis
  - management, 131
- antihypertensives, 110
  - choice of, 112
- aortocaval compression, 30
- arterial blood gas, 131
- aspiration of gastric contents, 25
  - difficult intubation and, 50
- baseline fetal heart rate, 80
- basic life support algorithm, 14
- bimanual uterine compression, 160
- bladder catheterisation, 160
- bladder filling, 223
- blood products, 141
- blood tests
  - arterial blood gas, 131
  - clotting studies, 131
  - C-reactive protein (CRP), 131
  - full blood count, 130
  - liver function, 131
  - maternal sepsis management, 130–1
  - renal function, 131
  - serum lactate, 130
- B-Lynch suture technique, 165
- brachial plexus injury (BPI), 211
- breech presentation, 230, *See also*
  - vaginal breech birth
  - consequences, 230–2
  - definition, 230
  - evidence and national recommendations, 233
  - predisposing factors, 230
- Burns–Marshall technique, 239
- caesarean section
  - anteartum haemorrhage management, 150
  - general anaesthesia, 48
  - perimortem caesarean pack, 36

## Index

---

- caesarean section (cont.)
  - shoulder dystocia prevention, 194
  - twin births, 247
- carboprost, 162
- cardiac arrest, 30, *See also* maternal collapse
  - aftermath, 44
  - community setting, 32
  - local anaesthetic-induced, 63
  - management, 32–43
    - medication, 43
    - post-resuscitation care, 44
    - recognition of heart rhythms, 38–41
    - team leader role, 36–7
  - potentially reversible causes, 42–3
- cardiac disease, 23–5
  - risk factors, 23
- cardiac tamponade, 43
- cardiorespiratory changes during pregnancy, 30–2
- cardiotocography (CTG), 79–92
  - accelerations, 82
  - antenatal, 94
    - classification, 95
  - baseline rate, 80
  - baseline variability, 81
  - decelerations, 82
  - interpretation, 86–8
  - normal intrapartum CTG, 79
  - pathological, 88
    - fetal blood sampling, 88–92
  - suspicious, 88
  - twin birth, 253
- cell salvage, 142
- cerebrovascular event (CVE), 22
- chest compression, 278
- chest pain, 24
- clavicular fractures, 212
- clotting abnormalities
  - blood tests, 131
  - maternal haemorrhage and, 137
  - with pre-eclampsia, 115
- communication, 5–7
  - with woman and birth partner/relatives, 8
- continuous positive airway pressure (CPAP), preterm birth, 287
- cord clamping, 154
  - hypoxic neonate, 272
  - preterm birth, 287
  - twin births, 256
- cord presentation, 218
- cord prolapse, 218
  - breech presentation, 240
- debrief of parents, 225
- documentation, 225
- management, 220–5
  - assessment for birth, 224
  - birth plan, 223
  - bladder filling, 223
  - call for help, 221
  - contraction reduction, 222
  - digital elevation of presenting part, 222
  - early recognition, 221
  - fetal monitoring, 223
  - maternal positioning, 222
  - neonatal resuscitation, 225
  - post birth, 225
  - training, 225
- perinatal complications, 219
- prevention, 219
- risk factors, 218
- cord traction, 154
- C-reactive protein (CRP), 131
- deep vein thrombosis (DVT), 21
  - prophylaxis with maternal sepsis, 133
- delayed cord clamping. *See* cord clamping
- diabetes mellitus, 23, 193
- difficult intubation, 49
  - optimal positioning, 51

## Index

- pre-oxygenation, 52
- risk factors, 49
- disseminated intravascular coagulation (DIC), 22, 131, 137
  - with pre-eclampsia, 115
- dizygotic twins, 245, *See also* twin births
- documentation
  - cord prolapse, 225
  - eclampsia, 106
  - newborn resuscitation, 283
  - postpartum haemorrhage, 167
  - shoulder dystocia, 208
  - uterine inversion, 267
- drug reactions, 25, 43
- eclampsia, 101
  - documentation, 106
  - incidence, 102
  - management, 102
    - basic support, 102
    - call for help, 102
    - community setting, 108–10
    - eclampsia box, 103
    - hypertension control, 110–12
    - magnesium sulfate emergency regimen, 105
    - plan for labour/birth, 116–17
    - post-birth care, 117
    - seizure control, 104–6, 112
  - monitoring, 112–15
  - presenting features, 102
  - seizures and coma, 22
- electronic fetal monitoring (EFM), 66–8, *See also* fetal monitoring
  - standards, 77–8
  - technical considerations, 77
  - vaginal breech birth, 235
  - with cord prolapse, 223
- emergency response triggers, 15, *See also* maternal collapse
- episiotomy, shoulder dystocia
  - management, 201
- Erb's palsy, 211
- ergometrine, 155
- external cephalic version, 254
- extubation, 55
- failed intubation, 48–55
  - definition, 48
  - management, 50–5
    - algorithm, 52
- fetal blood sampling (FBS), 91–2
  - with pathological CTG, 88–92
- fetal heart rate (FHR)
  - accelerations, 82
  - baseline, 80
  - decelerations, 82
  - tachycardia, 80
  - variability, 81
- fetal monitoring, 66–8, *See also*
  - electronic fetal monitoring (EFM)
  - risk management, 68
  - standards and quality, 71–8
    - informed choice, 73
    - intermittent auscultation in labour, 73–6
    - intrapartum risk assessments, 71
  - training, 69
- fetal oxygen supply, 70–1
  - hypoxia risk factors, 69
  - influences, 71
  - responses to impeded supply, 70–1
- fetal scalp stimulation (FSS), 90–1
- fluid balance, pre-eclampsia and, 113
- fluid resuscitation, 132
  - major obstetric haemorrhage, 138–9
    - blood products, 140–1
    - cell salvage, 142
    - intravenous access, 140
    - point-of-care testing, 142
    - volume to be infused, 140
- forceps use, vaginal breech birth, 239
- full blood cell count, 130

## Index

---

- general anaesthesia, 48
  - indications, 48
- glucose management with newborn
  - resuscitation, 281
- glycogenolysis, 70
- haemorrhage. *See* maternal
  - haemorrhage
- heart disease. *See* cardiac disease
- heart rate, fetal. *See* fetal heart rate
- heart rhythms during cardiac arrest,
  - 38–41
  - non-shockable rhythms, 40–1
  - shockable rhythms, 39–40
- high regional block, 56–9
  - management, 58–9
  - presentation, 56–7
  - risk factors, 57
- humeral fractures, 212
- hydrostatic method for uterine
  - inversion, 265
- hyperglycaemia, 23
- hyperkalaemia, 42
- hypermagnesaemia, 42
- hypertension, 108, *See also* eclampsia;
  - pre-eclampsia
  - management, 110–12
  - antihypertensives, 110
- hypocalcaemia, 42
- hypoglycaemia, 23, 42
- hypotension, fluid resuscitation, 132
- hypothermia, 42
  - therapeutic, 280
- hypovolaemia, 21, 42, 137
- hypoxia
  - fetal responses to impeded oxygen
    - supply, 70–1
  - fetal risk factors, 69
  - maternal cardiac arrest, 42
  - neonatal, 270, *See also* newborn
    - resuscitation
    - cord clamping, 272
    - physiology, 270–1
- hysterectomy, postpartum haemorrhage
  - management, 166
- induction of labour, shoulder dystocia
  - prevention, 195
- intensive care unit transfer, 186
- intermittent auscultation (IA), 73–6
  - abnormal fetal heart rate guidance,
    - 75
  - indications for changing to
    - continuous EFM, 76
  - optimal timing, 73–4
  - practice recommendations, 74
- internal iliac artery ligation, 166
- internal podalic version, 254
- internal rotational manoeuvres, 201,
  - 205
- interventional radiology, postpartum
  - haemorrhage, 165
- intrapartum haemorrhage, 137
- intravenous access, 140
- intubation. *See* difficult intubation;
  - failed intubation
- ischaemic heart disease. *See* cardiac
  - disease
- Klumpke's palsy, 211
- labetalol, 112
- laparotomy, postpartum haemorrhage
  - management, 164
- leadership, 8
- legal claim analyses (LCAs), 294
- liver function test, 131
- local anaesthetic toxicity, 59–63
  - follow-up, 63
  - management, 61–3
    - specific treatment, 63
  - signs and symptoms, 59

## Index

- macrosomia, 192
- magnesium sulfate, 104
  - emergency regimen, 105
  - neuroprotection in preterm birth, 288
  - seizure prevention, 112
  - toxicity, 106
    - emergency protocol, 106
- major obstetric haemorrhage. *See* maternal haemorrhage
- maternal collapse, 14, 15, *See also* cardiac arrest
  - basic life support algorithm, 14
  - causes, 20–7
    - air embolism, 27
    - amniotic fluid embolism (AFE), 26–7
    - anaphylactic or toxic reactions, 25
    - cardiac disease, 23–5
    - cerebrovascular event (CVE), 22
    - disseminated intravascular coagulation (DIC), 22
    - eclamptic seizures and coma, 22
    - haemorrhage, 21
    - hypo- or hyperglycaemia, 23
    - pulmonary aspiration of gastric contents, 25
    - pulmonary thromboembolism, 20–1
    - sepsis, 22
  - management, 16–19
    - continuing treatment, 18–19
    - initial management, 17–18
    - primary obstetric survey, 18
    - secondary obstetric survey, 19
- maternal critical care, 174–5
  - equipment, 186
  - indications, 176
  - investigations, 180
  - long term impacts of near-miss maternal morbidity, 187
  - provision location, 180
  - recognition of the critically ill woman, 176
    - MOEWS charts, 177
  - regular structured review, 180
  - transfer to ICU, 186
  - unique features of, 175–6
- maternal critical care chart, 182
- maternal critical care structured review sheet, 182
- maternal death
  - haemorrhage, 136
  - sepsis, 120
- maternal haemorrhage, 21, 136
  - antenatal risk assessment, 143–4
    - anaemia, 143
    - haemorrhagic disorders, 143
    - maternal weight, 143
    - MOEWS chart use, 144
    - placenta praevia and accreta, 144
    - women who decline blood products, 144
  - anteperium. *See* anteperium haemorrhage
  - intrapartum haemorrhage, 137
  - ongoing care, 167
  - pathophysiology, 137
  - postpartum. *See* postpartum haemorrhage
  - protocol, 138–42
    - blood products, 140–1
    - cell salvage, 142
    - fluid resuscitation, 138–9
    - intravenous access, 140
    - point of care testing, 142
    - volume to be infused, 140
  - situational awareness, 170
- maternal sepsis, 22, 120–1
  - imaging, 133
  - management, 126–33
    - actions within 1 hour, 126–9
    - antibiotics, 131
    - blood cultures, 129

## Index

- maternal sepsis (cont.)
  - blood tests, 130–1
  - call for help, 129
  - clinical examination, 129
  - DVT prophylaxis, 133
  - fluid resuscitation, 132
  - monitoring, 132
  - multi-professional approach, 133
  - oxygen administration, 129
  - removal of source of infection, 133
- mortality, 120
- prevention, 121
- recognition of, 122
- risk factors, 125
- signs and symptoms, 122–5
  - genital tract sepsis, 122–3
  - non-obstetric sepsis, 124–5
- Mauriceau–Smellie–Veit manoeuvre, 239
- McRoberts' manoeuvre, 198–9
- meconium-stained liquor, 77, 279
- metabolic acidosis, 70, 131
- misoprostol, 163
- modified obstetric early warning score (MOEWS) charts, 144, 177
- monozygotic twins, 246, *See also* twin births
- neonatal O-negative blood, 280
- newborn assessment, 92–4, 272–5
- newborn resuscitation, 270, 272–80
  - airway, 275–6
  - assessment at birth, 272–5
  - breathing, 276–7
  - chest compression/circulation, 278
  - cord clamping, 272
  - documentation, 283
  - emergency medication, 279
  - emergency neonatal O-negative blood, 280
  - equipment preparation, 271–2
  - meconium management, 279
  - oximetry and supplementary oxygen, 278
  - post-resuscitation care, 280–1
    - glucose, 281
    - therapeutic hypothermia, 280
  - warmth, 272
- nifedipine, 112
- non-shockable heart rhythms, 40–1
- nuchal arms, 240
- obesity, 193
- obstetric haemorrhage. *See* maternal haemorrhage
- oxytocin, 154, 159
  - twin birth, 253
- patient-reported outcome measures (PROMs), 296
- perimortem birth, 31, 36
  - caesarean pack, 36
- perinatal mortality
  - cord prolapse, 219
  - multiple births, 247
- placenta accreta, 144, 154
- placenta praevia, 144
- positive end-expiratory pressure (PEEP),
  - preterm birth, 287
- posterior arm delivery, 201, 203
- postpartum haemorrhage (PPH), 137, 151–67, *See also* maternal haemorrhage
  - causes, 146–54
  - documentation, 167
  - major haemorrhage management, 155–67
    - bimanual uterine compression, 160
    - bladder catheterisation, 160
    - B-Lynch suture technique, 165
    - call for help, 157
    - continuing management, 167
    - emergency box, 157
    - examination under anaesthetic, 161

## Index

- hysterectomy, 166
- immediate actions, 157
- internal iliac artery ligation, 166
- interventional radiology, 165
- keeping the mother warm, 162
- laparotomy, 164
- manual removal of retained products, 161
- medications, 159–60, 162–3
- ongoing care, 167
- rapid evaluation, 158
- tear repair, 161
- unrelenting haemorrhage, 162
- uterine balloon tamponade, 164
- uterine massage, 159
- uterine vessel ligation, 166
- prevention, 154
  - active management of third stage of labour, 154–5
  - physiological management of third stage of labour, 155–4
- primary, 137
- risk factors, 151
  - intrapartum, 152
  - pre-labour, 152
- secondary, 137
- situational awareness, 170
- with uterine inversion, 261
- pre-eclampsia, 100
  - fetal complications, 101
  - management
    - community setting, 108–10
    - hypertension control, 110–12
    - plan for labour/birth, 116–17
    - post-birth care, 117
    - seizure prevention, 112
    - severe pre-eclampsia, 108
  - maternal complications, 100
  - monitoring, 112–15
    - clotting abnormalities, 115
    - fluid balance, 113
    - pulmonary oedema, 114
  - predisposing factors, 101
- preterm birth, 286–8
  - cord clamping, 287
  - counselling parents, 286
  - CPAP and PEEP, 287
  - neuroprotection with magnesium sulfate, 288
  - surfactant therapy, 288
  - thermal care of the preterm infant, 286
- process measures, 293
- pulmonary aspiration of gastric contents, 25
- pulmonary oedema, 114
- pulmonary thromboembolism, 20–1
- quality indicators (QIs), 293–5
- quality of care, 299
  - definition, 292
  - information presentation, 297
  - measurement, 291–2
    - clinical quality indicators (QIs), 293–5
    - culture of unit, 296
    - patient reported outcome measures (PROMs), 296
    - process measures, 293
    - source data, 295–6
- recombinant factor VIIa, 163
- renal function test, 131
- respiratory distress syndrome (RDS), 288
- safety culture, 296
- Saving Babies Lives Care Bundle, 69
- seizures
  - eclampsia, 102
    - magnesium sulfate emergency regimen, 105
    - management, 104–6
    - prevention, 112



## Index

- sepsis, 121, *See also* maternal sepsis
  - prevention, 121
  - recognition of, 122
  - signs and symptoms, 122–5
    - genital tract sepsis, 122–3
    - non-obstetric sepsis, 124–5
- serum lactate, 130
- Severity Index (SI), 293
- Sexton Safety Attitudes Questionnaire, 297
- shockable heart rhythms, 39–40
- shoulder dystocia, 190–1
  - consequences, 210–12
    - acidosis, 210
    - brachial plexus injury, 211
    - humeral and clavicular fractures, 212
- documentation, 208
- incidence, 191
- management, 195–207
  - after the birth, 208
  - all-fours position, 199–200
  - call for help, 197
  - episiotomy need evaluation, 201
  - gaining internal vaginal access, 202
  - internal rotational manoeuvres, 205
  - McRoberts' manoeuvre, 198–9
  - posterior arm delivery, 203
  - suprapubic pressure, 200
  - tertiary manoeuvres, 206
  - time limit, 206
  - what to avoid, 207–8
- prevention and antenatal counselling, 194–5
  - caesarean section, 194
  - labour induction, 195
- recognition of, 197
- risk factors, 191–3
  - gestational age, 193
  - macrosomia, 192
  - maternal diabetes, 193
  - obesity, 193
  - operative vaginal birth, 198
  - previous occurrence, 192
- situational awareness, 9–11
- source data, 295–6
- suprapubic pressure, 200
- surfactant therapy, 288
- symphysiotomy, 206
- Syntometrine, 154, 159
- tachycardia, fetal, 80
- teamwork, 2
  - communication, 5–7
  - definition, 3
  - high reliability and resilience, 3–4
  - leadership roles and responsibilities, 8
  - situational awareness, 9–11
  - training, 2
    - local training, 3, 4–5
    - under pressure, 11
- tension pneumothorax, 43
- therapeutic hypothermia, 280
- thromboembolism, 43, *See also*
  - deep vein thrombosis (DVT);
  - pulmonary thromboembolism
- timely cord clamping. *See* cord clamping
- tocolysis, uterine inversion
  - management, 264
- training
  - cord prolapse management, 225
  - fetal monitoring, 69
  - teamwork, 2
    - local training, 3, 5
- tranexamic acid (TXA), 159–60
- transfer or critically ill woman, 186
- twin births, 245–7
  - management, 250–3
    - first stage of labour, 250
    - preparation, 252
    - second stage of labour, 252
    - third stage of labour, 256
  - mode of birth, 247–9



## Index

- presentation at birth, 247
  - procedure, 253–6
    - delayed cord clamping, 256
    - external cephalic version, 254
    - internal podalic version, 254
    - length of inter-twin birth interval, 256
  - timing of birth, 249
  - umbilical cord prolapse. *See* cord prolapse
  - uterine atony, 153, 159
  - uterine balloon tamponade, 164
  - uterine compression, 160
  - uterine inversion, 260
    - definition, 260
    - diagnosis, 261
    - documentation, 267
    - management, 262–7
      - continuing management, 266
      - debriefing, 267
      - hydrostatic method, 265
      - immediate action, 262
      - manual replacement, 264
      - resuscitation, 263
      - surgical management, 266
      - treatment, 263
      - uterine relaxants, 264
    - risk factors, 260
  - uterine massage, 159
  - uterine rupture, 153
  - uterine vessel ligation, 166
  - uterus, manual displacement, 31
  - vaginal breech birth, 230, *See also*
    - breech presentation
    - complications, 240–1
      - cord prolapse, 240
      - failure to assist birth of after-coming head, 240
    - fetal risks, 241
    - head entrapment during preterm birth, 240
    - nuchal arms, 240
  - management, 232–9
    - assisted manoeuvres, 236–9
    - electronic fetal monitoring, 235
    - first stage of labour, 234–5
    - second stage of labour, 235
  - types of, 232
- Weighted Adverse Outcome Score (WAOS), 293
- white blood cell count (WBC), 130
- Zavanelli manoeuvre, 206