

Index

- absorption
 - RF power, 118
 - due to Provtotorov, 119
- acceptance drift
 - geometric and dead time, 439
- acceptance of detector, 284
 - in each kinematic bin, 436
- acceptor, 100
- acoustic phonon
 - emitted energy flux, 340
 - velocity of, 231
- acoustic velocity, 83
- adiabatic
 - transition, 29
- adiabatic condition
 - ignoring relaxation, 442
- adiabatic demagnetization, 33
 - cooling by, 35
 - in rotating frame, 35, 38, 95
- adiabatic passage
 - in rotating frame, 39, 96
- ADRF, 38, *See adiabatic demagnetization*
- alcohol, 288
- alignment, 17, 19, 20
- alkanes, 288
- amine, 288
- aminoxy
 - 4-hydroxy-TEMPO, 148
 - OH-TEMPO, 148
 - oxo-TEMPO, 148
 - TEMPO, 148
 - TEMPOL, 148
- ammonia
 - second-order spin couplings, 225
- ammonia beads
 - irradiation, 312
 - preparation by freezing liquid droplets, 311
 - preparation by slow freezing and crushing, 311
- ammonium borohydride, 288
- analyte
 - molecule under study, 309
- angular momentum
 - classical definition of, 2
 - intrinsic, 3
 - law of conservation, 2
 - orbital, 3
 - quantum mechanical definition, 3
 - vector, 2
- anhydrous magnesium sulphate
 - water removal, 303
- anisotropic hyperfine tensor, 227
- anomalous magnetic moment, 4, 98
- APRF, 40, *See adiabatic passage in rotating frame at constant entropy*, 443
- density matrix with two temperatures, 443
- high-temperature models, 444
- magnetic ordering
 - loss of reversibility, 444
- mixing field, 443
- of polarized proton targets, 460
- polarization loss mechanisms, 444
- APRF reversal of deuteron spins, 445
- APRF reversal of proton polarization
 - superradiance, 445
- Arrhenius law, 293
- attenuation
 - in oversized waveguide, 400
 - rectangular guide, 400
- azephenylenyls, 148
- beam normalization, 439
- beam polarization, 437
- beam transmission, 290
- Bessel function, 401
 - root of, 401
- BHHA-Cr(V), 149
 - proton hyperfine spectrum, 228
- binomial coefficients, 108
- biradical, 147, 309
 - in cross effect, 183
 - TOTAPOL, 147
- bisdiphenylallyl (BDPA), 148

510

Index

- bisdiphenylene-b-phenylallyl, 148
- Bloch equations, 87
- Bloch function
 - of conduction electron, 235
- Bloch state
 - of conduction electron, 235
- Bohr magneton, 4
- boiling heat transfer
 - at low pressure, 344
 - correlation for ^4He , 345
 - Cu-dilute solution, 345
 - dryout heat flux limit, 346
 - forced convection, 346
 - nucleate, 345
 - pool boiling, 346
 - propanediol- ^3He , 345
- bolometer
 - microwave detector, 140
 - used for study of FM, 193
- Boltzmann constant, 18, 279
- Boltzmann distribution, 18, 24, 59, 66, 246
- Boltzmann factor, 18
- Boltzmann ratios, 189
- Born approximation, 452
- boron compounds, 288
- Bose-Einstein condensation
 - ^4He , 481
- Bose-Einstein statistics, 80
- Breit-Rabi polarimeter, 431
- bremssstrahlung
 - energy loss by, 286
- Brillouin function, 19, 21, 29, 30
- brute force polarization
 - ^{27}Al , ^{93}Nb , ^{165}Ho , 422
 - metal hydrides, 422
 - of nuclei in metals, 422
 - solid ^3He , 423
 - solid HD, 423
- BWO, 463, *See* backward wave oscillator
- canonical
 - density of protons, 115
 - distribution, 24
- CGS Gaussian system of units, 471
- charged glassy beads
 - discharging, 307
- chemical DNP, 425–427
- chemical potential
 - of ^4He , 367
- chemical shift, 55, 477
 - anisotropic, 56
 - magic angle spinning (MAS), 229
- chemical spin polarization
 - atomic hydrogen, 425
 - CIDNP, 427
 - parahydrogen induced polarization, 427
- Clausius-Clapeyron equation, 381
- coaxial line
 - characteristics, 503
 - electrical length, 250
 - phase transition in PTFE, 267
 - propagation constant, 250
 - semi-rigid, 267
 - thermal drift of attenuation, 267
 - thermal drift of length, 267
- cobalt-iron alloy
 - magnet pole pieces, 409
- coherent scattering
 - polarized slow neutrons, 290
 - coherent slow neutron scattering, 452
- cold neutron beam
 - polarization by magnetic supermirrors, 452
- commutation relations, 5
- commutation rules, 32
 - exponential operators, 12
- COMPASS collaboration
 - triple target, 450
- concentrated stream
 - energy balance, 358
- conduction electrons
 - cooling of, 36
 - paramagnetic, 100
- configurational degrees of freedom
 - loss of, 292
- contamination
 - by water frost, 331
- convectiveal heat transfer
 - hydrodynamic surface area, 347
 - in dilute solution, 346
- cooling cycle
 - dilution refrigerator, 355
 - evaporation refrigerator
 - ^3He , 350
 - ^4He , 350
- cooling of the rotating sample
 - in MAS NMR, 463
- cooling power
 - evaporation refrigerator, 351
- counterflow heat exchanger
 - effectiveness, 375
- coupling constant
 - spin-orbit, 51
- coupling resistance, 249
- Cr(V) complexes
 - in stable pure form, 304
 - preparation with diols, 302
- Cr(V)-diol reaction
 - sensitivity to light and water, 302
- critical temperature
 - for magnetic ordering, 460
- critical velocity
 - for zero viscosity, 481

- cross effect
 - for MAS-DNP, 183
 - simulation using TEMPO, 184
- cross effect DNP, 463
- cross relaxation, 46, 79
 - flip-flop terms, 133
 - inhomogeneous broadening, 167
 - nuclear spins, 115
 - rate parameter, 184
 - role of nuclear spins, 135
 - spin pair flip-flop transitions, 133
- cross relaxation parameter, 183
- cross section
 - difference for opposite orientations, 284
 - unpolarized, 284
- cross-relaxation
 - between different nuclear spin species, 239
- crystallization
 - hindered nucleation, 292
- Curie law, 21, 78
 - correction to, 22
- Curie temperature, 22, 23, 32
- Curie-Weiss law, 22
- cut-off wavelength
 - rectangular guide, 399
- damping resistance
 - resonant circuit, 252
- Debye
 - cut-off, 126
 - frequency, 231
 - model, 80
 - temperature, 80, 132, 341
- Debye frequency, 479
- Debye model
 - phonons in glassy materials, 231
- Debye temperature, 479, 480
- density matrix, 23, 24, 73, 155
 - high-temperature approximation, 25
 - linearized, 25
 - off-diagonal elements, 25
- deuterated ammonia
 - post-irradiation, 314
- deuteron NMR signal
 - errors in asymmetry, 269
- differential solid effect, 463
- diffusion
 - nuclear spin, 230
 - of nuclear spin polarization, 242
- diffusion barrier
 - for nuclear spins, 242
- diffusion constant
 - in dilute stream, 367
- dilute solubility line
 - of phase diagram, 482
- dilute solution
 - degenerate Fermi fluid, 366
 - isenthalpic expansion, 366
- dilute stream flow
 - limited by diffusion, 366
- dilution factor
 - for pure chemical substance, 286
 - of PT material, 284
 - of target
 - mean over kinematic bins, 437
- dilution refrigerator
 - cooling power, 357
 - counterflow heat exchanger, 357
 - design of flow channels, 362
 - enthalpy in the dilute stream, 483
 - evaporator and condenser, 355
 - expansion device, 355
 - main heat exchanger, 356
 - maximum cooling power, 357, 361
 - mixing chamber, 356
 - optimum flow, 360
 - optimum flow of ^3He , 357
 - phase boundary, 356
 - principle of operation, 355
 - still, 355
 - diol, 288
- diphenyl picryl hydrazyl (DPPH), 148
- dipolar broadening
 - fourth moment, 207
 - of EPR line, 111
 - of NMR line, 206
 - second moment, 206
- dipolar energy
 - quantum mechanical, 45
- dipolar field, 32
 - local, 112
- dipolar frequency, 69, 72, 113
- dipolar interaction
 - energy, 24
- dipolar lineshape
 - at high polarization, 207
 - moments at high polarization, 208
- dipolar spin-spin interactions
 - secular part, 164
- dipolar temperature, 61, 133
- dipolar width, 113, 114
 - EPR line, 115
- Dirac term
 - scalar interaction, 184
- direct process
 - relaxation rate due to, 84
- dispersion relation
 - for ^4He superfluid, 481
- dissolution-DNP (dDNP), 183, 465

512

- dissolution NMR
 - DNP before rapid melting, 309
- Dittus and Böltner correlation, 376
- divacancy, 100
- DNP, 155
 - by cross effect, 182
 - by Overhauser effect, 187
 - cross effect at low temperatures, 183
 - diffusion barrier, 180
 - dynamic cooling of the spin-spin interactions, 155
 - enhancement of MAS NMR signal, 309
 - in fluorene
 - using photoexcited phenanthrene, 321
 - in irradiated $\text{Ca}(\text{OH})_2$, 319
 - in irradiated CaF_2 , 319
 - in irradiated glassy butanol, 320
 - in scintillating plastics, 308
 - irradiated deuterated butanol, 321
 - leakage factor, 179
 - of hyperfine nuclei, 184
 - Overhauser effect, 155
 - in liquid state, 190
 - in solids, 190
 - solid effect, 155
 - time evolution, 174
 - using cross effect, 309
 - via electron spin locking, 429
 - with frequency modulation, 191
 - with inhomogeneously broadened EPR line, 167
- DNP in crystalline materials
 - using substitutional paramagnetic ions, 322
- DNP MAS NMR spectroscopy, 462
- donor, 100
- double balanced mixer, 249
- doublet
 - non-Kramers, 125
- D-state probability
 - deuteron, 201
- dynamic cooling
 - inhomogeneous broadening, 163
 - of electron dipolar interactions
 - at high temperature, 161
- dynamic cooling of nuclear spins
 - phenomenology, 156
- dynamic nuclear polarization. *See DNP*
- effective damping
 - frequency sensitivity, 252
- effective field, 38
- effectiveness method
 - design of heat exchangers, 374
- EHBA-Cr(V), 149
 - preparation, 304
 - proton hyperfine spectrum, 228
- eigenfunction, 25
 - of spin operator, 6
- eigenvalues
 - of spin 1 Hamiltonian, 216

Index

- EIO, 463, *See extended interaction oscillator*
- elastic coherent scattering
 - forward peaked, 455
- elastic peak
 - for free protons, 289
- elastic scattering
 - elastic peak, 439
- electric field
 - asymmetry parameter, 49
 - gradient, 49
- electric quadrupole moments
 - of nuclei, 200, 203
- electromagnetic compatibility (EMC), 279
- electromagnetic interference (EMI)
 - control of, 279, 382
- electron dipolar interactions
 - neglect of, 172
- electron-nucleus spin interaction
 - dipolar, 227
 - hyperfine, 227
- Ellis-Jaffe sum rule, 450
- EMC solenoid
 - main and trim coils, 418
- EMI control
 - design of signal paths, 280
 - hardening by circuit design, 280
 - RF interferences, 279
- energy conservation
 - in dynamic cooling, 165
- energy reservoirs
 - electron dipolar, 156
 - electron Zeeman, 156
 - nuclear Zeeman, 156
- enthalpy
 - dilute solution, 357
 - helium isotopes and their mixtures, 483–485
 - of helium mixtures, 483
 - of the concentrated stream, 484
 - osmotic, 483
- enthalpy-pressure diagram
 - Joule-Thomson cycle, 351
- entropy, 28, 31, 35
 - additive quantity, 28
 - equilibrium, 27
 - excess between liquid and crystal, 293
 - in high magnetic field, 29, 30
 - loss of, 39
 - maximum for a spin system, 28
 - measurement of, 29
 - of ^4He , 367
 - of a composite system, 28
 - of a spin system, 24
 - quantum statistical definition, 28
 - relation with polarization, 460
 - with dipolar interactions, 32
- entropy theory
 - glass transition, 293

- EPR saturation, 172
- EPR spectrometer
 - high field, 138
 - V-band, 139
 - X-band, 135
- EPR spectroscopy, 135
 - in situ*, 140
- EPR spectrum
 - from NEDOR signal, 146
- equilibrium thermodynamics
 - in a glass, 292
- equivalent surface-to-volume
 - heat transfer surface, 339
- equivalent surface-to-volume diameter
 - flow resistance parameter, 340
- European Muon Collaboration (EMC)
 - CERN NA2 twin target, 448
- evaporation refrigerator
 - with ^3He - ^4He mixture, 353
- exchange interaction, 40, 55, 58
- exchange narrowing, 118
- Fabry-Perot interferometer
 - semiconfocal, 139
- false asymmetry
 - estimation and mitigation, 439–441
- fast neutron beam
 - polarization using polarized proton target, 290
- F-center, 100
- F-centers
 - in glassy organic materials, 149
- Fermi contact interaction, 54
- Fermi distribution function, 236
- Fermi pseudopotential, 453, 458
- Fermi surface, 57
 - of conduction electrons, 236
- Fermi-Dirac statistics, 481
- fibre-optic links
 - replacing galvanic lines, 280
- FID. *See* free induction decay
 - time scale, 88
- FID envelope, 67
- figure of merit
 - for a scattering experiment, 436
 - material dependent part, 285
 - maximization in intense beam, 285
 - of PT, 284
 - of the target, 437
- filling factor
 - of probe coil, 261
- film boiling
 - critical heat flux, 380
- finite element (FE)
 - numeric simulation, 413
- flash evaporation, 353
- flip-flop term, 46
- fountain pressure
 - in ^4He superfluid, 481
- free electron
 - trapped by solute molecule, 150
 - trapped in a cavity, 150
- free induction decay, 62
- free precession signal
 - non-exponential, 88
- free radical, 147
 - radiolytic, 99
 - stable, 99
- free radicals
 - table, 475
- free-volume model
 - glass transition, 293
- Fremy's salt, 148
- frequency modulation
 - during DNP, 463
 - effect on EPR spectrum, 143
 - effect on NEDOR signal, 146
 - repopulation of hyperfine lines, 195
- frozen spin operation
 - coolant density measurement, 440
 - during field rotation, 379
 - transverse field mode, 379
- fusion with polarized fuel, 290
- gamma function, 498, 500
- gaseous helium
 - specific heat, 486
- Gaussian function
 - normalized, 71, 497
- gel matrix
 - immobilizing radicals, 465
- general relativity, 2
- g*-factor
 - anisotropic, 101, 114
 - coupling with the lattice, 101
 - Dirac value of, 4, 98
 - free nucleon, 199
 - quark
 - in naïve quark model, 200
 - structural, 4
- Gibbs free energy, 28
- glass
 - organic, 132
- glass former
 - characteristics, 292
- glass transition
 - 1,2-propanediol, 299
 - microcalorimetry, 296
- glass transition temperature, 292
- glassy material, 70, 115
 - delayed heat release, 299
 - mould casting, 308
- glassy materials
 - chemically doped, 291
 - properties of the matrix, 292

514

Index

glassy target beads
 preparation of, 306
 Glebsch-Gordan coefficients, 49, 201
 gluon, 2
 contribution to nucleon spin, 200
 Gomberg's triphenylmethyl radical, 148
 Gorter's formula
 for relaxation time, 236
 gradient tensor
 electric field, 48
 group velocity
 rectangular guide, 399
 g -shift
 paramagnetic electron, 100
 unpaired electron, 101
 gyromagnetic factor, 45
 gyromagnetic ratio, 3, 21, 199
 sign of, 5
 gyrotron, 463
 hadron, 2
 Hamiltonian, 47
 dipolar, 31, 45
 secular part, 33
 electron exchange, 58, 118
 Hermitian, 69
 phonon, 82
 secular part, 47
 spin, 13, 17
 spin orbit, 51
 Zeeman, 17, 33, 45
 heat capacity
 of spin system, 131
 heat exchanger
 continuous-flow, 358
 heat flush effect, 355
 for measurement of osmotic pressure, 485
 heat transfer
 in sintered sponge, 359
 sintered powder and liquid, 492
 heat transport
 by evaporating superfluid film, 349
 subcooled ^4He superfluid, 349
 Heaviside-Lorentz system of units, 473
 helium gas purifiers, 386–388
 operation, 388
 helium pump
 leak testing, 386
 Root's blower, 384
 rotary blade pump, 385
 helium recirculation systems
 active purifiers, 385
 adsorbent purifiers, 386
 pump oil purge, 385
 shaft seal, 386
 high-intensity beam
 direct heating of target, 438
 radiation damage, 438

high-resolution NMR technique
 pulsed NMR, 462
 high-temperature approximation
 interacting spin system, 31
 horn antenna
 coupling structure, 463
 hydraulic surface area, 370
 hydrodynamic surface area, 493
 hydrogen
 free molecule wavefunction, 223
 solid spin conversion, 224
 hyperfine constant
 isotropic, 107
 hyperfine interaction, 54
 dipolar term, 184
 neglect in DNP, 173
 hyperfine splitting, 107
 hyperfine tensor, 107
 anisotropic, 54, 107, 109, 114
 isotropic, 54
 hyperpolarization, 421, 460
 deuterated glucose, 466
 impedance
 rectangular guide, 399
 incomplete inner shells, 98
 indirect coupling between nuclear spins, 55
 indirect nuclear spin interaction
 pseudo-dipolar, 226
 pseudo-exchange, 226
 indirect spin interaction
 NMR lineshape, 222
 integral theorem of Cauchy, 64
 integrated beam fluxes
 through target, 436
 integrated NMR signal
 linearity, 270
 International System of units (SI), 471
 inverse temperature, 34
 irradiated ammonia
 preparation, 311
 irradiated crystalline materials
 F-centers, 151
 H-centers, 151
 irradiated lithium hydride
 structure and properties, 314
 irradiated materials
 radical yield, 310
 irradiated PT materials, 309
 irradiation
 safety aspects with liquid N_2 , 322
 irradiation damage
 bleaching by light, 326
 in irradiated ammonia, 327
 repair by annealing, 325
 irradiation effects
in situ, 324

- isospin mirror nuclei, 201
- isotropic hyperfine tensor, 227
- isotropic space, 2, 6
- ITS-90 scale
 - possible improvement using NMR, 278
 - vapor pressure of helium, 276
- ITS-90 temperature scale, 381
- Kapitza conductance, 171, 492
 - Cu-³He, 341
 - Cu-⁴He, 341
 - deuterated butanol-dilute solution, 344
 - epoxy-³He, 343
 - FEP-dilute solution, 343
 - glassy butanol-⁴He, 343
 - paramagnetic salts-⁴He, 342
- Kapitza conductances, 341
- Kapitza resistance
 - acoustic phonon mismatch, 340
 - heat transfer, 306
 - solid ortho-hydrogen, 224
 - thermal boundary resistance, 340
- kinematical bin, 436
- Knight shift, 55, 56, 229, 477
- Korringa constant, 36, 238
 - effect of impurities, 238
 - with weak coupling, 239
- Korringa law, 422, 428
- Korringa relation
 - between relaxation and Knight shift, 238
- Kozeny-Carman equation, 370
- Kramers doublet
 - in YES, 424
- Kramers-Krönig equations, 247
- Kramers-Krönig relations, 64, 65, 76
- Kramers-Krönig transform, 499
- Lagrange multipliers, 34
- lambda-transition, 481
- Landé g-factor, 51, 101
- large superconducting magnets
 - examples, 417
- Larmor frequency, 17
 - chemical shift, 229
- Larmor precession, 11
 - classical treatment, 12
 - frequency of, 21, 246, 477
- lattice interactions, 76
- lattice stress
 - interstitial radiolytic centers, 314
- leakage factor, 173
- leakage mechanism
 - for nuclear spin polarization, 94
- level gauge
 - for liquid helium, 384
- Ley's radical, 182
- linear distortion
 - of wide NMR signal, 269
- linear response
 - approximation, 119
 - theory, 60, 162, 204
- lineshape
 - Gaussian, 134
 - Lorentzian with cut-offs, 114
 - quadrupolar, 49, 50, 210
 - in high field, 210
 - lineshape functions, 496
 - moments of, 67, 68, 496
- linewidth
 - EPR contribution of nuclear spins, 115
- Liouville formalism, 78
- liquid nitrogen
 - explosion following irradiation, 322
- lithium deuteride ⁶LiD
 - irradiated, 288
- lithium hydrides
 - DNP results, 315–319
 - high radiation length, 288
- LMN
 - crystal growth, 323
- local field, 33, 44, 72
- Lorentzian function, 71, 498
- Mach number
 - at freezing radius of atomic beam, 430
- magnet pole pieces
 - shimming, 412
- magnet system
 - CEN-Saclay Nucleon-Nucleon experiments, 419
- magnetic cooling, 29
- magnetic dipole interaction, 43–48
 - energy of, 17
- magnetic dipole moment
 - of nuclei, 199, 200
 - of nucleon, 199
- magnetic dipole operator
 - for a nucleus, 200
- magnetic energy levels, 9
- magnetic field
 - transverse, 48
- magnetic induction, 43, 113
- magnetic moment
 - density, 20
- magnetic moments
 - table, 476
- magnetic ordering
 - due to exchange interaction, 58
 - loss of entropy, 460
- magnetic resonance, 58
- magnetic scalar potential, 43
- magnetization, 21
 - evolution of, 16
- static, 20, 246

516

Index

main heat exchanger
 axial conduction, 365
 viscous heating, 365
 MAS-DNP
 sensitivity enhancement, 229
 mass diffusion
 of dilute solutions, 490
 mass flowmeters
 calorimetric, 383
 matrix representation
 of spin operator, 7–9
 spin 1 vector, 216
 Maxwell-Boltzmann distribution, 157, 159
 mechanical vacuum model, 491
 memory functions, 78
 metallo-organic complexes, 477
 metallo-organic compounds, 100, 149
 methane, 288
 spin conversion in solid, 224
 microwave counter
 with harmonic downconversion, 408
 microwave power
 measurement by calorimeter, 408
 microwave source
 backward wave oscillator (BWO)
 carcinotron, 393
 extended interaction oscillator (EIO), 394
 IMPATT diode source, 395
 klystron, 394
 measurement of frequency and power, 408
 miniature gas turbines
 for MAS NMR, 462
 mixing field, 34
 modified Lorenzian function
 moments of, 499
 modified Lorenzian functions, 498
 molecular dioxygen (O_2), 147
 molecular field effect, 23
 molecular spin effects, 57
 molecular spin isomers, 222
 MRI contrast enhancement, 464
 MRI sensitivity enhancement, 464
 multimode cavity, 463
 multiple scattering, 285, 437, 438
 multiple targets in same beam
 mitigation of drift sources, 448
 mutual friction
 in dilute solution below 0.25 K, 491
 in dilute stream, 366
 ND_3
 irradiated, 288
 NEDOR spectroscopy, 145
 nitronyl nitroxides, 148
 NMR circuit
 crossed-coil, 272

NMR coil
 Q-factor, 252
 NMR line broadening
 Van Vleck formulas, 227
 NMR measurement of polarization
 principles, 245
 NMR saturation
 by polarization measurement
 neglect of, 172
 NMR signal
ab initio size, 253
 calibration for polarization measurement, 275
 continuous-wave (CW), 260
 saturation, 256
 signal-to-noise ratio, 254
 non-equilibrium phenomenon
 liquid-glass transition, 292
 normalized Gaussian function, 497
 nuclear magnetic cooling, 35
 nuclear magnetic ordering, 36, 39, 40, 460
 in Ag, 40
 in $AuIn_2$, 40
 in $Ca(OH)_2$, 41
 in CaF_2 , 41
 in Cu, 40
 in LiH, 41
 in Rh, 40
 nuclear magnetic phases
 by neutron diffraction experiments, 461
 nuclear magneton, 477
 nuclear Overhauser effect
 NOE sequence
 I. Solomon, 429
 nuclear spin polarization
 by brute force, 421
 nuclear spin relaxation
 in dielectric solids, 235
 in metals, 235
 polarization dependence, 232
 nuclear spin-lattice relaxation
 shell-of-influence model, 230
 nucleation of crystal growth
 effect of impurities, 297
 in slow cooling, 296
 nucleon, 2
 Nusselt number, 374, 375
 operation in high beam flux
 frozen spin mode, 373
 operation in high beam fluxes
 choice of refrigeration method, 372
 operator
 exponential, 12
 ladder, 6
 optical pumping
 by collisional spin-exchange (SEOP), 432

- by metastability-exchange (MEOP), 431, 432
- ground-state and metastable atoms and
 - ions, 431
 - in dense gas, 430
- optically pumped ^3He
 - MRI contrast medium, 291
- optimum filter
 - pre-filtering and cross-correlation, 446
- Orbach expansion
 - lattice potential, 124
- orbital moment
 - nuclear, 53
 - quenched, 53
- orbital motion
 - contribution to nucleon spin, 200
- oriented single crystal
 - ellipsoidal, 461
- ortho-hydrogen, 223
- osmotic pressure
 - loss in dilute stream, 366
 - of dilute solutions, 485
- Overhauser effect, 427, 464
 - DNP, 463
 - in liquid phase using TEMPO, 464
 - in metals and liquids, 427
- para-hydrogen, 223
- paramagnetic compounds, 147
- paramagnetic ion
 - Nd^{3+} in LaAlO_3 , 149
 - Nd^{3+} in LaF_3 , 149
 - Nd^{3+} in LMN, 149
 - Tm^{2+} in CaF_2 , 149
- paramagnetic oxygen
 - magnetic liquid, 147
- paramagnetic substance
 - diluted, 98
- paramagnetic susceptibility
 - static, 21, 99
- paramagnetism, 98
- partition function, 24, 28, 34
- parton, 2
- Pascal's triangle, 108
- Penning trap, 4, 477
 - single-electron, 98
- perdeuterated propanediol
 - reduction of potassium dichromate, 303
- permeability
 - of free space, 20
- perturbation theory
 - time-dependent, 16
- phase diagram of $^3\text{He}/^4\text{He}$ mixtures, 481
- phase sensitive detection
 - real part of Q-meter signal, 249
- phase separator
 - ^4He liquid-gas, 378
- phase velocity
 - rectangular guide, 399
- phonon, 80
 - ballistic, 130
- phonon bottleneck, 130
 - DNP speed limiting, 165
 - Kochelaev model, 132
 - with frequency modulation, 132
- phonon energy, 82
- phonon spectrum, 479
- phonons
 - in ^4He superfluid, 481
- Planck constant, 2
- PLTS-2000
 - provisional low temperature scale, 486
 - provisional temperature scale, 277
- point defect
 - interstitial, 100
 - substitutional, 100
- Poiseuille flow
 - laminar region, 369
- Poisson's equation
 - vector form, 412
- polarization, 17, 18
 - adiabatic reversal of, 34
 - measurement, 22, 60
 - measurement of, 21
 - tensor. *See* alignment
 - vector, 17
- polarization asymmetry
 - in scattering experiments, 283
- polarization by spin selection
 - in atomic beam, 430
- polarization of nuclei in large biomolecules, 454
- polarization reversal
 - by adiabatic passage in rotating frame (APRF), 442
 - by DNP, 441
 - by field rotation, 450
 - by spin rotation, 441
 - diabatic, 37
 - efficiency of, 39
 - optimum filter theory, 446
- polarization reversal frequency
 - optimization, 445
- polarized target refrigerator
 - SMC double cell target, 379
 - Virginia-Basel-SLAC target, 377
- polarizing agent (PA)
 - in MAS-DNP, 309
 - generic name for a free radical, 464
- porphyrroxide (PX), 148
- porphyrindene (PB), 148
- powder
 - of small crystals, 70

518

Index

- power absorption
 - due to spin flips, 59
- power supplies
 - separated analog and digital parts, 280
- Poynting vector, 400
- Prandtl number, 374
 - for gaseous helium, 491
- precooling heat exchangers
 - pressure drop, 376
- pressure measurement
 - capacitive gauge, 382
 - McLeod gauge, 383
 - Pirani gauge, 383
 - vacuum gauges, 383
- probe coil
 - filling factor, 258
 - reduced current, 265
 - sampling function, 257
 - size compared with wavelength, 258
- proton NMR spectrum
 - hyperfine protons, 228
- Provotorov equations, 38, 60, 63, 72, 74, 75, 88, 160, 161
 - generalized, 78
- pseudo-dipolar coupling, 57
- pseudo-exchange coupling, 57
- pseudomagnetic field
 - due to strong interaction, 459
- pseudomagnetism, 457
- PT refrigerator
 - target holder insert, 379
 - target insert, 378
- PTB-2006 scale
 - forthcoming new temperature scale, 486
 - provisional scale, 382
- pulse response
 - Fourier transform, 205
- pycril-N-aminocarbazyl (PAC), 148
- Q*-curve
 - drift of, 255
- Q-meter
 - circuit optimization, 265
 - complex circuit theory, 257
 - Liverpool circuit, 249
 - optimum tuning, 266
 - parallel tuned, 249
 - probe coil
 - series tuned, 256
 - Q-curve, 251
 - series tuned, 249
 - signal expansion, 251
- quadrupole energy
 - sign of, 214
- quadrupole interaction, 48–50
 - asymmetry parameter, 214
 - coupling parameters, 221
 - in low field, 215
- quadrupole moments
 - table, 476
- quadrupole resonance
 - due to magnetic interaction, 215
- Quantum Chromodynamics (QCD)
 - testing based on nucleon spin, 200
- quantum fluids, 480
- quantum statistical treatment, 16, 120
- quantum statistics, 27
 - of a system of spins, 17
- quark, 2
- quark model, 199
- quenching of the angular momentum, 52
- quenching rate
 - glass formation, 292
- radiation damage
 - chemically doped hydrocarbons, 325
 - in lithium hydrides, 329
- radiation length, 286
- radiation resistance
 - of PT materials, 324
- radicals
 - alcoxy, 150
 - in irradiated alcohols, 150
 - in irradiated alkanes, 150
 - in irradiated ammonia, 151
 - in irradiated aryls, 150
 - radiolytic centers, 147
 - radiolytic free radical, 466
 - radiolytic radicals
 - in glassy organic materials, 149
- Raman process
 - relaxation rate due to, 84
- rare earth group, 477
- rare spin species
 - polarization measurement, 276
- rectangular waveguide
 - characteristics, 501
- refrigerator for polarized target
 - precooling heat exchangers, 373
- relative permeability
 - magnet material, 410
- relaxation
 - Blume and Orbach process, 128
 - cross-relaxation, 90
 - dipolar, 87
 - direct leakage, 165
 - direct process, 124, 128
 - effective rate with phonon bottleneck, 132
 - electron spin-lattice, 3, 101, 122
 - electron spin-spin, 122
 - nuclear spin, 230
 - nuclear spin-lattice, 94
 - of transverse magnetization, 87
 - Orbach process, 127, 128
 - Orbach-Aminov process, 127

- Raman process, 125
- spin-lattice, 18, 76, 78
 - experimental, 128
- Waugh and Slichter process, 128
- relaxation measurement
 - NEDOR method, 129
 - pulse recovery method, 128
- relaxation mechanisms, 16
- relaxation of the dipolar energy
 - loss of entropy, 460
- relaxation time
 - determined from pulse recovery, 480
 - dipolar
 - from saturated lineshape, 138
 - from pulse recovery, 137
 - from saturation spectrum, 137
 - from signal height, 137
 - in situ* NEDOR measurement, 146
- reluctance
 - of magnetic circuit, 410
- Remotely Enhanced Liquids for Image Contrast (RELIC), 464
- reservoir
 - dipolar, 38
 - Zeeman, 38
- residual gas analyzer, 384
- resonator impedance, 249
- reversal of nuclear spin polarization, 94
- Reynolds number, 374
- RF field
 - transverse, 25
- RF filters
 - in power feeds, 280
- rotating frame, 15, 23, 161
 - spin resonance in, 12
 - transformation, 13
- rotation operators, 5
- rotons
 - in ${}^4\text{He}$ superfluid, 481
- round waveguide, 404
 - circular polarization, 401
 - dominant mode, 401
 - guide wavelength, 404
 - phase velocity, 404
 - propagation constant, 404
 - propagation impedance
 - TE_{11} mode, 404
 - TM_{01} mode, 405
- round waveguides, 503
- Rudermann-Kittel interaction, 40
- saturation, 16, 47, 72, 155
 - BPP phenomenological model, 119
 - effect of frequency modulation on, 193
 - effect on DNP, 174
 - electron spin resonance, 118
 - error in polarization measurement, 261
- function, 119
- NMR lineshape, 76
- Provotorov equations, 119
- strong, 133
- time constant, 260
- saturation decay
 - experimental
 - deuteron NMR signal, 264
- saturation error
 - deuteron NMR signal measurement, 262
- saturation function, 119
- saturation of NMR signal
 - error due to, 256
- saturation parameter
 - in cross effect, 182
- scattering amplitude
 - neutron beam traversing a polarized target, 458
- Schrödinger equation, 13
 - time-dependent, 9
- scintillating detector
 - polarized by DNP, 308
- second (thermal) sound waves, 481
- second quantization, 83
- self-inductance
 - definition, 257
- semi-rigid coaxial cables
 - replacing flexible cables, 280
- series Q-meter
 - capacitively coupled, 272
 - design for deuteron NMR, 268
 - design for proton NMR, 268
 - improvements, 271
 - probe coil design, 270
 - using a hybrid bridge, 274
 - with quadrature mixer, 274
- shock wave
 - at atomic beam skimmer, 430
- sintered heat exchanger
 - design and evaluation, 369
 - thermal penetration, 371
- sintered heat exchangers
 - continuous approximation, 371
- sintered powder
 - extended surface, 359
- skin depth, 257, 400
 - anomalous, 400
- slow drift
 - beam and acceptance, 448
- SMA connectors, 280
- small-angle neutron scattering
 - time resolved, 456
- SMC magnet system
 - solenoid and dipole coils, 419
- SMC polarized target
 - microwave isolator, 380
 - mixing chamber, 380
 - sintered copper heat exchanger, 380

520

Index

SMC solenoid
 main coil and trim coils, 418
 solenoid winding
 gap filling technique, 417
 solid effect, 47, 174
 cavity losses, 179
 differential, 181
 diffusion barrier, 180
 leakage factor, 179
 phonon bottleneck, 181
 rate equations, 175
 resolved, 175
 semi-phenomenological model, 174
 transition rates, 175
 solid effect DNP, 463
 solutions of ${}^3\text{He}$ in ${}^4\text{He}$
 spontaneous separation in two phases, 355
 sources of noise
 RF oscillator, 279
 thermal, 279
 spallation neutron source, 457
 specific heat, 33, 36
 of a spin system, 29
 specific heat anomaly
 at glass transition, 292
 specific heat of liquid ${}^3\text{He}$, 483
 spectral diffusion, 135
 spectral redistribution
 cross relaxation, 159
 spherical polar coordinate system, 43
 spin
 components of, 2
 effective, 3
 intrinsic, 2
 maximum projection of, 2
 quantum number, 7
 temperature, 17
 vector, 2
 spin contrast variation, 290
 small-angle scattering of neutrons, 454
 spin filter
 transmission of neutrons, 451
 transmitted neutron polarization, 451
 spin filtering, 290
 spin filtering of neutron beams, 450
 spin filtering of neutrons
 by Sm, ${}^3\text{He}$, TiH_2 , 452
 spin flip coil
 neutron beam polarization reversal, 455
 spin packet, 135
 dipolar width, 135
 spin packets
 quadrupolar lineshape, 212
 with individual Zeeman temperatures, 183

spin pairs
 like, 46
 unlike, 46, 47
 spin polarization
 matrix, 25
 spin refrigerator, 463
 by rotating YES crystals, 424
 polarized target, 323
 spin temperature, 25
 in low effective field, 23
 in rotating frame, 25
 inverse, 21
 measurement, 28
 spin-orbit coupling, 106
 spin-orbit interaction, 50
 split-coil solenoid magnet
 SLAC E143 experiment, 419
 stable atomic hydrogen
 as polarized electron target, 433
 stable isotopes, 476
 standard waveguides, 502
 state vector
 spin 1 exact solution, 220
 static field
 internal, 21
 statistical accuracy
 dominated by systematic errors, 438
 of polarization asymmetry, 436
 statistical error
 of target asymmetry, 437
 statistical fluctuations
 multiplicative and additive components, 446
 statistical uncertainty
 minimization, 438
 storage ring
 muon, 4
 strong saturation
 transverse field, 159
 substructure, 98
 superconducting magnet, 413
 rectangular conductor, 413
 round conductor, 413
 with split coils, 413
 superconducting magnets
 control and protection, 417
 superconducting solenoid
 end compensation, 415
 field on axis, 414
 winding accuracy, 415
 supercooled liquid, 292
 superfluid ${}^3\text{He}$, 481
 superfluid film creep, 349, 353, 481
 superfluid state
 ${}^4\text{He}$, 481
 superisolation, 330

- superradiance
 - of NMR circuit, 271
- surface-to-volume diameter, 370, 493
- susceptibility
 - complex, 66
 - static, 20, 67, 119
- systematic error
 - due to slow drift
 - beam and acceptance, 438
- Taler and Taler correlation, 376
- target asymmetry, 436
- target cavity
 - bolometer, 407
 - coupling slots, 407
 - microwave field strength, 407
 - multimode structure, 407
- target dilution factor, 437
- target length, 437
- target materials
 - storage and handling, 330
- target thickness, 437
- target thickness variation
 - in dilution refrigerator, 440
 - in evaporative cooling, 440
- Taylor expansion, 25
- Taylor series
 - of density matrix, 31
- temperature
 - dipolar, 159
 - Zeeman, 159
- temperature measurement
 - during NMR signal calibration, 381
 - of helium bath, 276
 - using resistance thermometers, 382
- tetramethylsilane marker, 229
- thermal conductivity
 - in gaseous helium, 492
 - of helium isotope mixtures, 487
- thermal equilibrium, 18
 - between spins and lattice, 275
 - calibration, 22
- thermal exchange
 - effective surface area, 494
- thermal mixing, 36, 39, 93
- thermal penetration depth, 494
 - dilute solution, 371
- thermodynamic functions, 27–28
- thermodynamics
 - at high polarization, 34
 - second law of, 28
- thermomolecular effect, 353
- toluene
 - glass former, 301
- transition metal ions, 100
- transition metals, 98, 477
- transition probability, 16, 119
- transport properties
 - liquid-glass transition, 293
- transverse field, 38
 - strong limit, 94
- transverse susceptibility, 246
 - absorption part, 59, 76
 - complex, 246
 - dispersion part, 59
 - general features
 - complex, 205
 - in NMR, 204
- triplet state
 - optically induced, 100
- triradical, 309
- tris(2,4,6-trichlorophenyl)methyl radical
 - TTM, 148
- trytyl radicals
 - Finland-D36, 148
 - OX063, 148
 - triphenylmethyl, 148
- tune shift
 - attenuation in coaxial line, 267
 - coaxial line, 252
- twin target
 - acceptance drift elimination, 450
- two-fluid model
 - for ^4He superfluid, 481
- two-phase boiling
 - heat transfer, 306
- two-phase region
 - liquid and gas, 480
- units
 - CGS Gaussian, 1
 - MKSA, 1
 - Système International (SI)*, 1
- unpolarized cross section, 437
- unpolarized dummy
 - target tests, 441
- V_1 -center, 100
- vacuum-insulated dewar
 - for biological samples
 - low loss, 330
- Van Vleck formulas
 - electron-nucleus interaction, 227
- Van Vleck's
 - second moment, 113
- Van Vleck's method of moments, 68
- vapor pressure
 - of ^3He , 276
- ITS-90 temperature scale, 486

522

Index

- viscosity
 - glass former, 293
 - in gaseous helium, 492
 - of liquid helium isotope mixtures, 489
- von Neumann-Liouville equation, 23, 67
- vortex pinned to walls
 - flow of ^3He in ^4He , 368
- vortex tangle
 - of superfluid ^4He , 368
- wave vector
 - of phonon, 81
- waveguide
 - complex propagation constant, 398
 - hollow metallic, 396
- propagation modes, 396
- rectangular, 396
- round, 401
- single mode band, 396
- single mode propagation constant, 398
- waveguide components, 405
- waveguide window, 405, 406
- Wigner-Eckart theorem, 49, 52
- yttrium ethyl sulphate (YES)
 - spin refrigerator, 424
- Zeeman energy, 18, 29, 48
- Zeeman temperature, 61, 133
 - inverse, 73