

---

## Index

- $2 \times 2$  table, 80, 101, 124, 201  
   odds ratio, 56, 81, 82, 84, 89  
   sampling scheme, 81  
   test, 83, 88, 89  
 adjacency matrix (definition), 210  
 affine independence (definition), 24  
 algorithm for MLE computation  
   Baum–Welch, 161  
   ECM, 154  
   EM, 46, 96, 150–155, 157, 161, 162,  
     242, 253, 255, 263, 264  
   Fisher scoring, 45, 127, 153, 171  
   iterative proportional scaling, 233  
   MCMC-ML, 46, 214–218, 222, 224,  
     225, 251, 252  
   Newton–Raphson, 45, 127, 153, 171,  
     233  
   pseudo-ML, 215, 216, 224, 225, 251  
   PX-EM, 163  
   quasi-Newton, 45, 153  
   TM, 46  
 ancillarity  
   definition, 136  
   distribution-constant, 73  
   Efron–Hinkley statistic, 139  
   exact, 124, 137–140, 142, 266  
   extended, *see* S-  
   first-order, 60  
   local, 138, 141  
   nonunique, 138  
   S-, 47–49, 51, 53, 95, 137, 141, 190,  
     206, 235, 249  
 ANOVA model, one-way, 209, 231  
   area–interaction model, 253  
 Barndorff–Nielsen’s formula, 73, 139  
 Basu’s theorem, 59, 60, 79, 142, 199  
 Behrens–Fisher model, 119, 127, 131  
 Bernoulli graph (definition), 211  
 binary data, 19, 97, 166, 169, 241, 248,  
   256  
 binary fission, 123  
 binomial inverse theorem, 163  
 Boltzmann constant, 18  
 Boltzmann’s law, 18, 100, 102, 115, 244  
 Box–Cox transformation model, 185  
 Bradley–Terry model, 114–117, 243  
 Bradley–Terry/Mallows model, 244  
 calibration of items, 231  
 calibration, multivariate, 259  
 canonical link (definition), 167  
 censored data, 122, 126, 127, 143  
 coefficient of variation, when known, 26,  
   118, 138  
 complete family, *see also* Basu’s  
   theorem, *see* exponential family,  
   completeness  
   definition, 59  
 concentration matrix, 12, 121, 195  
 conditional distribution, 12  
 conditional distribution given canonical  
   component, 40, 41, 48, 49, 77, 88  
 conditional distribution given canonical  
   statistic, 4, 101  
 conditional independence structure  
   for contingency table, 201–205  
   for Gaussian data, 195–200  
   for mixed data, 207–209  
   under symmetry, 199  
 conditional inference, *see also* principle  
   ancillarity in curved family, 136  
   example, 48, 51, 53, 55–57, 79, 136,  
     142, 196, 211, 217, 224, 232, 249  
   with nuisance parameter, 46, 67, 77, 85  
 conditionality principle, *see* principle  
 confidence region, 44, 67, 86, 93, 108,  
   119, 126, 216, 222, 265

- relation with test, 76
- consistency,  $\sqrt{n}$ -, 65, 66, 69, 149, 150, 155
- contingency table, *see also*  $2 \times 2$  table, 9, 83, 165, 193, 201–205, 234, 239
  - large sample test, 96, 113, 204
- convolution model, 145
- correlation
  - as single parameter, 124, 131, 135
  - partial, 198–200
  - Spearman rank, 78, 108, 244
- covariance lemma, by conditioning, 196
- covariance selection model, *see also*
  - conditional independence structure, Gaussian data, 13, 55, 121, 191, 195
- Cox regression, 186
- Cramér–Rao inequality, 61–63
- curvature of model (definition), 129
- curved model, *see* exponential family
- cut, *see also* ancillarity, S-
  - definition, 47
- Darmonis, Koopman and Pitman, 26
- decomposability, 194–209
- deviance, *see also* LR test
- deviance residuals, 174, 179
- deviance statistic, 90, 92, 173, 204, 239
- diagnostic model check, 5, 77, 247, 266
- directional data, 16
- dispersion parameter, 175, 177–179, 182
- distribution
  - beta, 15, 22
  - binomial, 4, 34
  - Boltzmann, 18, 20
  - Cauchy, 27, 147, 154
  - CG, *see* conditionally Gaussian
  - compound Poisson, 182
  - conditional, *see* conditional
    - distribution
    - conditionally Gaussian, 205–209
    - exponential, 5, 10, 53, 98, 169, 177
    - Fisher’s, for directions, 17
    - folded binomial, 147
    - folded normal, 144, 150
    - gamma, 10, 53, 72, 74, 98, 169, 177, 178
    - Gaussian, *see* normal
    - geometric, *see* time to first success
    - hypergeometric, 83
    - hypergeometric, generalized, 56
    - inverse Gaussian, 22, 35, 58, 72, 181
    - Ising, *see* Ising model
    - Lévy, 23, 35
    - logarithmic, 22, 147
    - marginal, *see* marginal distribution
    - Maxwell–Boltzmann, 17
    - microcanonical, 100, 101
    - mixture, *see* mixture models
    - multinomial, 24
    - negative binomial, 21, 57, 146, 154, 180
    - normal
      - multivariate, 12, 13, 55, 88, 195, 205
      - univariate, 1, 11, 41, 48, 54, 78, 169
    - Poisson, 5, 169
    - Rayleigh, 22
    - skew-logistic, 31
    - time to first success, 10
    - truncated, 143, 260
    - von Mises, for directions, 16, 108, 142
    - Weibull, 27
    - wrapped normal, 147
  - distribution-constant statistic, *see*
    - ancillarity, definition
  - dose–response, 56, 119, 166
  - edge problem, 253
  - Efron–Hinkley ancillary, 139
  - EM, *see* algorithm for MLE
  - embedding, 14, 71, 85, 105, 107, 266
  - entropy
    - canonical, 111
    - microcanonical, 110, 112
  - Erdős–Rényi model, 211, 217
  - ERGM, *see* exponential random graph
  - exact test
    - example, 78–84, 97, 198, 234, 244
    - generally, 76–110, 127
  - exponential dispersion model, 181, 189
  - exponential distribution, *see* distribution
  - exponential family characteristics
    - completeness, 59, 85
    - convexity of  $\Theta$ , 27
    - convexity of  $\log C(\theta)$ , 29
    - cumulants, 30
    - curved, 26, 76, 118–142, 155, 168, 200, 226, 241
    - full (definition), 26
    - higher-order moments, 30
    - Legendre transform, 36
    - linear, 35, 39, 147, 164, 167, 169, 180–183
    - minimal, *see also* minimal
      - representation

- minimal (definition), 24
- moment-generating function, 30
- nonregular, 28, 33, 35, 36, 251
- order (definition), 24
- partition function (definition), 19
- prime (definition), 35
- regular (definition), 28
- semi-invariants, 30
- steep, 28, 34, 35
- structure function (definition), 4
- exponential random graph, general, 225
- exponential tilting, 14, 36, 85, 105
- F-test for variance ratio, 80
- factor analysis
  - binary data, 241
  - Gaussian model, 161
  - regression scores, 163
- Fisher's exact test, 80
- frailty models, 146
- gamma distribution, *see* distribution
- gamma function, 53, 270
- generalized linear model, 7, 164–190, 239, 241, 245, 249
  - dynamic, 256
- geographic information system, 249
- Gibbs model, 253
- graphical Gaussian models, 14
- graphical model, *see* conditional independence structure
- grouped data, 143
- Hammersley–Clifford theorem, 193, 208, 212, 252
- Hardy–Weinberg equilibrium, 156, 258
- hidden Markov model, 153, 159, 255
- hierarchical log-linear model, 202, 203
- HMM, *see* hidden Markov model
- hyperparameter, 255
- incidental parameters, 49, 67, 231, 232
- incomplete data, 123, 143–255
- information inequality, *see* Cramér–Rao information orthogonality, 42, 43, 181, 247
- insurance pricing, 187
- inverse Gaussian, *see* distribution
- inversion of partitioned matrices, 268
- Ising model, 18, 19, 21, 219
- isostatistical region, 101
- item response function, 241
- iterative MLE computation, *see* algorithm for MLE computation
- Jacobian matrix (definition), 269
- Kalman filter/smoothing, 255
- Laplace transform, 14, 30, 59, 71
- latent trait model, 241, 242
- latent variables, 159, 161, 227, 241, 255
- Lehmann–Scheffé theorem, 85
- Lexis ratio, 97
- life-time data, 122, 123, 126, 143, 145
- likelihood concepts, definition
  - marginal, 255
  - orthogonality, 42
  - partial, 186
  - profile, 43
  - pseudo-, 214
  - quasi-, 183
- likelihood ratio test, *see* LR test
- linear exponential family, *see* exponential family, linear
- linear model, normal, 11, 49, 58, 164, 184
- link function (definition), 164, 167
- logistic regression, 7, 34, 50, 51, 56, 166, 180, 214, 242
- logit function (definition), 6
- logit/probit or logit/normal model, 241
- LR test, *see also* deviance, 95, 155
- Lugannani–Rice formula, 70
- Mallows'  $\theta$ -model, 244
- marginal distribution
  - canonical component, 40, 41, 48, 49, 57, 77, 87, 194, 205, 206
  - canonical statistic, 3
  - for incomplete data, 148
  - parameter-free, *see* Basu's theorem
  - test statistic, 79, 199
- Markov chains, 15
- matched pairs study, 51
- Maxwell–Boltzmann model, 17
- MCMC-ML, *see* algorithm for MLE
- microcanonical distribution, 100, 101
- minimal representation, 24–26
- minimal sufficiency, 26
- missing data, 146
- mixture model
  - normal, 205
- mixture models, 209
  - Cauchy, 147
  - HMM, 159
  - negative binomial, 147
  - normal, 145, 150, 155

- moment method of estimation, 54
- multivariate normal, *see* distribution
- multivariate regression model, 119, 120
- negative binomial, *see* distribution
- Nelder and Wedderburn, 164
- Neyman structure, 85, 128
- Neyman–Pearson lemma, 85
- Neyman–Scott ML inconsistency, 49, 67, 231
- nonlife insurance, 187
- odds ratio, 56, 81, 82, 84, 89
- orthogonality
  - information, 42, 47, 139, 181, 247
  - likelihood, 42
- overdispersion, 146, 180, 184, 189
- $p^*$  formula, 73, 139
- parameterization type
  - canonical, 3
  - mean value, 38
  - mixed, 76
  - natural, 3
- partial likelihood, 34, 186
- Pearson  $\chi^2$  test, 88
- Per Martin-Löf’s philosophy, 100
- phase transition, 21, 217, 252, 254
- pivot, 76
- Poisson model
  - distribution, *see* distribution, Poisson
  - multiplicative, 9, 21, 57, 99
  - spatial, 246
  - two samples, 48, 98, 136, 140
- Poisson trick, 95–98, 154, 157, 165, 236, 239
- positron emission tomography, 262
- precision index, 136, 190, 266
- prediction, 108, 163
- principle
  - conditioning, 47, 53, 77, 95, 136, 137, 217, 232, 249, 251, 266
  - exact test, 108
  - repeated sampling, 5, 265
  - sufficiency, 4, 266
  - weak likelihood, 266
- probit function, 119, 166, 242
- probit regression, 119, 128, 166, 168, 180
- profile likelihood, 43, 54, 67, 131, 185, 187, 252
- proportional hazards, 186
- Pseudo-ML, *see* algorithm for MLE
- quadrat counting test, 247
- R package/function, 197, 245
  - BradleyTerry2, 116
  - RSIENA, 226
  - clogit, 51
  - eRm, 233
  - ergm, 225
  - glm, 183, 184, 240
  - spatstat, 246, 247
  - for HMMs, 159
- rank data models, 243
- Rao’s test, *see* score test
- Rao–Blackwell theorem, 59
- Rasch models, 34, 96, 228–242
- rating factors, 188
- redundancy
  - canonical, 112
  - microcanonical, 110
  - Shannon’s, 112
- REML, 49
- reparameterization lemma, 37, 39, 43, 56, 58, 67, 73, 125, 131
- repetitive structure, 3, 102
- S-ancillarity, *see* ancillarity, S-
- saddlepoint approximation, 14, 70–72, 74, 87, 105, 124, 139, 140
- sample (definition), 3
- saturated model (definition), 173
- scan statistic, 80
- score test
  - example, 88, 89, 97, 98, 199, 234–238
  - theory, 86–88, 91, 95, 133, 155, 174
- seemingly unrelated regressions, *see* SUR model
- shaved dice example, 123, 144
- similar test, 85, 86, 128
- Simpson’s paradox, 201
- Spearman’s model, *see* correlation
- star of edges (definition), 212
- Stirling’s formula, 72–74, 141, 270
- Strauss model, 35, 41, 219, 246, 249–254
- SUR model, 119–121, 131, 200
- survival data, 122, 126, 186
- symmetric polynomials, 232
- test for  $H_0$ :
  - common binomial, 97
  - common Poisson, 98
  - conditional independence, 198, 204
  - exponential density, 98
  - fit of Rasch model, 234–240
  - Gaussian mean value, 79
  - Gaussian variance, 78

- homogeneity, 80, 81, 88, 96
- independence, 96
- independence in  $2 \times 2$  table, 80, 81, 88
- lack of correlation, 80
- Poisson multiplicativity, 80, 81, 88, 99
- specified coefficient of variation, 98
- specified multinomial, 98
- variance equality, 79
- total time at risk, 123
- triplets interaction process, 253
- Tweedie model, 182, 189
- unidentifiability, 144, 145, 150
- uniform density family, 15
- variation independent parameters, 42, 43, 47, 77, 137, 141, 235, 247
- Wald test
  - example, 199
  - theory, 92, 95, 128, 155
- Yule process, 123
- Yule–Simpson effect, 201