

Index

- Π_1^1 comprehension, 462
- α -equivalence, 21, 352
- β -reduction, 303, 354
- η -reduction, 354
- \exists -free formula, 386
- ι -reduction, 376
- ι -reduction (iota), 376
- λ -saturated model, 136
- ω -consistent, 330
- n -type, 136
- absolute formula, 134
- absorption laws, 66
- Ackermann function, 212
- admissible rule of inference, 35, 40
- algebraically closed field, 347
- almost negative formula, 386
- analysis, 440, 449
- analytic hierarchy, 426
- analytic set, 427
- ancestors of a node of a tree, 12
- application (in the lambda calculus), 302
- applicative structure, 372
- arithmetic
 - definability, 241
 - hierarchy, 241
- arithmetic comprehension axiom, 441
- arithmetic transfinite recursion, 464
- arity, 2
- assumption rule, 38, 139
- automorphism, 3
- axiom
 - of choice, 397, 412, 423, 477
 - of dependent choice, 423
 - of infinity, 6, 424, 476
- Baire space, 312
- Beth model, 80
 - generalized, 79
- Beth's definability theorem, 180
- binder, x, 21, 84
- boldface definability, 427
- Boolean algebra, 66
- Boolean-valued model, 123
- bootstrapping, 219
- Borel set, 427
- bound variable, 21
- bounded
 - lattice, 66
 - quantifier, 197, 222, 241
- branch of a tree, 13
- c. e. set, 290
- calculus of inductive constructions, 479
- Cantor space, 312, 459
- Cantor's pairing function, 200
- cardinality
 - of a language, 133
 - of a model, 133
- categoricity, 130
- characteristic function, 196
- children of a node of a tree, 12
- choice axiom, *see* axiom of choice
- Church's thesis, 296, 390
- Church–Rosser property, 303, 356
- classical logic, x, 28
- clause, 55
- closed term, 19
- co-analytic set, 427
- coding, 199
- collection schema, 239, 249
- combinatory algebra, 372
- compactness
 - failure in second-order logic, 414
 - of a topological space, 459
 - of first-order logic, 128
 - of propositional logic, 62
- complement in a lattice, 66
- complete
 - Boolean algebra, 123
 - Heyting algebra, 123
 - lattice, 77
 - set of connectives, 47
 - theory, 322
- complete diagram, 133
- complete induction, 223, 238
- completeness
 - of classical first-order logic, 113, 124
 - of classical propositional logic, 61, 67
 - of classical second-order logic with Henkin semantics, 413

- of intuitionistic first-order logic, 121, 126
- of intuitionistic propositional logic, 73, 75, 77
- composition, 193
- comprehension schema, 410
- computable
 - function, 281
 - functional, 310
 - relation, 281
 - set, 281
- computable analysis, 440, 450
- computably enumerable set, 290
- computably inseparable sets, 293, 314, 331
- computation sequence, 298
- conditional function, 197
- configuration of a Turing machine, 298
- conflict driven clause learning (CDCL), 56
- confluence, 303, 356
- congruence closure, 96, 109
- conjunctive normal form (CNF), 47
- conservative extension, 40, 102
- consistent set, 61, 113
- constant domain schema, 122
- constructive logic, *see* intuitionistic logic
- context, 37, 379, 479
- contraction, 353
- convergent rewrite system, 356
- conversion rule, 480
- corner quotes, 260
- course-of-values recursion, 205, 223
- covering notion, 79, 162
- Craig's theorem, 334
- Curry's paradox, 339
- Curry–Howard correspondence, 379
- currying, 353
- cut
 - in a theory of arithmetic, 444
 - rank, 151
 - rule, 138, 139, 145
- cut elimination
 - for first-order logic, 151
 - for second-order logic, 417
- cut-free proof, 139
- Davis Putnam procedure, 56
- de Bruijn index, 25
- De Morgan's laws, 45
- decision problem, 286
- deduction theorem, 34
- definability, 127
 - arithmetic, 241
 - boldface, 427
 - implicit, 180
 - of a class of structures, 127
 - of a relation in a structure, 127
 - second-order, 414
- defined term, 107, 280
- definite description, 101, 474
- definition by cases, 197
- definitional equality, 479
- definitional extension, 104
- degree
 - many-one, 309
 - Turing, 309
- denotation, 110
- denotational semantics, 297
- dense linear order, 342
- dependent function type, 481
- dependent type theory, 478
- derivation, 31
- derived rule, 40
- descendents of a node of a tree, 12
- descriptive set theory, 427, 440
 - effective, 450
- Diaconescu's trick, 474
- diagonal set, 287
- diagonalization, 210
- diagram, 133
- Dialectica interpretation, 401
- Diller–Nahm interpretation, 404
- disjunction property, 166, 172
- disjunctive normal form (DNF), 47
- distributive lattice, 66
- distributivity, 77
- double-negation
 - shift, 97, 122
 - translation, 51, 97, 240
- dovetailing, 200
- downward persistent formula, 134
- DPLL procedure, 56
- dual lattice, 67
- duality (for Boolean algebras), 67
- effectively continuous function, 312
- effectively open set, 312
- Ehrenfeucht–Fraissé game, 130
- eigenvariable condition, 86, 169
- elementarily equivalent
 - structures, 130
- elementary embedding, 133
- elimination rule, 36, 480
- embedding, 3, 133
- empty type, 381
- endomorphism, 3
- entailment, 60, 112
- epsilon calculus, 190, 191
- equational fragment of first-order logic, 92
- equational logic, 94
- equisatisfiable, 55
- equivalence
 - of formulas, 43
 - up to renaming, 21
- essentially \exists -free, 386
- essentially undecidable, 333
- evaluation, 18, 210
- ex falso sequitur quodlibet, 30
- exclusive or, 47

- existential
 - quantifier, x
 - formula, 89
 - quantifier, 84
- expansion, 132
- explicit definability property, 166, 172, 390
- explicit definition, 194
- extension, 218, 369
- extensional equality, 202
- extensionality, 369, 373, 374, 410, 476
- F (second-order lambda calculus), 431
- fast-growing hierarchy, 211
- filter, 68
- finite character, 34
- finite separation principle, 257
- finite type arithmetic, 378
- finite types, 351
- finitistic reasoning, 323
- Fitch diagram, 41
- fixed-point theorem, 293
- forcing
 - classical, 81, 123
 - in a Kripke model, 71
- formation sequence, 7
- foundation axiom, 476
- free variable, 23
- freely generated, 16
- Frege system, 55
- fresh variable, 86
- Friedman–Dragalin translation, 101, 256
- full second-order model, 413
- function extensionality, 369
- Glivenko’s theorem, 53
- Gödel sentence, 335
- graph, 128
- greatest common divisor, 224
- greatest lower bound (in a lattice), 65
- halting computation sequence, 298
- halting problem, 286, 322
- Harrop formula, 174, 391
- Heine–Borel principle, 460
- Henkin
 - constant, 113
 - set, 113
 - structure, 413
- Herbrand normal form, 188
- Herbrand’s theorem, 166
- hereditarily
 - effective operations, 375, 377
 - extensional objects, 374, 473
 - recursive operations, 374, 377
 - undecidable theory, 333
- Heyting algebra, 75
- Heyting arithmetic, 237
 - second-order, 422
- Heyting-valued model, 123
- higher types, 351
- higher-order arithmetic, 438
- higher-order logic, 437
- Hilbert epsilon function, 474
- Hilbert–Bernays–Löb provability conditions, 337
- Hintikka set, 150
 - second-order, 418
- homomorphism, 3
- identity of indiscernibles, 411
- impredicative definition, 7
- incompleteness theorems, 322
 - first, 331
 - second, 338
- inconsistent set, 61
- indefinite description, 105, 180
- independence of premise, 122, 390
- index set, 288
- induction schema, 238, 247
- inductive type, 380, 481
- inductively defined set, 4
- infinitesimal, 131
- infinity axiom, *see* axiom of infinity
- initial segment
 - of a finite sequence, 10
 - of a function, 312
 - of a set, 311
- integer linear arithmetic, 344
- intension, 218, 369
- interpolant, 174
- interpolation theorem, 166, 174
 - Lyndon’s strengthening, 178
- introduction rule, 36, 480
- intuitionistic logic, x, 28
- invertible rule, 147
- irreducible term, 356
- isomorphism, 3
- join (in a partial order), 66
- judgmental equality, *see* definitional equality
- König’s lemma, 314
- Kleene
 - T* and *U*, 283
 - equality, 107, 280
 - normal form theorem, 283
- Kleene–Brouwer order, 15, 428
- Kolmogorov translation, 53
- König’s lemma, 13, 457
- Kripke model, 118
 - exploding, 74
 - for propositional logic, 71
 - normal, 121
 - pointed, 74
- Kuroda translation, 98
- lambda abstraction, x, 302
- lambda calculus, 301
 - second-order, 430

- language, 2
 - of arithmetic, 237
 - of first-order logic, 83
 - of propositional logic, 11, 28
 - of second-order logic, 410
- lattice, 65
 - law of the excluded middle, 43, 145
 - least fixed point
 - of a monotone operator, 461
 - of a monotone set operator, 8
 - least upper bound (in a lattice), 65
 - least-element principle, 223
- Leibniz equality, 411
- lexicographic order, 209
- linear arithmetic, 344
- literal, 47
- Löb's theorem, 339
- locally confluent rewrite system, 363
- logic of partial terms, 108
- logic, nature of, 83
- logical consequence, 60, 112
- logical equivalence, 43
- Łoś–Vaught test, 136
- low basis theorem, 315
- low degree, 315
- Löwenheim–Skolem theorems, 133
- MacNeille completion of a lattice, 125
- main premise, 151
- many-one degree, 309
- many-one reducible, 286
- many-sorted first-order logic, 106
- Markov's principle, 392
- maximally consistent set, 63, 115
- meet (in a partial order), 66
- metalanguage, 1
- metatheory, 1
- metavariables, 9
- middle sequent theorem, 169
- minimal logic, 28
- model, 2, 110
- modified realizability, 394
 - with truth, 397
- modulus of uniform continuity, 460
- monotone set function, 7, 461
- monotonicity
 - in a Kripke model, 71, 119
 - of the provability relation, 34
- mutual recursion, 204
- nand, 47
- natural deduction
 - for first-order logic, 86
 - for propositional logic, 40
 - for second-order logic, 411
- negation normal form, 48, 96
- negative formula, 51, 97, 174, 386
- negative subformula, 178
- neutral term, 358, 432
- Newman's lemma, 362
- no-counterexample interpretation, 402, 405
- nonstandard analysis, 131
- nonstandard model of arithmetic, 131
- nor, 47
- normal
 - form, 303, 356
 - proof, 140
 - term, 356
- normalizing
 - relation, 356
 - term, 356
- numeral, 304
- operational semantics, 297
- order filter, 125
- order ideal, 125
- order relation, 127
- pairing function, 200
- parallel reduction, 303
- parameters, 111, 132
- parsing, 11
- partial computable function, 280
- partial term, 107
- path through a tree, 13, 314
- Peano arithmetic, 237
 - second-order, 422
- permutative conversion, 381
- Pi type, 381, 481
- polymorphism, 430
- positive arithmetic operator, 462
- positive subformula, 178
- Post's theorem, 312
- prenex formula, 96
- Presburger arithmetic, 344
- preservation under increasing chains, 135
- primitive recursion, 193
- primitive recursive
 - arithmetic, 214
 - function, 192
 - functional, 376
- principal
 - formula, 151
 - order filter, 125
 - order ideal, 125
- product type, 482
- projection function, 192
- proof complexity, 55
- proof mining, 440
- proof theory, 323
- proper filter, 69
- propositional equality, 479
- propositional logic, 58
- propositions as types, 379, 480
- provability conditions, 337
- provably total computable function, 398
 - of PA or HA, 430

- pseudocomplement, 75
- pseudodvision, 347
- pseudohierarchy, 468
- Q (Robinson's theory), 323
- quantifier, x
- quantifier elimination, 341
- quantifier-free
 - fragment of a theory, 109
 - fragment of first-order logic, 95
 - instance of a formula, 166
 - logic, 95
- quotient structure, 4, 116
- real closed field, 348
- realizability, 383
 - with truth, 391
- recursion
 - course-of-values, 205
 - mutual, 204
 - simultaneous, 204
 - structural, 20
 - well founded, 208
 - with a measure, 206
 - with changeable parameters, 205
- recursion theorem, 293
 - for freely generated inductive structures, 17
- recursive comprehension axiom, 447
- recursor, 376
- redex, 353
- reducible term, 356
- reduct, 132
- reductio ad absurdum, 38
- reduction
 - β -, 303, 354
 - η -, 354
 - many-one, 286
 - truth-table, 317
 - Turing, 309
- reduction property, 246, 430
- reference, 369
- relative definability, 242
- relativized computation, 308
- relativized quantifier, 106
- renaming of variables, 142, 163
- replacement axiom, 477
- representability, 323
- representable
 - function, 327, 373
 - relation, 326
- representation, 199
- resolution rule, 56
- resolution theorem proving, 190
- reverse mathematics, 440, 450
- rewrite system, 356
- Rice's theorem, 288
- Rosser sentence, 336
- Russell's iota, 474
- s-m-n theorem, 283
- SAT solver, 56
- satisfaction
 - for propositional formulas, 60
 - in a model, 112
- saturated model, 136
- saturated set
 - of first-order sentences, 120
 - of propositional formulas, 73
- schema, 30
- second-order
 - arithmetic, 422
 - logic, 410
- self-realizing sentence, 388
- semantic consequence, 60, 112
- semidecidable, 291
- sense, 369
- sentence, 23
- separation axiom, 476
- separation property, 246, 430
- sequent calculus, 138
 - for second-order logic, 417
- set theory, 475
- Sheffer stroke, 47
- shmovable, 336
- shortening of cuts, 445
- side formula, 151
- Sigma type, 381, 483
- signature, *see* language
- simple term, 358
- simple type, 351
- simple type theory, 472
- simply typed lambda calculus, 351
 - second-order, 430
- simultaneous recursion, 204
- Skolem
 - axiom, 187
 - function, 187
 - normal form, 187
- Skolem's paradox, 134
- SMT solver, 109
- sort, 105
- soundness
 - of classical first-order logic, 112, 124
 - of classical propositional logic, 61, 67
 - of classical second-order logic, 413
 - of intuitionistic first-order logic, 119, 126
 - of intuitionistic propositional logic, 72, 75, 77
- standard element of a nonstandard model, 131
- standard interpretation of second-order logic, 413
- strictly positive subformula, 172
- strong collection schema, 258
- strong induction, 238
- strongly computable term, 358
- strongly normalizing
 - rewrite system, 356
 - term, 356
- structural induction, 20
- structure, *see* model

- subformula
 - generalized, 141
 - property, 142
 - strictly positive, 172
- subject reduction, 354
- substitution, 19
- substructure, 132
- subterm, 19
- subtraction, *see* truncated subtraction
- successor, 192
- sum type, 380, 482
- superlow basis theorem, 317
- symbol, 1
- T (Gödel's theory), 378, 399
- tableau, 150
- Tarski–Lindenbaum algebra, 68
- Tarski–Vaught test, 132
- tautology, 60
- term, 9
 - algebra, 18
 - model, 117
 - of first-order logic, 83
- termination proof, 295
- tertium non datur, 43
- theory, 89
 - axiomatized by a set of sentences, 89
 - of a model, 130
 - universally axiomatized, 89
- tree, 11
 - finitely branching, 13
- truncated subtraction, 195, 216
- truth assignment, 60
- truth value, 60
- truth-table reduction, 317
- Tseitin transformation, 56
- Turing degree, 309
- Turing machine, 295
- Turing reducible, 309
- type, 351
 - finite, 351
 - simple, 351
 - formation rule, 480
 - level, 352
 - preservation, 354
- type (in model theory), 135
- type 2 computability, 450
- type 2 functional, 310
- ultrafilter, 69
- unbounded search, 280
- uncurrying, 353
- undecidability, 284, 329, 349
 - of second-order logic, 415
- undefined term, 107, 280
- uniformization property, 247
 - of Π_1^1 relations, 429
 - of Σ_1^1 relations, 429
- unique factorization theorem, 203, 227
- unique readability, 11
- unit propagation, 56
- unit type, 381
- universal
 - axiom, 135
 - computable binary tree, 317
 - computably inseparable c.e. sets, 317
 - formula, 89, 95
 - partial computable function, 282
 - quantifier, x, 84
- universal closure of a formula, 86
- universal fragment of a theory, 109
- universally axiomatized theory, 89, 229
- universe of a model, 2, 110
- upward persistent formula, 134
- validity
 - in first-order logic, 112
 - in propositional logic, 60
- weak König's lemma, 457
- weak reduction, in combinatory logic, 365
- weakening, 34, 139
- well-founded
 - part of a tree, 462
 - relation, 129
 - tree, 12
- well-order
 - computable, 428
 - definable, 428
- well-orders, comparability of, 429
- word problem, 95
- Zorn's lemma, 63