

PALEOZOOLOGY AND PALEOENVIRONMENTS

Paleozoology and Paleoenvironments outlines the reconstruction of ancient climates, floras, and habitats on the basis of animal fossil remains recovered from archaeological and paleontological sites. In addition to outlining the ecological fundamentals and analytical assumptions attending such analyses, Tyler Faith and Lee Lyman describe and critically evaluate many of the varied analytical techniques that have been applied to paleozoological remains for the purpose of paleoenvironmental reconstruction. These techniques range from analyses based on the presence or abundance of species in a fossil assemblage to those based on taxon–free ecological characterizations. All techniques are illustrated using faunal data from archaeological or paleontological contexts.

Aimed at students and professionals, this volume will serve as a fundamental resource for courses in zooarchaeology, paleontology, and paleoecology.

J. Tyler Faith is Curator of Archaeology at the Natural History Museum of Utah and Assistant Professor of Anthropology at the University of Utah. His research addresses the relationships between Quaternary mammal communities, environmental change, and human–environment interactions, with an emphasis on eastern and southern Africa. This is his first book.

R. Lee Lyman is Emeritus Professor of Anthropology at the University of Missouri, Columbia. A scholar of late Quaternary paleozoology and human prehistory of the Pacific Northwest United States, he is author of *Vertebrate Taphonomy* (Cambridge, 1994), *Quantitative Paleozoology* (Cambridge, 2008), and *Theodore E. White and the Development of Zooarchaeology in North America* (2016).





PALEOZOOLOGY AND PALEOENVIRONMENTS

FUNDAMENTALS, ASSUMPTIONS, TECHNIQUES

J. TYLER FAITH

University of Utah, Salt Lake City and

R. LEE LYMAN

University of Missouri, Columbia





CAMBRIDGE UNIVERSITY PRESS

University Printing House, Cambridge CB2 8BS, United Kingdom

One Liberty Plaza, 20th Floor, New York, NY 10006, USA

477 Williamstown Road, Port Melbourne, VIC 3207, Australia

314-321, 3rd Floor, Plot 3, Splendor Forum, Jasola District Centre, New Delhi - 110025, India

79 Anson Road, #06-04/06, Singapore 079906

Cambridge University Press is part of the University of Cambridge.

It furthers the University's mission by disseminating knowledge in the pursuit of education, learning, and research at the highest international levels of excellence.

www.cambridge.org

Information on this title: www.cambridge.org/9781108480352

DOI: 10.1017/9781108648608

© Cambridge University Press 2019

This publication is in copyright. Subject to statutory exception and to the provisions of relevant collective licensing agreements, no reproduction of any part may take place without the written permission of Cambridge University Press.

First published 2019

Printed in the United Kingdom by TJ International Ltd, Padstow, Cornwall

A catalogue record for this publication is available from the British Library.

Library of Congress Cataloging-in-Publication Data

Names: Faith, J. Tyler, author. | Lyman, R. Lee, author.

Title: Paleozoology and paleoenvironments: fundamentals, assumptions,

techniques / J. Tyler Faith, R. Lee Lyman

Description: Cambridge ; New York : Cambridge University Press, 2019. |

Includes bibliographical references and index.

Identifiers: LCCN 2018039941 | ISBN 9781108480352 (hardback) |

ISBN 9781108727327 (pbk.)

Subjects: LCSH: Paleontology. | Paleoecology.

Classification: LCC QE761.F35 2019 | DDC 560-dc23

LC record available at https://lccn.loc.gov/2018039941

ISBN 978-1-108-48035-2 Hardback

ISBN 978-1-108-72732-7 Paperback

Cambridge University Press has no responsibility for the persistence or accuracy of URLs for external or third-party internet websites referred to in this publication and does not guarantee that any content on such websites is, or will remain, accurate or appropriate.



CONTENTS

| List of Figures | page ix |
|---|---------|
| List of Tables | xiii |
| Acknowledgments | XV |
| 1 WHY A BOOK ON PALEOENVIRONMENTAL. | |
| RECONSTRUCTION FROM FAUNAL REMAINS? | I |
| A Bit More History | 3 |
| Volume Structure | 5 |
| What We Do Not Do | 9 |
| Final Comments | 10 |
| 2 FUNDAMENTALS OF ECOLOGY AND BIOGEOGRAPHY | 12 |
| Historical Sketch | 14 |
| Biogeography | 14 |
| Ecology | 16 |
| Summary | 21 |
| Ecological Tolerances | 21 |
| Environments and Niches | 26 |
| Ecological Succession and Climax | 31 |
| What Is a Species to Do When the Environment Changes? | 32 |
| Top-Down or Bottom-Up Ecology | 33 |
| Resolution and Scale | 36 |
| Spatial and Temporal Scale | 37 |
| Kinds of Paleoenvironmental Change | 41 |
| Adaptation (Evolutionary Genetic) or Adjustment | |
| (Phenotypic Plasticity) | 45 |
| Conclusion | 46 |
| 3 ANALYTICAL ASSUMPTIONS | 48 |
| Assumption 1: Uniformitarianism | 50 |
| Assumption 2: Nearest Living Relative | 53 |
| Assumption 3: Ecological Tolerances | 55 |
| Assumption 4: Taxonomic Presence/Absence | 57 |

v



vi CONTENTS

| Assumption 5: Faunal Composition | 59 |
|---|------------|
| Assumption 6: Taxonomic Identification | 62 |
| Assumption 7: Sample Size and Taphonomy | 64 |
| Assumption 8: Small Bodies or Large Bodies | 66 |
| Assumption 9: Temporal Resolution | 68 |
| Assumption 10: Ecotone | 72 |
| Discussion | 75 |
| BACKGROUND OF SELECT PALEOZOOLOGICAL | |
| SAMPLES | 77 |
| Boomplaas Cave | 78 |
| Paleoenvironmental Summary | 82 |
| Homestead Cave | 83 |
| Paleoenvironmental Summary | 90 |
| Summary | 90 |
| • | |
| THE PRESENCE/ABSENCE OF TAXA | 0.2 |
| | 92 |
| A Bit of History | 93 |
| What about the Absence of a Taxon? | 94 |
| From Faunal List to Paleoenvironmental Reconstruction | 95 |
| Ordination | 99 |
| Correspondence Analysis | 99 |
| Detrended Correspondence Analysis | 103 |
| Other Ordination Techniques | 105 |
| One Species at a Time (Usually) Indicator Taxa | 107 |
| | 107 |
| Climatograph Multiple Taxa | 108 |
| Area of Sympatry | 113 114 |
| Mutual Climatic Range (Coexistence Approach) | 114 |
| The UTM-MCR Technique | 119 |
| Discussion | 121 |
| Summary | 121 |
| • | |
| 5 ENVIRONMENTAL RECONSTRUCTION BASED ON | |
| TAXONOMIC ABUNDANCES | 123 |
| Taxonomic Presences or Abundances? | 126 |
| Is the Analysis of Taxonomic Presences a Safer Alternative? | 129 |
| How to Gain (or Lose) Confidence | 134 |
| Abundances of a Few Taxa | 134 |
| Abundances of Indicator Taxa | 135 |
| Abundance Indices | 138 |
| Discussion | 142 |



| | | CONTENTS | vii |
|---|--|----------|-----|
| | History of Analyzing Taxonomic Abundances | 142 | |
| | Abundances of All Taxa in Assemblages | 145 | |
| | Ordination of Taxonomic Abundances | 145 | |
| | Other Techniques to Examine Abundances of All Taxa | 149 | |
| | Summary | 153 | |
| 7 | TAXON-FREE TECHNIQUES | 155 | |
| | Taxon Free: What It Is and What It Is Not | 156 | |
| | A Brief History | 158 | |
| | Autecological Approaches | 160 | |
| | Ecomorphology | 160 | |
| | Paleodietary Reconstruction | 164 | |
| | Synecological Approaches | 171 | |
| | Habitat Metrics | 173 | |
| | Ecometrics | 184 | |
| | Summary | 196 | |
| 8 | ENVIRONMENTAL INFERENCES BASED ON | | |
| | TAXONOMIC DIVERSITY | 197 | |
| | What Is Diversity? | 198 | |
| | How Is Diversity Quantified? | 199 | |
| | Taxonomic Richness | 201 | |
| | Taxonomic Heterogeneity | 213 | |
| | Taxonomic Evenness | 215 | |
| | Which Metric Should I Choose? | 217 | |
| | Practical Considerations with Fossil Data | 217 | |
| | Taxonomic Scope | 218 | |
| | Recovery and Analytical Methods | 219 | |
| | Agent of Accumulation | 220 | |
| | Differential Fragmentation | 221 | |
| | Differential Skeletal Element Representation | 222 | |
| | Time Averaging | 222 | |
| | Paleoenvironmental Implications of Taxonomic Diversity | 222 | |
| | The Varied Response of Diversity to Environmental | | |
| | Gradients | 223 | |
| | Paleoenvironmental Inferences | 228 | |
| | Case Study | 229 | |
| | Differential Identifiability | 230 | |
| | Differential Fragmentation | 230 | |
| | Skeletal Element Representation | 230 | |
| | Time Averaging | 231 | |
| | Agent of Bone Accumulation | 232 | |
| | Interpretation | 232 | |
| | Summary | 233 | |



viii CONTENTS

| 9 | TRANSFER FUNCTIONS AND QUANTITATIVE | |
|-------|---|-----|
| | PALEOENVIRONMENTAL RECONSTRUCTION | 234 |
| | How It Works | 235 |
| | The Analytical Toolkit | 238 |
| | Thackeray's Method | 238 |
| | Quantitative Bioclimatic Models | 242 |
| | Dental Ecometrics | 246 |
| | Tree Cover from Taxon-Free Characterizations | 251 |
| | Future Prospects | 256 |
| | The No-Analog Problem | 257 |
| | Further Cause for Concern | 260 |
| | Summary | 265 |
| 10 | SIZE CLINES AS PALEOENVIRONMENTAL INDICATORS | 266 |
| | Size Clines in Modern Organisms | 268 |
| | Bergmann's Rule: More than Just Temperature | 268 |
| | The Island Rule | 272 |
| | Summary | 273 |
| | Assumptions, Why Measure, and Method Basics | 273 |
| | Assumptions | 274 |
| | Why Measure? | 276 |
| | Method Basics | 277 |
| | Size Clines as a Paleoenvironmental Proxy | 278 |
| | One Way to Go about It | 279 |
| | Bergmann's Rule and Past Temperature Change | 288 |
| | Forage Availability and Predation in Large Herbivores | 291 |
| | Summary | 299 |
| 11 | SOME FINAL THOUGHTS | 301 |
| | Conservation Paleozoology | 303 |
| | Environments of Human Biological and Cultural Evolution | 305 |
| | What Next? | 307 |
| | Final Thoughts | 308 |
| Gl | ossary | 311 |
| Rę | ferences | 319 |
| Index | | 395 |



FIGURES

| 2.1 | A model of Merriam's life zones superimposed on the | |
|------|--|---------|
| | San Francisco Peaks, Arizona | page 18 |
| 2.2 | A model of world biome types in relation to mean annual | |
| | precipitation and temperature | 19 |
| 2.3 | Models of variability in the abruptness of community boundaries | |
| | along an environmental gradient | 20 |
| 2.4 | Ecological tolerance curve | 22 |
| 2.5 | A model of how tolerances of individual organisms of a species | |
| 2.6 | produce a fundamental niche or a species-level tolerance curve A model of a two-dimensional or two environmental gradient | 24 |
| | niche for a species population | 25 |
| 2.7 | A model of presences and abundances of nine species along an | |
| | environmental gradient of moisture | 26 |
| 2.8 | A model of species' niches, non-analog faunas, and species | |
| | assemblage change over time as a result of environmental change | 29 |
| | A model of the trophic pyramid | 34 |
| 2.10 | A model of covariation of spatial and temporal scales of biological | |
| | phenomena and processes | 39 |
| 2.11 | A model of the approximate spatial and temporal scales provided by typical fossil assemblages of terrestrial vertebrates from different | |
| | depositional settings | 40 |
| 2.12 | A model of the spatial and temporal scales of selected weather | |
| | and climatic episodes | 40 |
| 2.13 | Categories of environmental change along space and time | 42 |
| 2.14 | Environmental fluctuation over time | 43 |
| 3.I | One model of time averaging | 70 |
| 3.2 | Another model of time averaging | 70 |
| 3.3 | Models of the distribution of a species | 73 |
| 5.I | Correspondence analysis and plot of axis 1 scores of taxonomic | |
| | presences, Boomplaas Cave ungulates | 102 |
| 5.2 | Abundances and correspondence analysis of taxa along a | |
| 5.3 | hypothetical environmental gradient spanning nineteen sites Correspondence analysis and detrended correspondence analysis | 104 |
| | of Homestead Cave small mammal assemblages | 106 |
| | | |

ix



x FIGURES

| 5.4 | Modern distribution of Columbian ground squirrel in northwestern | |
|------|--|-------|
| | North America | III |
| 5.5 | Climatograph for Columbian ground squirrel | 112 |
| 5.6 | Bioclimatic range graphs for Lemmiscus curtatus and Mictomys borealis | 113 |
| 5.7 | Schematic illustration of how an area of sympatry and mutual | |
| | climatic range is determined for a fossil fauna | 116 |
| 5.8 | Areas of sympatry for three mammalian faunas of different ages from | |
| | the Cherokee Site | 118 |
| 6.I | Relative abundances of chisel-toothed kangaroo rats, | |
| | Homestead Cave | 124 |
| 6.2 | Relative abundances of small mammals from pellets of three owl | |
| | species, central Oregon | 127 |
| 6.3 | Species abundance distribution for small mammals from Stratum IV, | |
| | Homestead Cave | 130 |
| 6.4 | Percentage of taxa associated with mesic environments, Homestead | |
| | Cave Stratum IV | 131 |
| 6.5 | Absolute abundances of small mammal taxa, Wasden Site-Owl Cave | 137 |
| 6.6 | Relative abundances of small mammal taxa, Wasden Site-Owl Cave | 138 |
| 6.7 | Ratio abundances of pocket gophers and rabbits, Wasden | |
| | Site-Owl Cave | 139 |
| 6.8 | Relative abundances of some ungulate taxa, Boomplaas Cave | 146 |
| 6.9 | Correspondence analysis and plot of axis 1 scores of taxonomic | |
| | abundances, Boomplaas Cave ungulates | 147 |
| 6.10 | Chord distances for adjacent pairs of stratigraphic units, Homestead | |
| | Cave sequence | 153 |
| 7. I | Distribution of mammalian taxa across locomotor and dietary classes, | |
| | Congo Basin and Serengeti Plains | 156 |
| 7.2 | Example of linear dimensions measured to study the ecomorphology | |
| | of the bovid astragalus | 163 |
| 7.3 | Temporal variation in the proportion of bovid astragali assigned to | |
| | habitat categories in the Shungura Formation | 165 |
| 7.4 | Dental microwear surfaces of a typical grazer and a typical browser | 167 |
| 7.5 | Schematic of dental microwear surfaces | 168 |
| 7.6 | Mean dental microwear complexity vs. anisotropy for modern | |
| | African bovids and fossil taxa from Laetoli, Tanzania | 169 |
| 7.7 | Schematic of mesowear variables | 170 |
| 7.8 | Univariate mesowear scores for maxillary third molars of modern | |
| | Australian marsupials compared with fossil Macropus giganteus titan | 171 |
| 7.9 | Distribution of species across body size classes, locomotor regimes, | |
| | and dietary classes across modern habitats relative to the Upper | |
| 7.10 | Ndolanya Beds (Laetoli) P. elationship between woody cover and the first axis of a CA | 175 |
| /.10 | Relationship between woody cover and the first axis of a CA of taxonomic presences in modern Serengeti raptor roosts | T 0 0 |
| | or taxonomic diesences in modern Sciengen fablor foosis | 180 |



| 7.11 | Habitat spectra for Serengeti raptor roosts across a gradient from | |
|------|--|-----|
| | open to wooded habitats | 182 |
| 7.12 | Mammal communities associated with Kalahari thornveld | |
| | vegetation and moist tropical forest | 187 |
| 7.13 | Interpretive algorithm for cenograms | 188 |
| 7.14 | Cenograms for Boomplaas Cave and the relationship between | |
| | richness and slope for large and small mammals | 191 |
| 7.15 | Average hypsodonty index of mandibular third molars for 133 | |
| | extant ungulates | 194 |
| 7.16 | Changes in mean hypsodonty index and relative abundance of | |
| | ungulate grazers, Boomplaas Cave | 195 |
| 8.1 | Four hypothetical faunal assemblages of varied richness and evenness | 200 |
| 8.2 | Relationship between sample size and metrics of taxonomic | |
| | richness. Homestead Cave Stratum IV | 203 |
| 8.3 | Relationship between sample size and NTAXA for the Boomplaas | Č |
| | Cave ungulates | 204 |
| 8.4 | Relationship between sample size and NTAXA assumed by | |
| | Margalef's richness index and Menhinick's richness index | 206 |
| 8.5 | Sampling-to-redundancy curves showing cumulative taxonomic | |
| | richness and heterogeneity across six annual field seasons | 208 |
| 8.6 | Rarefaction curves for Boomplaas Cave ungulates | 210 |
| | Rarefaction curves for mammal remains, Marmes site | 211 |
| 8.8 | Two different relationships between sample size and NTAXA, | |
| | Homestead Cave and Samwell Cave | 212 |
| 8.9 | Relationship between sample size and metrics of taxonomic | |
| | heterogeneity and evenness, Homestead Cave Stratum IV | 214 |
| 8.10 | Comparison of Shannon index for Boomplaas Cave ungulates | |
| | to values observed when assemblages are standardized to an | |
| | equivalent sample size | 215 |
| 8.11 | Relationship between mean annual precipitation and species | - |
| | richness for sub-Saharan African ungulates | 224 |
| 8.12 | Relationship between mean annual precipitation and species | · |
| | richness for Australian mammals | 225 |
| 8.13 | Assemblage evenness and the abundance of kangaroo rats, | - |
| | Homestead Cave | 226 |
| 8.14 | Assemblage evenness and the abundance of reindeer, Grotte XVI | 226 |
| | Changes in evenness and richness in micromammal assemblage, | |
| | Boomplaas Cave | 227 |
| 8.16 | Changes in evenness and richness in ungulate assemblage, | , |
| | Boomplaas Cave | 228 |
| 8.17 | Relationship between richness and abundance of isolated teeth, | |
| , | abundance of skull elements, time averaging, and anthropogenic | |
| | bone modifications in the ungulate assemblage from Boomplaas Cave | 231 |

xi

FIGURES



xii FIGURES

| 9.1 | Ranking of temperature reconstructions from coolest to warmest | |
|------|---|-----|
| | for Boomplaas Cave micromammals | 241 |
| 9.2 | Ranking of temperature and precipitation reconstructions for | |
| | Boomplaas Cave rodents | 246 |
| 9.3 | Ranking of temperature and precipitation reconstructions for | |
| | Boomplaas Cave large herbivores | 250 |
| 9.4 | Schematic representation of tree cover categories | 252 |
| 9.5 | Relationship between faunal community composition | |
| | and %heavy tree cover | 255 |
| 9.6 | Faunal PCI, %heavy tree cover, and ranking of %heavy tree cover | |
| | reconstructions for Boomplaas Cave large mammals | 256 |
| 9.7 | Range of temperatures in which genus Fraxinus (ash trees) are | |
| | found in North America today | 259 |
| 9.8 | Antarctic ice core records of changes in atmospheric CO ₂ , | |
| | past 800,000 years | 262 |
| 9.9 | Predicted superiority of C ₃ and C ₄ photosynthetic pathways as | |
| | a function of daytime growing-season temperature and atmospheric | |
| | CO ₂ concentrations | 262 |
| 9.10 | Reconstructed mean annual precipitation for the Great Plains | |
| | of North America, past ~25 Ma | 264 |
| IO.I | Geometric bases of Bergmann's rule, and of Allen's rule | 267 |
| 10.2 | Changes in adult body mass of moose individuals across a | |
| | latitudinal gradient, Sweden | 269 |
| 10.3 | Frequency distributions of mandibular alveolar lengths in | |
| | bushy-tailed woodrats | 286 |
| 10.4 | Mean mandibular alveolar length of bushy-tailed woodrats, | |
| | Gatecliff Shelter, Nevada | 287 |
| 10.5 | Mandibular alveolar lengths and relationship with mean annual | |
| | temperature, modern East African mole-rats and Holocene | |
| | assemblages | 289 |
| 10.6 | Changes in astragalus size in white-tailed deer relative to changes in | |
| | summer insolation, past 10,000 years | 292 |
| 10.7 | Heuristic model showing influence of net primary productivity and | |
| | population density on white-tailed deer growth rates and body size | 294 |
| | Changes in weight of deer harvested 1971–2005, Fort Hood, Texas | 295 |
| 10.9 | Changes in Great Plains bison calcaneal length, past 37,000 years | 297 |



TABLES

| 4.I | Stratigraphy and chronology of Boomplaas Cave | page 80 |
|------------|--|---------|
| 4.2 | Taxonomic abundances (MNI) for Boomplaas Cave microfauna | 81 |
| 4.3 | Taxonomic abundances (NISP) for Boomplaas Cave macromammals | 84 |
| 4.4 | Stratigraphy and chronology of Homestead Cave | 87 |
| 4.5 | Taxonomic abundances (NISP) for Homestead Cave small mammals | 88 |
| 5. I | The presence of ungulate species through time at Boomplaas Cave | 97 |
| 5.2 | Modern climate data for Columbian ground squirrel | IIO |
| 5.3 | Small mammal faunas from the Cherokee archaeological site | 117 |
| 6.1 | Taxonomic abundances (NISP) of small mammals from Homestead | |
| 6.2 | Cave Stratum IV and randomly sub-sampled assemblages Taxonomic abundances (NISP) and adjusted residuals for | 132 |
| | Homestead Cave small mammals | 151 |
| 7. I | Preferred habitats of Cherokee archaeological site stenotopic | |
| | micromammals | 172 |
| 7.2 | Niche models for Serengeti rodents | 181 |
| 7.3 | Climatic zones and corresponding vegetation biomes used in | |
| | bioclimatic model | 183 |
| 7·4 8.1 | Example of the bioclimatic model technique The number of non-overlapping taxa for five hypothetical | 185 |
| | assemblages of Odocoileus spp. | 202 |
| 8.2 | Sample size and various indices of richness for the Boomplaas | |
| | Cave ungulates | 205 |
| 8.3 | Taxonomic list and diversity indices for Henderson Island | |
| | fish remains | 220 |
| 8.4 | Diversity indices for small mammal assemblages accumulated by | |
| | three owl species in Oregon | 221 |
| 9.1 | Factor I loadings and mean annual temperature of South African | |
| | rodent taxa | 239 |
| 9.2 | Reconstructions of past climate for Boomplaas Cave micromammals | 240 |
| 9.3 | Comparison of observed mean annual temperatures with values | |
| | reconstructed using transfer functions for rodents | 243 |
| 9.4 | Hypsodonty score and lophedness for Boomplaas Cave primates | |
| | and ungulates | 2.48 |

xiii



xiv TABLES

| 9.5 | Mean hypsodonty score and lophedness of large mammalian | |
|------|--|-----|
| | herbivores and reconstructions of past climate, Boomplaas Cave | 249 |
| 9.6 | Ecological groups for mammals | 252 |
| 9.7 | Percentage of large mammal species (>1 kg) across ecological | |
| | categories in modern communities | 254 |
| IO.I | A selection of mammalian taxa for which size clines across | |
| | spatial environmental gradients have been used to infer | |
| | paleoenvironmental change | 280 |



ACKNOWLEDGMENTS

This volume is the outcome of years of reflection and discussion with our mentors, collaborators, colleagues, and friends, and we sincerely thank the many people who contributed directly or indirectly to the ideas presented here. We are particularly indebted to those who responded to our questions, comments, and requests while writing this volume: Andrew Barr provided a brief but important discussion concerning measurements in ecomorphology; Larisa DeSantis sent us dental microwear photosimulations; Andrew Du supplemented our limited coding skills and offered insightful comments on taxonomic diversity; Christine Janis and John Damuth sent us raw data concerning precipitation estimates for the Great Plains; Julien Louys clarified points concerning a methodological technique; Dan Peppe directed us to some useful paleobotanical literature; Steve Wolverton provided raw data concerning the size of deer from Texas. René Bobe and an anonymous reader inspired useful fine-tuning in later drafts of this volume. A portion of Faith's contribution to this volume was supported by an Australian Research Council DECRA fellowship at the University of Queensland.

