

Index

- Abbreviated Injury Scale (AIS), 33
- antiepileptic drug (AED) therapy
- behavioral effects from, 176
 - brivaracetam, 170
 - carbamazepine, 169–70
 - concussive
 - convulsions, 93–5
 - eslicarbazepine, 170
 - gabapentin, 170–1
 - lacosamide, 171
 - lamotrigine, 171
 - levetiracetam, 171–2
 - methylphenidate, 175
 - oxcarbazepine, 172
 - perampanel, 172
 - phenobarbital, 169–70
 - phenytoin, 169–70
 - post-traumatic epilepsy (PTE) in children, 86–7
 - risk factors and side effects, 175
 - rufinamide, 172–3
 - tiagabine, 173
 - topiramate, 173
 - valproate, 169–70
 - vigabatrin, 173–4
 - zonisamide, 174
- apathy, 131–2
- blast-induced traumatic brain injury model (bTBI)
- mechanism of, 13–14
 - realistic test for military-type explosions, 14
- brain pathologies from traumatic brain injury (TBI)
- age and recovery, 17
 - age related, 18
 - inflammatory signaling pathways, 17
 - multifaceted nature of, 15
 - neurogenesis from, 17
 - and role in post-traumatic epilepsy (PTE), 18
 - spontaneous seizures from, 16
 - superoxide dismutase (SOD) changes, 17–18
- brivaracetam, 170
- carbamazepine, 169–70
- cerebral salt wasting (CSW), 68–9
- cognitive behavioral therapy (CBT), 115–16, 152
- cognitive consequences of epilepsy
- age and, 108–9
 - antiepileptic drug (AED) effects, 110
 - brain structures affected, 108, 109
 - cognitive impairments and, 109
 - depression, 109–10
 - surgical effects, 110
- cognitive consequences of post-traumatic epilepsy (PTE)
- cognitive behavioral therapy (CBT), 115–16
 - as double hit after head injury, 104, 110–11
 - fatigue, 116–17
 - stigma of epilepsy, 116
- cognitive consequences of traumatic brain injury (TBI)
- brain structures affected, 105
 - case study, 106–8
 - diffuse axonal injury (DAI), 105
 - emotional recognition, 105
 - social relationship breakdown, 105–6
- cognitive rehabilitation strategies for post-traumatic epilepsy (PTE)
- attention and processing speed, 113–14
 - executive function, 114
 - fatigue, 116–17
 - memory, 114
 - social cognition, 115
- Collaborative European neuroTrauma Effectiveness Research in Traumatic Brain Injury study (CENTER-TBI), 47
- computerized tomography (CT), 85
- concussive convulsions
- defined, 90, 91
 - diagnosis, 93–4
 - epidemiology, 92
 - history of, 90
 - management, 93–5
 - semiology, 91
- continuous electroencephalography (cEEG), 85
- controlled cortical impact model (CCI)
- epileptic rate from, 13
 - memory and cognitive deficits from, 13
 - procedure, 13
- copper, 14
- deep brain stimulation (DBS), 174–5
- default mode network (DMN), 142
- depression
- after traumatic brain injury (TBI), 128–30
 - cognitive consequence of epilepsy, 109–10
- diffuse axonal injury (DAI), 105
- diffusion tensor imaging (DTI), 141–2
- early post-traumatic seizures
- defined, 157
 - epidemiology, 158
 - studies, 161–5
- electroencephalogram (EEG)
- concussive convulsions, 93, 95
 - continuous, 85
 - enacted stigma (epilepsy), 117
- epilepsy

- cognitive consequences of, 108–10
 defined, 30–1
 diagnosis, 30, 31
 etiology, 32
 neurostimulation for, 174–5
 rehabilitation, 117–18
 surgery, 174
- epilepsy accidents and injuries
 motor vehicle
 accidents, 98–9
 prevalence of, 97–8
 seizure related, 99–100
- epileptic seizure
 classification, 31, 32
 defined, 29, 30
- epileptogenesis after traumatic brain injury (TBI)
 clinical challenges, 159–61
 clinical practice
 implications, 165–6
 factors influencing, 158–9
 prevention studies, 161–5
- epileptogenesis and fluid percussion injury (FPI), 2–12
- eslicarbazepine, 170
- Feeney's weight-drop
 model, 14
- felt stigma (epilepsy), 117
- fluid percussion injury (FPI)
 model
 age-related results
 differences in, 2
 behavioral
 comorbidities, 2–11
 incidence of epilepsy
 after, 12
 limitations of, 12–13
 procedure, 2
 role in epileptogenesis, 2–12
 spike-wave discharges, 12
- gabapentin, 170–1
- Glasgow Coma Scale (GCS), 33
- hyponatraemia, 68–9
- Integrative Cognitive Model (ICM), 142
- International Classification of Disease (ICD), 33
- International Initiative for Traumatic Brain Injury Research (InTBIR), 47
- intracranial pressure (ICP)
 control, 70–1
- lacosamide, 171
- lamotrigine, 171
- late post-traumatic seizures
 defined, 157
 epidemiology, 158
 studies, 165
- levetiracetam, 171–2
- low income countries,
 PTE in
 epidemiology, 179, 181
 history of, 179
 management, 182–3
 mortality and morbidity
 prevalence, 179
 pathophysiology, 181–2
- low income countries, TBI in
 epidemiology, 180
 falls and, 180–1
 road traffic accidents, 180
 substance abuse, 180
 violence and, 180
- magnetic resonance imaging (MRI), 85
- Marmatou's impact
 acceleration model, 15
- Mayo classification system, 49
- methylphenidate, 175
- microdialysis, 73
- motor vehicle accidents
 epilepsy accidents and
 injuries, 98–9
 and traumatic brain injury (TBI) in low income
 countries, 98–9
- near infrared spectroscopy (NIRS), 73
- neuroimaging and convulsive
 convulsions, 93, 95
- neuroprotection for traumatic
 brain injury
 intracranial pressure (ICP)
 control, 70–1
 osmotherapy, 71
 pharmacological, 72–3
 seizure control, 72
 surgical intervention, 72
 temperature control, 71
- Operation Brain Trauma
 Therapy (OBTB), 18–19
- osmotherapy, 71
- oxcarbazepine, 172
- pediatric weight-drop
 model, 15
- penetrating brain injury model
 brain changes from, 14
 limitations of, 14
 procedure, 14
- perampanel, 172
- phenobarbital, 169–70
- phenytoin, 169–70
- post-traumatic amnesia (PTA), 132–4
- post-traumatic epilepsy (PTE)
 after traumatic brain injury
 (TBI), 1, 18
 cognitive consequences of,
 110–11
 diagnosis, 157–8
 epidemiology, 29–30, 34
 frequency of, 157
 neuropsychiatric problems,
 135–6
 post-traumatic seizures
 and, 32–3
 rehabilitation for, 111–18
 spontaneous seizures in
 preclinical trials, 16
 traumatic brain injury
 evidence for, 34–44
- post-traumatic epilepsy (PTE)
 and post-traumatic stress
 disorder (PTSD)
 background of, 149
 clinical implications, 151–2
 epidemiology, 150–1
 management, 152–4
 neurobiology, 149–50
 pharmacological
 management, 153
- post-traumatic epilepsy (PTE)
 animal models, 2–15
- post-traumatic epilepsy (PTE)
 in children
 assessment and
 diagnosis, 85–6
 clinical presentation, 85
 epidemiology, 78
 management, 86–7
 mortality and morbidity
 prevalence, 78
 pathophysiology, 78–84
 risk factors for, 79

- post-traumatic epilepsy (PTE)
 in low income countries
 epidemiology, 179, 181
 history of, 179
 management, 182–3
 mortality and morbidity
 prevalence, 179
 pathophysiology, 181–2
- post-traumatic stress disorder (PTSD)
 after traumatic brain injury (TBI), 131
 and post-traumatic epilepsy (PTE), 149–54
 and psychogenic non-epileptic seizures (PNES), 144–5
- psychogenic nonepileptic seizures (PNES)
 defined, 140
 etiology, 140–1
 as neural network disorder, 142–3
 traumatic brain injury (TBI)
 perpetuating factor, 145–6
 traumatic brain injury (TBI)
 psychological effects and, 143–5
 traumatic brain injury (TBI)
 structural effects and, 141–3
- psychosis, 132
- rehabilitation for epilepsy
 driving inability, 117–18
 enacted stigma, 117
 felt stigma, 117
- rehabilitation for post-traumatic epilepsy (PTE)
 cognitive strategies, 113–15
 emotional comorbidities, 115–18
 future research in, 118
 as interactive process, 111
 interdisciplinary approach, 111–12
 SMART goals, 112–13
- responsive neurostimulation (RNS), 174–5
- rodent developmental stages, 1–2
- rodent models for post-traumatic epilepsy (PTE)
 blast-induced traumatic brain injury model (bTBI), 13–14
- controlled cortical impact model (CCI), 13
 findings summary, 3
 fluid percussion injury (FPI)
 model, 2–13
 penetrating brain injury
 model, 14
 weight-drop models, 14–15
- rufinamide, 172–3
- Shohami's weight-drop model, 15
- SMART goals, 112–13
- social cognition therapy, 115
- sports-related
 concussion (SRC)
 occurrence of, 90
 pathophysiology, 91–2
 post-traumatic epilepsy (PTE), 91
- substance abuse
 and TBI in low income countries, 180
 from traumatic brain injury, 135
- superoxide dismutase (SOD), 17–18
- tiagabine, 173
- Time Pressure Management (TPG), 113
- topiramate, 173
- Transforming Research and Clinical Knowledge in Traumatic Brain Injury (TRACK-TBI), 47
- traumatic brain injury (TBI)
 age and, 56
 behavioral
 comorbidities, 2–11
 brain pathologies, 15–17
 cognitive consequences of, 105–8
 epidemiology, 48
 epileptogenesis after, 158–61
 evidence for post-traumatic epilepsy (PTE), 34–44
 Operation Brain Trauma Therapy (OBT), 18–19
 pathology, 63–4
 and post-traumatic epilepsy (PTE), 1
 prevention, 54–6
 and psychogenic non-epileptic seizures (PNES), 141–6
- rehabilitation, 116
 risk factors for, 54
 severity classification, 33–4, 48–9
- traumatic brain injury (TBI) in low income countries
 epidemiology, 180
 falls and, 180–1
 motor vehicle accidents, 180
 substance abuse, 180
 violence and, 180
- traumatic brain injury (TBI)
 management
 acute, 50–4
 clinical assessment, 66–7
 hematological, 68
 immediate prehospital and emergency room, 64–5
 metabolic and endocrine, 68–9
 neuromonitoring, 73–6
 neuroprotection, 70–3
 programs, 47
 sedation, 69–70
 systemic, 66–8
- traumatic brain injury (TBI)
 neuropsychiatric problems
 anxiety disorders, 130–1
 apathy, 131–2
 long-term personality changes, 134–5
 mood disorders, 128–30
 post-traumatic amnesia (PTA), 132–4
 psychosis, 132
 substance abuse, 135
 summary of, 128–9
- uncinate fasciculus (UF), 141–2
- unprovoked seizure. *See* late post-traumatic seizures
- vagal nerve stimulation (VNS), 174–5
- valproate, 169–70
- vigabatrin, 173–4
- weight-drop models
 Feeney's, 14
 Marmatou's impact acceleration model, 15
 pediatric, 15
 Shohami's, 15
- zonisamide, 174