
Index

- Castelnuovo–Severi inequality, 29, 35
- Character
 - cyclotomic, 75, 111
 - Dirichlet, 73, 74, 76, 79, 84, 92
 - Hecke, 90, 92, 93, 102, 107, 133
- Chow group, 58, 63–65
- Cohomology
 - l -adic, 67, 70, 122, 125, 134, 143, 153
 - Betti, 67, 70, 142, 143
 - crystalline, 67, 157
 - de Rham, 59, 67, 142
 - Weil, 53, 57, 60, 66, 68, 70, 129, 134, 143, 154
 - classical, 67, 153, 155
- Conjecture
 - Artin, 100, 143, 166
 - Tate, 45, 164
 - weight-monodromy, 135
- Conjectures
 - standard, 70, 151, 154, 155, 160
 - Weil, 21, 44, 54, 56, 57, 68, 143
- Functional equation, 7, 13, 18, 20, 25, 27, 53, 56, 68, 69, 93, 100, 102, 104, 107, 120, 125, 133, 140, 159, 161–163, 180
- Good reduction, 108, 135, 152
- l -adic sheaf, 107, 110–112, 114, 115, 119, 166
- L -function
 - Artin, 96
 - Dirichlet, 74
 - Hecke, 92, 99, 105
 - of an l -adic sheaf, 114, 119
- Numbers
 - Betti, 55, 56, 68, 70
 - Weil, 55, 129, 135
- Riemann hypothesis, 7, 15
 - for abelian varieties, 52
 - for curves, 20, 21, 25, 29, 101, 163
 - for smooth projective varieties, 56, 70, 71, 105, 131, 135, 156, 162
- Riemann–Hurwitz–Hasse formula, 95
- Theorem
 - l -adic monodromy, 135, 139
 - Dirichlet’s, 2, 76, 80, 81, 83
 - Riemann–Roch, 1, 20, 26, 28, 35, 37, 94
- Zeta function
 - Dedekind, 15, 18, 22, 75, 97
 - Hasse–Weil, 104, 106, 109, 133
 - of a (pure) motive, 109, 143, 163
 - of a scheme of finite type over $\mathbf{Z}$, 20, 52, 106
 - of an endomorphism, 159, 179
 - Riemann, 6, 19, 93