

## Index

- 'A Matter of Balance' program, 311
- accelerometers, 212
  - fall risk assessment and, 218
  - gait stability measurements and, 54
  - turning measurements and, 65
- acetylcholinesterase inhibitors, 476
- Action Cycle (of the KTA Framework), 438, 440–53
- acute illness, 185, 242
- alcohol consumption, risk of falls and, 238
- Alternate Step Test, 32, 228
- Alzheimer's disease, 144
  - cerebral amyloid/tau and, 136
  - peripheral sensation and, 92
- analgesics, as risk factors for falls, 193, 197
- ankle strategy, 36
- ankle-foot orthoses, 350
- anti-anxiety agents, 243
- anti-arrhythmic medications, 198, 244
- anticoagulants, 198
- antidepressants, 195, 243
- antiepileptics, 244
- antihypertensives, 197, 243
- antipsychotics, 196, 243
- anxiety
  - cognitive-specific risk factors for falls and, 148
  - as a consequence of falls, 163
  - exercise and, 165
  - fear of falling and, 161
  - risk factors for falls and, 163, 242
- anxiolytics, 243
- arrhythmia, 331
  - anti-arrhythmic medications and, 198
  - cerebral hypoperfusion and, 181
- arthritis, 242
- assistive devices and self-care aids, 360
- AST (Alternate Step Test), 228
- backward falls, 114
- balance
  - assessment of, 33, 325
  - exercises for, 290, 297
  - as impacted by footwear, 122
  - risk of falls and, 239
  - white matter hyperintensities and, 132
- balance control
  - anxiety and, 164
  - balance confidence and, 160–3, 302
  - brain changes and, 131
  - cognitive impairment risk factors and, 145
  - cognitive-motor training and, 304
  - exercise for preventing falls and, 252, 260
  - exercises for, 470
  - orthoses and, 350
  - standing and, 105–18
  - step training and, 272
  - vestibular disorders and, 177, 243
- balance range, 28
- base of support, 23, 28, 53
  - defined, 105
  - exercise and, 252
  - foot problems and, 120
  - shoes and, 123
  - step training and, 271
- behavioural and psychological symptoms (BPS), 147
- behavioural risk factors, 205, 470
- benzodiazepines, 194, 475
- beta-amyloid, 136
- beta blockers, 197, 244
- biomechanics, 470
  - balance-related falls and, 105–18
  - obstacle negotiation and, 61
  - slipping and, 71, 73
  - stair negotiation and, 66
- BPS (behavioural and psychological symptoms), 147
- brain function, 130–43
  - brain atrophy and, 130
  - functional near-infrared spectroscopy and, 138
  - magnetic resonance imaging and, 137
- brain hypoperfusion, 181
- brain injury, 411
- brain networks, 137
- camera detection systems, 212
- camera risk assessment systems, 216
- cardiac syncope, 331

**482 Index**

- cardiovascular medications, risk of falls and, 194, 197, 244
- cardiovascular problems, 181
- Care Home Falls Screen, 234
- carotid sinus hypersensitivity, 243
- carotid sinus syndrome, 330
- cataracts
  - cataract surgery and, 341
  - as a risk factor for falls, 173
- centre of mass (centre of gravity), 105
  - footwear and, 122
  - predictive adaptation and, 275
- cerebellar disorders, 179
- cerebral amyloid, 136
- cervical spine, 176, 183
- cholinergic denervation, 136
- clinical research/implementation
  - gait monitoring for fall risk assessment and, 219
  - new technology devices and, 220, 222
- Cochrane review of exercise and falls, 148–51, 251
- cognition, 130, 470
  - Alzheimer's disease and, 136
  - dancing and, 297
  - gait and, 58–64
  - infarcts and, 136
  - interactive cognitive-motor training and, 301
  - microbleeds and, 136
  - white matter hyperintensities and, 132, 136
- cognitive behavioural interventions, for fear of falling/fall risk, 165, 311–21
  - 'A Matter of Balance' program and, 311
  - outcomes and, 314–18
- cognitive impairment, 11, 131, 144–59
  - older people in residential care facilities and, 419
  - preventing falls and, 148–51
  - risk factors for falls and, 144–8, 242
- cognitive-motor training
  - dual-tasking/multi-tasking and, 287, 288, 295
  - efficacy of, 289–90
  - future research perspectives and, 476
  - mode of action of, 303
  - vs. traditional exercise, 302
- cognitive processing speed, impairment of, 130, 147
- cognitive stepping, 298
- compliant flooring, 381, 478
- consequences of falls, 14–16
- coordinated stability, 29
- costs of falls/fall-related injuries, 16, 460–8, 474
  - future perspectives for decision-making and, 466
  - methodological limitations and, 465
  - trial-based vs. model-based economic evaluations and, 462–5
- crutches, 351
- daily-life gait, fall risk assessment and, 55, 218
- daily-life movements, fall risk assessment and, 220
- dancing, 290, 296
- delirium, 150, 402
- dementia, 180
  - defined, 144
  - Donepezil anti-dementia drug and, 149
- exercise for preventing falls and, 261
  - non-cognitive manifestations of, 147
  - older people in residential care facilities and, 419
  - risk groups for falls and, 322
  - types of, 144
- depression, 181
  - antidepressants and, 195
  - cognitive-specific risk factors for falls and, 148
  - as a consequence of falls, 163
  - exercise and, 165
  - fear of falling and, 161, 164
  - risk factors for falls and, 163, 242
- dermatological conditions, foot problems and, 184
- detecting falls, 4, 211–16
  - new technology's real-world utility and, 215
  - stages of falls detected, 211
- diabetes
  - foot problems and, 184
  - peripheral sensation and, 176
- diabetic peripheral neuropathy, wearable devices and, 353
- digoxin, 198, 244
- diuretics, 198
- dizziness, risk factors for falls and, 243
- "drop attacks", 328
- dual-tasking/multi-tasking, 59
  - cognitive-motor training and, 287, 288, 295, 302, 304
  - dual-tasking during gait, cognitive impairment risk factors and, 146
- endurance, 95, 96, 240
- the environment, three levels of, 202
- environmental devices
  - for fall detection, 212
  - for fall risk assessment, 216
- environmental factors, 470
- environmental interventions to prevent falls, 360–77
  - environmental risk screens/assessments and, 367
  - reducing hazards and, 360
  - sample solutions for, 369, 370
- environmental risk factors for falls, 202–10
- epidemiology of falls/fall-related injuries, 3–18
  - incidence of falls and, 5–12
  - location of falls and, 12
  - older people in residential care facilities and, 411
- executive function
  - balance control and, 146
  - cognitive-motor training and, 304
  - gait and, 62
  - impairment of, 130, 147
  - as a risk factor for falls, 242
- exercise, as physical activity for healthy older people, 265
- exercise, for preventing falls, 148, 238, 251–70, 471
  - vs. cognitive-motor training, 302
  - combined with 'A Matter of Balance' program, 313
  - cost considerations and, 461, 465
  - multifactorial benefits of, 165

**483 Index**

- older people's perception of, 428
- residential care facilities and, 417
- step training and, 272
- strong evidence supporting, 251
- types of exercise and, 251, 260
- exergames, 290, 299
  - cognitive-motor training and, 289
  - future research perspectives and, 476
  - volitional step training and, 274
- external perturbations, 35–9
- eyeglasses, 360
  - bifocal/multifocal and, 68, 245, 343
  - first time/new prescription and, 342–5
  - risk of falls and, 245, 328
  - unifocal, 328
- fall efficacy, 160–3
- fall prevention, 148–51, 410–24, 471–4
  - adherence strategies for, 427–35, 473
  - behaviour change strategies/techniques for, 429, 477
  - costs of falls/fall-related injuries and, 460–8, 474
  - fall prevention interventions for the visually impaired, 341–7
  - fall risk assessment and, 216–22
  - footwear/devices and, 348–59
  - from research to practice and, 436–59
  - safety education interventions and, 149, 151
  - social support and, 431
- fall-related injuries
  - defined, 5
  - epidemiology of, 3–18
  - fractures and. *see* fractures
- falls
  - consequences of, 14–16
  - defined, 3–5
  - detection/risk assessment for, 211–26
  - epidemiology of, 3–18
  - in hospitals, 11, 396
  - risk of falling again and, 11, 410
- FDG PET, 137
- fear of falling, 15, 160–3, 164, 241
  - cognitive behavioural interventions for, 165, 311–21
  - gait changes and, 55
  - interactive cognitive-motor training and, 302
- femur, hip protectors and, 378
- flooring, compliant for preventing falls, 381, 478
- fluorodeoxyglucose (FDG) PET, 137
- fNIRS (functional near-infrared spectroscopy), 138
- foot drop, wearable devices and, 353
- foot orthoses, 350
- foot problems, 119–29, 184
  - podiatry and, 125
  - prevalent types of, 119
  - risk of falls and, 242
- footwear, 121–4, 184, 470
  - podiatry and, 125
  - potential detrimental effects on balance/gait and, 122
  - risk of falls and, 124, 245, 348
  - socks and, 400
- forward falls, 113
- fractures, 417
  - impact of exercise on, 263
  - multiple interventions and, 393
  - older people in residential care facilities and, 411
- frailty, risk of falls and, 11
- frames, as walking aids, 351
- functional impairment, risk factors for falls and, 147
- functional magnetic resonance imaging, 137
- functional reach, 28
- gait, 51–86, 470
  - Alzheimer's disease and, 136
  - assessment of, 325
  - brain changes and, 131
  - cognition and, 58–64
  - cognitive impairment risk factors and, 146
  - dementia and, 180
  - fall risk assessment and, 216–22
  - as impacted by footwear, 122
  - level walking and, 51–8
  - perturbations to, 110
  - risk of falls and, 239
  - slipping and, 70–3
  - stair negotiation and, 65–8
  - stroke and, 178
  - tripping and, 68–70
  - turning and, 64
  - variability of, 52
  - vestibular function and, 177
  - walking aids and, 351
  - white matter hyperintensities and, 132
- gait adaptability, 62
- gait changes, 55
- gait speed
  - interactive cognitive-motor training and, 302
  - level walking and, 52
  - medications and, 192
  - non-modifiable factors and, 58
  - predicting falls and, 53
- glaucoma, as a risk factor for falls, 175
- grey matter, 130
- HAROLD (Hemispheric Asymmetry Reduction in Older Adults) model, 137
- health coaching, 432
- hearing impairment, risk of falls and, 89, 241
- heel elevation, footwear choices and, 122
- Hemispheric Asymmetry Reduction in Older Adults (HAROLD) model, 137
- high-risk groups
  - exercise for preventing falls and, 261, 264
  - future research into falls and, 476
  - older people in residential care facilities and, 410–24
- the hindbrain, 181
- hip protectors, 378
  - effectiveness/potential adverse effects of, 379
  - providing soon after admission to residential care, 416

**484**      **Index**

- hip strategy, 36
- the home, risk factors for falls and, 202
- hospitals
  - equipment/devices and, 400
  - older patients, fall risk assessment for, 231
  - prevention of falls in, 150, 396–409, 473
  - recently released patients, exercise for preventing falls and, 261
- hypertension
  - medications and, 192
  - stroke and, 178
  - white matter hyperintensities and, 132
- ICMT (interactive cognitive-motor training), 299, 304
- implementation science, 474
  - described, 437
  - future perspectives and, 453
  - theoretical approaches of, 438
- inactivity, cognitive impairment risk factors and, 147
- incidence of falls, 5–12
- incontinence, 185, 242, 418
- infarcts, 135
- injuries
  - fall-related injuries, epidemiology of, 3–18
  - future research perspectives and, 477
  - impact of exercise and, 263
- insomnia, 164, 195
- interactive cognitive-motor training (ICMT), 299, 304
- internal perturbations generated by functional tasks, 33
- interventions, 148–51, 260
  - cognitive behavioural, for fear of falling/fall risk, 311–21
  - costs of falls/fall-related injuries and, 461
  - environmental, to prevent falls, 360–77
  - future research perspectives and, 477
  - multiple. *see* multiple interventions
  - single, hospital patients and, 400–3
  - types of, defined, 386
- judgement error, risk factors for falls and, 148
- Kellogg International Working Group on the Prevention of Falls in the Elderly, falls defined by, 3
- kinematic gait patterns, 54
- Knowledge-to-Action (KTA) Framework
  - Knowledge Translation and, 437
  - two key phases of, 438
- Knowledge Creation Phase (of the KTA Framework), 439
- Knowledge Inquiry component (of the KTA Framework), 439
- Knowledge Synthesis component (of the KTA Framework), 440
- Knowledge Tools and Products component (of the KTA Framework), 440
- KT (Knowledge Translation), 437
- laboratory tests, fall risk assessment and, 327
- lateral falls, 113
- lateral reach, 28
- leaning, 26, 33
- leg extensor power, 95
- level walking, 51–8
  - gait changes and, 55
  - gait speed and, 52
- limbs, peripheral sensation and, 91
- location of falls, 12
- the “long lie”, 15
- low- /middle-income settings, future research into fall prevention and, 477
- lower limb, 91
  - footwear and, 122
  - peripheral sensation and, 91
  - proprioception and, 176
  - sensory abnormalities and, 92
  - strength/weakness and, 95
- machine learning-based fall detection, 214
- macular degeneration, as a risk factor for falls, 175
- magnetic resonance imaging (MRI), 137
- mechanoreceptors, 176
- medical conditions, as risk factors for falls, 11, 172–91, 242
  - acute illness and, 185
  - diseases affecting central processing and, 177–82
  - diseases affecting sensory input and, 173–7
  - fall prevention strategies and, 427
  - foot problems and, 184
  - peripheral arterial disease and, 183
  - suggestions/examinations and, 325
  - urinary incontinence and, 185
- medical management of risk groups for falls, 322–40
  - laboratory tests and, 327
  - risk assessment and, 323–7
  - treatments and, 327–32
- medications for fall prevention, 475
- medications, as risk factors for falls, 181, 192–201, 243, 322, 470
  - cardiovascular drugs and, 197
  - centrally acting medications and, 146, 193, 332
  - cognitive impairment risk factors and, 146
  - depression/anxiety and, 163, 165
  - Donepezil anti-dementia drug and, 149
  - medication review/assessment and, 332, 403
  - older people in residential care facilities and, 418
  - orthostatic hypotension and, 182
  - polypharmacy and, 198, 244
  - psychotropic medications and, 193–7
- mediolateral dynamic stability, 55
- methodological considerations for research into falls, 3–5
- methylphenidate, 476
- microbleeds, 135
- middle-aged people, future research into falls and, 474
- midsole cushioning, footwear choices and, 122
- mobility

**485 Index**

- assessment of, 325
- risk of falls and, 239
- mobility impairment, 16, 59
- foot problems and, 120, 184
- infarcts and, 136
- microbleeds and, 136
- peripheral arterial disease and, 183
- white matter hyperintensities and, 136
- MRI (magnetic resonance imaging), 137
- multi-direction reach, 28
- multiple interventions, 386–95, 472, 473
  - cost considerations and, 461, 465
  - fall-related fractures and, 393
  - for older people living in residential care facilities, 150
  - hospital patients and, 400
  - multiple, 386–95
  - types of, defined, 386
- muscle strength, 94, 240
- muscle synergies, 36
- myelopathy, 183
- myostatin inhibitors, 475
- neurally mediated syncope, 329
- neurological conditions, risk factors for falls and, 242
- new technology, for fall detection and risk assessment, 113, 211–26, 244, 470
  - behavioural factors/habits and, 205
  - future perspectives and, 216, 221
  - future research perspectives and, 474
  - hospitals and, 400
  - interactive cognitive-motor training systems and, 299
  - low-tech assessment tools and, 216
  - real-world utility of, 215
  - types of technology and, 212
- NSAIDs (non-steroidal anti-inflammatory drugs), 197, 244
- obstacle negotiation, 60–4
  - obstacle avoidance and, 62
  - step training and, 271, 277
- older people
  - defined, 5
  - perception of falls by, 427
- older people in hospitals
  - fall risk assessment for, 231
  - prevention of falls and, 150, 396–409, 473
  - recently released patients, exercise for preventing falls and, 261
- older people living at home (community dwellers)
  - environmental interventions to prevent falls and, 360–77
  - exercise for preventing falls and, 251, 261, 264
  - fall risk screening and assessment for, 228
  - incidence of falls and, 5–10
  - interventions for, 471
  - reactive step training and, 280
- older people living in apartment-style retirement villages, incidence of falls and, 11
- older people living in intermediate care facilities, incidence of falls and, 10
- older people living in residential care facilities
  - exercise for preventing falls and, 263
  - fall risk assessment for, 231, 413
  - incidence of falls and, 10
  - reactive step training and, 280
  - reducing risk of falling and, 410
  - risk factors and, 412
  - as a risk group for falls, 322
- opiates, 197, 244
- orthoses, 350
- orthostatic hypotension, 147, 181
  - medications and, 192
  - older people in residential care facilities and, 419
  - risk factors for falls and, 243
  - treatment and, 329
- osteoarthritis, 182, 184
- overground perturbation systems, for reactive step training, 277
- pain, foot pain/foot problems and, 119
- Parkinson's disease, 11, 179, 242, 470
  - cholinergic denervation and, 136
  - exercise for preventing falls and, 261
  - freezing of gait and, 64
  - older people in residential care facilities and, 419
  - reactive step training and, 280
  - risk groups for falls and, 322
  - wearable devices and, 354
- patient sitters, 401
- pattern recognition fall detection, 214
- peer leaders/peer support, fall prevention strategies and, 432
- Peninsula Health Falls Risk Assessment Tool, 231
- peripheral arterial disease, 183
- peripheral sensation, risk of falls and, 91, 175
- peripheral vascular disease, foot problems and, 184
- perturbations
  - external, responses to, 35–9
  - internal, generated by functional tasks, 33
  - reactive step training and, 274–81
  - translational, 36
- PET (Position Emission Tomography), 136
- pharmacodynamics, 192
- pharmacokinetics, 192
- physical ability
  - gait changes and, 55
  - preventing falls and, 149
  - risk factors for falls and, 148
- physical activity for healthy older people, 265
- physical fatigue, gait patterns and, 58
- physical restraints, preventing falls and, 354, 419
- physiological factors, risk of falls and, 240
- physiological function, 469
- physiological profile assessment (PPA), 136, 228
- plantarflexor power, 95
- podiatry, role in preventing falls and, 125, 185
- Poisson distribution, incidence of falls and, 4
- polypharmacy, 198, 244
- Position Emission Tomography (PET), 136

- postural hypotension. *see* orthostatic hypotension
- postural stability, 23–39, 470
  - assessment of, 325
  - defined, 23
  - foot problems and, 185
  - medical conditions and, 172, 176, 179, 182–5
  - medications and, 192–201
  - risk of falls and, 239, 240
- postural stress test, 35
- postural sway, 24–6
  - assessing, 106–9
  - leaning and, 26
  - perturbations and, 33–8
  - standing and, 24–6
  - stepping and, 29–33
- PPA (physiological profile assessment), 136, 228
- predicting falls, 469
  - coordinated stability and, 29
  - functional reach and, 28
  - lateral reach/multi-directional reach and, 28
  - maximum balance range and, 29
  - postural sway and, 25
  - standing and, 26
  - stepping and, 31
- prescribing medications, polypharmacy and, 198, 244
- Prevention of Falls Network Europe (ProFaNE)
  - fall-related injury definitions/classifications and, 5
  - falls, defined by, 3, 386
- private residences, risk factors for falls and, 202
- ProFaNE (Prevention of Falls Network Europe)
  - fall-related injury definitions/classifications and, 5
  - falls, defined by, 3, 386
- proprioception, 176
- protective stepping, 29
- psychoactive medications, 470
  - exercise for preventing falls and, 261
  - future research perspectives and, 475
  - low alcohol use and, 239
  - older people in residential care facilities and, 418
  - as a risk factors for falls, 193, 243, 403
- psychological factors, 160–71, 470
  - gait changes and, 55
  - risk of falls and, 241
- psychotropic medications, as risk factors for falls, 193–7
- public places
  - environmental interventions to prevent falls and, 360–77
  - risk factors for falls and, 205
- QuickScreen fall risk assessment tool, 230, 327
- reach, 28
- reactive step training, 271, 274–81
- reflex syncope
  - carotid sinus syndrome and, 330
  - vasovagal syncope and, 329
- research
  - methodological considerations for, 3–5
  - perspectives for the future and, 474–8
- residential care facilities
  - incidence of falls in, 10
  - prevention of falls in, 149, 410–24, 473
    - fall prevention interventions and, 416–19
    - managing and running/starting fall prevention programs and, 415
    - physical restraints and, 419
    - risk factors for falls and, 203, 412
- risk assessment, 216–22, 227–36, 323–7, 471
  - environmental profile and, 361
  - hospital patients and, 231, 397, 405
  - limitations of, 222
  - multifactorial fall risk assessment and, 228
  - older people in residential care facilities and, 231, 413
  - questions to ask, 323
- risk factors for falls, 144–8
  - extrinsic, 119, 348, 417
  - four-level rating system for, 237
  - intrinsic, 119, 206, 221, 348
  - older people in residential care facilities and, 412
  - relative importance of, 237–48
  - socio-demographic factors and, 237
- risk groups for falls, medical management of, 322–40
- risk screening, 227–36
- risk-taking behaviour, 242
- safety education interventions
  - hospital patients and, 151, 401
  - older people living at home (community dwellers) and, 149
  - visual impairment and, 345
- seasonal variations in the frequency of falls, 8
- secular trends in fall-related injuries, 9
- sedatives, 194, 243
- seizure, 328
- sensorimotor function, 87–104
  - physiological system risk assessment and, 230
  - risk of falls and, 130, 240
- sensory organization test, 35
- serotonin inhibitors, 195
- Short Physical Performance Battery (SPPB), 228
- single interventions, 473
  - environmental adaptations and, 362
  - exercise and, 261, 417
  - hospital patients and, 400–3
  - older people in residential care facilities and, 150
  - older people living at home and, 471
- sit-to-stand ability, 33, 58, 228
- epidemiology of falls/fall-related injuries
  - and, 411
  - fall risk assessment and, 216, 220
  - stroke patients and, 178
- sleep, quality of, 238
- slipping, 70–3
  - balance and, 112
  - footwear choices and, 123
  - step training and, 272, 275–80

**487**      **Index**

- socio-demographic factors, risk of falls and, 237
- socio-economic factors, low- /middle-income settings and, 477
- spectacles, 360
  - bifocal/multifocal and, 68, 245, 343
  - first time/new prescription and, 342–5
  - risk of falls and, 245, 328
  - unifocal, 328
- SPPB (Short Physical Performance Battery), 228
- stability limits, 23
- staff, 150, 151
  - compliant flooring and, 381
  - education/training and, 234
  - hip protector use and, 380
  - in hospitals, 397–405
  - multiple interventions and, 392
  - in residential care facilities, 410, 415, 416
- stair negotiation, 65–8, 206, 470
- standing, 24–6
  - balance control and, 105–10
  - reactive step training and, 279
- step length, 52, 69, 123, 218
- step training, 271–86
  - defined, 271
  - future research perspectives and, 475
  - impact for preventing falls and, 271
- step width, 52, 53, 105, 123, 217
- stepping, 29–33
  - internal perturbations and, 33
  - risk of falls and, 239
- stepping exercises, 290, 298
- stepping strategy, 36
- striatal dopaminergic transporter activity, 136
- stroke, 177, 242, 328
  - anticoagulants and, 198
  - exercise for preventing falls and, 261
  - older people in residential care facilities and, 419
  - risk groups for falls and, 322
  - wearable devices and, 353
- subcortical infarcts, 135
- sway. *see* postural sway
- syncope, 172, 327, 328
- tau, 136
- threshold-based fall detection, 213
- timed-up-and go test, 32, 34
  - peripheral sensation and, 92
  - as a simple screening tool, 228
  - turning and, 65
- transient self-limited loss of consciousness (TLOC), 323, 327, 328
- translational perturbations, 36
- treadmills, reactive step training and, 276
- tripping, 60, 68–70
  - balance and, 110–13
  - step training and, 272, 273–81
- turning, 64, 220
- vasovagal syncope, 329
- vestibular disorders, risk factors for falls and, 242
- vestibular function, 176, 243
  - components of, 176
  - risk of falls and, 93, 240
- vision
  - examinations/treatment and, 328
  - obstacle negotiation and, 63
  - stair negotiation and, 67
- visual impairment
  - exercise for preventing falls and, 261, 345
  - fall prevention interventions and, 341–7
  - residential care facilities and, 418
  - as a risk factor for falls, 89, 173, 240
  - safety education interventions and, 345
- visuospatial function, 137, 147
- vitamin D, 473
  - deficiency in, 8, 95
  - older people in residential care facilities and, 419
  - supplementation and, 150, 333
- volitional step training, 271–4
  - using exergames, 274
  - using targets, 272
- voluntary stepping, 29–33
- walking. *see* gait
- walking aids, 351, 470
- walking sticks, 351
- wearable devices/aids, 381, 470
  - for fall detection, 212, 216
  - for fall prevention, 353, 360
  - for risk assessment, 217
  - hip protectors and, 378, 379, 416
- white matter hyperintensities (WMHs), 132, 135, 242
- white matter integrity, 134
- WHO (World Health Organization), falls defined by, 3
- WMHs (white matter hyperintensities), 132, 135
- World Health Organization (WHO), falls defined by, 3
- Z-drugs, 195