

Index

- ACA. *See* Australian Cricket Association
- acamprosate, 76
- ACL injury. *See* anterior cruciate ligament injury
- ADHD. *See* attention deficit hyperactivity disorder
- adjustment disorder, from concussion, among rugby players, 104
- adolescent-focused psychotherapy, for anorexia nervosa, 119
- alcohol abuse
- concussion and, among rugby players, 100
 - among cricket players, 43
 - sleeping disorders and, 92, 95
- alcohol use disorder, European football culture and, 68–83
- with ACL injury, 68
 - in medical history, 69
- alcohol withdrawal syndrome, manifestations of, 78–83
- central nervous system irritability, 78
 - delirium tremens, 78–81
 - pharmacological therapy for, 81
- case formulation for, 71
- family history and, 69
- medical history and, 69
- medication history and, 69
- personal history and, 69–70
- collateral history, 70
 - forensic history, 70
- presenting complaints, 68–9
- psychiatric manifestations of, onset of, 77
- risk assessment, 71
- for comorbidity, 77
- screening investigations, 71
- answers in, 73–7
 - AUDIT tool, 73–5
 - CIWA-Ar, 71, 78–81
 - questions, 72–3
 - scoring from, 73, 74–5
- social history and, 69
- STIs and, 71
- testing scales
- MMSE scores, 70–1
 - MSE scores, 70–1
 - transtheoretical change model and, stage of, 70–1
- treatment plans and strategies, 71–2, 76
- alcohol support groups and, 75
- cognitive behavioural therapy, 75, 76
- pharmacotherapy in, 75–6
- alcohol withdrawal syndrome, manifestations of, 78–83
- central nervous system irritability, 78
 - delirium tremens, 78–81
 - pharmacological therapy for, 81
- alcohol support groups, 75
- amenorrhea, RED-S and, 21
- cognitive behavioural therapy for, 23–4
- primary, 21
 - screening for, 21–2
 - secondary, 21
- amphetamines, for ADHD, 62
- anabolic androgenic steroids, 64
- anorexia nervosa
- among adolescent swimmers, mental health emergencies from, 117–18
 - adolescent-focused psychotherapy for, 119
 - cognitive behavioural therapy for eating disorders, 119
- anterior cruciate ligament (ACL) injury, alcohol use disorder and, 68
- anxiety disorders. *See also* attention deficit hyperactivity disorder
- catastrophising, 94
 - core symptoms, 94
 - generalised anxiety disorder, 92
- atomoxetine, 61
- attention deficit hyperactivity disorder (ADHD), cycling and
- age of onset, 60
 - core symptoms, 59–60
 - diagnostic criteria for, 59, 60–1
 - hyperactive/impulsive type, 60–1
 - hyperkinetic disorder and, 60
 - inattentive type, 60
 - patient history, 55–7
 - collateral history, 56–7
 - family history, 55
 - forensic history, 56
 - medical history, 55
 - medication history, 55–6
 - past psychiatric history, 55
 - personal history, 56
 - pre-morbid personality, 56
 - social history, 55–6
- planning strategies for, formulation of, 58–65
- of assessment criteria, 59–60
- presenting complaint, 54–5
- risk assessment for, 57–8
- screening investigations for, 58
- testing scales, with MSE, 57
- treatment strategies for, 61–5
- cognitive behavioural therapy, 62–3
 - hormonal therapy, 64
 - pharmacotherapy in, 61–3

- (ADHD) (cont.)
 therapeutic use
 exemption, 54, 61, 63–5
 under World Anti-Doping
 Agency guidelines, 60–5
 AUDIT tool, for alcohol use
 disorder, 73–5
 Australian Cricket Association
 (ACA), 51–2
- benzodiazepines, 78
 bipolar affective disorder
 age of onset, 50–1
 boxing and, 36
 clinical presentation of,
 51
 comorbidity risks, 51
 cricket and, 45, 47, 48,
 50–1
 lithium treatment for,
 48–9, 50
 long-term prognosis, 51
 mortality rates, 51
- blood tests
 exercise addiction and, 20
 for mental health
 emergencies, among
 adolescent
 swimmers, 114
- bone health, RED-S and, 22
 boxing. *See* low mood disorders
- CAMHS. *See* Child and
 Adolescent Mental
 Health Services
- catastrophising, 94
 Cavaday, Naomi, 130
 CBT. *See* cognitive behavioural
 therapy
- CBT-E. *See* cognitive
 behavioural therapy for
 eating disorders
- central nervous system (CNS)
 irritability, 78
- Child and Adolescent Mental
 Health Services
 (CAMHS), for
 adolescent swimmers,
 mental health
 emergencies among,
 109, 115–16
- chlordiazepoxide, 78
 chronic PTSD, 130
 chronic traumatic
 encephalopathy (CTE),
 7, 8–11
- assessment and diagnosis
 of, 8–9
- chronic traumatic
 encephalopathy
 neuropathic change
 (CTE-NC), 9, 10–11
- citalopram, 94
- Clinical Institute Withdrawal
 Assessment for
 Alcohol – revised
 version (CIWA-Ar), 71,
 78–81
- CNS irritability. *See* central
 nervous system
 irritability
- cognitive behavioural
 therapy (CBT)
 for alcohol use disorder,
 75, 76
 for amenorrhea, from RED-
 S, 23–4
 for attention deficit
 hyperactivity disorder,
 with cycling, 62–3
 CBT-E, 119
 for concussion, from
 rugby, 102
 e-CBT, 76
 for exercise addiction, 24–5
 for gambling disorder, 38
 boxing and, 33
 for mental health
 emergencies, from
 cricket, 49
 for sleeping disorders, 93–4
 for trauma, from tennis, 127
- cognitive behavioural therapy
 for eating disorders
 (CBT-E), 119
- cognitive impairment,
 American football and
- chronic traumatic
 encephalopathy, 7, 8–11
 assessment and diagnosis
 of, 8–9
 CTE-NC, 9, 10–11
 lab investigations for, 5–6
 patient history, 3–4
 collateral history, 4
 medication history, 3
 past medical history, 3
 personal history, 4
 social history, 3
 physical examination for, 5
 planning strategies for,
 formulation of, 6–12
- presenting complaint, 1–2
 risk assessment for, 5
 testing scales for, with
 MSE, 4–5
 treatment strategies for,
 through psychiatric
 medications, 12
- CogSport tests, in rugby,
 105
- Compulsive Exercise Test,
 118–19
- concussion, from rugby
 alcohol use issues, 100
 case formulation with,
 102
 collateral history, 101
 diagnosis of, 104–5
 adjustment disorder
 and, 104
 major depressive disorder
 and, 104
 injury history, 98
 investigations for, 102
 medical history, 100
 medication history, 100
 mental health after. *See*
 mental health
- personal history issues,
 100–1
- post-concussive symptoms,
 104–5
- pre-morbid personality,
 101
- presenting complaints, 98–9
 history of, 99
 protocols for, 98
 psychiatric history with,
 99–100
- recovery from, 105
 social history for, 100
 testing scales for, 105
 CogSport tests, 105
 MSE, 101–2
 SCAT5, 105
- treatment plans and
 strategies, 102–7
 cognitive behavioural
 therapy, 102
 pharmacotherapy in,
 103
- Cotterill, David, 82
- cricket. *See* mental health
 emergencies,
 cricket and
- CTE. *See* chronic traumatic
 encephalopathy

- CTE-NC. *See* chronic traumatic encephalopathy neuropathic change
- cycling. *See* attention deficit hyperactivity disorder
- deliberate self-harm (DSH), 44–5
 risk assessment for, 45
- delirium tremens (DTs), 78–81
- depression. *See* major depressive disorder
- dexamphetamine, 62
- disordered eating, 23
- disulfiram, 76
- DSH. *See* deliberate self-harm
- DTs. *See* delirium tremens
- duloxetine, 94
- eating disorders
 among adolescent swimmers, mental health emergencies and
 anorexia nervosa, 117–18
 family history of, 115
 food diaries and, 111
 male presentation of, 118
 multi-factorial causes of, 118
 screening tools for, 118–19
- anorexia nervosa
 among adolescent swimmers, mental health emergencies from, 117–18
 adolescent-focused psychotherapy for, 119
 cognitive behavioural therapy for eating disorders, 119
- disordered eating, 23
- RED-S and, 23
 perfectionism and, 23
- screening tools for
 Compulsive Exercise Test for, 118–19
- Eating Disorders Examination Questionnaire, 118–19
- SCOFF questionnaire, 118
- Eating Disorders Examination Questionnaire (EDE-Q), 118–19
- e-CBT. *See* electronic cognitive behavioural therapy
- EDE-Q. *See* Eating Disorders Examination Questionnaire
- electronic cognitive behavioural therapy (e-CBT), 76
- Epstein-Barr virus, 21
- escitalopram, 94
- European football culture. *See* alcohol use disorder
- exercise addiction. *See also* relative energy deficiency in sport
 overtraining syndrome and, 19, 23, 24
- patient history, 17
 medical history, 17
 medication history, 17
 pre-morbid personality, 17–18
 personal history, 17
 social history, 17
- perfectionism and, 23
- plan management for, formulation of, 19–23
- presenting complaint, 15–16
 history of, 15–17
- risk assessment, 18, 24
- screening investigations, 18–19
 blood tests, 20
- testing scales, with MSE, 18
- treatment approaches and strategies, 24–5
- fluoxetine, 36, 50, 94, 103
- food diaries, 111
- gambling disorder, 37
 boxing and, 33
 cognitive behavioural therapy approach to, 33
- diagnostic criteria for, 37
- treatment strategies for, 38
- cognitive behavioural therapy, 38
- motivational enhancement therapy, 38
- motivational interviewing, 38
- generalised anxiety disorder, 92
- golf. *See* sleeping disorder
- hormone therapy
 for attention deficit hyperactivity disorder, 64
- RED-S and, 22
- hormone replacement therapy, 22
- hyperactive attention deficit hyperactivity disorder, 60–1
- hyperkinetic disorder, 60
- hypomania, 36
- impulsive attention deficit hyperactivity disorder, 60–1
- inattentive-type attention deficit hyperactivity disorder, 60
- insomnia, 95
- International Society for Sports Psychiatry (ISSP), 50
 curriculum for, 12
- International Trauma Questionnaire (ITQ), 128–30
- interpersonal psychotherapy (IPT), 49
- intramuscular Pabrinex, 81
- IPT. *See* interpersonal psychotherapy
- ISSP. *See* International Society for Sports Psychiatry
- ITQ. *See* International Trauma Questionnaire
- Korsakoff syndrome, 81
- lamotrigine, 50
- low mood disorders, boxing and
 bipolar affective disorder and, 36
- concussions and, 28
- gambling disorder and, 33
- cognitive behavioural therapy approach to, 33
- hypomania and, 36

- low mood disorders (cont.)
 major depressive disorder,
 35–6. *See also* major
 depressive disorder
 mania and, 36
 patient history, 30–2
 family history, 30
 forensic history, 32
 medical history, 30
 past psychiatric history, 30
 personal history, 31
 pre-morbid
 personality, 32
 social history, 31
 plan management for,
 formulation of, 33–8
 presenting complaint, 29–30
 history of, 29–30
 risk assessment for, 32
 screening investigations
 for, 33
 testing scales for, with MSE/
 MMSE, 32
 lurasidone, 50
- major depressive disorder
 boxing and, 35–6
 diagnostic guidelines
 for, 35–6
 among rugby players, 106
 diagnosis of, 104
 sleeping disorders and,
 among golfers, 92
 symptoms of, 35
 somatic, 35–6
 treatment strategies
 for, 36–7
- mania, 36. *See also* hypomania
- mental health, after
 concussions, through
 rugby
 collateral history, 101
 family history, 100
 medication history and,
 100
 past psychiatric history,
 99–100
 personal history issues,
 100–1
 pre-morbid personality, 101
 prevalence rates, of common
 depressive systems, 106
 retirement and, 106
 risk assessment of, 102
 mental health emergencies,
 cricket and
- bipolar affective disorder
 and, 45, 47, 48, 50–1
 lithium treatment for,
 48–9, 50
 for deliberate self-harm, risk
 assessment for, 45
 patient history, 42–3
 for alcohol and drug
 use, 43
 collateral, from other
 sources, 44
 family history, 42–3
 medical history, 42–3
 medication history, 43
 personal history, 43
 plan management for,
 formulation of, 45–9
 psychological
 interventions as part
 of, 46
 player associations' response
 to, 51–2
 presenting complaints, 42
 history of, 42
 risk assessment of, 45
 testing scales for, with
 MSE, 44–5
 treatment strategies
 with cognitive behavioural
 therapy, 49
 International Society for
 Sports Psychiatry
 guidelines, 50
 with interpersonal
 psychotherapy, 49
 with lithium, 48–9, 50
 pharmacotherapy
 approaches, 49, 50
- mental health emergencies,
 among adolescent
 athlete swimmers
 CAMHS and, 109, 115–16
 eating disorders and
 anorexia nervosa, 117–18
 family history of, 115
 food diaries and, 111
 male presentation of, 118
 as multi-factorial, 118
 screening tools for, 118–19
 patient history, 112–13
 collateral history, from
 other sources, 113
 family history, 112
 food diary, 111
 medical history, 112
 personal history, 112–13
- social history, 112
 plan management for,
 formulation of, 115–19
 blood testing as part
 of, 114
 CAMHS interventions as
 part of, 115–16
 pre-morbid personality, 113
 presenting complaint,
 110–11
 history of, 110–11
 risk assessment for, 114–15
 testing scales, with MSE, 113
- Mental State
 Examination (MSE)
 alcohol use disorder
 and, 70–1
 for attention deficit
 hyperactivity disorder,
 with cycling, 57
 for cognitive impairment,
 from American
 football, 4–5
 for concussion, from rugby,
 101–2
 for exercise addiction, 18
 for low mood disorders,
 among boxers, 32
 for mental health
 emergencies, among
 adolescent
 swimmers, 113
 for sleeping disorders,
 among golfers, 89
- methylphenidate, 62
- MI. *See* motivational
 interviewing
- Mini Mental State Examination
 (MMSE)
 alcohol use disorder
 and, 70–1
 for low mood disorders,
 among boxers, 32
 motivational enhancement
 therapy, 38
 motivational interviewing
 (MI), 38
- MSE. *See* Mental State
 Examination
- naltrexone, 38, 76
- National Society for the
 Prevention of Cruelty to
 Children, 132
- New Zealand Cricket Players
 Association, 51–2

- obstructive sleep apnoea (OSA), 95–6
- olanzapine, 50
- ondansetron, 76
- OSA. *See* obstructive sleep apnoea
- overtraining syndrome, 23
- exercise addiction and, 19, 23, 24
- perfectionism and, 23
- symptoms of, 23
- paroxetine, 94
- PCA. *See* Players Cricket Association
- perfectionism, 23
- Players Cricket Association (PCA), 51
- post-traumatic stress disorder (PTSD)
- chronic PTSD as distinct from, 130
- trauma, from tennis, 127, 130
- diagnostic criteria, 129–30
- pharmacological therapy strategies, 131
- treatment plans for, 127
- pregabalin, 94
- primary amenorrhea, 21
- psychotherapy
- adolescent-focused, for anorexia nervosa, 119
- interpersonal psychotherapy, 49
- PTSD. *See* post-traumatic stress disorder
- quetiapine, 50
- relative energy deficiency in sport (RED-S), 19, 21–5
- amenorrhea and, 21
- cognitive behavioural therapy for, 23–4
- primary, 21
- screening for, 21–2
- secondary, 21
- causes of, 22
- bone health, 22
- hormone production, 22
- under-fuelling, 22–3
- clinical presentation, 21–2
- disordered eating, 23
- eating disorders and, 23
- perfectionism and, 23
- Epstein-Barr virus and, 21
- hormone health and, 22
- hormone replacement therapy, 22
- overtraining syndrome, 23
- perfectionism and, 23
- symptoms of, 23
- return to play recovery from, 25
- treatment strategies for, 22
- hormone replacement therapy, 22
- return to play (RTP) recovery, 25
- Rowe, Luke, 64
- RTP recovery. *See* return to play recovery
- rugby. *See* concussion; mental health
- SCAT5. *See* Sports Concussion Assessment Tool
- SCOFF questionnaire, for eating disorders, 118
- secondary amenorrhea, 21
- sertraline, 36, 94
- sexual abuse, in tennis, 131–2
- National Society for the Prevention of Cruelty to Children guidelines, 132
- safeguarding from, 131–2
- sexually transmitted infections, 71
- sleeping disorders, golf and alcohol abuse and, 92, 95
- classifications of, 95–6
- generalised anxiety disorder and, 92
- insomnia, 95
- major depressive disorder and, 92
- obstructive sleep apnoea, 95–6
- patient history, 87–9
- collateral history, 89
- family history, 87
- medication history, 87
- past medical history, 87
- personal history, 88
- social history, 88
- planning strategies for, formulation of, 90–6
- presenting complaint, 85–7
- risk assessment for, 89
- screening investigations for, 90
- testing scales for, with MSE, 89
- treatment strategies for, 93–4
- cognitive behavioural therapy, 93–4
- pharmacotherapy approaches, 94
- soccer. *See* alcohol use disorder
- Sports Concussion Assessment Tool (5th edition) (SCAT5), 105
- suicide. *See* deliberate self-harm
- swimming. *See* mental health emergencies, among adolescent athlete swimmers
- tennis. *See* trauma
- testing scales
- for alcohol use disorder
- MMSE scores, 70–1
- MSE scores, 70–1
- transtheoretical change model and, stage of, 70–1
- for concussion, from rugby, 105
- CogSport tests, 105
- MSE, 101–2
- SCAT, 5, 105
- for cricket, for mental health emergencies, 44–5
- for low mood disorders, among boxers, 32
- for trauma, in tennis
- ITQ, 128–30
- MSE, 126
- therapeutic use exemption (TUE), 54, 61, 63
- under World Anti-Doping Agency guidelines, 63–5
- transtheoretical change model, for alcohol use disorder, 70–1
- trauma, tennis and case formulation, 127
- patient history
- family of, 124
- medical, 124
- for medications, 124

140	Index	
trauma, tennis (cont.) <ul style="list-style-type: none">of presenting complaints, 123–4psychiatric issues, 124social history, 124	to Children <ul style="list-style-type: none">guidelines, 132safeguarding from, 131–2	valproate, 49, 50
presenting complaints, 123–4 <ul style="list-style-type: none">history of, 123–4	testing scales for <ul style="list-style-type: none">ITQ, 128–30MSE, 126	vortioxetine, 94
PTSD and, 127 <ul style="list-style-type: none">diagnostic criteria, 129–30pharmacological therapy, 131treatment plans for, 127	treatment plans and strategies, 127–32	WADA. <i>See</i> World Anti-Doping Agency
risk assessment for, 126	cognitive behavioural therapy, 127	WE. <i>See</i> Wernicke’s encephalopathy
sexual abuse, 131–2 <ul style="list-style-type: none">National Society for the Prevention of Cruelty	pharmacological therapy, 131	Wenger, Arsene, 82
	for PTSD, 127	Wernicke’s encephalopathy (WE), 81
	TUE. <i>See</i> therapeutic use exemption	women athletes, psychiatric case studies for. <i>See</i> mental health emergencies, among adolescent athlete swimmers
	under-fuelling, RED-S and, 22–3	World Anti-Doping Agency (WADA), 60–5