

Index

- accumulating data 44–47
- additive rules of probability 231–233
 - disjoint (mutually exclusive) events 231–233
 - Second Additive Rule 232–233
- adjusted R^2 squared values 523
- Akaike, Hirotugu 612–613
- Akaike information criterion (AIC) 612, 684–685
- alternative hypothesis 297
- analysis of variance (ANOVA) 297, 414–416
 - assumptions of 419–422
 - Bonferroni adjustment 444–446
 - computations for 417–420, 469–474
 - development of 414
 - effect size 426–431
 - F distribution 425
 - F-ratios 427
 - index of separation for 416, 419
 - Kruskal–Wallis analysis of variance 639
 - Levene’s test 451
 - mean squares 417–420
 - multiple t -tests and 412–414
 - one-tailed tests 303
 - regression 518–520
 - robvar** command 425
 - summary table 425
 - t -tests 416
 - two-tailed test 333
 - in two-way analysis of variance 465
 - variability between/within groups 416
 - variance estimators 417
 - see also* two-way analysis of variance
- anova** command 429
 - t -distribution and 420
- Anscombe.dta* 524–525, 673
- arithmetic mean
 - comparing with mode and median 84–85
 - of dichotomous variable 82
 - weighted 83–84
- arithmetic operators 104
- assumptions 622, 679–683
 - of analysis of variance 419–420, 466–467
 - of chi-square test of independence 633
 - of independent samples 340–342
 - of paired samples 354–359
- asymmetric curves 95–99
- average *see* arithmetic mean
- average squared deviation 205
- average treatment effect 398, 399, 404
 - b -weight equation 556
 - first-order effect variables 598–599
 - in multiple regression 558
 - as partial slope in multiple regression 568–571
 - statistical significance 563–564
 - testing for statistical significance 514–518
- b, β coefficients, with non-correlated predictors 555–556
- b -weight 590, 594
- balanced factorial designs 467–468
- bar graphs 56
 - clustered for bivariate relationships 184–187
 - frequency and percent distributions of nominal data in Stata 28
- Bartlett’s test 429
- basketball.dta* 673
- Bayes, Sir Thomas 239
- Bayesian information criterion (BIC) 684–685
- Bem Sex-Role Inventory* 145
- best-fitting linear equations
 - average squared deviation in 205
 - least squares criterion 205
 - Pearson correlation coefficient and 202–209
 - prediction lines 205
 - regression lines 205
 - residuals 204
 - standard error of estimate in 205
- between-group variance 416
- bias
 - populations and 285–286
 - selection 405
- bimodal distributions 75
- bimodality 75
- binomial probability model
 - applicability 249–254
 - approximation of using normal distribution 260
 - distribution in 244–254
 - exercises 261–268
 - factorial notation 248
 - identical trials 244, 249
 - independent trials 244, 249
 - law of large numbers 251
 - normal probability model and 254–261

INDEX

717

- binomial tail function 254
- bivariate contingency table 630
- bivariate normal distribution 509, 513
 - homoscedasticity property 503–506
 - linearity property 503–506
- Blood 673
- Bonferroni adjustment 444–446
 - multiple measures 446–447
- bootstrapping 360–368
- Brainsz.dta* 674
- causal studies 391–395
- centering 125
- central tendency 74
- chi-square test of independence, assumptions 633
- Cohen's *d* 311
- command summary for chapter 4–5, 614–616
- continuous variables 8–9
- correlation scale 171
- Court TV 75
- criterion variables 512
- Cronbach's alpha 392
- cross-sectional design
- cross-tabulation 630, 632
- crossed factorial design
- cumul** command 45
- cumulative binomial probability value 252, 253
- cumulative percent distributions 44–45
- Currency 674
- data analysis
 - computers in 3
 - context setting in 11–14
- data management, Stata .do-file 147–150
- data reliability 174
- data sets
 - Anscombe.dta* 524–525, 673
 - Hamburg.dta* 159, 160, 676–677
 - Ice Cream.dta* 106, 506, 559, 560, 679–683
 - Impeach.dta* 679–683
 - Learndis.dta* 112, 154, 160, 375, 378–379, 678
 - Likert.dta* 125, 126, 144
 - ManDext.dta* 417
 - NELS.dta* 11, 13, 26, 43, 44, 74, 159, 236, 511
 - Statisticians.dta* 25
 - Stepping.dta* 499
 - TEMP.dta* 87
- data transformation 54
 - nonlinear 135–142
 - recoding variables 144–147
- data value
- de Moivre, Abraham 282
- degrees of freedom
 - in ANOVA 420–422
 - chi-square distribution 623–639
 - independent samples 340
 - t*-distribution 320–321
 - in univariate distributions 26
 - dependent samples 337–338
- dependent variables 202, 217, 402, 427
- describe** command 108
- descriptive statistics 2, 228, 269
 - defined 2
- descriptive studies 391–395
- detail** option 79, 424
- diagnostics 524–529, 533–536
- dichotomous independent variables 536
 - interaction and 603–610
 - multiple regression 571–576
- dichotomous variables 4, 244
 - defined 82
 - mean values 82
 - phi coefficient 184
 - Point Biserial correlation coefficient 176–185
 - simple linear regression and 214–217
- discrepancy 531
- discrete probability 228–230
- discrete variables 8–9, 26, 47, 244
- disjoint events 230
 - additive rules of probability 231–233
- disordinal interaction 461
- dispersion *see* spread
- display** command 96, 168, 246–247, 253, 256, 302, 331, 423
- distribution
 - asymmetric curves 95–99
 - shapes of 95–99
 - z*-scores 133–134
- distribution location 74–86
 - arithmetic mean as 80
 - comparing mean, mode, and median 84–85
 - mean 80–84
 - median 78–80
 - mode 74–78
 - selecting measures of 99
 - weighted mean 83–84
- distribution-tied methods 622
- division, in linear transformation 119
- Doctrine of Chances* (de Moivre) 282
- doedit** command 147, 148
- drop** command 108
- Dynarski, Susan 410
- edit** command 19
- effect size
 - hypothesis testing and 305–306, 310
 - independent sample *t*-test 349–353
 - one sample *t*-test 333–336
 - one-way analysis of variance 426–431
 - paired sample *t*-test 359–360
 - partial eta squared 476–477
 - R* 520–524
 - simple linear regression and 217
 - two-way analysis of variance 476–477
- Efron, Bradley 360
- egen** command 34, 78, 129, 143, 166
- egen mode** command 77

- endogeneity 402
- equations
 - for *b*-weights 556
 - for hypothesis tests 369
 - independent variables 210
 - standardized regression equation 210
 - for *t*-test values 369
 - unstandardized regression equation 573, 594, 606
 - z*-values 298
- estimators
 - biased and unbiased 285
 - variance 339, 340
- estpost** command 656
- events 228
- Excel
 - correlation matrices 661–663
 - regression output 663–668
 - summary statistics tables 656
 - working directory 655–656
- expected values 400
- experimental mortality 405, 406
- experiments
 - agricultural 414
 - natural 410
 - probability 244
 - simple 228
- Exploratory Data Analysis* (Tukey) 55
- external validity 404
 - for quasi-experimental design 404, 405–406
- F*-distribution 420–422
 - descriptive statistics 435
 - Levene's test 427
 - one-way analysis of variance 425, 427
- robvar** command 427
 - two-way ANOVA 469
- factorial designs 467–468
- factorial notation 248
- family-wise error rate (α_{FW}) 413, 431, 444
- firearms 79
- first additive rule of probability 231,
 - 232, 236
- first-order effects 594, 599–600
- Fisher, Ronald A. 301, 414
- Fisher's exact test 635, 637–638
 - hypergeometric distribution 635
 - Stata for 635
- Fisher *r* to *Z* transformation 509
- Fisher's least significant difference (LSD) 432
- five-number summary 51–52
- fixed-effects models 479
- fixed factors 479
- forval** command 98
- Framingham* data set 17, 62,
 - 113, 379
- Framingham Heart Study 675
- freq** option 46
- frequency distribution table 27, 32, 56
- frequency distributions 26
- bar charts 28
- boxplots 51–56
- five-number summaries 51–52
- graphical selection summary 56
- grouped frequency distribution 34
- histograms 38–39
- line graphs 40–42
- mutually exclusive and exhaustive categories and 28
- ogive curve 46
- percentile ranks 46–51
- percentiles 46–51
- pie charts 31
- relative frequency 28
- skewness of 95–99
- stem-and-leaf displays and 35
- Galton, Sir Francis 165, 206–207
- gen** command 120, 145, 146
- generate **random** command 273
- global hypothesis 444
- global variables 98
- goodness-of-fit
 - chi-square and 625–632
 - regression equations and 524
- Gossett, William Sealy 320
- grand mean 474–475
- graph bar** command 57
- graph box** command 53, 55, 88, 100–102
- graph hbar** command 28
- graph twoway** command 208
- graphs
 - graphical displays of variables 56
 - modifying appearance of 56
- grouped frequency distribution 34
- Hamburg.dta* 159, 160, 676–677
- Handbook of Experimental Psychology* 3
- help** command 30, 46, 51, 79, 107, 120
- heterogeneity *see* spread
- heteroscedasticity 527
- hierarchical analyses 595
- hierarchical multiple regression 595
- histogram** command 39, 276, 278, 325
- histograms 38–39, 56, 57
- homoscedasticity 506
- honestly significant difference (HSD) 431
- hypergeometric probability distribution 635, 637–638
- hypothesis testing 296
 - decreasing standard error of means 311
 - effect size 306–307, 310
 - equations for 369
 - interval estimation and 305–306
 - Neyman and Pearson null 301
 - nondirectional hypothesis test 306
 - one sample *t*-test 327
 - pitfalls of 312
 - power of the test 307–312
 - significance level 310

INDEX

719

- two-way analysis of variance 466–467
 - type I and II errors and 307–312
- Ice Cream.dta* 106, 506, 560, 679–683
- identical trials 244
- Impeach.dta* 679–683
- inconsistency *see* spread
- independent samples 338–340
 - assumptions of 340–342
 - degrees of freedom 340
- independent samples *t*-test 373
 - ANOVA as extension 416
 - effect size 349–353
 - flowchart 371
- independent trials 244
- independent variables 558
 - dichotomous 536
 - relative importance of in multiple regression 565–566
 - simple linear regression 214–217
- index of separation 416
- indirect linear relationship 166
- inferential statistics 228
 - defined 2
 - random sampling and 270–273
 - sampling and 269, 275
- infinite populations 274
- influential observations 529–530
- information criterion 684–685
- instrumental variables 402
- instrumentation 405
- interaction 460–465, 591
 - dichotomous and continuous independent variables 603–610
 - disordinal 462
 - ordinal 462
 - statistically significant two-way interaction 600–603
- interaction models 592
- internal validity 404
- interpretation 14
- interquartile range (IQR) 50, 52, 86, 89–91
- interval estimation 312
 - central limit theorem and 293
 - confidence interval and 293
 - hypothesis testing and 305–306
 - interval estimation 293–296
- interval-leveled variables 175, 437
- interval measurement of variables 5–6
- inverse normal probability function 259
- invnormal(*p*)** function 259, 261
- IQ scale 10, 75
- joint distribution 503
- “Justice at work and reduced risk of coronary heart disease among employees” (article) 394
- Kaplan, E. L. 312
- kdensity** command 41
- Kruskal–Wallis analysis of variance 639, 647–648
- kwallis** command 648
- label define** command 126, 145
- label value** command 126, 145
- law of large numbers 240, 251, 253
- Learndis.dta* 112, 154, 160, 375, 378–379, 678
- least interval-leveled variables 159–166, 596, 597, 604
- least significant difference (LSD) 432
- least squares criterion 205
- legend option 46, 575
- Levene’s test 439
 - F*-distribution 438
 - one-way analysis of variance 438, 439
- robvar** command 439, 494
- two-sample designs and 346, 351
- two-way analysis of variance 494
- variances 340, 341, 346
- leverage 529–530
 - Stata for 530–531
- Likert scale 10
- Likert.dta* 125, 126, 144
- line graphs 40–42, 56, 57
 - Stata 601
- linear composite variables 554
- linear regression
 - b*-weight for 514–518
 - equation 202
 - influential outliers 529–530
 - leverage 529–530
 - residual plots 524–529
 - statistical significance 516
- linear regression equation 503
 - best-fitting 202–209
 - criterion variables 512
 - dichotomous independent variables and 214–217
 - prediction accuracy with 56–58, 512
- linear regression model
 - R* as measure of fit of 210–217
 - scatterplot importance for 217–218
- linear relationships 503
- linear transformation 119–121
 - defined 118
 - effect of summary statistics of a distribution 121–123, 151–153
- effect on measures of central tendency 123
- effect on measures of shape 123
- effect on measures of spread 123
- exercises 151–153, 154–156
- Pearson correlation coefficient and 170
- reflection as 126
- shape of a distribution and 121
- standard scores 126–127
- translation as 125
- z*-scores 127–128
- list** command 78
 - median as 78
- location characteristics
 - mode as 74
 - of a univariate distribution 74

- log file, in Stata 13, 18–19
- log-linear analysis 639
- logarithms 135
- logical operators 104
- longitudinal designs
- lwidth** option 41, 42, 57
- main effects 460
- Mandex 679
- ManDext.dta* 417
- Mann–Whitney *U*-test 642–647
- marginal counts 183
- marginal effect 558
- margins** command 481, 482, 600, 602, 606
- marginsplot** command 601, 602
- Marijuana.dta* 260, 676–677
- marker option 53
- math achievement of tenth-graders in southern US 3
- matlist** command 660
- matrices 169
 - in Excel 661–663
- maturity, in quasi-experimental design 405
- mean 80–84
 - column marginal mean 460
 - comparing with mode and median 84–85
 - dichotomous variables 82
 - grand mean 474–475
 - one-way analysis of means (ANOVA) 414–416
 - row marginal mean 460
 - two-way analysis of variance
 - weighted mean 83–84
- mean squares 418
- measurement of variables
 - data transformation 54
 - location in 99
 - spread in 99
- median 78–80
 - comparing with mode and mean 84–85
- median splits 145–146
- metrics 102, 118
- mixed-effects models 479
- mode 74–78
 - bimodal 74–78
 - comparing with mean and median 84–85
 - trimodal 76
- moderator variables 591
- monotonic transformations 118
- multiple regression
 - additive models 590
 - b*-weight as partial slope in multiple regression 568–571
 - basic equation with two predictors 554–555
 - correlated predictors 556–558
 - dichotomous independent variables 571–576
 - first-order effect variables 599–600
 - hierarchical 595
 - interaction 596
 - introduction 553–554
 - model selection 610–613
 - R*-squared significance 566–568
- summary of important principles of 556–558
- multiple *t*-tests 412–414
- multiplication, in linear transformations 119
- multiplicative rule of probability 233–236
- mutually exclusive events 230
- NELS.dta* 11–14, 19, 26, 43, 44, 74, 159, 178, 236, 511, 536, 634
- nestreg** command 595, 597
- Neyman, Jerzy 296, 301
- nodots** option 362
- noise components 337
- nominal variables 4, 26–32
 - chi-square goodness-of-fit test 623–639
 - graphical display of 56
- non-randomized controlled studies 395
 - Simpson’s paradox and 397
- nondirectional hypothesis testing 306, 331
- nonlinear transformation 118
 - logarithm and square root transformations 135–142
 - ranking variables 142–144
- nonparametric methods 639–649
 - ordinal dependent variables and 639–649
 - Spearman rank correlation coefficient 640
 - Wilcoxon sign test 3, 640
- norm** command 325
- normal distribution 255
- normal probability model 254–261
 - approximation of a binomial probability 260
 - complement rule 255
- normality 322, 325
- null hypothesis 297
- observational studies 394
- ogive curves 45–46
- omnibus *F*-test 432
- one sample *t*-test 320–321
 - confidence intervals 323–333
 - degrees of freedom 320–321
 - effect size 333–336
 - equation 321–323
 - exercises 376–378
 - flowchart 370
 - hypothesis testing 330–333
 - parent population normal deviation and 322–323
 - Stata commands 372
- one-tailed tests 303
- one-way analysis of variance (ANOVA) 414–426
 - assumptions 419–420
 - computation 417–420
 - effect size 426–431
 - post-hoc multiple comparison tests 431–432
 - summary table 425
- oneway** command 33, 423, 428, 430, 432, 433, 438
- ordinal interaction 462
- ordinal measurement 4–5

INDEX

721

- ordinal variables
 - chi-square test of independence 623–639
 - graphical display of 56
 - nonparametric methods 639–649
 - Spearman rank correlation coefficient and 174–176
 - orthogonal factorial designs 467–468
 - outcome variable 393
 - outliers 44, 53, 603
 - bivariate
 - influential 529–530
 - z -score detection of 130–134
 - over** commands 58–61
 - Stata commands
 - p*-values 299, 437, 472
 - bootstrap technique 374
 - chi-square test of independence 630
 - Fisher's exact test 635
 - in *t*-distribution 322, 333, 345
 - paired samples *t*-test 353–360
 - flowchart 371
 - panel design 399
 - parameters 269
 - parametric methods 622
 - parent populations 275, 282
 - parsimony 52
 - partial eta squared effect size 477
 - partial regression coefficients 558
 - partial slope 568–571
 - partialed product 594, 605
 - partitioning total sum of squares, two-way analysis of variance 468–469
 - Pearson
 - Egon 25, 301
 - Karl 25, 165, 626
 - Pearson product moment correlation coefficient 165, 167, 185, 217, 503, 509–512
 - best-fitting linear equation 202–209
 - bivariate normal distribution and 506–509
 - causation and 172
 - correlation strength of linear relationship and 168
 - data reliability and 174
 - dichotomous variables and 176–185
 - direct linear relationships 166
 - indirect linear relationships 166
 - linear regressions and 206, 209–210
 - linear transformations effect and 172–173
 - ordinal variables and 174–176
 - perfect negative linear relationships 168
 - perfect positive linear relationships 168
 - phi coefficient 176, 181–185
 - Point Biserial correlation coefficient 176–185
 - restriction of range and 173–174
 - shape of underlying distributions and 174
 - Spearman rank correlation coefficient 174–176, 640
 - Stata for 167, 184, 510, 511
 - strength of relationship 168, 176
 - zero value for 506–509
 - per-comparison error rate 413, 444
 - percent distribution table 27, 33
 - percentiles 46–51, 58
 - ranks 46–51
 - percents 28
 - perfect linear relationships 168
 - phi coefficient 176, 181–185
 - contingency tables 183
 - marginal counts 183
 - pie charts 31–32
 - planned comparison tests, Bonferroni adjustment 444–446
 - Point Biserial correlation coefficient 176–185, 216, 349
 - point estimation 291
 - population
 - bias and 285
 - estimators for 285, 291
 - infinite 274
 - parent populations 282
 - population correlation coefficient 507, 509–512
 - population standard deviation 275, 291, 293, 412
 - sampling of 269–270
 - standard error of prediction 513
 - target 393, 405–406
 - population mean estimation 291
 - post-estimation statistics 168
 - post-hoc multiple comparison tests, one-way analysis of variance 435
 - post-hoc multiple means test, two-way analysis of variance 479–481
 - post option 482
 - power, in hypothesis testing 307–312
 - power analysis 538–540, 613–614
 - multiple regression 580–582
 - power** command 448
 - pre-post design 353
 - predict** command 211
 - prediction model
 - predictions
 - accuracy of with linear regressions 209–210
 - standard error of estimate 513
 - standard error of prediction 513
 - predictor variable 512
 - prefix** command/s 361
 - preserve** command 108
 - pretest to posttest design 354, 399, 405, 406
 - probability
 - additive rules 231–233
 - complement rules 230
 - conditional 236–239
 - discrete 228–230
 - experiments 244–254
 - law of large numbers 240, 251, 253
 - total 239
 - probability models
 - binomial probability model 244–254, 260
 - binomial probability model applicability 249–254
 - normal probability model 254–261
 - theoretical probability model 244
 - "Probable error of the mean" (article, Gossett) 320

- propensity score matching 402
- putexcel** command 656
- pwcompare** command
- pwcorr** command 179
- pwmean** command 433
- quasi-experimental designs 395, 404
 - experimental mortality 406
 - external validity threats to 406–408
 - instrumentation in 405
 - internal validity threats to 404
 - maturity 405
 - selection bias in 405
- R 520–524, 558
 - effect size 520
 - effect size for simple regression 217
 - linear regression model and 210–217
 - values 511, 512
- R squared
 - adjusted 523
 - change
 - r* to *Z* relationship 509
 - statistical change 566–568
 - simple regression 520–522
 - statistical significance 522–523
 - true population 523
- random assignments, in causal studies 395–396
- random factors 479
- random number generator 271–272
 - obtaining a simple 273
 - with and without replacement 273–275
- random sampling 270–273
- range and interquartile range 89–91
- rank function 143
- ratio measurement of variables 6
- raw total SS 475
- RDD *see* regression discontinuity designs (RDD)
- recode** command 126
- recoding variables 144–147
- rectangular distributions 76
- redundancy 553, 558
- reflection 126
- regress** command 208, 219, 509, 515, 560, 563, 567, 569, 571
- regression coefficients 514, 515, 531
- regression diagnostics 524–529, 533–536
- regression discontinuity designs (RDD) 403
- regression lines 203
- regression models
 - Excel output 663–668
 - see also* linear regression; multiple regression
- regression to the mean 206
- regressn** command 606
- regressors
- relational operators 104
- relational studies 391–395
 - construct measures 391–395
- relative frequency 28
- relative frequency distributions 28, 182, 240, 254, 505
- residual option 211
- studentized 531
- reliability 391
- replace** command 145
- residuals
 - best-fitting linear equations 204
 - sum of squares due to 518, 519
 - variances for 418–419
- resistant statistics 89, 94
- restore** command 108
- return list** command 168, 429
- r(max)** return value 98
- r(min)** return value 98
- robvar** command 340, 346, 425, 438, 479, 494
- row marginal mean 460
- runiform function 273
- S curves 45
- sampling, random 270–273
 - independent and dependent samples 337–338
 - inferential statistics and 269–270
 - of populations 269–270
 - with and without replacement 273–275
- sampling distribution of means 275–280
 - central limit theorem 281–285
 - empirically 275–280
 - exercises 287–290, 319
 - theoretically 280–282
- sampling with replacement 234
- SAT Critical Reading score 159
- scatterplot varying color and pattern 222
- scatterplots 185, 222, 526
 - between two variables 160–171
 - direction 163
 - linear regression models and 217–218
 - regression line superimposed 508
 - residual 526
 - shape and 160–171
 - strength 163
- school effectiveness in the United States 11–14
- Schwartz, Gideon E. 613
- seed** option 362
- selection bias 405
- set seed** command 279, 283
- shortened form of commands 120
- sign test 640
- signal components 337
- significance level 300
- signtest recordwt** command 641
- simple effects 482
- simple experiment 228–230
 - defined 228
- simple linear regression 512–513
 - dichotomous independent variables and 214–217, 536
 - effect size and 217
 - exercises for interval or ratio variables 219–227
- simple slopes 600

INDEX

723

- Simpson's paradox 397
- skewness 95–99
 - calculating 96–97
 - defined 43
 - skewness statistic 95
 - Stata for 96–97, 98
 - summskew** 98, 99, 102, 344
 - summskewcell** 325, 466, 467
 - t*-distribution 333
- Skulls data set 683
- socioeconomic status 77
- sort cum** command 46
- Spearman rank correlation coefficient 174–176, 640
- spread 86–95, 133
 - range and interquartile range 89–91
 - selecting measures of 99
 - standard deviation 93–95
 - variance 91–93
- square root transformations 135
- standard deviation 93–95
 - see also *z*-scores
- standard error of estimate 514
- standard error of the mean 281
- standard error of mean differences 339
- standard error of prediction 514
- standard normal distribution 255
- standard scores see *z*-scores
- standardized regression equation 210
- Stata
 - anova** command 429, 486
 - arithmetic operators 104
 - as balanced option
 - bar charts for frequency and percent distributions of nominal data in 28
 - bar graph height sorting 30
 - binomial tail function 254
 - bootstrap** command 276, 278, 361, 429, 597
 - bootstrap estimation 517
 - boxplots 100–102
 - browse** command 19
 - by option 101, 102, 343
 - capture** command 278
 - cd** (change directory) command 278
 - centile** command 50
 - chi-square goodness-of-fit test
 - raw data 628
 - tabulated data 627
 - chi-square independence test 630, 635
 - chi2tail** function 625, 627
 - clear** command 108, 277
 - clustered bar graph** commands 184
 - codebook** command 108, 627
 - command line input 13
 - command line syntax 106
 - command options 107
 - command summary for Chapter 2 56–58
 - command summary for Chapter 3 104–105
 - command summary for Chapter 4 150
 - command summary for Chapter 5 189
 - command summary for Chapter 6 219
 - command summary for Chapter 8 260
 - command summary for Chapter 9 286–287
 - command summary for Chapter 10 313–314
 - command summary for Chapter 11 372–374
 - command summary for Chapter 13 450
 - command summary for Chapter 14 540–541
 - command summary for Chapter 16 582
 - command summary for Chapter 17 614–616
 - command summary for Chapter 18 649
 - Cook's distance 531
 - correlate** command 179, 211
 - csgof** command 627, 628
 - cumul** command 45
 - cumulative binomial probability value 252, 253
 - describe** command 108
 - detail** option 79, 424
 - discrepancy 531
 - display** command 96, 98, 168, 246–247, 252, 253, 256, 257, 259, 302, 331, 423
 - .do-files 147–150
 - doedit** command 147, 148
 - drop** command 108
 - edit** command 19
 - egen** command 34, 78, 129, 143, 166
 - egen mode** command 77
 - erase** command 278
 - estat bootstrap** command 429, 430
 - Excel and 656
 - exp** function for denoting exponential constant *e* 511
 - expand** command 628
 - Explore procedure 100
 - F*-distribution tables and Stata functions 420–422
 - finding *p*-value associated with a chi-squared goodness-of-fit test 627
 - for Fisher's exact test 635
 - forval** command 98
 - freq** option 46
 - frequency and percent distributions 27, 33
 - gen** or **generate** command 120, 146, 147, 355, 593
 - generate random** command 273
 - global k = r(max)** command 98
 - global m = r(max)** command 98
 - graph bar** command 30, 57
 - graph box** command 88, 101, 133
 - graph hbar** command 31
 - graph matrix** command
 - graph twoway** command 208
 - help** command 30, 41, 49, 52, 79, 107, 120
 - helpful hints 105
 - histogram** command 38–39, 41, 276, 278, 325
 - if** in command line 106
 - immediate** command 633
 - in** in the command line 106
 - independent samples *t*-test 373–374, 538
 - interface structure 2–12
 - jitter option 610
 - kdensity** command 41
 - kwallis** command 648

- Stata (cont.)
 - label define** command 126, 145, 195
 - label value** command 126, 145–146, 195
 - lcolor** option 41
 - legend** option 46, 575
 - Levene's test 426–431
 - leverage** 530–531
 - line graphs 40–42
 - list** command 78
 - log files 18–19
 - logging out 25
 - logical operators 104
 - lwidth** option 41, 42, 57
 - margins** command 481, 482, 600, 602, 606
 - marginsplot** command 601
 - marker option 53
 - mysm(Oh)** option 610
 - nestreg** command 595, 597
 - nodots** option 361, 364
 - obtain influence 531–533
 - one sample *t*-test commands 372
 - oneway** command 33, 423, 428, 429, 433, 438
 - online resources 106
 - opening data file 108
 - operators and expressions 104
 - over** commands 58–61
 - paired samples *t*-test commands 374
 - pcorr** command 565
 - Pearson product moment correlation 168, 184, 510, 511, 541
 - post-estimation statistics 168
 - post option 482
 - power** command 447, 538–540
 - predict** command 211, 532, 541
 - leverage 531, 541
 - predict predicted** command 222
 - predict residual** command 211, 219, 569
 - predict stres rstudent** command 531, 541
 - predict y1 predict x1orthx2resid**, residuals 527, 541
 - predict yhat** command 211, 493
 - predicted** command 536
 - predit yhat** command 516
 - prefix** command 361
 - preserve** command 108
 - product term 594
 - pwcompare** command 486
 - pwcorr** command 179, 511
 - pwmean** command 433
 - qnorm** command 325
 - rank function 143
 - recode** command 126
 - regress** command 208, 219, 509, 515, 560, 563, 567, 569, 571, 593
 - regression analysis 514–515
 - renaming variable in a bootstrap procedure 517
 - replace** command 144–147
 - restore** command 108
 - return list** command 168, 429
 - r(max)** 98
 - r(min)** 98
 - robvar** command 340, 346, 425, 479
 - runiform** function 273
 - save** command 19, 25, 147
 - seed** option 362, 364, 517
 - set seed** command 279, 283
 - significance testing 593–598
 - sigttest recordwt** command 641
 - skewness 96–97, 98
 - sort** command 90, 95
 - sort cum** command 46
 - stem pertak** command 35
 - studentized residual values 531
 - summarize** command 80, 82, 90, 93, 105, 106, 108, 207, 324, 342, 424
 - summarize detail** command 98
 - summarize prod** command 168
 - summarize ses** command 49, 58, 79, 81
 - summskew** command 98, 99, 102, 344
 - summskewcell** command 324, 466, 467
 - tab1** command 28
 - tabi** command 633
 - table** command 133, 338
 - tabstat** command 49, 50, 80, 81, 91, 105, 424
 - tabulate** command 28, 34, 77, 82, 131, 133, 184, 231, 233, 234, 236, 276, 428, 493, 632, 636
 - tabulated data 627
 - test** command
 - testing interaction for statistical significance 593–598
 - ttail** command 331
 - ttest** command 326, 332, 346, 364
 - twoway** command 41, 46, 160, 177, 179, 255, 541, 575
 - twoway line** command 41, 46
 - twoway scatter** command 507, 533
 - use** command 19, 149
 - variable creation 569
 - varname** command 99
 - xlabel** option 177
 - statistical significance
 - b*-weight for 563–564
 - of effect size 306–307
 - hierarchical multiple regression approach 595
 - of interactions 593–598
 - of *p*-values 345
 - of *R*-values 522–523
 - Statisticians.dta* 25
 - descriptive 3, 26
 - post-estimation statistics 168
 - purpose of 1–2
 - summary statistics 86, 98
 - statistics software 2
 - status quo 391–395
 - stem and leaf displays 35–38, 57
 - stem pertak** command 35
 - Stepping.dta* 499, 684
 - strength
 - in Pearson product moment correlation coefficient 168
 - in scatterplots 163
 - studentized range statistic 427

INDEX

725

- studentized residual value 531
- sum of squares due to residual 519
- studentized range distribution 431
- sum of squares due to the regression (SSREG) 519
- summarize** command 49, 58, 79, 80, 81, 82, 90, 93, 99, 106, 108, 168, 207, 324, 342, 656
- summary statistics 86, 98, 106, 121–123
 - in Excel 656
 - of temperature 90
- summskew** command 98, 99, 102, 105, 344
- summskewcell** command 324, 466, 467
 - installing 324
- symmetric distributions, 42
- symmetric index 506
- t*-statistic 320
- t*-test
 - independent samples 340–342
 - paired samples 353–360
 - protected 432
 - Type I error 413
 - see also* paired samples *t*-test; Stata, independent samples *t*-test
- tab1** command 28, 634
- table** command 133, 373
- tables and Stata functions 420–422
- tabstat** command 49, 50, 80, 81, 91, 105, 424
- tabulate** command 28, 34, 77, 82, 131, 133, 184, 231, 233, 234, 236, 276, 428, 493, 633, 635
- target 392–393
- target populations 393
 - alternative hypotheses and 320
 - bootstrap 363
 - degrees of freedom 320–321
 - t*-distribution 320, 340
 - z*-statistics 319
- TEMP.dta* 87
- test** command
 - theoretical probability model 244
- total probability 239
- transformations 118
- translation 125
- trimodal distributions 76
- true control groups 398–401
- true experiment 395–396
- ttest** command 326, 332, 346, 364
- Tukey, John 55, 431
- two-factor designs 398, 457–460
- two-sample designs 336–337
 - assumptions of 337–338
 - independent samples 353
 - Levene's test 346, 351
 - noise components in 337
 - repeated-measures design 353
 - signal components in 337
- two-way analysis of variance
 - assumptions 466–467
 - balanced and unbalanced factorial designs 467–468
 - column main effect test 465
 - computation by hand 469–474
 - crossed factorial designs 458
 - degrees of freedom 469
 - effect size 477
 - F*-test 469
 - fixed vs. random factors 479
 - grand mean 460, 475
 - hypothesis testing 465
 - interactions in 460–465
 - joint effects 475–476
 - line graph 461
 - modeling scores 475
 - partitioning total sum of squares 468–469
 - post-hoc multiple means test 479–481
 - randomized block designs 458
 - row main effect test 465
 - simple effects 482–484
 - two-factor design 457–460
- twoway** 609
- twoway** command 42, 46, 162, 177, 179, 255, 533, 575
- twoway line** command 42, 46
- Type I error 413
- unbalanced factorial designs 467–468
- unbiased estimators 285
- ungrouped frequency distribution 35, 47
- uniform distributions 76
- uniform random number generator 273
- unimodal distributions 77
- univariate distributions 26, 32
 - bimodal 75
 - data value occurrences 26
 - mean 80–84
 - median 78–80
 - mode in 74–78
 - range of 89–91
 - shape of 74
 - spread of 74, 86–95
 - unpredictability *see* spread
 - unrelated samples *see* independent samples
- unstandardized regression equations 208
- use** command 19, 149
- validity 391
 - external for quasi-experimental design 404, 405–406
 - internal for quasi-experimental design 404
- values
 - cumulative binomial probability value 252, 253
 - expected 400
- variability *see* spread
- variables
 - choosing scale of measurement of 6–7
 - combining variables 147
 - interval measurement of 5
 - linear composites 554
 - measurement 3–5
 - nominal measurement of 4, 26–32
 - ordinal measurement of 4

- variables (cont.)
 - ratio measurement of 6
 - recoding variables 144–147
 - redundant 553
 - variance 91–93
 - between group 416, 417
 - residual variance 426
 - test of means and 414–416
 - within group 416, 417
- varname** command 99
- Wages.dta* 613–614
- weighted mean 83–84
- Whitney, D. Ransom 639
- Wilcoxon sign test 3, 640
- Wilcoxon’s Rank-sum test 642
- within-group variance 416
- xlabel option 177
- Yates’ correction 628
- z-scores 127–128, 295, 300, 319
 - comparing different distributions and 133–134
 - outlier detection and 130–134
 - percentile ranks and 135
- z-test 299
- zero-order correlations 555