

Index

316L steel	source of profuse bleeding	question, 307–8
property conferred by the chromium	answer, 46–47	muscle incised during posterolateral
content	question, 39	approach
answer, 456	Achilles tendinopathy	answer, 347
question, 443	description of the reactive phase	question, 329
abdominal aorta	answer, 92–93	muscle overlying the tibia in a
vertebral level of bifurcation and	question, 85	posterolateral approach
termination	treatment for non-insertional	answer, 348
answer, 349	Achilles tendinopathy	question, 330
question, 330	answer, 94	nature of a painful lump
abductor muscles	question, 85	answer, 426
force required to maintain a single	Achilles tendon ruptures	question, 416
leg stance	features of	nature of a painless soft tissue lump
answer, 45	answer, 94	answer, 426
question, 38	question, 85	question, 416
above knee amputation	acid-fast bacilli in joint fluid	weakness and pain in a child
cause of gait problem	identification of the organism	answer, 315
answer, 236	answer, 377	question, 307
question, 226–29	question, 369	ankle arthrodesis
above knee amputee with a prosthesis	adductor hallucis	ideal position of the ankle
cause of a medial whip	nerve supply	answer, 102
answer, 437	answer, 91–92	question, 89
question, 432	question, 85	ankle arthroscopy
acetabular anteversion	adhesive wear	cause of persistent numbness
conditions associated with	description of	complication
answer, 50	answer, 380	answer, 100
question, 41	question, 370	question, 88
acetabular dysplasia in children	aging	ankle-brachial pressure index
features of	effects on articular cartilage	indication of vascular injury
answer, 300	answer, 402–3	answer, 220
question, 290	question, 397	question, 202
acetabular fracture	allograft bone	ankle foot orthosis
structure found during ilioinguinal	properties of	proper positioning in a 6-year-old
approach	answer, 362	patient with forefoot
answer, 345	question, 353	abduction
question, 328	anatomical snuffbox	answer, 437–38
acetabular protrusio	structure near to	question, 432
conditions not associated with	answer, 343	ankle fracture
answer, 45	question, 327	management to allow early weight
question, 38	Angoff method of standard setting,	bearing
definition from anteroposterior	18–19	answer, 231
radiograph	ankle	question, 223
answer, 50	location of neurovascular structures	plating for medial malleolus fixation
question, 41	during an anterior approach	answer, 236
acetabular screw placement	answer, 347	question, 229
description of quadrant	question, 329	ankle fracture-dislocation
answer, 47	management of septic arthritis in a	application of an external fixator
question, 39–40	child	answer, 221
	answer, 307–8	question, 204

ankle injury <ul style="list-style-type: none">cause of increased tibiofibular clear space<ul style="list-style-type: none">answer, 101question, 88non-specific pain and occasional swelling<ul style="list-style-type: none">answer, 101–2question, 88–89pronation external rotation injury<ul style="list-style-type: none">pattern<ul style="list-style-type: none">answer, 101question, 88worst position to sustain a pilon fracture<ul style="list-style-type: none">answer, 102question, 89	question, 135	arm injury after a fall <ul style="list-style-type: none">upper arm injury with difficulty moving wrist and fingers<ul style="list-style-type: none">answer, 216–17question, 197arm traction injury<ul style="list-style-type: none">signs of preganglionic injury<ul style="list-style-type: none">answer, 163–64question, 158–59
ankle instability <ul style="list-style-type: none">ligament that is most likely damaged<ul style="list-style-type: none">answer, 100–1question, 88	anterior cervical spine surgery <ul style="list-style-type: none">preparation for revision procedure<ul style="list-style-type: none">answer, 155question, 143anterior cruciate ligament reconstruction<ul style="list-style-type: none">cause of post-operative numbness of the foot<ul style="list-style-type: none">answer, 81question, 76cause of re-rupture<ul style="list-style-type: none">answer, 81question, 76exercises to avoid in the first 4 to 6 weeks<ul style="list-style-type: none">answer, 72question, 64graft option for elite sports player<ul style="list-style-type: none">answer, 69question, 62in a child<ul style="list-style-type: none">answer, 234–35question, 228in skeletally immature patients<ul style="list-style-type: none">answer, 72–73question, 65post-operative inability to fully extend the knee<ul style="list-style-type: none">answer, 81question, 76technique<ul style="list-style-type: none">answer, 72question, 65tunnel malposition<ul style="list-style-type: none">answer, 68–69question, 61–62	arthrititis of the hip <ul style="list-style-type: none">management of previously placed dynamic hip screw<ul style="list-style-type: none">answer, 57question, 44
ankle reflex <ul style="list-style-type: none">root value<ul style="list-style-type: none">answer, 146question, 136	in a child <ul style="list-style-type: none">answer, 234–35question, 228	articular cartilage <ul style="list-style-type: none">biomechanical changes in osteoarthritis<ul style="list-style-type: none">answer, 360question, 352changes in<ul style="list-style-type: none">answer, 358question, 351changes with age<ul style="list-style-type: none">answer, 358–59question, 351collagen composition<ul style="list-style-type: none">answer, 359question, 351function of<ul style="list-style-type: none">answer, 357question, 351structure of<ul style="list-style-type: none">answer, 357question, 351treatment options for defect in<ul style="list-style-type: none">answer, 358question, 351
ankle swellings <ul style="list-style-type: none">management in a child<ul style="list-style-type: none">answer, 314question, 306	anterior cruciate ligament sport injury <ul style="list-style-type: none">benefits of ACL reconstruction<ul style="list-style-type: none">answer, 71question, 64anterior cruciate ligament surgery<ul style="list-style-type: none">cause of intraoperative mistake<ul style="list-style-type: none">answer, 80–81question, 75–76anterior lumbar interbody fusion<ul style="list-style-type: none">procedure<ul style="list-style-type: none">source of significant haemorrhage<ul style="list-style-type: none">answer, 129question, 113	autologous chondrocyte implantation <ul style="list-style-type: none">use in knee cartilage defects<ul style="list-style-type: none">answer, 458question, 445
ankle twisting injury <ul style="list-style-type: none">Lauge-Hansen fracture classification<ul style="list-style-type: none">answer, 222question, 204–5	anterior cruciate ligament surgery <ul style="list-style-type: none">cause of intraoperative mistake<ul style="list-style-type: none">answer, 80–81question, 75–76	autosomal recessive B-glucocerebrosidase deficiency <ul style="list-style-type: none">physal zone affected by<ul style="list-style-type: none">answer, 393question, 389
ankylosing spondylitis <ul style="list-style-type: none">diagnostic criteria<ul style="list-style-type: none">answer, 153question, 141radiology<ul style="list-style-type: none">answer, 147question, 136–37	anterior cruciate ligament surgery <ul style="list-style-type: none">cause of intraoperative mistake<ul style="list-style-type: none">answer, 80–81question, 75–76	axillary artery <ul style="list-style-type: none">where it becomes the brachial artery<ul style="list-style-type: none">answer, 332question, 323
ankylosing spondylitis with kyphosis <ul style="list-style-type: none">pedicle subtraction osteotomy<ul style="list-style-type: none">answer, 147question, 136	anterior lumbar interbody fusion <ul style="list-style-type: none">procedure<ul style="list-style-type: none">source of significant haemorrhage<ul style="list-style-type: none">answer, 129question, 113	axonal injury <ul style="list-style-type: none">features of<ul style="list-style-type: none">answer, 459question, 445–46
annulus fibrosus <ul style="list-style-type: none">structure of<ul style="list-style-type: none">answer, 363question, 353–54	antibiotics <ul style="list-style-type: none">mechanism of hearing loss side-effect<ul style="list-style-type: none">answer, 402question, 397	back pain <ul style="list-style-type: none">action following MRI scan findings<ul style="list-style-type: none">answer, 132question, 119
antalgic gait pattern <ul style="list-style-type: none">features of<ul style="list-style-type: none">answer, 433question, 428	anticoagulant medication <ul style="list-style-type: none">mechanism of action<ul style="list-style-type: none">answer, 395question, 391	

Index

- back pain (cont.)
 and bilateral leg pain, treatment
 answer, 128
 question, 111
 and bilateral radiculopathy
 answer, 153–54
 question, 141–42
 and weight loss in a 9-year-old boy
 answer, 127
 question, 109
 cervicothoracic pain after a fall in a
 60-year-old man
 answer, 127
 question, 109
 histology of biopsy
 answer, 133
 question, 122
 investigation in diabetic pneumonia
 patient
 answer, 129
 question, 114–15
 ongoing pain with positive Mantoux
 test
 answer, 129–30
 question, 115
 paraspinal spasm in a 15-year-old
 boy
 answer, 126–27
 question, 109
 radiating into the foot
 answer, 459
 question, 445
 severe pain in patient with a history
 of metastatic prostate cancer
 answer, 133
 question, 123
 severe pain in patient with a history
 of renal cell carcinoma
 answer, 133
 question, 123
 severe thoracic pain and T8 lesion
 answer, 133
 question, 122
 treatment for chronic back pain
 answer, 147
 question, 137
See also low back pain; spine.
 back pain and gait disturbance
 with other findings in a diabetic
 patient
 answer, 154
 question, 142
 Baumann procedure
 features of
 answer, 297–98
 question, 288
 below knee amputation
 effect if the heel of the prosthesis is
 too rigid
 answer, 435
 question, 430
 prosthetic knee with variable
 cadence
 answer, 436
 question, 431
 terminal device suitable for running
 answer, 435
 question, 430
 biologically active particles
 size of
 answer, 52–53
 question, 42
 biomechanics
 classes of levers
 answer, 403
 question, 399
 biopsy
 aspects of use
 answer, 424
 question, 413–14
 bisphosphonates
 conditions used for
 answer, 422
 question, 410
 mechanism of action of nitrogen-
 containing formulations
 answer, 402
 question, 396
 blood coagulation test
 indication for blood product
 administration
 answer, 457–58
 question, 444
 blood transfusion
 patients who refuse on religious
 grounds
 answer, 470–71
 question, 464–65
 reduction of requirement in
 Jehovah's witness THA patient
 answer, 47–48
 question, 40
 Bloom's taxonomy of cognitive
 categories, 3–4, 10–12
 Blount's disease
 features of
 answer, 300–1
 question, 290
 bone
 blood supply
 answer, 360
 question, 352
 collagen structure
 answer, 363
 question, 353
 distinction between cortical and
 cancellous bone
 answer, 364
 question, 354
 rate dependency loading of cortical
 bone
 answer, 365–66
 question, 355
 bone banks
 processing of bone for
 answer, 403
 question, 399
 bone defect
 appropriate material to fill
 answer, 455
 question, 443
 bone graft
 mechanism of integration
 answer, 393–94
 question, 390
 strength of the graft over time
 answer, 403
 question, 398
 bone lesions
 radiolucent lesions with thin
 sclerotic rim and ground-glass
 matrix
 answer, 420
 question, 408
 bone morphogenetic protein (BMP)
 link to patient presentation
 answer, 395
 question, 391
 bone remodelling
 features of
 answer, 362
 question, 353
 bone screws
 way to increase the working length
 answer, 453–54
 question, 442
 ways to increase their pull-out
 strength
 answer, 214–21
 question, 195
 bone tumours
 classical appearances and
 locations
 answer, 418
 question, 406–7
 brachial plexus
 shoulder muscles innervated from
 answer, 333
 question, 323
 brachial plexus injury
 indicator of level of injury
 answer, 251
 question, 242
 level of injury suggested by trauma
 patient symptoms
 answer, 251
 question, 242

- management of later shoulder problems
 - answer, 251
 - question, 242
 - brachialgia
 - underlying pathology
 - answer, 129
 - question, 112–13
 - breast cancer
 - management of a metastatic femoral lesion
 - answer, 424–25
 - question, 414
 - Brodie’s abscess
 - features of
 - answer, 377
 - question, 368
 - bunion
 - appropriate surgical intervention
 - answer, 96
 - question, 86
 - diagnosis related to pain location
 - answer, 96–97
 - question, 86
 - burner nerve injury
 - unlikely cause of
 - answer, 151
 - question, 139
 - buttock
 - diagnosis of enlarging painless mass
 - answer, 418
 - question, 406
 - calcaneal squeeze test
 - diagnosis when pain occurs
 - answer, 93
 - question, 85
 - calcaneum
 - diagnosis when pain occurs on
 - palpation of the medial plantar aspect
 - answer, 93
 - question, 85
 - calcaneus
 - amputation that preserves and fuses it to the distal tibia
 - answer, 435
 - question, 430
 - calcified zone
 - feature of
 - answer, 459
 - question, 445
 - calcium
 - role in skeletal muscle contraction
 - answer, 403
 - question, 398
 - candidate guidance
 - applying to take the exam, 5
 - Bloom’s level 1, 2 and 3 questions, 3–4
 - build-up to the exam, 5
 - exam format, 1
 - example questions from the JCIE
 - website, 2–3
 - extended matching item (EMI)
 - questions, 1
 - general advice, 4–5
 - higher-order judgement, 3–4
 - higher-order thinking, 3–4
 - multilogical thinking, 4
 - overview of the FRCS (Tr & Orth)
 - exam, 1–3
 - Parts 1 and 2 of the FRCS (Tr & Orth) exam, 1
 - relevance of the FRCS (Tr & Orth)
 - exam, 3
 - resources for preparation and revision, 5–8
 - test centres, 1
 - the week of the exam, 8
 - See also* SBA writing process.
- cane
 - utilisation for ipsilateral hip arthritis
 - answer, 435
 - question, 430
 - carpal bones
 - sequence of ossification
 - answer, 254
 - question, 245–46
 - carpal tunnel
 - anatomy of
 - answer, 343
 - question, 327
 - carpal tunnel decompression
 - choice of local anaesthetic
 - answer, 278
 - question, 260
 - location of nerve
 - answer, 338
 - question, 325
 - prognostic indicator
 - answer, 252
 - question, 243
 - structure most at risk of injury
 - answer, 253
 - question, 244
 - cartilage
 - effects of aging on articular cartilage
 - answer, 402–3
 - question, 397
 - major component of articular cartilage
 - answer, 458
 - question, 445
 - cartilage defects in the knee
 - use of autologous chondrocyte implantation
 - answer, 458
 - question, 445
 - cartilaginous tumours
 - features of
 - answer, 419
 - question, 407
 - case-control study
 - feature of the study design
 - answer, 485
 - question, 474–79
 - cavovarus with positive Coleman block test
 - suitable orthotic for
 - answer, 435
 - question, 429
 - cement
 - antibiotic-loaded bone cement
 - answer, 384
 - question, 372
 - use of antibiotic-loaded bone cement
 - answer, 384
 - question, 372–73
 - cemented femoral stem fracture
 - risk factors
 - answer, 56–57
 - question, 44
 - ceramic
 - mechanical property that predisposes to fracture
 - answer, 50
 - question, 41
 - cerebral palsy
 - contraindication for selective dorsal rhizotomy
 - answer, 317–18
 - question, 310
 - features of the Baumann procedure
 - answer, 297–98
 - question, 288
 - gait compensatory mechanism
 - answer, 316
 - question, 308
 - hip conditions related to
 - answer, 302
 - question, 291
 - orthopaedic management
 - answer, 303
 - question, 291
 - procedures to help crouch gait and fixed knee deformity
 - answer, 317
 - question, 310
 - radiographic evaluation
 - answer, 302
 - question, 291
 - surgical management of CP hips
 - answer, 303
 - question, 291
 - total hip arthroplasty

Index

- cerebral palsy (cont.)
 - answer, 299
 - question, 289
- cervical disc replacement
 - indication for
 - answer, 154
 - question, 142
- cervical discectomy
 - muscles affected by injured nerve
 - answer, 349
 - question, 331
- cervical spine
 - structure encountered during
 - anterior approach
 - answer, 349
 - question, 330
- cervical spine fracture-dislocation
 - ASIA grade and neurological level of injury
 - answer, 130
 - question, 115
 - best management option
 - answer, 231
 - question, 223
- cervical spine injuries
 - use of halo traction
 - answer, 436
 - question, 431
- cervical spine surgery
 - left scapula winging after surgery
 - answer, 164
 - question, 159
- child having recurrent falls
 - indicated action to investigate
 - answer, 126
 - question, 107–8
- child with short neck, low hairline and reduced neck range of motion
 - nature of the condition
 - answer, 155
 - question, 143
- choke syndrome
 - likely cause with a below knee prosthesis
 - answer, 434
 - question, 429
- chondrosarcoma
 - features of
 - answer, 425
 - question, 415
- chondrosarcoma from the femur
 - treatment
 - answer, 424
 - question, 412
- Chopart amputation
 - deformity that would result in
 - answer, 434
 - question, 429
- chordoma
 - likely form of treatment
 - answer, 424
 - question, 413
- chromosomal breakage test
 - expected finding in a child with a positive test
 - answer, 458
 - question, 444–45
- clavicle
 - structure at surgical risk in right medial clavicle excision
 - answer, 166
 - question, 161
- clavicle fracture
 - indications for operative fixation
 - answer, 165
 - question, 161
 - Neer type 1 lateral third clavicle fracture
 - answer, 165
 - question, 161
 - nerve supply to muscle
 - answer, 342
 - question, 326
 - operative versus non-operative management
 - answer, 233–34
 - question, 226
- clavicle injury
 - operative versus non-operative treatment
 - answer, 218
 - question, 199
- clinical trials
 - assessment of relative risk
 - answer, 489
 - question, 479–80
 - combined versus single modality treatment
 - answer, 489
 - question, 479
 - comparison of patient groups
 - answer, 487
 - question, 476
 - comparison of two treatments
 - answer, 489
 - question, 479
 - control of patient variables
 - answer, 487
 - question, 476
 - correlation between variables
 - answer, 490
 - question, 480
 - data analysis
 - answer, 486
 - question, 476–80
 - forest plot interpretation
 - answer, 490–91
 - question, 481
 - graphical presentation of results
 - answer, 486
 - question, 475–76
 - intention to treat analysis
 - answer, 489–90
 - question, 480
 - interpretation of odds ratios
 - answer, 487
 - question, 477
 - interpretation of results
 - answer, 486
 - question, 475
 - minimising bias and confounding
 - answer, 487
 - question, 476
 - odds ratio comparison between groups
 - answer, 487–88
 - question, 477
 - power of the study
 - answer, 488–89
 - question, 478–79
 - procedural risk assessment
 - answer, 488
 - question, 477
 - sensitivity and specificity of screening tests
 - answer, 490
 - question, 481
 - statistical significance of results
 - answer, 486
 - question, 475
 - use of logistic regression
 - answer, 490
 - question, 480
 - use of multivariate linear regression
 - answer, 490
 - question, 480
- collagen
 - predominant type in healing tissue two weeks after ankle sprain
 - answer, 394
 - question, 391
 - predominant type in knee cartilage
 - answer, 394
 - question, 390
 - structure in bone
 - answer, 363
 - question, 353
 - type produced during recovery from knee soft-tissue stress injury
 - answer, 393
 - question, 389–90
- common femoral artery
 - surface anatomy within the femoral triangle
 - answer, 344
 - question, 328
- compartment concept

identifying an extracompartmental site answer, 427 question, 417	role of vitamin E added to answer, 448 question, 439	answer, 301–2 question, 291
compartment syndrome	cubital fossa	DEXA T-score and Z-score
dual incision fasciotomy of the leg answer, 234 question, 226–27	structure near to answer, 336 question, 324	use in diagnosis answer, 395 question, 391–92
confidence interval	cubital tunnel syndrome	diabetes
feature of answer, 484 question, 473	bracing position to assist relief of symptoms answer, 438 question, 432	considerations for surgical patient with answer, 385–86 question, 373–74
congenital scoliosis	data analysis	diabetes mellitus
appropriate management of a 9-month-old child answer, 125 question, 106	correct application of statistical tests answer, 485 question, 474	involvement in midfoot collapse answer, 103 question, 89
embryonic tissue involved in answer, 124–25 question, 106	data distribution	diabetes mellitus type 2
gene mutation associated with answer, 125 question, 106	interpretation of answer, 485–86 question, 474–75	management of midfoot collapse answer, 103 question, 89
congenital talipes equinovarus (CTEV)	debridement and exchange of the femoral head	diabetic foot ulcer
management in children answer, 294–95 question, 286–87	patient suitability for answer, 384–85 question, 373	negative predictor of wound healing answer, 103–4 question, 89
congenital thoracic scoliosis	deep peroneal nerve	predictor of associated osteomyelitis answer, 104 question, 89
aim of surgery answer, 125–26 question, 107	clinical signs of injury answer, 348 question, 329–30	diabetic patient
appropriate management of a 12-year-old child answer, 125 question, 106	degenerative listhesis	identifying unresponsive afferent nerve fibres answer, 394 question, 390
likely cause of progressive deformity answer, 124 question, 106	management answer, 155 question, 143–44	diabetic THA patient
likely outcome without treatment answer, 125 question, 107	degenerative listhesis at L45	reducing prosthetic joint infection risk answer, 48 question, 40
potential curve progression answer, 125 question, 107	likely presentation answer, 153 question, 141	diathermy
consent	degenerative lumbar discs	description of answer, 468 question, 462
assessing a young person’s capacity for consent answer, 470 question, 464	histological features answer, 395 question, 391	digital nerve transection
requirement when carrying out clinical trials answer, 469 question, 463–64	Denis Browne bar brace	first modality to return after repair answer, 393 question, 389
consent process	appropriate position of the feet answer, 435 question, 430	direct anterior approach (DAA) to the hip
GMC guidance for answer, 469 question, 463	developmental dysplasia of the hip (DDH)	vessel that is sacrificed answer, 52 question, 42
Cronbach’s alpha, 20	clicking hips in babies answer, 301 question, 290	disproportionate dwarfism in a child
cross-linked UHMWPE	ongoing management answer, 317–18 question, 309–10	likely cause of answer, 393 question, 389
	ongoing management of unilateral DDH answer, 322 question, 313	distal biceps rupture
	radiographic measurement	neurologic complication from 2-incision surgical approach answer, 176 question, 168
		distal biceps tendon repair

Index

- distal biceps tendon repair (cont.)
 - most likely subsequent neurological deficit
 - answer, 162
 - question, 158
- distal biceps tendon rupture
 - complication with 2-incision surgical repair
 - answer, 163
 - question, 158
 - strength loss with nonoperative management
 - answer, 163
 - question, 158
- Down's syndrome
 - hip conditions related to
 - answer, 303
 - question, 291
- duty of candour
 - disclosure of information relating to patient harm
 - answer, 470
 - question, 464
- Eichenholz Stage III midfoot collapse
 - suitable orthotic for a diabetic patient
 - answer, 437
 - question, 431–32
- elastic nailing
 - technique
 - answer, 222
 - question, 205
- elastic stable intramedullary nailing (ESIN)
 - principles
 - answer, 454
 - question, 442
- elastic TENS nails
 - principles of
 - answer, 214
 - question, 195
- elastin
 - loaded strain capability
 - answer, 402
 - question, 397
- elbow
 - carrying angle
 - answer, 186
 - question, 173
 - cause of nerve injury in supracondylar fracture repair
 - answer, 317
 - question, 309
 - effects of nerve transection during anterior approach
 - answer, 342–43
 - question, 326
 - nerve injury complication from supracondylar fracture repair
 - answer, 317
 - question, 309
 - tenderness following ciprofloxacin treatment for olecranon bursitis
 - answer, 175–76
 - question, 167
 - elbow dislocation
 - clicking elbow joint 6 months after surgery
 - answer, 166
 - question, 161
 - elbow fracture
 - football injury in a child
 - answer, 178–79
 - question, 169–70
 - ongoing problems 4 years later
 - answer, 181–82
 - question, 170–71
 - reconstruction using a double loop tendon graft technique
 - answer, 178
 - question, 168
 - elbow fracture-dislocation
 - management of pulseless white hand
 - answer, 219
 - question, 201
 - elbow fracture from a fall
 - treatment in an elderly patient
 - answer, 217
 - question, 197–98
 - elbow fracture in a child
 - features of
 - answer, 213–14
 - question, 194
 - elbow injury
 - recurrent popping, clicking, clunking and locking
 - answer, 182–83
 - question, 171
 - elbow injury from a fall
 - most likely deficit caused by
 - answer, 163
 - question, 158
 - technical objectives of surgery
 - answer, 209–10
 - question, 192
 - elbow injury from a motorbike accident
 - definitive treatment
 - answer, 217–18
 - question, 198
 - elbow injury in a child
 - consequences of management problems
 - answer, 178
 - question, 168–69
 - elbow injury recovery
 - improving DASH score and function
 - answer, 180–81
 - question, 170
 - elbow injury while lifting overhead weights
 - diagnosis
 - answer, 165
 - question, 160–61
 - elbow injury with no radial pulse
 - management in a 6-year-old child
 - answer, 165
 - question, 159–60
 - elbow lateral mass fracture
 - late-onset functional deficit in a child
 - answer, 163
 - question, 158
 - elbow mass
 - management of a lipomatous tumour
 - answer, 425
 - question, 414
 - elbow pain
 - chronic lateral pain in a tennis player
 - answer, 176–77
 - question, 168
 - in javelin athlete
 - answer, 183–84
 - question, 171–72
 - management of medial pain in a child
 - answer, 175
 - question, 167
 - structure involved in gym injury
 - answer, 176
 - question, 167–68
 - tennis elbow management
 - answer, 177
 - question, 168
 - elbow pain and sensitivity
 - considerations during surgical repair
 - answer, 183
 - question, 171–72
 - elbow pain with effusion and click
 - MRI findings
 - answer, 181
 - question, 170
 - elbow stiffness with soft tissue calcification
 - answer, 422–24
 - question, 410
 - enchondroma
 - histology in the centre of
 - answer, 402
 - question, 396
 - epidemiological testing
 - statistical interpretation of results
 - answer, 467
 - question, 461
 - epiphysis separation
 - findings of Salter and Harris

answer, 321–22 question, 312–13	reduction of a sub-trochanteric fracture answer, 234 question, 226	management of an asymptomatic bone lesion in a child answer, 298–99 question, 288–89
extended matching item (EMI) questions, 1	traction device as temporary splint for trauma patient answer, 437 question, 431	peri-prosthetic fracture answer, 35 question, 29
extracellular matrix effect of overall negative charge answer, 458 question, 445	femoral fracture in a child most common complication answer, 220 question, 202	potential complications of fracture surgery answer, 33–34 question, 28
Factor V Leiden mutation consideration for surgical patient with answer, 385 question, 373	femoral fracture through a lytic lesion management in a breast cancer patient answer, 426–27 question, 416	reason for expansion as patients get older answer, 456 question, 443
fall investigation of subsequent restriction of arm movement answer, 157 question, 157	femoral head stainless steel versus cobalt-chrome answer, 447 question, 439	risk of intraoperative periprosthetic femoral fracture (IOPFF) answer, 34 question, 28
fall in a child effects of neurological injury answer, 218 question, 200	femoral head collapse management answer, 34 question, 28	FGFR3 gene mutation likely cause of spinal stenosis in patient with answer, 457 question, 444
femoral acetabular impingement changes on hip MRI answer, 55–56 question, 43	femoral head materials testing of properties answer, 447 question, 439	part of the bone affected by answer, 402 question, 396
femoral and popliteal masses answer, 426 question, 415	femoral lesion in a 10-year-old boy features of answer, 425–26 question, 415	fibula tumour comprising small round blue cells answer, 424 question, 412
femoral B-cell lymphoma treatment answer, 426 question, 415	femoral nailing avoiding complications in antegrade nailing answer, 236 question, 229	finger accidental injection with a high-power grease gun answer, 276 question, 259
femoral circumflex vessels structure associated with answer, 54–55 question, 43	femoral neck fracture Garden 4 fracture answer, 35 question, 29–30	alternative to non-operative management for PIP joint osteoarthritis answer, 459 question, 446
femoral components amount of proximal bone loss due to stress shielding answer, 52–54 question, 42	hemiarthroplasty versus total hip arthroplasty answer, 234 question, 227	cause of little finger numbness answer, 128 question, 112
femoral fracture contraindication for using Gallow's traction in a child answer, 316 question, 308	femur arterial components of the trochanteric anastomosis answer, 345–46 question, 328	damage caused by a climbing injury answer, 248 question, 239
distal femoral physeal fracture in an adolescent answer, 294 question, 286	Dorr classification, 35	damage in Type 1 flexor tendon injury answer, 249 question, 240
fatigue failure of antegrade nailing answer, 231–34 question, 227	Gruen zones answer, 32 question, 27	definitive management of deformity in a patient with rheumatoid arthritis answer, 282–83 question, 261
nail entry point and fixation varus deformity answer, 235 question, 228	Hartofilakidis Type B answer, 37 question, 31	diagnosis of excised exophytic proximal phalangeal lesion

Index

- finger (cont.)
 answer, 418
 question, 406
 digital clawing with altered sensation
 answer, 283
 question, 261
 inability to flex after tendon injury
 answer, 233
 question, 225
 injury sustained while playing rugby
 answer, 207
 question, 189–90
 management of a closed fracture-
 subluxation
 answer, 268
 question, 257
 management of a metacarpal neck
 fracture with puncture wound
 answer, 378
 question, 369
 management of closed mallet
 injuries
 answer, 267–68
 question, 257
 management of flexion contractures
 answer, 272
 question, 258
 management of metacarpal neck
 fracture
 answer, 284
 question, 262
 management of painful football
 injury from two weeks earlier
 answer, 269–70
 question, 257–58
 management of painful,
 erythematous dog bite
 answer, 281–82
 question, 261
 movement difficulty along with
 swelling and bruising
 answer, 251
 question, 242
 painful snapping sensation after
 punch injury
 answer, 278–79
 question, 260
 post-operative management of flexor
 tendon repairs
 answer, 264
 question, 256
 splint for extensor tendon slip
 rupture
 answer, 436–37
 question, 431
 structures affected by flexion
 contracture
 answer, 276
 question, 259
 testing before surgery for
 benign fibro-proliferative
 disorder of the palmar
 fascia
 answer, 248
 question, 238
 tingling and numbness in body
 builder
 answer, 184
 question, 158–59
 treatment of partial amputation
 answer, 252–53
 question, 243–44
 zone of extensor tendon injury
 answer, 254
 question, 245
 finger pain at PIP joint
 management to optimise joint
 function
 answer, 255
 question, 246
 fingertip accidental amputation
 reconstruction technique
 answer, 278
 question, 260
 fingertip pain
 diagnosis of mass lesion
 answer, 253
 question, 244
 fingertip saw injury
 management of
 answer, 249–50
 question, 240
 first metatarsal phalangeal joint
 treatment of effects of long-standing
 sport injury
 answer, 99
 question, 87
 first metatarsal phalangeal joint
 arthroplasty
 treatment of later osteolysis and
 valgus deformity
 answer, 98
 question, 87
 fixed flat foot deformity
 forefoot deformity associated with
 answer, 96
 question, 86
 flat feet
 indicator of congenital vertical talus
 answer, 320
 question, 312
 flexor digitorum longus
 structure immediately dorsal or deep
 to it
 answer, 95
 question, 86
 flexor digitorum profundus
 consequences of avulsion
 answer, 233
 question, 225
 flexor tendon injury
 flexor zone of the injury
 answer, 339
 question, 325
 flexor tendon pulleys
 preservation to avoid bowstringing
 and weakness
 answer, 343
 question, 327
 flexor tendon repair
 Kleinert protocol
 answer, 435
 question, 430
 flexor zone III of the hand
 structure and its nerve supply
 answer, 341
 question, 325–26
 floating toe iatrogenic complication
 procedure associated with
 answer, 98
 question, 86–87
 fluid-film lubrication
 hip prosthesis design that will allow
 it
 answer, 451–52
 question, 441
 foot
 anatomical structure in the 3rd layer
 answer, 90
 question, 84
 diagnosis of a painful plantar lesion
 answer, 422
 question, 410
 injury from a fall from a height
 answer, 207–8
 question, 190
 management of cavo varus in a
 neonate
 answer, 316
 question, 309
 management of midfoot collapse
 answer, 103
 question, 89
 most common malignancy
 answer, 102–3
 question, 89
 muscle that plantar flexes the 1st
 metatarsal
 answer, 90
 question, 84
 procedure to benefit deformity from
 PMP22 gene duplication
 answer, 319
 question, 311
 treatment for rigid plantarflexion
 deformity of 40°
 answer, 319

- question, 311
- underlying process involved in
 - unilateral midfoot collapse
- answer, 103
- question, 89
- foot lesion
 - biopsy diagnosis
- answer, 96
- question, 86
- foot pathology
 - in-shoe orthotic associated with
- answer, 95–96
- question, 86
- foot preposition
 - muscle that enables
- answer, 393
- question, 389
- foot prosthesis
 - effect of placing too far inset
- answer, 433
- question, 428
- forearm
 - complication following Henry’s
- approach
- answer, 340
- question, 324
- description of anatomy
- answer, 254
- question, 245
- extensor compartment of ruptured tendon
- answer, 340–41
- question, 325
- investigation of an enlarging mass on the volar aspect
- answer, 272–73
- question, 258
- muscles of the posterior deep compartment
- answer, 339–40
- question, 325
- forearm and wrist pain and heaviness
- investigation
- answer, 247–48
- question, 238
- forearm fracture
 - approach for surgical repair
- answer, 231
- question, 224
- retention of forearm plates in children
- answer, 297
- question, 288
- forearm fracture-dislocation
 - management of recurrent dislocation
- answer, 234
- question, 227
- forearm muscles
- nerve roots
- answer, 336
- question, 324
- superficial flexors
- answer, 336–37
- question, 324
- fracture fixation
 - design of an ideal plate
- answer, 456
- question, 443
- fracture fixation plate
 - working length
- answer, 453
- question, 442
- fractures
 - element of a locking plate construct that reduces axial forces
- answer, 394
- question, 390
- healing by a high strain fracture environment
- answer, 209
- question, 191
- friction between two surfaces
 - features of
- answer, 457
- question, 444
- gait
 - fastest gait pattern
- answer, 434
- question, 428
- nature of a vaulting gait
- answer, 437
- question, 432
- sequence of events in a 3-point gait
 - utilising a walker
- answer, 434
- question, 429
- gait disturbances
 - patient with metastatic prostate cancer
- answer, 150
- question, 139
- gene mutation
 - associated with hip deformity and café au lait spots
- answer, 394
- question, 391
- General Medical Council (GMC), 1
- genetic defect
 - identification from patient
- presentation
- answer, 395
- question, 391
- genu valgum
 - treatment options in a child
- answer, 316
- question, 308
- genu varum
- other findings in an 8-year-old boy
- answer, 317
- question, 309
- gluteal region
 - features of
- answer, 349
- question, 330
- graphical presentation of results
 - interpretation
- answer, 488
- question, 477–78
- great toe arthrodesis
 - nerve encountered on dorsomedial approach
- answer, 348
- question, 330
- greater sciatic foramen
 - structure which exits it and does not re-enter the lesser sciatic foramen
- answer, 48
- question, 40
- groin and knee pain in a child
 - predisposing factors
- answer, 303
- question, 292
- groin pain in endurance athlete
 - diagnosis and management
- answer, 50
- question, 41
- growth deficit in an infant
 - blood test finding
- answer, 362
- question, 353
- growth plates
 - order of closure
- answer, 320–21
- question, 312
- Gruen zones
 - answer, 32
 - question, 27
- haemophilia
 - description of the two best known types
- answer, 404
- question, 399
- inheritance pattern
- answer, 387
- question, 374
- haemophilia A
 - clotting factor affected
- answer, 387
- question, 374
- hallux sesamoids
 - structure attached to the plantar surface
- answer, 91
- question, 84

Index

- hallux
 - iatrogenic complication of excision of sesamoids
 - answer, 86
 - question, 86
- hallux valgus deformity
 - force created by the abductor hallucis on the proximal phalanx
 - answer, 98
 - question, 87
- halo orthosis
 - checks before application
 - answer, 154
 - question, 143
- halo orthosis application
 - complications
 - answer, 155
 - question, 143
- hand
 - altered sensation in digits
 - answer, 254
 - question, 245
 - annular pulley identification
 - answer, 341
 - question, 326
 - condition associated with pain and difficulty straightening digits
 - answer, 248
 - question, 239
 - diagnosis of altered sensation in digits and thenar eminence
 - answer, 283–84
 - question, 261–62
 - finger swelling and multiple small bone osteolytic lesions
 - answer, 427
 - question, 416–17
 - immunisation consideration in wound management
 - answer, 378
 - question, 369
 - management of an asymptomatic lytic lesion
 - answer, 250
 - question, 241
 - metacarpal fracture through a lesion
 - answer, 423
 - question, 412
 - muscle assessment and evidence of weakness
 - answer, 344
 - question, 327
 - numbness and progressive loss of manual dexterity
 - answer, 183
 - question, 171
 - palmar nodule in line with the little finger
 - answer, 273
 - question, 258
 - tender palmar lump at the base of the ring finger
 - answer, 248
 - question, 239
 - hand deformity in a child
 - explanation of the cause
 - answer, 247
 - question, 238
 - hand injury
 - antibiotic treatment following flexor sheath washout
 - answer, 247
 - question, 237
 - assessment and management in elderly patient with dementia
 - answer, 247
 - question, 237
 - cause of problem after flexor digitorum profundus repair
 - answer, 252
 - question, 243
 - damage caused by a fall on an outstretched hand
 - answer, 249
 - question, 239–40
 - indications for emergent surgical treatment of a puncture wound
 - answer, 247
 - question, 237
 - infectious organism from a cat bite
 - answer, 254
 - question, 245
 - initial management of a perilunate dislocation in a trauma patient
 - answer, 247
 - question, 237–38
 - management of deep tissue damage over the thenar eminence
 - answer, 277
 - question, 260
 - management of multiple metacarpal fractures
 - answer, 251–52
 - question, 242
 - management of multiple trauma-related fractures
 - answer, 284–85
 - question, 262
 - management of pain and swelling from a puncture wound
 - answer, 248
 - question, 238–39
 - management of painful PIPJ and intra-articular fracture
 - answer, 252
 - question, 243
 - management of palm injury
 - answer, 253
 - question, 244
 - management of palmar knife laceration
 - answer, 263
 - question, 256
 - pain and swelling following a puncture wound
 - answer, 247
 - question, 238
 - structure damaged by a fall on an outstretched hand
 - answer, 249
 - question, 240
 - tendon injury from palm laceration
 - answer, 249
 - question, 240
 - hand pain and grip weakness
 - investigation in a 76-year-old female patient
 - answer, 252
 - question, 243
 - hand surgery
 - post-operative rehabilitation regimen
 - answer, 250
 - question, 241
 - hand tumours
 - features of
 - answer, 427
 - question, 417
 - hand weakness and altered sensation
 - likely site of nerve pathology
 - answer, 253
 - question, 244
 - hand with altered sensation
 - findings from nerve conduction testing
 - answer, 247
 - question, 238
 - hands
 - decreasing hand function in patient with rheumatoid arthritis
 - answer, 129
 - question, 113–14
 - findings with morning stiffness and polyarthropathy
 - answer, 277
 - question, 260
 - hands with painful, swollen and deformed DIP joints
 - x-ray findings
 - answer, 254
 - question, 245
 - hangman's fracture
 - feature of
 - answer, 150–51
 - question, 139

- Hardinge approach to the hip
 - safe length of split of the gluteus medius
 - answer, 51–52
 - question, 42
- heel
 - non-trauma pain, swelling and erythema in a child
 - answer, 319
 - question, 311
- heel strike phase of the gait cycle
 - muscle that contracts eccentrically
 - answer, 91
 - question, 84
- heterotopic ossification
 - prophylaxis
 - answer, 386
 - question, 374
 - risk factors
 - answer, 386
 - question, 374
 - treatment planning
 - answer, 386–87
 - question, 374
- higher-order judgement, 3–4
- higher-order thinking, 3–4, 12, 427
- highly cross-linked polyethylene (HCLPE)
 - mechanical properties
 - answer, 57
 - question, 44
- hip
 - clicking hips in babies
 - answer, 301
 - question, 290
 - conditions related to cerebral palsy
 - answer, 302
 - question, 291
 - conditions related to Down’s syndrome
 - answer, 303
 - question, 291
 - diagnosis of non-specific bilateral hip problem
 - answer, 365
 - question, 354
 - features of windblown hips
 - answer, 300
 - question, 290
 - major structure at risk during posterior approach
 - answer, 346
 - question, 329
 - management of bilateral hip pain in a 7-year-old child
 - answer, 319
 - question, 310–11
 - management of hip clicking in a 3-month-old baby
 - answer, 315
 - question, 308
 - muscle exposed during posterior approach
 - answer, 345
 - question, 328
 - muscle retraction during anterior approach
 - answer, 344–45
 - question, 328
 - retroverted acetabular component
 - answer, 32–33
 - question, 27
 - Waldenstrom stages of Legg–Calve–Perthes disease
 - answer, 317
 - question, 310
- hip and groin pain in late pregnancy
 - answer, 33
 - question, 27
- hip angle identification
 - answer, 36
 - question, 30
- hip arthrodesis
 - optimal position
 - answer, 45
 - question, 38
 - risk of ipsilateral knee pain
 - answer, 45
 - question, 38
- hip arthroplasty
 - follow-up
 - answer, 33
 - question, 28
 - injury during
 - answer, 35
 - question, 29
- hip dislocation
 - risk factors
 - answer, 57
 - question, 44
- hip dysplasia
 - Hartofilakidis Type B
 - answer, 37
 - question, 31
 - hip dysplasia (Crowe type IV)
 - leg length correction in THA
 - answer, 57
 - question, 44
- hip fracture
 - features of fractures in children
 - answer, 299
 - question, 289
 - management
 - answer, 37
 - question, 30
 - management of fractures in children
 - answer, 296–97
 - question, 290
- hip fracture fixation
 - management of post-operative complications
 - answer, 385
 - question, 373
- hip fracture in an elderly person
 - muscles that produce the deformity of procurvatum and varus
 - answer, 219
 - question, 201
- hip fusion position
 - answer, 35
 - question, 29
- hip impingement
 - diagnosis
 - answer, 37
 - question, 31
 - testing
 - answer, 32
 - question, 27
- hip injury from a fall
 - appropriate treatment
 - answer, 220
 - question, 202–3
 - management considerations
 - answer, 218–19
 - question, 200–1
- hip injury from a motor vehicle accident
 - management stages
 - answer, 218
 - question, 200
- hip joint capsule
 - features of
 - answer, 55
 - question, 43
- hip joint pathology
 - nerve impingement mimic of
 - answer, 55
 - question, 43
- hip osteonecrosis
 - cause of
 - answer, 382–83
 - question, 371
 - management
 - answer, 382
 - question, 371
 - staging according to the Ficat classification
 - answer, 382
 - question, 371
- hip pain
 - diagnosis
 - answer, 45
 - question, 38
 - ‘double-line sign’ on MRI scan
 - answer, 381
 - question, 371

Index

- hip pain (cont.)
 - history of pain and epiphyseal lytic lesion
 - answer, 426
 - question, 415–16
 - management in a patient with polymyalgia rheumatica
 - answer, 381–82
 - question, 371
 - test for contracture
 - answer, 36
 - question, 30
- hip pain associated with osteonecrosis
 - management
 - answer, 52
 - question, 42
- hip pain in a child
 - management of
 - answer, 377
 - question, 369
- hip replacement
 - location of vulnerable nerve
 - answer, 34
 - question, 28
 - with femoral nailing
 - answer, 33
 - question, 27–28
- hip resurfacing
 - follow-up
 - answer, 33
 - question, 28
 - imaging to detect heterotopic ossification
 - answer, 49
 - question, 40
 - patient suitability
 - answer, 33
 - question, 27–28
- hip scoring systems
 - answer, 32
 - question, 27
- hip screw placement
 - vulnerable structures
 - answer, 34
 - question, 28
- hip surgery
 - detection of nerve damage
 - answer, 33
 - question, 28
- humeral fracture
 - classification of lateral condylar fractures in children
 - answer, 295–96
 - question, 287
 - complications of lateral condylar fractures in children
 - answer, 297
 - question, 287–88
 - features of lateral condylar fractures in children
 - answer, 295
 - question, 287
 - management of lateral condylar fractures in children
 - answer, 296–97
 - question, 287
 - open reduction of a lateral condylar fracture in a child
 - answer, 314–15
 - question, 307
 - prediction of humeral head ischaemia
 - answer, 206–7
 - question, 189
 - surgical procedures for humeral nailing
 - answer, 165
 - question, 161
- humeral fracture complications
 - use of tendon transfer
 - answer, 250
 - question, 241
- humeral fracture from a fall
 - most appropriate treatment
 - answer, 218
 - question, 198–99
- humeral fracture in a child
 - management approach for best outcome
 - answer, 231–32
 - question, 224
- humeral lytic lesion and associated pathological fracture
 - management of
 - answer, 419
 - question, 407
- humeral lytic lesion with cortical erosion
 - likely diagnosis
 - answer, 419
 - question, 407
- humeral shaft
 - vessel at risk during proximal deep dissection
 - answer, 342
 - question, 326
- humeral shaft fracture
 - posterior approach and axillary nerve injury
 - answer, 235
 - question, 228–29
- hydroxyapatite-coated implants
 - factors affecting bony integration
 - answer, 455
 - question, 443
- idiopathic congenital clubfoot
 - consequences of non-operative management in a 4-week-old infant
 - answer, 316
 - question, 308–9
- idiopathic transient osteoporosis of the hip (ITOH)
 - answer, 33
 - question, 27
- iliopsoas impingement syndrome
 - answer, 32–33
 - question, 27
- imaging
 - pseudotumour associated with metal-on-metal hip resurfacing
 - answer, 46
 - question, 39
- implant removal in children
 - answer, 305
 - question, 293
- implant with perfect hydrodynamic lubrication
 - concern about longevity
 - answer, 455–56
 - question, 443–44
- implants
 - greatest cause of wear leading to loosening
 - answer, 379
 - question, 370
- infant with bilateral deformities of 4th and 5th toes
 - likely cause
 - answer, 100
 - question, 88
- infection reduction in arthroplasty surgery
 - answer, 383–84
 - question, 372
- infectious microorganisms
 - characteristics of
 - answer, 378
 - question, 369
- infraclavicular area
 - neurological consequences of a penetrating injury
 - answer, 164
 - question, 159
- infraclavicular fossa
 - recurrent painful mass
 - answer, 421
 - question, 409
- Injury Severity Score (ISS)
 - trauma patient (motorcycle accident)
 - answer, 222
 - question, 204
- Intercollegiate Examination in Trauma and Orthopaedic Surgery, 1
- intervertebral discs

age-related change in answer, 151 question, 140	answer, 388 question, 375	ligament injury diagnosis answer, 69 question, 62
degenerative process answer, 360 question, 352	structure found during posterior approach answer, 346 question, 329	knee jerk reflex primary afferent fibres which mediate answer, 394 question, 390
description of a normal disc answer, 404 question, 400	structures of the posteromedial corner answer, 79–81 question, 74	knee ligaments cause of a positive Pivot shift test answer, 67 question, 60
feature of answer, 151 question, 140	treatment options for genu valgum in a child answer, 316 question, 308	knee pain diagnosis of a lytic lesion within the metaphysis answer, 423 question, 411
function of the annulus fibrosus answer, 360 question, 352	validated outcome measures answer, 79 question, 74	distal femoral physeal fracture in an adolescent answer, 294 question, 286
structure of the annulus fibrosus answer, 361 question, 352	knee and ankle pain and swelling with a previous rash, in a child answer, 315 question, 307	imaging diagnosis of lateral knee pain answer, 80 question, 75
tissue collagen type answer, 124 question, 105	knee arthroplasty contraindications for medial (partial) unicompartmental arthroplasty answer, 68 question, 61	lateral femoral condyle lesion answer, 70–71 question, 63
intra-articular metaphysis joints that do not have answer, 376 question, 368	liner dislocation with unicompartmental knee arthroplasty answer, 70 question, 62	liner dislocation with unicompartmental knee arthroplasty answer, 70 question, 62
intramedullary nails torsional rigidity of different sizes of nail answer, 394 question, 390	pain and flexion instability after revision procedure answer, 72 question, 64	spontaneous onset of medial pain answer, 67 question, 60
Jehovah’s witness patient reduction of blood transfusion requirement in THA answer, 47–48 question, 40	knee arthroscopy continuing knee pain following answer, 79–80 question, 75	knee pain and swelling lytic lesion within the tibial plateau answer, 420 question, 408
Joint Committee on Intercollegiate Examinations (JCIE), 1	repair of portal placement complication answer, 80 question, 75	knee pain and swelling following minor trauma blood-stained aspirated joint fluid answer, 421 question, 409
Judet views assessment of pelvic fracture healing answer, 236 question, 230	knee dislocation ankle-brachial pressure index indication of vascular injury answer, 220 question, 201–2	knee pain associated with bony growth management of answer, 422–23 question, 411
knee features of the meniscomfemoral ligaments answer, 364–65 question, 354	sequence of management answer, 233 question, 226	knee pain of 6 months’ duration management of a tibial lytic lesion answer, 423 question, 411
management of pain and swelling after a fall in a child answer, 319 question, 311	treatment of multi-ligament injuries answer, 222 question, 204	knee pain with bone oedema likely diagnosis answer, 381 question, 371
posterolateral corner reconstruction answer, 35, 79 question, 74	knee injury cell type that facilitates regeneration answer, 395 question, 392	knee pain with swelling, stiffness and locking diagnosis
posteromedial approach answer, 79 question, 74	knee injury from motorbike accident	
source of osteochondral lesion		

Index

- knee pain with swelling, stiffness and locking (cont.)
 - answer, 421–22
 - question, 410
- knee pain, swelling and locking
 - management of patient with chronic history
 - answer, 83
 - question, 78
- knee pathology radiograph
 - diagnosis
 - answer, 387–88
 - question, 374–75
- knee prosthesis
 - features of the polycentric knee
 - answer, 434
 - question, 428–29
 - suitable type for paediatric patients
 - answer, 433
 - question, 428
- knee replacement
 - cause of later patellar crepitus and jumping of the patella
 - answer, 66
 - question, 59
 - femoral component sagittal measurement between sizes 4 and 5
 - answer, 66
 - question, 59
 - patellar clunk syndrome
 - answer, 66
 - question, 59
 - primary role of the femoral and tibial stems
 - answer, 450
 - question, 440
 - procedure with increased risk of periprosthetic fracture
 - answer, 82
 - question, 77
- knee replacement prosthesis
 - use of all-polyethylene tibia component
 - answer, 68
 - question, 61
- knee snapping and locking
 - management
 - answer, 80
 - question, 75
- knee swelling
 - management of patient with acute non-trauma swelling
 - answer, 83
 - question, 78
- knee swelling and intermittent locking
 - diagnosis
 - answer, 79
 - question, 74–75
- knees
 - recurrent swelling in a young male
 - answer, 387
 - question, 374
- Kocher’s criteria for septic arthritis of the hip
 - answer, 376
 - question, 368
- leg
 - contents of the anterior compartment
 - answer, 213
 - question, 194
 - leg compartment
 - muscle associated with, 329
 - answer, 347
 - leg compartments
 - muscular origin of innervation
 - answer, 346–47
 - question, 329
 - leg fasciotomy
 - structures at risk in the posterior compartments
 - answer, 346
 - question, 329
 - leg length discrepancy
 - management in a 5-year-old child
 - answer, 318–19
 - question, 310
 - leg pain
 - treatment for ongoing lateral pain
 - answer, 129
 - question, 113
 - leg weakness
 - patient with a history of breast cancer
 - answer, 133
 - question, 122–23
- Legg–Calvé–Perthes disease
 - Waldenstrom stages
 - answer, 317
 - question, 310
- levels of evidence
 - answer, 467
 - question, 461
- ligaments
 - comparative strength of
 - answer, 51–53
 - question, 41
 - stress strain curve compared with tendons
 - answer, 365
 - question, 355
- limb length discrepancy (LLD)
 - arithmetical method for determining
 - answer, 304
 - question, 292
- Lisfranc injuries
 - radiology for
 - answer, 208
 - question, 191
- Lisfranc joint anatomy
 - answer, 208
 - question, 191
- Lisfranc ligament
 - anatomical description
 - answer, 90–91
 - question, 84
- Lisfranc ligament injuries
 - features of
 - answer, 94–95
 - question, 85
- local anaesthetic
 - mechanism of action on nerves
 - answer, 402
 - question, 396
- locking bolts
 - purpose when stabilising a femoral fracture
 - answer, 447
 - question, 439
- low back pain
 - and fever in a child
 - answer, 146
 - question, 135
 - and MRI lesion in patient with history of breast cancer
 - answer, 133
 - question, 121–22
 - and pain radiating down the leg
 - answer, 127–28
 - question, 109
- disc prolapse and inability to micturate
 - answer, 130
 - question, 115
- leg symptoms associated with
 - answer, 126
 - question, 108
- management of lytic spondylolisthesis
 - answer, 126
 - question, 109
- management of patient with neurogenic claudication
 - answer, 155
 - question, 143–44
- management of patient with neurogenic claudication and gait disturbance
 - answer, 145–46
 - question, 135
- vertebral step in lytic spondylolisthesis
 - answer, 126–27
 - question, 108–9
- with small lumbar scoliosis

answer, 132–33	question, 109	answer, 491
question, 121	step between lumbar vertebrae	question, 482
low molecular weight heparin	answer, 126–27	metal
thromboprophylactic effect	question, 108–9	features of the S-N curve
answer, 404	symptoms of	answer, 455
question, 400	answer, 126	question, 442–43
Lower Extremity Assessment Project	question, 108	methods of hardening
(LEAP)		answer, 456
findings	malignant transformation	question, 443–44
answer, 222	condition with the highest risk of	property displayed with increasing
question, 205	answer, 426	cyclical loading
lower limb amputation	question, 416	answer, 449–50
increase in metabolic demand	conditions with the potential to	question, 440
answer, 433	undergo	metal hypersensitivity to orthopaedic
question, 428	answer, 423	implants
lower-order thinking, 12	question, 411	answer, 45
lubrication	McKellop’s modes of wear	question, 38
types of	polythene wear in a hip joint	metal-on-metal hip resurfacing
answer, 457	answer, 450	imaging pseudotumour associated
question, 444	question, 440–41	with
lumbar canal stenosis	medial approach to the hip	answer, 46
classical finding	intermuscular plane	question, 39
answer, 153	answer, 54	metal-on-polythene hip joint surface
question, 141	question, 42	mechanism of wear leading to
lumbar compression fractures	medial patellofemoral ligament	osteolysis
diagnosis based on T-scores and Z-	(MPFL) reconstruction	answer, 449
scores	answer, 70	question, 440
answer, 395	question, 62	metastasis
question, 392	median nerve	conditions with the potential to
features of	muscle weakness caused by	undergo
answer, 149	damage to	answer, 423
question, 138	answer, 338–39	question, 411
lumbar interlaminar region	question, 325	metatarsal fracture
compressive soft tissue structure	meniscal repair techniques	appropriate treatment
answer, 124	for anterior horn meniscal tear	answer, 97
question, 105	answer, 70	question, 86
lumbar lordosis	question, 63	microdiscectomy (conventional
feature of	meniscal transplantation	discectomy)
answer, 148	contraindications for	comparison with tubular discectomy
question, 137	answer, 69–70	answer, 156
lumbar myelomeningocele	question, 62	question, 144
management for increasing leg pain	meniscus	Miller’s pyramid of levels of
and progressive foot deformity	changes following twisting knee	assessment, 10
answer, 319–20	injury	monoclonal antibody medication
question, 311–12	answer, 80	mechanism of action
lumbar spine	question, 75	answer, 133
signs and symptoms of nerve injury	effects of medial meniscectomy	question, 122
during anterior retroperitoneal	answer, 366–67	Monteggia fracture-dislocation
approach	question, 356	management of recurrent dislocation
answer, 350	most important function	answer, 234
question, 331	answer, 366	question, 227
lumbar spine flexion distraction injury	question, 355	MRI scan
in a trauma patient	meniscus tear patterns	scanning an MRI safe ex-fix on a
appropriate fixation method	prognosis for meniscal	knee
answer, 460	repair	answer, 456
question, 446	answer, 70	question, 443
lytic spondylolisthesis	question, 63	MRI with contrast enhancement
ongoing management of	meta-analysis from a systematic review	mode of action of gadolinium
answer, 126	of studies	contrast agent

Index

MRI with contrast enhancement (cont.) answer, 402 question, 397	management of an elderly patient answer, 215–16 question, 196	high-specificity injuries answer, 212 question, 193
MRSA infection protein responsible for antibiotic resistance answer, 394 question, 391	neck pain and gait disturbance management answer, 154 question, 142–43	non-ossifying fibroma conditions associated with answer, 422 question, 410
multilogical thinking, 4, 12	needlestick injury with HIV patient risk of transmission answer, 385 question, 373	normal distribution feature of answer, 484 question, 473
multiple myeloma investigation answer, 424 question, 413	nerve conduction studies findings from answer, 403 question, 398	occipitocervical dissociation (OCD) indicators of answer, 130 question, 115–16
muscle innervation brachial plexus root answer, 333–34 question, 323	nerve damage from hip surgery detection answer, 33 question, 28	odontoid fracture anterior pin placement for a Halo device answer, 210 question, 192
matching muscle and nerve answer, 54 question, 43	nerve fibres main determinant of healing after surgical repair answer, 394 question, 390	olecranon fracture surgical repair muscle function at risk answer, 179 question, 170
myelopathy causes of answer, 149–50 question, 139	nerve impingement mimic of hip joint pathology answer, 55 question, 43	Orthobullets website, 7–8
myxofibrosarcoma treatment of a high-grade tumour answer, 425 question, 414	nerve injury from posterior hip dislocation root value of the nerve answer, 393 question, 389	osteoarthritis biomechanical changes in articular cartilage answer, 360 question, 352
nail anatomy structures involved in fingertip laceration answer, 343–44 question, 327	nerve repair suitability for end-to-end repair answer, 459 question, 446	comparison with aging effects answer, 359 question, 351–52
National Board of Medical Examiners (NBME) item-writing manual, 9	nerve root injury identification from clinical signs answer, 344 question, 327	response of chondrocytes answer, 360 question, 352
neck C1 right side injury status of the transverse ligament answer, 130 question, 116	nerve that lies just lateral to the axillary artery answer, 334 question, 323–24	osteochondritis dissecans feature of answer, 388 question, 375
neck injury anterior and posterior arch fracture answer, 150 question, 139	nerve transection clinical signs caused by answer, 341–42 question, 326	incorrect statement about answer, 388 question, 375
comparison of treatment options answer, 131 question, 116	neurofibromatosis type 1 patient with enlarging painful thigh mass answer, 421 question, 409–10	knee lesions answer, 388 question, 375
treatment of odontoid fracture answer, 130–31 question, 116	neutralisation plate role in fracture fixation answer, 455 question, 442	surgical management answer, 388 question, 375
neck pain diagnosis and treatment answer, 128–29 question, 112	non-accidental injury (NAI) to a child	osteochondroma features of answer, 418–19 question, 407
neck pain after a fall initial management answer, 215 question, 196		osteochondromatosis features of answer, 425 question, 414
		osteocytes

- factor which decreases sclerostin expression
 - answer, 363
 - question, 353
- osteonecrosis
 - management of associated hip pain
 - answer, 52
 - question, 42
- osteoporosis
 - mechanism of action of bisphosphonates
 - answer, 362
 - question, 353
- osteoporotic fracture
 - vertebroplasty treatment
 - answer, 146
 - question, 135
- osteosarcoma
 - characteristics of high-grade tumours
 - answer, 424
 - question, 413
 - poor prognostic factors
 - answer, 423
 - question, 411
 - treatment options for a 9-year-old boy
 - answer, 423
 - question, 412
- Oxford knee replacement
 - need for more surgery than the patient has consented
 - answer, 83
 - question, 78
 - pre-surgery patient examination
 - concerns
 - answer, 83
 - question, 78
- p value
 - feature of
 - answer, 484–85
 - question, 473
- paediatric fractures
 - physis layer they occur through
 - answer, 215
 - question, 196
- Paget’s disease
 - pathophysiology of
 - answer, 404
 - question, 401
- paired sample t-test
 - appropriate use of
 - answer, 484
 - question, 473
- patella
 - primary role of
 - answer, 450
- question, 440
- patellar instability
 - considerations before tibial tubercle osteotomy
 - answer, 71
 - question, 64
 - medial patellofemoral ligament (MPFL) reconstruction
 - answer, 70
 - question, 62
- patellofemoral joint
 - effect of body weight when climbing stairs
 - answer, 79
 - question, 74
- pathological fracture
 - criteria for intramedullary nailing
 - answer, 232
 - question, 224–25
- patient confidentiality
 - sharing of patient information
 - answer, 470
 - question, 464
- patient warming
 - appropriate action for emergency surgical patient
 - answer, 468
 - question, 462
- Pavlik harness
 - principle of
 - answer, 436
 - question, 430–31
 - transient femoral nerve palsy related to
 - answer, 301
 - question, 290
- pectoral girdle
 - muscle identification
 - answer, 335
 - question, 324
- pelvic fracture
 - best Judet view to assess healing
 - answer, 223
 - question, 230
- pelvic incidence
 - features of
 - answer, 145
 - question, 134
- pelvic injury from a fall
 - appropriate surgical approach
 - answer, 221
 - question, 203
- pelvic pain
 - expanded bone area with soft tissue mass
 - answer, 423
 - question, 411–12
- pelvic planes and angles
 - answer, 34–35
 - question, 28–29
- pelvis
 - identify tear drop structures on anteroposterior radiograph
 - answer, 49–50
 - question, 41
 - management of motorbike accident trauma
 - answer, 211
 - question, 193
 - pelvic parameter that does not change with posture
 - answer, 148
 - question, 137
 - structure which does not enter via the lesser sciatic foramen
 - answer, 48
 - question, 40
- pelvis radiograph
 - structure to identify
 - answer, 219
 - question, 201
- per rectal examination
 - afferent nerve involved in penile response
 - answer, 394
 - question, 390
- peripheral nerve fibres
 - type that transmits vibration sense to the spinal cord
 - answer, 376
 - question, 368
- periprosthetic fracture
 - treatment strategy
 - answer, 233
 - question, 226
- periprosthetic hip fracture
 - management of different Vancouver types
 - answer, 379
 - question, 370
 - management of Vancouver type A
 - answer, 378–79
 - question, 370
 - management of Vancouver type B2
 - answer, 379
 - question, 370
 - Vancouver classification
 - answer, 378
 - question, 369–70
- periprosthetic infection
 - diagnostic criteria
 - answer, 384
 - question, 372
- Perthes disease
 - effects of proximal femoral varus osteotomy (PFVO)
 - answer, 294

Index

Perthes disease (cont.) question, 286 potential need for surgical intervention answer, 314 question, 306 technical issues in proximal femoral varus osteotomy (PFVO) answer, 294 question, 286 pes planus caused by navicular collapse diagnosis answer, 381 question, 371 physeal injuries in children features of answer, 299 question, 289 pilon fracture worst ankle position to sustain answer, 102 question, 89 placebo-controlled trial interpretation of results answer, 485 question, 474 plagiocephaly with torticollis management in a baby answer, 314 question, 306–7 plantar fasciopathy common aetiological factor answer, 92 question, 85 first line treatment answer, 93 question, 85 plating system material property answer, 456–57 question, 444 polyethylene manufacturing process answer, 460 question, 446 polymer cross-linking type of bond between monomer chains answer, 451 question, 441 polymethylmethacrylate (PMMA) cement function of zirconium dioxide answer, 451 question, 441 mechanical properties answer, 451 question, 441	polymethylmethacrylate (PMMA) cement powder components of answer, 451 question, 441 Ponseti casting viscoelastic material property utilised by answer, 450 question, 440 posterior cruciate ligament reconstruction use of x-ray guidance answer, 80 question, 75 posterior spinal fusion post-surgery pain and paraesthesia in a diabetic patient answer, 151–52 question, 140 Postgraduate Orthopaedics website, 8 post-operative wound infection likely infectious organism answer, 402 question, 396 pre-operative check-up rheumatoid patient with chronic neck pain answer, 155 question, 143 pre-operative patient skin preparation answer, 468 question, 462 primary bone healing plate fixation to achieve answer, 454 question, 442 prostate cancer management of metastatic spread answer, 132 question, 119–21 prosthetic joint infection diagnostic testing answer, 383 question, 372 most common causative organism answer, 383 question, 372 proteoglycan polymer identification answer, 459 question, 445 proteoglycans function in the extracellular matrix answer, 458 question, 445 proximal femoral varus osteotomy (PFVO) effects in Perthes disease	answer, 294 question, 286 technical issues in Perthes disease answer, 294 question, 286 quadriplegic spastic cerebral palsy management of problems related to spinal shape answer, 126 question, 108 radial fracture implant removal in children answer, 305 question, 293 likely impairments of thumb movement answer, 250 question, 241 nerve injury during fixation answer, 337–38 question, 324–25 plating for a distal fracture answer, 208–9 question, 191 post-operative investigation and management of nerve injury answer, 270–71 question, 258 rupture associated with undisplaced distal fracture answer, 214 question, 195 surgical approach and structure most at risk of damage answer, 248 question, 239 way to increase rigidity of the unilateral external fixator answer, 394 question, 390 wrist pain and swelling following distal fracture answer, 255 question, 246 radial head fracture complication of the Kaplan surgical approach answer, 217 question, 197 fixation of answer, 210–11 question, 192–93 radial head replacement management of nerve palsy complication answer, 271 question, 258
--	---	---

- radial neck fractures
 - features of
 - answer, 183
 - question, 171
- radial nerve
 - location of
 - answer, 350
 - question, 331
 - recovery after injury
 - answer, 235
 - question, 229
- radial nerve palsy
 - definitive management plan
 - answer, 273–74
 - question, 259
- radius
 - extensor tendon just medial to the dorsal tubercle
 - answer, 344
 - question, 327–28
 - volar (Henry) approach
 - answer, 336–37
 - question, 324
- randomised double-blind trial
 - feature of
 - answer, 485
 - question, 474
- reamed nailing
 - features of
 - answer, 363–64
 - question, 354
- recombinant human BMP (rhBMP)
 - licensed bone graft substitute
 - answer, 395
 - question, 391
- renal impairment
 - development of tertiary hyperparathyroidism
 - answer, 404
 - question, 400–1
- rheumatoid joint pathology
 - appropriate management to improve function
 - answer, 254
 - question, 245
- sacral giant cell tumour
 - mechanism of action of monoclonal antibody medication
 - answer, 133
 - question, 122
- sacroiliac joint fixation
 - symptom of screw penetration of the anterior cortex
 - answer, 221–22
 - question, 204
- SBA writing process
 - advantages of SBAs, 13
 - aims of, 9
 - aims of the SBA format, 26
 - Angoff method of standard setting, 18–19
 - basic rules for writing SBAs, 24
 - Bloom's taxonomy of cognitive categories, 10–12
 - blueprinting the curriculum, 20
 - constructing SBAs, 20–22
 - example of a poorly-constructed SBA, 21–22
 - lead-in, 21
 - options, 21
 - stem, 20
 - deriving the eligibility to proceed mark, 18–19
 - disadvantages of SBAs, 13
 - educational theory, 10–12
 - flaws related to irrelevant difficulty, 22–23
 - content of alternatives/distractors, 23
 - general guidelines, 23
 - language or structure is not homogeneous or parallel, 23
 - none of the above is used as an option, 23
 - numerical data are not consistently presented, 22
 - overly complicated or long questions, 22
 - stems contain negative phrasing, 23
 - terms in the options or the stem are vague, 23
 - true/false format, 23
 - unnecessarily complicated stems, 23
 - window dressing, 23
 - flaws related to testwiseness, 22
 - grammatical cue, 22
 - logical cue, 22
 - longest answer is 'single best answer', 22
 - presence of convergence, 22
 - presence of grouped or collectively exhaustive options, 22
 - repeated words, 22
 - use of absolute terms, 22
 - higher-order thinking, 12
 - highly discriminating questions, 12
 - item analysis, 13–18
 - difficult questions, 14
 - difficulty index (DIF I), 15, 18
 - discrimination index (DI), 15–18
 - distractor efficiency (DE), 18
 - easy questions, 14
 - good performing SBAs, 15–17
 - ideal SBA, 15–17
 - poorly performing SBAs, 14–16
 - lower-order thinking, 12
 - MCQ question writing committee, 23–24
 - Miller's pyramid of levels of assessment, 10
 - multilogical thinking, 12
 - NBME item-writing manual, 9
 - presentation of questions for candidates, 26
 - Question Writing Checklist, 26
 - reliability of a test, 19–20
 - standard error of measurement (SEM) and the GMC, 19
 - standard setting, 18–19
 - technical item flaws, 22–23
 - usefulness of understanding for candidates, 9
 - utility model to analyse assessment tools, 13
 - validity of a test, 20
 - writing guidelines - dos and don'ts, 24–26
- scapula fixation
 - nerve damage during Judet approach
 - answer, 218
 - question, 199–200
- scapular neck fracture
 - surgical approach
 - answer, 393
 - question, 389
- scapular region
 - effects of nerve damage on movement
 - answer, 332
 - question, 323
- scapular winging
 - root of injured nerve
 - answer, 335–36
 - question, 324
- scarf osteotomy
 - location of the distal transverse cut
 - answer, 100
 - question, 87–88
- Schatzker II tibial fracture
 - damage frequently associated with
 - answer, 236
 - question, 229
- void filler
 - answer, 236
 - question, 230
- sciatic nerve
 - anatomical variation exiting the greater sciatic foramen
 - answer, 48–49
 - question, 40
- scoliosis
 - in a patient with cerebral palsy
 - answer, 148

Index

- scoliosis (cont.)
 - question, 137
 - management in a patient with Duchenne’s muscular dystrophy
 - answer, 148
 - question, 138
 - suitable brace for a child
 - answer, 436
 - question, 431
- scoliosis correction
 - pedicle screw construction
 - answer, 124
 - question, 105
- scoliosis correction rods
 - comparison of material properties
 - answer, 124
 - question, 105
- scoliosis in a child
 - need for a full spine MRI scan
 - answer, 145
 - question, 134–36
 - pain more prevalent at night
 - answer, 145
 - question, 134
- screw fixation
 - factor with most impact on screw pull-out
 - answer, 404
 - question, 400
- second metatarsal phalangeal joint
 - diagnosis of pain and stiffness
 - answer, 98–99
 - question, 87
- septic arthritis
 - common infectious organism in patients with sickle cell disease
 - answer, 377
 - question, 368
- septic arthritis of the hip
 - Kocher’s criteria for
 - answer, 376
 - question, 368
- septic hip
 - best approach for open arthrotomy
 - answer, 377
 - question, 368
- shins
 - medial pain down both shins
 - answer, 128
 - question, 110–11
- shock
 - ATLS classification
 - answer, 220
 - question, 203
- short stature and bowed legs in a child
 - diagnosis
 - answer, 393
 - question, 389
- short stature and limb deformity in a child
 - laboratory finding
 - answer, 362–63
 - question, 353
- shoulder
 - cause of periprosthetic fracture in 71-year-old lady
 - answer, 185
 - question, 172
 - features of rotator cuff tears
 - answer, 184–85
 - question, 172
 - imaging investigation of weakness and deep-seated pain
 - answer, 162
 - question, 157
 - pain and stiffness in a 71-year-old patient
 - answer, 162
 - question, 157
- shoulder and neck pain
 - location of compressed nerve root
 - answer, 128
 - question, 112
- shoulder arthroplasty infection
 - common infective organism
 - answer, 383
 - question, 372
- shoulder arthroscopy
 - structure at risk of injury with posterior portal
 - answer, 332–33
 - question, 323
- shoulder dislocation
 - cause of recurrent instability
 - answer, 163
 - question, 158
 - test for deficit following reduction
 - answer, 162
 - question, 157
- shoulder dislocation surgery
 - consequences of nerve injury during graft harvesting
 - answer, 164–65
 - question, 159
- shoulder injury
 - traumatic injury irreducible in the ED
 - answer, 214
 - question, 195–96
- shoulder injury from a fall
 - Types of injuries
 - answer, 210
 - question, 192
- shoulder instability
 - findings in non-traumatic instability
 - answer, 166
 - question, 161
- shoulder pain
 - management of glenohumeral joint osteoarthritis
 - answer, 165
 - question, 160
 - management of pain associated with movement
 - answer, 185
 - question, 172–73
 - treatment for severe pain at night with shoulder weakness
 - answer, 185–86
 - question, 173
 - worsening pain with loss of shoulder movement
 - answer, 180–86
 - question, 173
- shoulder replacement
 - biomechanical advantages of reverse polarity shoulder prosthesis
 - answer, 163–64
 - question, 159
 - deficit related to McKenzie approach
 - nerve injury
 - answer, 162
 - question, 157
 - features of reverse polarity shoulder replacement
 - answer, 452–53
 - question, 441
 - mechanism of the reverse polarity procedure
 - answer, 404
 - question, 400
 - structure most at risk of damage
 - answer, 342
 - question, 326
- shoulder surgery
 - regional anaesthesia side-effect signs in an elderly patient
 - answer, 164
 - question, 159
- sickle cell disease
 - THA complication associated with
 - answer, 50–51
 - question, 41
- skeletal muscle
 - features of the actin and myosin functional units
 - answer, 403
 - question, 398
 - role of calcium in muscle contraction
 - answer, 403
 - question, 398
- slipped upper femoral epiphysis (SUFE)
 - complications of surgical pinning
 - answer, 304
 - question, 292

considerations with unilateral SUFE answer, 321 question, 312	answer, 154 question, 142	question, 225
fixation with a 'growing' screw answer, 320–21 question, 312	spinal cord injury complications with T4 injury and paraplegia answer, 148 question, 137	spine infection mode of spread answer, 153 question, 141
percutaneous in-situ fixation options answer, 320 question, 312	prognosis of different types of injury answer, 130 question, 115	spine-pelvis-hip relationship answer, 34–35 question, 28–29
risk of osteonecrosis answer, 304 question, 292	spinal deformity in a child indication for a brace answer, 146–47 question, 136	spino-pelvic alignment description of normal alignment answer, 124 question, 105
surgical management answer, 304 question, 292	spinal infection imaging findings answer, 146 question, 135	spontaneous osteonecrosis of the knee (SONK) features of answer, 67 question, 60
use of an unthreaded fixation device answer, 314 question, 306	spinal nerve injury complication of iliosacral screw fixation answer, 345 question, 328	spring ligament anatomy of answer, 90 question, 84
Smith-Petersen approach to the hip vessel that is sacrificed answer, 52 question, 42	spinal trauma likely location of spinal injury answer, 150 question, 139	statistics choice of appropriate statistical test answer, 471 question, 465
Smith Robinson approach features of answer, 146 question, 135	significance of return of the bulbocavernous reflex answer, 150 question, 139	stem cement fretting mode of wear answer, 380 question, 370
soft tissue reconstruction in open fractures answer, 235–36 question, 229	spine diagnostic injection for multilevel pathology answer, 128 question, 111	stem loosening types of answer, 380 question, 370
soft tissue sarcoma treatment answer, 424 question, 413	location of compressed nerve root answer, 128 question, 112	sternal and rib pain with osteopaenia diagnosis answer, 418 question, 406
type most likely to develop lymph node metastases answer, 425 question, 415	management of lateral recess stenosis answer, 128 question, 110–11	stinger nerve injury unlikely cause of answer, 151 question, 139
spastic diplegia gait management options answer, 316 question, 308	paediatric spinal trauma answer, 147 question, 136	stress relaxation definition answer, 402 question, 397
spina bifida features of answer, 304–5 question, 293	pathology of lateral recess stenosis answer, 128 question, 111	stress strain curve features of answer, 365 question, 355
spinal cord anatomy of the afferent and efferent tracts answer, 403 question, 398	thoracic or lumbar compression fractures answer, 149 question, 136–38	for different materials answer, 453 question, 441–42
lumbar level of termination answer, 350 question, 331	spine anatomy answer, 147–48 question, 137	study design approach to answer, 471 question, 465
predominant blood supply answer, 154 question, 142	spine fracture indications for posterior stabilisation answer, 232	dealing with lost data answer, 488 question, 478
structures found in a superficial to deep approach		

Index

- study design (cont.)
 - types of error
 - answer, 471
 - question, 465
- subchondral bone
 - substance released to aid formation of new cartilage
 - answer, 458
 - question, 445
- submicron particles
 - generation in a metal on polythene THA
 - answer, 58
 - question, 44
- subscapularis muscle
 - innervation
 - answer, 335
 - question, 324
- superior gemelli
 - nerve supply to
 - answer, 48
 - question, 40
- supinator muscle
 - nerve branch within
 - answer, 343
 - question, 326–27
- supracondylar fracture
 - indications for urgent surgical treatment
 - answer, 209
 - question, 191
 - lack of a pulse after surgical repair
 - answer, 179–80
 - question, 170
 - neuropraxia finding after cast application
 - answer, 177–78
 - question, 168
 - no feeling in little finger post-surgery
 - answer, 182
 - question, 171
- supracondylar fracture surgery
 - post-operative complications in a 5-year-old child
 - answer, 165
 - question, 160
- survival analysis study
 - statistical effect of data adjustment
 - answer, 467
 - question, 461
- survival data
 - interpretation of the graph
 - answer, 491
 - question, 482–83
- synapse
 - ion concentration changes in an action potential
 - answer, 404–5
 - question, 401
- synovial fluid
 - non-Newtonian characteristics
 - answer, 404
 - question, 399
- synovial joint lubrication
 - most important form in prolonged stance
 - answer, 447
 - question, 439
- synovium
 - cell types and functions
 - answer, 404
 - question, 399
- t-test
 - interpretation of p value
 - answer, 484
 - question, 473
- talar body
 - main blood supply
 - answer, 91
 - question, 84
- talar fracture-dislocation
 - approach for fixation
 - answer, 233
 - question, 225
- talar injuries
 - location of
 - answer, 101–2
 - question, 88–89
- talar neck fracture
 - blood supply to the talar neck
 - answer, 211–12
 - question, 193
 - risk of talar avascular necrosis
 - answer, 102
 - question, 89
- taper slip cemented stem fixation
 - optimisation
 - answer, 55
 - question, 43
- TB
 - expected microscopy finding
 - answer, 402
 - question, 397
- teicoplanin
 - mechanism of action
 - answer, 383
 - question, 372
- tendon tensile testing
 - shape of the curve in the elastic region
 - answer, 459
 - question, 446
- tendon transfer
 - for humeral fracture complications
 - answer, 250
 - question, 241
- tendons
 - proteoglycan that regulates tendon diameter
 - answer, 394
 - question, 391
- tenosynovial giant cell tumour
 - feature of
 - answer, 421–22
 - question, 410–11
- testing
 - sensitivity and specificity of a test
 - answer, 488
 - question, 477
- theatre
 - reducing the risk of airborne contamination
 - answer, 467–68
 - question, 462
- theatre design
 - acceptable level of bacteria for a ventilation system
 - answer, 467
 - question, 461–62
- theatre suite
 - zoning within
 - answer, 462
 - question, 462
- theatre ventilation
 - features of laminar flow systems
 - answer, 404
 - question, 400
- thigh
 - enlarging painful mass in patient with neurofibromatosis type 1
 - answer, 421
 - question, 409–10
 - enlarging painless mass with 10-year history
 - answer, 420–21
 - question, 408–9
 - features of the anterior thigh
 - answer, 348
 - question, 330
 - painful mass with shooting pain to the foot
 - answer, 421
 - question, 409
- thigh mass
 - diagnosis of rectus femoris lesion
 - answer, 422
 - question, 410
- thigh pain following fever
 - histiocyte-rich femoral lesion
 - answer, 419–20

- question, 408
- thigh pain in an adolescent
 - longstanding pain that is worse at night
 - answer, 419
 - question, 407
- third class levers
 - features of
 - answer, 365
 - question, 352–55
- thoracic compression fractures
 - features of
 - answer, 149
 - question, 138
- thoracic kyphosis
 - in a teenage patient
 - answer, 152
 - question, 140
- thoracic spine
 - structure most at risk during
 - left-sided transthoracic approach
 - answer, 349
 - question, 330
- thoracolumbar burst fracture
 - when surgery is not indicated
 - answer, 148–49
 - question, 138
- Thoracolumbar Injury
 - Classification and Severity Score
 - components of
 - answer, 149
 - question, 138
- thoracolumbar spine
 - flexion distraction injury
 - answer, 149
 - question, 138–39
- thumb
 - damage caused by forced abduction injury
 - answer, 250–51
 - question, 241–42
 - inability to extend following distal radial fracture
 - answer, 279
 - question, 260
 - management of accidental amputation
 - answer, 269
 - question, 257
 - management of displaced intra-articular fracture
 - answer, 276
 - question, 259–60
 - management of extension restriction in a child
 - answer, 254–55
 - question, 246
 - management of injury shown on radiograph
 - answer, 285
 - question, 262
 - management of palmar laceration
 - answer, 263–64
 - question, 256
 - management of unilateral small thumb in a child
 - answer, 271–72
 - question, 258
 - treatment of partial amputation
 - answer, 253
 - question, 244
- tibia
 - diagnosis of a bone dysplasia
 - answer, 361
 - question, 352–53
 - distal tibia physis closure process
 - answer, 320
 - question, 312
 - structure found during anterior approach
 - answer, 345
 - question, 328
 - unilateral bowing in a 6-month-old child
 - answer, 315
 - question, 306–7
- tibial fracture
 - continuing discomfort four months after intramedullary nailing
 - answer, 218
 - question, 199
 - criterion for non-emergency operation
 - answer, 221
 - question, 203
 - damage frequently associated with Schatzker II fracture
 - answer, 236
 - question, 229
 - features of high-energy open fractures in children
 - answer, 298
 - question, 288
 - initial management of traumatic injury
 - answer, 214
 - question, 195
 - management of an open tibial shaft fracture
 - answer, 377
 - question, 368
 - MESS score of patient with gunshot wound
 - answer, 215
 - question, 196
 - soft tissue reconstruction in open fractures
 - answer, 235–36
 - question, 229
 - surgical technique for intramedullary nailing
 - answer, 206–7
 - question, 189–90
 - void filler for Schatzker II fracture
 - answer, 236
 - question, 230
- tibial lesion in a 14-year-old male
 - appropriate management of
 - answer, 418
 - question, 406
- tibial lytic lesion with soap bubble appearance
 - answer, 424
 - question, 413
- tibial pain in a 6-year-old boy
 - mode of investigation
 - answer, 423
 - question, 411
- tibial plateau
 - bony landmarks in an anterolateral approach
 - answer, 346
 - question, 329
- tibial tendon dysfunction
 - suitable orthotic
 - answer, 434
 - question, 429
- tibial tubercle osteotomy
 - choice of technique
 - answer, 71–72
 - question, 64
 - considerations before
 - answer, 71
 - question, 64
- tibialis anterior tendon rupture
 - management in a diabetic patient
 - answer, 434–35
 - question, 429
- toddler with irritability
 - injuries with high specificity for non-accidental injury (NAI)
 - answer, 212
 - question, 193
- toe deformities in an infant
 - condition suggested by
 - answer, 99
 - question, 87
- toe joint pain and deformity in an elderly patient
 - appropriate treatment
 - answer, 99
 - question, 87
 - ‘too many toes’ sign
 - diagnostic indication

Index

- ‘too many toes’ sign (cont.)
 - answer, 95
 - question, 85
- torsional profile
 - components of
 - answer, 317
 - question, 309
- total elbow replacement
 - distal humerus fracture in elderly patients
 - answer, 235
 - question, 228
- total hip arthroplasty
 - cause of failure after ten years
 - answer, 46–56
 - question, 43
 - cause of leg shortening
 - answer, 37
 - question, 31
 - cause of polythene wear
 - answer, 371
 - question, 370
 - cementing
 - answer, 380–81
 - question, 371
 - complication associated with sickle cell disease
 - answer, 50–51
 - question, 41
 - complications
 - answer, 35
 - question, 29
 - effects of a plus 4mm (neck length) head
 - answer, 56
 - question, 43
 - factors affecting the fluid film and lubrication system
 - answer, 457
 - question, 444
 - factors that increase primary arc range
 - answer, 57–58
 - question, 44
 - foot turning inwards one year later
 - answer, 46
 - question, 39
 - for high developmental hip dysplasia (Crowe type IV)
 - answer, 57
 - question, 44
 - generation of submicron particles in a metal on polythene THA
 - answer, 58
 - question, 44
 - in patients with cerebral palsy
 - answer, 299
 - question, 289
 - indication for a constrained liner following dislocations
 - answer, 51
 - question, 41–42
 - lubrication in a metal on polythene joint
 - answer, 46
 - question, 39
 - management of post-surgery
 - indication of loss of nerve function
 - answer, 57
 - question, 44
 - management of previously placed dynamic hip screw
 - answer, 57
 - question, 44
 - modes of femoral stem loosening
 - answer, 46–56
 - question, 43–44
 - muscle recovery after nerve injury
 - answer, 49
 - question, 40
 - optimal combination of implant position
 - answer, 51
 - question, 41
 - optimal combined version of acetabular cup and femoral stem
 - answer, 51
 - question, 41
 - osteolysis secondary to polythene wear debris
 - answer, 52
 - question, 42
 - post-surgery pain, numbness and weakness
 - answer, 37
 - question, 31
 - reducing prosthetic joint infection risk
 - answer, 48
 - question, 40
 - reduction of blood transfusion requirement
 - answer, 47–48
 - question, 40
 - revision following damage to
 - answer, 46
 - question, 39
 - revision materials
 - answer, 37
 - question, 31
 - sites of loosening pre-fracture
 - answer, 45–46
 - question, 39
 - use of a PMMA centraliser in a taper slip design
 - answer, 458
 - question, 444
- total hip replacement
 - advice regarding rheumatoid medication
 - answer, 385
 - question, 373
 - effect of increasing femoral head diameter
 - answer, 449
 - question, 440
 - heterotopic ossification
 - answer, 36
 - question, 30
 - increasing hip motion
 - answer, 35
 - question, 29
 - increasing the primary arc range of motion
 - answer, 449–50
 - question, 440
 - nerve injury risk
 - answer, 34–35
 - question, 29
 - prophylaxis for Brooker grade 3 changes
 - answer, 36
 - question, 30
 - prosthesis design to achieve hydrodynamic fluid-film lubrication
 - answer, 452–57
 - question, 441
 - reason for not continually impacting the implant
 - answer, 456
 - question, 444
 - wound complication
 - answer, 35–36
 - question, 29–30
- total knee arthroplasty
 - cause of a cam jump dislocation
 - answer, 435
 - question, 430
 - genetic risk factor for venous thromboembolism
 - answer, 394
 - question, 391
 - improving patellofemoral tracking
 - answer, 67
 - question, 60
 - issues with previous high tibial osteotomy (HTO)
 - answer, 68
 - question, 60–61
 - optimising tibial component rotation
 - answer, 66
 - question, 59

- radiographic findings that make it more difficult
 - answer, 83
 - question, 77
- sagittal balancing of the knee
 - answer, 69
 - question, 62
- surgical approach
 - answer, 68
 - question, 61
- total knee replacement
 - arthritic valgus knee
 - answer, 71
 - question, 64
 - balancing a valgus knee
 - answer, 82
 - question, 77
 - cause of later flexion instability
 - answer, 66–67
 - question, 59
 - flexion is loose but extension is ok
 - answer, 81–82
 - question, 76
 - flexion is ok but extension is loose
 - answer, 81
 - question, 76
 - management of later *S. epidermidis* infection
 - answer, 383
 - question, 372
 - management of post-operative osteolysis
 - answer, 82
 - question, 77
 - numbness after arthritic valgus knee
 - procedure
 - answer, 82–83
 - question, 77
 - patient states they were ‘never really happy’ with the new knee
 - answer, 82
 - question, 76–77
 - patient with previous ORIF of lateral tibial plateau
 - answer, 67
 - question, 60
 - procedure with risk of future patellar degeneration
 - answer, 82
 - question, 77
- total shoulder arthroplasty
 - biomechanics of reverse total shoulder arthroplasty
 - answer, 186
 - question, 173
 - strategies to avoid future functional deficits
 - answer, 186–87
 - question, 174
- total shoulder replacement
 - loss of range of motion 8 weeks after surgery
 - answer, 165
 - question, 39
 - neurological deficit caused by use of large soft tissue retractors
 - answer, 162
 - question, 157
 - pain and loss of range of motion 6 months after surgery
 - answer, 165
 - question, 160
- tourniquet application
 - comorbidity contraindications for
 - answer, 469
 - question, 463
 - patient with a high BMI
 - answer, 469
 - question, 463
- transradial amputation
 - features of
 - answer, 438
 - question, 432
- trapeziectomy
 - structures not at risk with a dorsal approach
 - answer, 253
 - question, 244
- trauma patient (adult)
 - likely location of spinal injury
 - answer, 150
 - question, 139
- trauma patient (car accident)
 - description of spinal injury
 - answer, 131
 - question, 117
 - management of spinal injury
 - answer, 131
 - question, 117–18
- trauma patient (fall from a height)
 - hypovolaemic shock classification (ATLS)
 - answer, 220
 - question, 203
- trauma patient (fall from a tractor)
 - description of the injury
 - answer, 131–32
 - question, 118
 - management of the injury
 - answer, 132
 - question, 118
- trauma patient (gunshot wound)
 - symptoms from a bullet lodged in the spinal cord
 - answer, 152
 - question, 141
- trauma patient (motor vehicle accident)
 - management of cervical subluxation
 - answer, 148
 - question, 137
 - odontoid fracture in a 70-year-old lady
 - answer, 152
 - question, 140
 - paediatric spinal trauma
 - answer, 147
 - question, 136
 - thoracolumbar flexion distraction injury
 - answer, 149
 - question, 138–39
- trauma patient (motorbike accident)
 - determination of adequate resuscitation
 - answer, 221
 - question, 203–4
 - Early Appropriate Care (EAC) protocol
 - answer, 215
 - question, 196
 - weakness in both legs and lower back pain
 - answer, 216
 - question, 197
- trauma patient (motorcycle accident)
 - femoral nailing in context of damage control orthopaedics (DCO)
 - answer, 235
 - question, 228
 - Injury Severity Score (ISS)
 - answer, 222
 - question, 204
 - management of head and spinal injuries
 - answer, 131
 - question, 116–17
 - spinal stability for transfer
 - answer, 131
 - question, 117
- trauma patient (odontoid fracture)
 - checks before halo orthosis
 - application
 - answer, 154
 - question, 143
- trauma patient (penetrating abdominal injury)
 - limb weakness after surgical repair
 - answer, 145
 - question, 134–35
- trauma patient (RTA) with ankylosing spondylitis
 - factor affecting outcome
 - answer, 145
 - question, 134
- trauma patient (rugby tackle)

Index

- trauma patient (rugby tackle) (cont.)
 - upper body nerve injury
 - answer, 151
 - question, 139
- trauma patient (stab wound to the back)
 - loss of fine touch in upper limb
 - answer, 152
 - question, 141
- tuberculous spondylodiscitis
 - features of
 - answer, 378
 - question, 369
- tubular discectomy
 - comparison with microdiscectomy (conventional discectomy)
 - answer, 156
 - question, 144
- tumour in posterior spinal elements of T10
 - diagnosis
 - answer, 152–53
 - question, 141
- tumour of the sacral midline
 - type of cell derived from
 - answer, 132
 - question, 118–19
- UCBL insert
 - condition suited for
 - answer, 437
 - question, 432
- ulnar nerve laceration
 - Sunderland Classification for nerve injury
 - answer, 249
 - question, 240
- ulnar nerve repair
 - splinting to correct deformity
 - answer, 436
 - question, 431
- unicameral bone cyst
 - features of
 - answer, 297
 - question, 288
 - management of a femoral lesion in a child
 - answer, 298–99
 - question, 288–89
- uncompartmental knee replacement
 - more damage discovered during the procedure
 - answer, 83
 - question, 78
- United Kingdom In-Training Examination (UKITE), 8
- upper limb prosthesis
 - for above elbow amputation
 - answer, 434
 - question, 429
 - type best indicated for heavy labour
 - answer, 434
 - question, 428
- viscoelastic materials
 - property of
 - answer, 393
 - question, 389
- viscoelasticity
 - feature of
 - answer, 449
 - question, 440
- Ward’s triangle description
 - answer, 52–53
 - question, 42
- windblown hips
 - features of
 - answer, 300
 - question, 290
- wound healing
 - cellular stages of
 - answer, 250
 - question, 241
- wrist
 - diagnosis of bilateral deformities and limited forearm rotation
 - answer, 274
 - question, 259
 - imaging of suspected scaphoid fracture
 - answer, 275
 - question, 259
 - management of a displaced proximal pole scaphoid fracture
 - answer, 274
 - question, 259
 - management of scaphoid waist fracture from high-energy injury
 - answer, 274–75
 - question, 259
 - structure providing stability to the distal radio-ulnar joint
 - answer, 252
 - question, 243
- wrist arthroscopy
 - best portal for access
 - answer, 253
 - question, 245
- wrist fracture and tendon rupture
 - reconstruction technique
 - answer, 233
 - question, 225
- wrist injury
 - management of later osteoarthritic changes
 - answer, 251
 - question, 242
 - management of undisplaced scaphoid waist fracture
 - answer, 255
 - question, 246
- wrist injury from a fall
 - pathoanatomical sequence of events
 - answer, 212
 - question, 193
 - pattern of injury
 - answer, 249
 - question, 240
 - volar approach to repair scaphoid fracture
 - answer, 212–13
 - question, 193–94
- wrist pain
 - cause of chronic pain in patient with Ehlers-Danlos syndrome
 - answer, 277–78
 - question, 260
 - central pain and stiffness of six months’ duration
 - answer, 280–81
 - question, 261
 - confirmation of scaphoid shift
 - diagnosis
 - answer, 267
 - question, 257
 - diagnosis in a patient with diabetes
 - answer, 254
 - question, 246
 - lytic lesion in the distal radius
 - answer, 420
 - question, 408
 - management of dorso-radial pain
 - answer, 248
 - question, 239
 - management of historic injury with scaphoid fracture non-union
 - answer, 268–69
 - question, 257
 - management of proximal lunate collapse
 - answer, 252
 - question, 243
 - radial-sided pain of eight months’ duration
 - answer, 279–80
 - question, 261
 - twelve-month history of worsening pain
 - answer, 277–81
 - question, 261
 - with triangular fibrocartilage complex (TFCC) lesion
 - answer, 265–66
 - question, 256

wrist pain after motorbike accident	yield point of a material	definition
nature of the injury	definition	answer, 402
answer, 266–67	answer, 448–49	question, 397
question, 257	question, 439–40	
wrist pain and reduced grip strength	Young’s Modulus	zone of injury
management in a 39-year-old	consideration in femoral stem	distal phalanx of the ring finger
patient	design	answer, 335
answer, 265	answer, 448	question, 324
question, 256	question, 439	