

Index

- ADHD. *See* attention deficit hyperactivity disorder
- adrenaline effects, of anxiety, 2
- ADVANCE programme, 78
- affective satisfaction, 25
- altered state
 through hypnosis, 54
 with procedure-induced anxiety, 14
- alternative point of entry, 45–6
- animation, in non-verbal communication, 17–24
- anxiety, 1–4. *See also specific topics*
 anxiolysis through
 premedication and, 60
 coping strategies for, 4–6
 cognitive approaches, 4
 internal distraction techniques, 4–13
 extreme, 2
 fear and, 2
 fight-flight response to, 2
 freeze-flight response to, 2
 management of, 4. *See also* disordered behaviour
 nature of, 2–13
 adrenaline effects, 2
 internal dialogue in, 3
 internal focus in, 3
 noradrenaline effects, 2
 physical symptoms, 3
 non-threats and, 3
 threats and, 3
 anxiety management services, 75
 anxiolysis, through
 premedication, 60–2
 anxiety and, 60
 apps, in elective management, 79–80
- ASD. *See* autistic spectrum disorder
- attention deficit hyperactivity disorder (ADHD), 65
- augmented reality, 79
- autistic spectrum disorder (ASD), 64
- the bind, 42–50
 Birdwhistell, Ray, 16
 British Medical Journal, 53–4
 Burgoon, Judee, 42–3. *See also* expected violation theory
- CBT. *See* cognitive behaviour therapy
- child-clinician interactions, elements of, 10–13
- coercion in, 12
- compassion, 11
 definition of, 11
- compliance in, 12
- restraint, 12–13
 confinement compared to, 12
 trust, 10–11
- clonidine, 61, 67
- coercion, in child-clinician interactions, 12
- cognitive appraisal, 51–2
- cognitive behaviour therapy (CBT), 76
- communication. *See also* non-verbal communication; verbal communication
 with procedure-induced anxiety, 15
 relational, 25
 satisfaction and, 25
 two-level communication theory, 29
- compassion
 in child-clinician interactions, 11
 definition of, 11
- compliance
 in child-clinician interactions, 12
 patient satisfaction and, 25
- confinement, restraint compared to, 12
- confusion, through overload, 44–5
- congruence, with procedure-induced anxiety, 14
- conscious awareness, 28–9
- content-free suggestions, 33–4
- conversational suggestion, 31–6
 complexity of, 31–2
 content-free suggestions, 33–4
 direct suggestions, 31
 embedded suggestions, 34
 imagination, 32–3
 implication in, 34–5
 indirect suggestions, 32–3
 interspersal of suggestions, 34
 language patterns in, 32
 metaphor and storytelling in, 33
 negative suggestions, 31
 negatives in, use of, 36
 permissiveness and, 32
 utilisation theory in, 29–35
- coping skills training, 78
- coping strategies, 4–6
 cognitive approaches, 4
 through enhanced communication, 41
 internal distraction techniques, 4–13
- decompression techniques, 55–6
- Dennig, M. D., 6
- de-potentiating conscious sets, 30
- dexmedetomidine, 61, 67
- direct suggestions, 31
- disordered behaviour
 anxiety management strategies, 64–7
 in community, 66
 on day of intervention, 66–7
 elective management, in advance of intervention, 66
 resources, 67
 for stress reduction, 67

- attention deficit
 - hyperactivity disorder, 65
- autistic spectrum disorder, 64
- extreme cases of, 67–8
- learning difficulties, 65
- premedication for, 67
- stereotypical characteristics, 64–5
- distraction techniques, 48
- dynamic transition, in verbal communication, 38–9
- elective management
 - anxiety management services, 75
 - through audiovisual material, 79
 - cognitive behaviour therapy, 76
 - through elective therapy, 75–6
 - with evolving technology, 79–80
 - apps, 79–80
 - augmented reality, 79
 - virtual reality, 79
 - family therapy, 76
 - grief therapy, 76
 - hospital tours and, 78–9
 - humanistic therapy, 76
 - hypnotherapy, 77
 - interpersonal therapy, 76
 - Neurolinguistic Programming, 76
 - play specialists, 76–7
 - play therapists, 77
 - preparation programmes, 77–8
 - ADVANCE programme, 78
 - coping skills training, 78
 - life skills training, 78
 - psychoanalytic therapy, 76
 - psychodynamic therapy, 76
 - theoretical approach to, 74
 - through therapeutic interventions, 76. *See also specific interventions*
 - transactional analysis, 76
 - written information on, 78–9
- elective therapy, 75–6
- embedded suggestions, 34
- enhanced communication
 - strategies
 - alternative point of entry, 45–6
 - the bind as, 42–50
 - through cognitive appraisal, 51–2
 - confusion and, through overload, 44–5
 - coping strategies through, 41
 - decompression techniques, 55–6
 - through distraction, 48
 - expectancy violations theory, 42–3
 - through guided imagery, 52–3
 - in storytelling, 53
 - through humour, 47–8
 - hypnosis in, 53–5
 - altered states, 54
 - in *British Medical Journal*, 53–4
 - highway, 54
 - incongruent behaviour, 45
 - for internal focus, 40
 - through magic, 55
 - negative perceptions in, challenges to, 46
 - obstacles in, 41
 - opportunities in, 41
 - pseudo-orientation in, 50
 - for rapport building, 40
 - release as part of, 50
 - successive approximation, 52
 - tag-team talking, 47
 - tension in, 50
 - touch, 43–4
 - validation through, 50–1
 - yes set techniques, 48–9
- Erickson, Milton, 29, 32, 35
- yes set techniques, 48–9
- expected violation theory (EVT), 19
- enhanced communication
 - strategies and, 42–3
- extreme anxiety, 2
- eye contact, 20–1
- family relationships,
 - procedure-induced anxiety influenced by, 5–6
 - academic approach to, 69–70
 - challenging behaviour as element of, 70
- post-traumatic stress
 - disorder and, 69
- family therapy, 76
- fear, anxiety and, 2
- fentanyl, 62
- fight-flight response, to anxiety, 2
- fixation of attention, 30
- flight distance, in non-verbal communication, 19
- freeze-flight-flight response, 2
- Global Malignant Anxiety Disorder, 6
- grief therapy, 76
- guided imagery, 52–3
 - in storytelling, 53
- Hall, Edward, 16–17
- Haptics. *See* touch
- highway hypnosis, 54
- hospital environment, anxiety as result of
 - ambient temperature, 73
 - clothing and, 71–3
 - hospital equipment, 71
 - from other children, 72–3
 - paediatric themes in, 72
 - personal protection equipment, 71–3
 - premedication taste and odour, 73
 - from tactile stimuli, 73
- humanistic therapy, 76
- humour, in enhanced communication
 - strategies, 47–8
- hypnosis, in enhanced communication
 - strategies, 53–5
 - altered states, 54
 - in *British Medical Journal*, 53–4
 - highway, 54
- hypnotherapy, 77
- hypnotic response, 30–1
- imagination, in conversational suggestion, 32–3
- incongruent behaviour, 45
- indirect suggestions, 32–3
- internal dialogue, 3
- internal distraction techniques, 4–13
- internal focus, 3, 40
- interpersonal therapy, 76

- interspersal of suggestions, 34
 intimate distance, in non-verbal communication, 17
- ketamine, 61, 67
 Kinesics, 16
- learning difficulties (LDs), 65
 life skills training, 78
 Linehan, Marsha, 51
 long-term operative procedure-induced anxiety, 8
- magic, as enhanced communication strategy, 55
 mass effect, 14
 Mehrabian, Albert, 27
 memories, procedure-induced anxiety and, 6
 metaphor, in conversational suggestion, 33
 microdynamics of suggestion, 29–31
 de-potentiating conscious sets, 30
 fixation of attention, 30
 hypnotic response, 30–1
 unconscious process, 30
 unconscious search, 30
 midazolam, 60–1
 mirroring, 23–4
 morphine. *See* oral morphine
- nasal sufentanyl, 62
 negative suggestions, 31
 negative vocabulary, 36–7
 Neurolinguistic Programming (NLP), 76
 non-threats, anxiety and, 3
 non-verbal communication. *See also* posture
 animation, 17–24
 definition of, 16–24
 expected violation theory and, 19
 eye contact, 20–1
 facial expression, 21
 flight distance, 19
 head movement, 21–2
 height and, 19–20
 intimate distance, 17
 Kinesics and, 16
 personal distance, 17
 proxemics, 16–19
 public distance, 18–19
 Western culture as influence on, 18–19
 rapport and, 19
 social distance, 18
 noradrenaline effects, of anxiety, 2
- operative procedure-induced anxiety (OPIA), 7–9
 consequences of, 7
 definition of, 1
 incidence rates, 7
 lifespan issues, 8–9
 long-term, 8
 post-operative dysfunctional behaviour and, 7
 procedure-induced psychological trauma and, 7
 short-term, 7–8
 oral morphine, 62
 overload, confusion through, 44–5
 Ozalin, 60–1
- paralinguistics, 27. *See also* Neurolinguistic Programming
 patient satisfaction. *See* satisfaction
 personal distance, in non-verbal communication, 17
 PIA. *See* procedure-induced anxiety
 PIPT. *See* procedure-induced psychological trauma
 play specialists, 76–7
 play therapists, 77
 PODB. *See* post-operative dysfunctional behaviour
 positive indirect conversational suggestion, 38
 positive pre-emptive interpretation, 37–8
 abstract alternatives, 37–8
 positive indirect conversational suggestion in, 38
 positive vocabulary, 36–7
 post-operative dysfunctional behaviour (PODB)
 definition of, 1
 operative procedure-induced anxiety and, 7
 UK statistics, 9–10
- post-procedure dysfunctional behaviour (PPDB), definition of, 1
 post-traumatic stress disorder (PTSD), 69
 posture, 22–4
 closed, 22
 frame of dominant orientation, 24
 interactive orientation, 23
 positions, 23
 mirroring, 23–4
 open, 22
 relaxation in, 22
 tension in, 22
 torso angle, 22
 PPDB. *See* post-procedure dysfunctional behaviour
 pre-conscious awareness, 28–9
 premedication
 acceptance of, 58–9
 administration of, 59–60
 anxiolysis through, 60–2
 anxiety and, 60
 clonidine, 61, 67
 dexmedetomidine, 61, 67
 fentanyl, 62
 ketamine, 61, 67
 midazolam, 60–1
 nasal sufentanyl, 62
 oral morphine, 62
 organisation of, 60
 Ozalin, 60–1
 refusal of, 58–9
 taste and odour of, anxiety as result of, 73
 timing of, 60
 procedure-induced anxiety (PIA). *See also* non-verbal communication; verbal communication
 age cohort for, 5
 assessment of, with State Trait Anxiety Inventory, 5
 communication with, 15
 core principles of, 14
 altered states, 14
 congruence, 14
 mass effect, 14
 resistance, 14
 definition of, 1
 family relationships as influence on. *See* family relationships

- management of, 6. *See also specific topics*
 through memory access, 6
 resources for, 6–7
 transgenerational effects of, 9
 procedure-induced
 psychological trauma (PIPT)
 definition of, 1
 operative procedure-induced
 anxiety and, 7
 proxemics, 16–19
 pseudo-orientation, in enhanced
 communication
 strategies, 50
 psychoanalytic therapy, 76
 psychodynamic therapy, 76
 PTSD. *See* post-traumatic stress
 disorder
 public distance, in non-verbal
 communication, 18–19
 Western culture as influence
 on, 18–19
- rapport
 in enhanced communication
 strategies, 40
 non-verbal communication
 and, 19
 relationships. *See* family
 relationships
 release, through enhanced
 communication
 strategies, 50
 resistance, in procedure-
 induced anxiety, 14
 restraint, in child-clinician
 interactions, 12–13
 confinement compared to, 12
 Rossi, Ernest, 29
 yes set techniques, 48–9
 Rudolph, K. D., 6
- satisfaction, of patients
 affective, 25
- communication and, 25
 compliance and, 25
 predictors of, 25
 Schefflin, Albert, 23–4
 short-term operative
 procedure-induced
 anxiety, 7–8
 social distance, in non-verbal
 communication, 18
 Sommer, Robert, 23
 State Trait Anxiety Inventory
 (STAI), 5
 storytelling
 in conversational suggestion,
 33
 guided imagery in, 53
 subconscious awareness, 28–9
 successive approximation, 52
 sufentanyl. *See* nasal sufentanyl
 suggestion. *See* conversational
 suggestion;
 microdynamics of
 suggestion
- TA. *See* transactional analysis
 tag-team talking, 47
 tension, in enhanced
 communication
 strategies, 50
 therapy. *See specific types*
 therapy-based verbal
 communication, 28
 threats, anxiety and, 3
 touch, in enhanced
 communication
 strategies, 43–4
Trancework, 53, 55
 transactional analysis (TA), 76
 transgenerational effects, of
 procedure-induced
 anxiety, 9
 trust, in child-clinician
 interactions, 10–11
 two-level communication
 theory, 29
- unconscious awareness, 28–9
 utilisation theory, 29–35
- validation, through enhanced
 communication
 strategies, 50–1
 verbal communication. *See also*
 conversational
 suggestion;
 microdynamics of
 suggestion
 conscious awareness,
 28–9
 definition of, 27
 dynamic transition in,
 38–9
 influence of, 27–8
 vocal intonation, 28
 negative vocabulary, 36–7
 paralinguistics, 27
 positive pre-emptive
 interpretation, 37–8
 abstract alternatives,
 37–8
 positive indirect
 conversational
 suggestion in, 38
 positive vocabulary, 36–7
 pre-conscious awareness,
 28–9
 subconscious awareness,
 28–9
 therapy-based, 28
 two-level communication
 theory, 29
 unconscious awareness,
 28–9
 virtual reality, 79
 vocabulary use. *See* negative
 vocabulary; positive
 vocabulary
- Weisz, J. R., 6
- yes set techniques, 48–9