

INDEX

- acceptance and commitment therapy (ACT), 77–8
- age and personality disorders, 42–3
- Age of Acceptance, 9
- Age of Discovery (460 BC–1700 AD), 1–4
- Age of Investigation, 7–8
- Age of Personality as a Substrate (1750–1950)
- Adolf Meyer, 4–5
- David Henderson, 5
- Gordon Allport, 5–6
- in literature, 6–7
- mental illness links to personality, 4
- Rolv Gjessing, 5
- Sigmund Freud, 4
- Allport, Gordon, 5–6
- Alternative Model for Personality Disorders (AMPD) (DSM-5), 21, 29
- anankastia
- as trait domain, 17
- treatment, 89
- Anatomy of Melancholy, The*, 4
- antisocial personality disorder, 43
- anxiety disorders
- health anxiety, 65–6
- pathoplasty model for treatment, 65
- and personality disorders, 60
- attention deficit hyperactivity disorder (ADHD), 62, 64
- Bateman, Antony, 72–3
- Beck, Tim, 75–6
- behavioural manifestations versus personality disorder, 16
- Bible, Old Testament
- book of Proverbs, 3
- and personality, 3
- book of Proverbs, Bible, 3
- borderline personality disorder
- diagnosis, 20
- ICD-11 definition, 20
- borderline personality disorder (BPD)
- classification problems with, 34–5
- clinical diagnosis of, 31–3
- clinical origins of, 31
- epidemiology, 43
- inconsistent clinical criteria for, 33–4
- integration into ICD-11, 113–14
- lack of evidence to make separate domain or factor, 35
- post-traumatic stress disorder (PTSD) and, 36
- severity levels, 38
- trauma and, 36
- treatment choices, 69–72
- brittle diabetes, 54–5
- characters (personality), 2–3
- cognitive analytic therapy (CAT), 76
- cognitive behaviour therapy (CBT), 75–6
- complication model, 64
- complication/scar model, 59
- consanguinity, 61
- culture and personality disorders, 43, 44–5, 47
- depression
- pathoplasty model for treatment, 65
- stigma attached to, 98
- detachment
- as trait domain, 17
- treatment, 88
- diagnosis
- borderline personality disorder, 20, 31–3, 81–2
- failed system, 111–12
- ICD-11, 13
- dialectical behaviour therapy (DBT), 73–4
- Die Psychopathischen Persönlichkeiten*, 7
- disinhibition
- as trait domain, 88–9
- treatment, 88–9
- dissociality
- as trait domain, 18
- treatment, 88
- DSM-III, 58
- DSM-5, 58
- eating disorders, 62
- externalising factor, 16
- Five Factor Personality Model, 7
- Freud, Sigmund, 4
- Galen, 1–2
- Galenic syndromes, 61
- gender and personality disorders, 42
- general neurotic syndrome, 61
- genetics and psychopathy, 19
- Gjessing, Rolv, 5
- good psychiatric management, 69
- habit, 7
- Hare, Robert, 17–19
- Hayes, Steven, 77–8
- health
- and personality disorders, 54–6
- and personality traits, 52–3
- Henderson, David, 5
- Hippocrates, 1
- Hofstede scale, 44–5
- humours (personality), 1–2
- ICD-11
- health professionals making personality status with every patient, 114–15
- personality disorder treatment, 81–2, 83–4
- post-traumatic stress disorder (PTSD) and, 36
- ICD-11
- personality disorder diagnosis assessment of severity, 13, 15

- personality disorder (cont.)
 borderline personality disorder, 20
 borderline personality disorder integration, 113–14
 definition, 15
 measurement, 20–4
 ICD-11
 trait domain
 anankastia, 17
 borderline personality disorder, 20
 detachment, 17, 88
 disinhibition, 17, 18, 88–9
 dissociality, 18
 and Five Factor Personality Model, 22–4
 measurement, 20–4
 negative affectivity, 17, 87–8
 personality difficulty, 16
 qualifiers, 15–18
 summary, 17
 immigration and personality disorders, 48–9
 individualism–collectivism
 cultural model for personality disorders, 44–5, 46–8
 internalising factor, 17

 Kernberg, Otto, 31–2, 74
 Knight, Robert, 31

 Level of Personality Functioning Scale (LPFS), 29
 Linehan, Marsha, 73–4
 literature and personality, 6–7

 mental illness and personality, heredity and, 4
 mental illness and personality disorders
 clinical implications, 67
 co-occurrence between DSM axis I and axis II, 58–9
 DSM-III impact, 58
 historical ages of, 1–9
 history of, 57
 mental illness and personality disorders examples
 anxiety disorders, 60
 attention deficit hyperactivity disorder (ADHD), 62, 64
 comorbid disorders, 66–7
 eating disorders, 62
 general neurotic syndrome, 61
 health anxiety, 65–6
 mood disorders, 60
 schizophrenia, 62
 substance use disorder, 61–2, 64, 67
 mental illness and personality disorders models
 complication/scar model, 59
 pathoplasty model, 60, 65
 spectrum model, 64–5
 spectrum/co-aggregation model, 59–60
 vulnerability and complication models, 63–4
 vulnerability/predisposition model, 59
 mentalisation based treatment (MBT), 72–3
 Meyer, Adolf, 4–5
 modernisation and personality disorders, 49
 mood disorders, 60

 National Morbidity Study, 2000 (UK), 26
 natural selection, 78
 negative affectivity
 as trait domain, 17
 treatment, 87–8
 Nottingham Study of Neurotic Disorder, 28

 pathoplasty model, 60, 65
Personality and Fate of Registered Prostitutes, The, 7
 Personality Assessment Questionnaire for ICD-11 (PAQ-11), 28
 personality assessment
 severity, 13, 15
 personality assessment stages, 10–12
 personality difficulty
 clinical ways to assess, 29
 frequency of, 28
 in ICD-11, 16
 ICD-11 coding, 26
 long-term implications, 30
 versus personality disorder, 26–8
 personality disorder
 diagnosis at any age, 115–16
 ICD-11 definition, 15
 versus personality difficulty, 26–8
 personality disorder classification
 borderline personality disorder and, 20, 31–6
 borderline personality disorder integration, 113–14
 changes from ICD-10 to ICD-11, 12–13
 Five Factor Personality Model, 7
 ICD-11 trait domain, 13–18
 new spectrum model of, 112–13
 PCL-R and, 17–19
 psychopathy and, 19
 personality disorder epidemiology
 age and, 42–3
 antisocial personality disorder, 43
 borderline personality disorder (BPD), 43
 cultural interactions, 46–7
 culture and, 43, 44–5
 gender and, 42
 immigration and, 48–9
 modernisation and, 49
 prevalence of personality problems, 39–41
 rates of individual disorders, 42
 social withdrawal and, 45
 Western influence in models, 49–50
 personality disorder historic ages
 age 1 – Age of Discovery (460 BC–1700 AD), 1–4
 age 2 – Age of Personality as a Substrate (1750–1950), 4–7
 age 3 – Age of Investigation and Intervention, 7–9
 age 4 – Age of Acceptance and Understanding, 9
 personality disorder outcome
 age range trends, 94
 background, 93–4
 individual and trait studies, 94–6

- personality disorder short scales
 Quick Personality Assessment Schedule (PAS-Q), 125
 Short Social Functioning Questionnaire (SSFQ), 128
 Standardised Assessment of personality (SAPAS), 122
 personality disorder treatment
 acceptance and commitment therapy (ACT), 77–8
 care for people not requesting treatment, 118
 cognitive analytic therapy (CAT), 76
 cognitive behaviour therapy (CBT), 75–6
 dialectical behaviour therapy (DBT), 73–4
 large number of, 69–72
 mentalisation based treatment (MBT), 72–3
 need to de-stigmatise, 118–20
 need to improve strategies, 116–17
 nidotherapy, 71, 78–80, 85, 88, 150, 154, 155
 Psychologically Informed Planned Environments (PIPEs), 81
 schema-focused therapy, 74
 social prescribing, 79–80
 summary, 71
 systems training for emotional predictability and problem solving (STEPPS), 76–7
 therapeutic communities, 80–1
 transference-focused therapy, 74
 personality disorder treatment (ICD-11)
 assessment and diagnosis, 81–2
 clinical utility of severity, 83–4
 pharmacology, 89–91
 severity as diagnostic criterion, 83
 personality disorders
 and comorbidity with other mental illnesses, 57–63
 failed diagnostic system, 111–12
 health and, 54–5
 physical health problems and, 55–6
 stigma of, 99–110
 personality function, 12
 Personality Inventory for ICD-11 (PiCD), 22, 28
 personality trait domains
 treatment
 challenge of, 84–5
 clinical assessment, 85
 clinical principles, 85
 as focus for psychiatric methods, 86
 for multiple traits, 92–3
 personality traits population studies, 95–6
 personality types, 3
 person-environment fit, 46
 post-traumatic stress disorder (PTSD)
 as anxiety disorder, 60
 borderline personality disorder (BPD) and, 36
 Prufrock syndrome, 78
 Psychologically Informed Planned Environments (PIPEs), 81
 Psychopathic States, 5
 psychopathy
 classification, 19
 genetics and, 19
 Psychopathy Check List Revised (PCL-R), 17
 Quick Personality Assessment Schedule (PAS-Q), 125
 Ryle, Anthony, 76
 schema-focused therapy, 74
 schizophrenia, 62
 Schneider, Kurt, 7–8
 Short Social Functioning Questionnaire (SSFQ), 128
 social prescribing, 79–80
 social withdrawal and personality disorders
 taijin kyofusho, 45
 Western versus Eastern cultural sensibilities, 45
 spectrum model, 64–5
 spectrum/co-aggregation model, 59–60
 Standardized Assessment of Severity of Personality Disorder (SASPD), 22
 background, 122
 and personality difficulty, 28
 questionnaire, 123–4
 Starting to Shrink, 30
 stigma
 and incorrect personality assessment, 12
 need to de-stigmatise personality disorders, 118–20
 in psychiatry, 98–9
 stigma of personality disorder health professionals' view, 99–100
 patients' view, 101–2
 prevalence of, 99
 psychiatrists' view, 102–10
 solution to, 110–11
 structured clinical management, 69
 substance use disorder and personality disorders, 61–2, 64, 67
 systems training for emotional predictability and problem solving (STEPPS), 76–7
 taijin kyofusho, 45
 Theophrastus, 2–3
 therapeutic communities, 80–1
 trait domain qualifiers (ICD-11), 22
 trait theory
 background of, 5–6
 and borderline personality disorder, 34–5
 Five Factor Personality Model, 7
 traits across cultures
 comparisons, 47–8
 prevalence of, 47
 traits and health, 52–3
 transference-focused therapy, 74
 trauma and borderline personality disorder (BPD), 36
 vulnerability model, 63–4
 vulnerability/predisposition model, 59
 Young, Jeffrey, 74
 Zettle, Robert, 77–8