

Index

- approximation
 - Moreau–Yosida, 225
 - test functions separated in all variables, 230
 - with polynomials in C^k norm, 230
- asynchronous exponential growth, 193
- bi-Lipschitz homeomorphism, 101
- Bochner integral, 269
- boundedness, 210
- cohort, 157
- commutator estimate, 85
- compactness, 210
 - local, 209
 - relative, 210
- cone of nonnegative measures, 14, 26, 238
- continuity
 - absolute (function), 273
 - absolute (measure), 239
 - α -Hölder, 226
 - Lipschitz (semigroup), 274
 - narrow, 14, 69
 - uniform, 38
 - weak*, 14
- convergence
 - narrow, 14, 243, 245
 - weak*, 14, 243, 245
- decomposition
 - Hahn, 12, 235
 - Jordan, 12, 236
 - polar, 241
- diagonalisation argument, 32
- diameter, 210
- dual problem, 64, 91, 92, 134
- Escalator Boxcar Train (EBT), 155
- essential growth bound, 190
- Euler scheme, 149
- flat metric, 16, 26
 - algorithm to compute, 152
- flow, *see* semigroup, two-parameter
- Gronwall inequality, 219
- isometry, 216
- measure
 - absolute continuity, 239
 - Borel, *see* measure, locally finite
 - finite, 13, 236
 - inner regular, 13, 238
 - locally finite, 13, 238
 - outer regular, 13, 238
 - pre-tight, 261
 - push-forward, 105, 237
 - Radon, 13, 238
 - regular, 13, 238
 - separable, 261
 - σ -additive, 12
 - σ -finite, 13, 236
 - signed, 12, 235
 - singular, 239
 - tight, 29, 251, 265
- measure solution
 - to linear model on metric space, 109
 - to linear model on $\mathbb{R}^+$, 63
 - to nonlinear model on metric space, 122
 - to nonlinear model on $\mathbb{R}^d$, 129
 - to nonlinear model on $\mathbb{R}^+$, 60, 61
- metric space, 209
 - uniformly discrete, 20
- completion, 216
- Polish, 212
- proper, 212

- metrizability, 209
- modulus of continuity, 102
- network, 205
- norm
 - Hölder seminorm, 197
 - Bielecki, 66, 94, 115
 - bounded Lipschitz, 11
 - operator, 11
 - total variation, 15, 236
- perturbation, 195
- Radon–Nikodým derivative, 240
- reconstruction
 - fixed-equal-mass, 180
 - fixed-location, 179
- Riemannian manifold, 233
- semigroup, 272
 - irreducible, 193
 - quasi-compact, 190
 - exponentially Lipschitz continuous, 274
 - Lipschitz-AC, 274
 - Lipschitz continuous, 274
 - Lipschitz-UC, 274
 - two-parameter, 273
- semigroup property, 68
- separability, 209
- separating family, 243
- solution
 - dual problem, 64
 - renormalised, 204
- space
 - bounded functions, 215
 - bounded Lipschitz functions, 11, 220
 - dual space, 11
 - finite signed Radon measures, 14, 15, 238, 239
 - metric, 209
 - positive linear functionals, 37
 - splitting
 - splitting operator technique, 82
 - splitting-particle method, 178
 - sum of semigroups, 89
 - supremum metric, 215
 - theorem
 - Arzelà–Ascoli, 217
 - Banach–Alaoglu, 218
 - Dini, 217
 - Hahn–Banach, *see* theorem, hyperplane separation
 - Hopf–Rinow, 234
 - hyperplane separation, 218
 - inverse mapping, 217
 - Lebesgue differentiation, 218
 - McShane–Whitney (extension), 226
 - Portmanteau, 246
 - Prokhorov, 31, 254
 - Rademacher, 222
 - Radon–Nikodým, 240
 - Riesz representation, 241
 - Stone–Weierstrass (compact version), 229
 - Stone–Weierstrass (locally compact version), 229
 - tightness, 29, 31
 - topology, 209
 - uniform integrability
 - of functions, 237
 - of measures, 49
 - upwind strategy, 149
 - variation, 13, 236
 - vertex, 205
- Wasserstein distance, 44
 - algorithm to compute, 151
 - generalised, 53
 - Wasserstein-type, 48