

Index

Entries in bold typeface denote figures, tables, and algorithms.

A

- ABC(DE) approach, 17–21
- abdominal pain
 - algorithm for, 234
 - characteristics/location of, 237–240
 - clinical management of, 240–244
 - pathophysiology of, 236–237
 - as symptom of serious illness, 32
- abdominal trauma
 - and complications due to pregnancy, 187–188
 - and primary/secondary survey procedures, 189–192
 - and trauma to uterus/haemorrhage, 188–189
- abortion, 13, 496–497
- Abortion Act (1967), 13, 496–497
- abruption, 167, 188
- absolute fluid loss, 40–41
- acquired heart disease, 248–253, **251**
- acute myocardial infarction (MI), 249–252, **251**
- adrenaline, 154
- adult respiratory distress syndrome (ARDS), 57, 228
- advanced directives, 313–314, 494–496
- advanced life support, 110, **108**, 114–115
- Advanced Life Support Group (ALSG), 3
- aftercoming head of breech, 440–441, 438
- airway
 - assessment of, 86–87
 - maintenance of, 90–96, **91**, **93**, **95**
 - procedure to open, 110–112
- airway devices, 90, **91**, 94–95
- airway management
 - advanced techniques for, 91–96, **93**, **95**
 - clearance of obstructed, 88–90, **89**
 - in neonatal resuscitation, 156–157
 - primary survey of, 17
- all-fours position, 382
- amniotic fluid embolism (AFE)
 - algorithm for, **120**
 - as cause of maternal death, 13
 - clinical manifestations of, 122–124, **124**
 - diagnosis of, 124–125 and fetal outcome, 127
 - incidence of, 121–122
 - major obstetric haemorrhage, 304
 - management procedure for, 125–126
 - and uterine trauma, 189
- AMPLE method, 190
- anaesthesia
 - as cause of maternal death, 13–14
 - complications from spinal/epidural, 450–457
 - failure of, 454–455
 - management and obstetric hemorrhage, 312–313
 - and manual removal of placenta, 335–336
 - and placenta accreta, 334
 - and pre-eclampsia, 291
 - and problems associated with administration of, **446**
- anal sphincter trauma
 - assessment of, 424, **425**
 - postoperative care, 431
 - repair of third and forth degree tears, **430**
 - repair of third and fourth degree tears, 426–430, **428**, **429**
 - training to improve recognition of, **432**, 432
- anaphylactic shock, 43, 48–49
- anaphylaxis, 43
- anencephaly, 390
- angiotensin-converting enzyme (ACE) inhibitors, **251**, 248
- antenatal care, 259–261, **260**
- antenatal steroids, 294
- anterior placenta, 325
- antidepressants, 280, 277–278
- antihypertensives, 290–291
- antiplatelet therapy, **251**
- antipsychotic drugs, 279
- anxiety, 33, **33**, 273–274
- aortic dissection, 252
- apnoea, 148, **141**, **142**, 141–143
- appendicitis (acute), 242
- Arrow Peripheral Emergency Infusion Device (EID), 66
- Arrow Rapid Infusion Catheter (RIC), 66, **67**
- arterial blood gases (ABG)
 - interpretation of measurements of, **35**, 34–35
 - and intravenous fluid administration, 77
 - and pulmonary embolisms, 133
- asystole rhythm, 114
- ATOMFC (thoracic injuries), 180
- at-risk women, 277–279
- automated external defibrillator (AED), 109, **107**, 113–115
- autoregulation, 40
- AVPU score, 26, 198
- awareness during anaesthesia, 447

B

- backache, 456
- bacteraemia, **52**
- balloon tamponade, 392
- basic life support, **106**, 109
- Battle's sign, 204
- beta-blockers, **251**, 290
- betamimetic treatment, 402
- bicarbonate, 154

502 MANAGING OBSTETRIC EMERGENCIES AND TRAUMA

- bipolar disorder, 277–278
- bladder filling, 392
- blood, 70, 72–75, **74**
- blood loss
 - clinical parameters of, **47**
 - clinical presentation/recognition of, 305
 - clinical signs of, 76–77, **76**
 - and fetal distress, 189 recognition of, 45
 - severity of during maternal haemorrhage, **46**
- blood pressure, 286–289, **286**
- blood transfusions
 - and major obstetric haemorrhage, 304
 - refusal of, 76, 313–315
 - and types of blood products for, 72–75, **74**
- B-Lynch suture, 309, **310**
- Bolam principle, 491
- brachial plexus impairment, 376
- Bracht technique, 407, 410, **411**
- bradycardia
 - and neurogenic shock, 212
 - in newborn babies, 140
- breathlessness, 31, **32**
- breech and external cephalic version, **400**, 401–405, **404**
- bupivacaine, 449 burns
 - algorithm for, 226
 - as cause of hypovolaemic shock, 44
 - electrical, 232
 - pathophysiology of, 227–229
 - primary survey, 229–231, **230**
 - secondary survey, 231
- button hole tear, **425**
- C**
- caesarean section
 - algorithm for, **316**, **317**, **318**
 - and cardiac arrest, 117–118
 - due to abdominal trauma, 193
 - incidence of, 319
 - and increase in occurrence of placenta accreta, 331–332
 - and mentoposterior position, 397
 - potential difficulties of, 323–327
 - rate of with external cephalic version, 403
 - in resource-poor areas, 441
 - and surgical technique for, 320–322
 - and twins, 417, 419, 421
 - and use of anaesthesia, 449
- calcium, 115
- calcium channel blockers, **251**
- cannulae, 66, **66**, 68
- capillary refill time (CRT), 45
- capnography, 113
- carbon monoxide, 228, 229
- carboprost, 309
- carboxyhaemoglobin, 228
- cardiac arrest, 114–115, 117–118
- cardiac disease
 - and acquired heart disease, 248–253
 - as cause of maternal death, 14
 - and congenital heart disease, 253–256
 - and intravenous fluid administration, 81
 - service provision for, 253–256
- cardiac output, 39, 41
- cardiac tamponade, 181
- Cardiff wedge, **209**
- cardiogenic shock, 42, 48
- cardiomyopathy, 248–249
- cardiopulmonary resuscitation (CPR)
 - and assessment of rhythms, 114–115
 - and eclampsia, 297
 - and major obstetric haemorrhage, 306–307
 - as part of primary survey, 18
 - and pregnancy considerations, 110–112, **111**, 116–118
 - procedure for, 110–115
 - and unresponsive patient, 19
 - and use of AED during, 113–114
 - and use of intravenous fluids, **77**
- cardiotocography, 288 cell salvage
 - and amniotic fluid embolism, 122
 - and blood loss, 75–76
- central venous pressure (CVP)
 - as measurement of circulation, 78
 - and monitoring for sepsis, 56
 - and need to monitor in pre-eclampsia, 296–297
 - and regulation of transfusion rate, 69
- Centre for Maternal and Child Enquiries (CMACE), 4
- cephalic replacement, 383
- cephalopelvic disproportion, 345, 395, 438
- cerebral malaria, 269
- cerebral perfusion, 196–197
- cerebral perfusion pressure (CPP), 196, 197
- cerebral thrombosis, **263**
- cerebral venous sinus thrombosis, **263**
- cerebrospinal fluid, **448**
- cervical collar, 215
- cervical spine immobilisation, 88, 214–215, **214**
- chest compressions, 112, 152–154, **153**
- chest drain insertion, 184–186, **186**
- chest radiograph (CXR), 132
- Chester v. Afshar (England), 489–490
- Children Act (1989), 495 chin life/head tilt, 88, **89**
- cholecystitis, 238
- cholecystitis (acute), 239, 242
- chorioamnionitis, 238
- circadian rhythm disturbance, 485–486
- circulation
 - and abdominal trauma, 190–191
 - assessment of, 18
 - and burns, 230, **230**
 - and neonatal resuscitation, 142–143, **142**, **143**
 - in newborn babies, 140
 - techniques to induce in neonatal resuscitation, 152–155, **153**
- citrate anticoagulation, 72–73
- cleidotomy, 380
- Clostridium septicum*, 53–54
- Coca Cola urine, 299 coincidental deaths, 15
- colicky pain, 237, 238, 240
- colonic neoplasms, 239
- colonic pseudo-obstructions, 239, 243

colour, 146–147
 communication
 during cardiopulmonary resuscitation, 119
 and human error, 480–482, **481**, **482**
 and major obstetric haemorrhage, 306
 with parents of new born baby, 162
 and patients refusing treatment, 313–314, **302**
 and transfer considerations, 468–469
 of value of external cephalic version, 403
 compartment syndrome, 223–224
 compressible haemorrhage, 47
 computed tomography pulmonary angiogram (CTPA), 133
 concussion, 201
 Confidential Enquiries into Maternal Deaths (CEMD), 4, 7–8
 Confidential Maternal and Child Enquiry (CMACE), 7
 confirmation bias, 484
 confusion, **263**
 congenital heart disease, 253–256
 consent
 capacity for, 492–496
 requirements for, 489–491
 and sufficient information, 491–492
 voluntary, 497–498
 continuous non-locking technique, 426
 continuous positive airway pressure (CPAP), 158, 160
 corticospinal tract, 210
 craniotomy, 440
 Crede's manoeuvre, 339
 Creutzfeldt-Jakob (CJD) prion, 73
 Crohn's disease, 238
 crossmatching, 73
 crush injuries, 221
 cryoprecipitate, **74**, 75, 126
 crystalloids, 71
 Cushing's response, 198

D

decapitation, 441
 deep vein thrombosis, 239
 depression, 273–274
 dermatomes, **211**
 destructive operations, **436**, 439–442
 dextrose solutions, 71
 diabetes
 classification of, 256
 and complications during pregnancy, 256–257
 gestational, 261
 service provision for, 258–261, **260**
 diabetic hypoglycaemia, **263**
 diabetic ketoacidosis, 246, 258
 diagnostic peritoneal lavage (DPL), 192
 diaphragmatic disruption, 182–183
 differential diagnosis, 124, 131, 266
 Difficult Airway Society guidelines, 446
 diffuse axonal injury, 201
 diffuse primary brain injury, 201
 dilated cardiomyopathy, 248
 distress, 33, **33**
 distributive shock, 42–44
 domestic abuse, 171–175, **178**

Domestic Violence Crime and Victims Act (2004), 174
 doppler probes, 79
 drug therapy
 and myocardial infarction, **251**
 and pre-eclampsia/eclampsia, 289–294
 and uterine inversion, 342
 dura mater, **448**

E

E. coli, 53–54
 echocardiography, 133
 eclampsia
 algorithm for, **282**
 definition of, 283
 and headaches, **263**
 and intravenous fluid administration, 80
 treatment of, 297–299, **298**
 ectopic pregnancy, 13
 Ehlers-Danlos Type IV disorder, 252
 electrocardiogram (ECG), 133
 encephalopathy, **263**
 endotracheal intubation, 92–93, **93**
 epidural abscess, 456
 epidural anaesthesia (block), **448**, 448–450
 epidural space, **448**
 epilepsy, **263**
 episiotomy
 and face presentation, 396
 and forceps delivery, 363
 and protocols for use of, 424
 and shoulder dystocia, 380
 and venthouse delivery, 360
 error chains, 478–480, **478**, **479**
 escharotomy, 231
Escherichia coli, 53–54
 ethnicity, 10, **10**
 express consent, **10**
 external cephalic version (ECV), 401–405, **404**, 418, **400**
 extracorporeal membrane oxygen treatment (ECMO), 62
 extradural haemorrhage, 202
 extubation, 96–97, 447

F

face presentation, 367, 395–398, **394**
 Family Law Reform Act (1969), 493
 fatigue, 485–486
 femoral vein access, 69
 fetal death
 and destructive procedures, 438
 and placental abruption, 311
 and shoulder dystocia, 376
 and trauma, 167
 fetal head
 bobbing of, 379
 entrapment of, 323–324, 412, 438
 fetal outcomes, 118, 127
 fetal position
 breech, 325
 and caesarian section, 321, 323–324
 and delivery type, 356
 and rotational delivery, 364
 and shoulder presentation, 326

504 MANAGING OBSTETRIC EMERGENCIES AND TRAUMA

and use of ventouse cup, 359
 fetal squames, 124 fetal wellbeing
 and abdominal trauma, 192 assessment of, 20–21
 and complications from anaesthesia, 457
 and placental abruption, 311–312
 and pre-eclampsia, 288
 and unconscious patient, 199
 fibroids, 304, 324
 first degree burns, 231
 flail chest, 181
 fluid management, 290–291, 295–296
 focal primary brain injury, 202
 Focused Assessment Sonography in Trauma (FAST) scan, 191
 forceps delivery, **352**
 after failure of ventouse cap, 368–369
 and aftercoming head, **409**
 and safety points for aftercoming head of breech baby, 367, **368**
 safety rules for, **363**
 and use in rotational delivery, 364–367
 and use of traction forceps, 363
 forceps placement
 and rotational delivery, 364–365 traction forceps, **364**
 fresh frozen plasma, 73, **74**
Fusobacterium necrophorum, 53–54

G

Gaskin manoeuvre, 382
 Generic Instructor Course, 4
 gestational diabetes, 261
 Gillick competent, 493
 Glasgow Coma Score (GCS), 19, 200–201, **200**
 glucose tolerance test (GTT), 261

H

haematocrit, 307
 haemorrhage
 as cause of maternal death, 11–12, **12**
 due to brain injury, 202–203
 and intravenous fluid administration, 79
 trauma-related, 189
 haemostasis, 322
 Hagen-Poiseuille equation, 65
 harm avoidance model, 490
 Hartmann's solution, 71
 Haultain's technique, 342
 head blocks/straps, 216
 head injuries, 201–204, **201**
 head tilt/chin lift, 88, **89** headache
 as complication from epidural anaesthesia, 455
 red flag symptoms, 31
 types of/causes of, **263**, 267–268
 heart rate, 44, 76, 147
 heart size, 140
 HELLP syndrome, 239, 283, 299
 HELPERR (mnemonic), 384 herpes zoster, 240
 high spinal anaesthesia (block), 452–454
 human factors
 and decision making, 486–487

and error chains, 478–480, **478**, **479**
 and fatigue, 485–486
 and miscommunication, 480–482, **481**, **482**
 and situation awareness, 483–485, **483**
 human factors training, 477
 Human Rights Act (1998), 495, 496
 Huntingdon's procedure, 342
 hydralazine, 290
 hydrostatic repositioning, 341
 hyperbaric anaesthesia, 449
 hypercarbia, 255
 hypertension, 283, 290–291
 hypertensive disease, 11, **11**
 hypertrophic obstructive cardiomyopathy (HOCM), 249
 hypoglycaemia, 257–258
 hypotension, 45, 450–451, 122
 hypothermia, 70–71
 hypovolaemic shock, 41, 44–45, 47
 hypoxia
 as cause of confusion, **263**
 and flail chest, 181
 in utero, 141–143
 hypoxic ischaemic encephalopathy, 127
 hysterectomy, 311, 334, 348

I

implied consent, 490
 incompetence, 493
 indirect death, 14–15, **14**
 inflation breaths, 151
 influenza, 62–63
 intermittent oxygenation, 93–94
 internal podalic version, 418, 421
 International Liaison Committee on Resuscitation (ILCOR), 94
 intertwin delivery interval, 418
 intestinal obstruction, 244
 intra-abdominal pathology, 235–236, **236**
 intracerebral laceration, 203
 intracranial neoplasms, 269
 intracranial pressure (ICP), 196, 197
 intraosseous technique (IO), 66–68, **67**
 intraperitoneal pain, 238
 intrathecal block, 448
 intrauterine tamponade, 309
 intravenous access, **64**, 66–70, **67**
 intravenous cutdown, 70
 intravenous fluid
 administration of, 70–71
 surveillance and management of, 78–81
 types of, 71–76, **74**
 and use in neonatal resuscitation, 155
 intubation
 difficulties associated with, 445–447
 endotracheal, 92–93, **93**
 failure of, **444**
 ischaemia, 223
 ischaemic heart disease, 249–252, **251**
 ischaemic skin loss, 213

J

jaw thrust, 88, **89**, **99**, **150**
Jehovah's Witnesses, 76, 313, **302**, 497–498
Johnson manoeuvre, 341

K

Kjelland forceps, 355–356, 365–367, **366**
Kleihauer test, 405

L

labetalol, 289, 290
lanugo, 124
laparotomy, 348
Laryngeal Mask Airways (LMA) device as alternative to bag mask, 94–95
 insertion procedures for, 99–101, **100**, **101**
 and use in neonatal resuscitation, 157
lateral tilt position, **42**
latex sensitivity, 43
learning disabilities, 492–493
levobupivacaine, 449
limb threatening injuries, 221–224
limb weakness, 199
lithium, 277, 279
Lloyd Davies position, 223
log roll, 210, **210**, 216–217, **217**
long-bone fractures, 220–221
Lovset's manoeuvre, **408**, 410
low-molecular-weight heparin (LMWH), 134–136, **134**, **251**

M

macrosomia, 257, 345
magnesium sulphate
 adverse effects of, 293, **294**
 and pre-eclampsia, 292–293
 and treatment of seizures, 297–298, **298**
 and use during cardiac arrest, 115
Magpie trial, 283, 292
major obstetric haemorrhage
 algorithm for, **301**
 and anaesthesia management, 312–313
 and antenatal and intrapartum care, 314–315
 as cause of maternal death, 303–304
 definition/cause of, 304
 general management of, 305–308
 tissue and trauma management, 311–312
 and tone management, 308–310
Managing Obstetric Emergencies and Trauma (MOET), 3–5, **5**, **4**
Marfan's syndrome, 181, 252
massive haemothorax, 181
massive pulmonary embolism, 131
material risk, 491
maternal collapse, 19, 41
maternal mortality
 and age as risk factor for, 10
 as caused by abortion, 13
 as caused by amniotic fluid embolism, 121–122, 13
 as caused by congenital heart disease, 253

 as caused by intra-abdominal pathology, 235–236, **236**
 as caused by suicide, 167
 and causes of direct deaths, 11–14, **11**, **12**
 global rate of, 3, 15–16, **16**
 and indirect deaths, 14–15
 and major obstetric haemorrhage, 303–304
 and pre-eclampsia, 437
 and pulmonary embolisms, 129–130
 reduction of during Second World War, 8–9
Mauriceau-Smellie-Viet manoeuvre, **409**, 408
MBRRACE-UK, 7
McRoberts' manoeuvre, 380–381, 412
mean arterial pressure (MAP), 196, 197
meconium aspiration, 150–151
medico-legal issues
 and domestic abuse, 174
 and postmortem caesarian section, 118
 and shoulder dystocia, 384–385
 vaginal breech delivery, 412
meningitis, **263**, 268, 456
Mental Capacity Act (2007), 494–496
Mental Health Act (1983), 495
mentoanterior position, 397
mentoposterior position, 397
MEOWS (Modified Early Obstetric Warning System), 12, 26–28, **27**
methicillin-resistant *S. aureus* (MRSA), 54
migraine, **263**
misoprostol, 309
MIST (abdominal injury), 189
Morganella morganii, 54
Mothers and Babies – Reducing Risk through Audit and Confidential Enquiries across the UK, 7
motor block, 451
moxibustion technique, 403
musculoskeletal trauma, **218**, 219–224
myocardial contusion, 182

N

naloxone, 155
nasopharyngeal airway device, 90
needle cricothyroidotomy, 103–104, **103**
needle placement, **67** needle thoracocentesis, 184
neonatal resuscitation
 algorithm for, **163**
 assessment of need to, 144–145
 clearance of airway, 149–151, **149**
 and creation of operative team, 162
 and decision to discontinue, 156
 equipment for, 143–144, **144**
 pathophysiology of, 141–143, **141**, **143**
 procedure for poor response to, 160–161
 ventilation in, 151–152, **152**
neonatal temperature maintenance, 145–146, 161–162
nerve block, 448
neurogenic shock, 212
neuropraxia, 456
neutral head position, **149**, 149
newborn babies

506 MANAGING OBSTETRIC EMERGENCIES AND TRAUMA

assessment of, 146–148
 life support algorithm for, **138**
 and need for temperature control, 145–146
 physiology of, 139–140
 nifedipine, 290
 non-compressible haemorrhage, 47 nonvertex
 presentation, 418

O

obesity
 as factor in ruptured uterus, 346
 as a factor in shoulder dystocia, **377**
 as factor in sudden adult death syndroms, 248
 as risk factor for maternal death, 10
 as risk factor for thromboembolism, 12
 obstetric and sphincter injuries (OASIS), 423
 obstetric injury, 9, **9**
 obstructive shock, 44
 oesophageal disruption, 183
 Ogilvie's syndrome, 243
 oliguria, 295–296
 open fractures/joint injuries, 222–224
 open pneumothorax, 181
 operative vaginal delivery (OVD), 353–356, 369–370
 opioids, 454 optic fundi, 288
 oropharyngeal (Guedel type) airway device, 90, **91**,
 98, **98**
 O'Sullivan's technique, 341
 oxygen administration, 151, 158–159
 oxygen saturation levels
 in newborn babies, 147–148, **148**
 and pre-eclampsia, 288 in preterm babies, **160**
 and pulmonary oedema, 296
 oxytocin, to augment contractions, 397

P

PACE communication tool, **482**, 482
 packed red cells, 72, **74**
 pain relief, 230
 Pajot's manoeuvre, **365**
 pancreatitis (acute), 240, 242–243
 Panton-Valentine leucocidin (PVL), 54
 paraplegia, 211
 pathophysiology
 burns, 227–229
 myocardial infarction, 249
 sepsis, 54–55
 Patwardhan's procedure, 326
 pelvic injuries, 220
 peptic ulceration, 239
 percutaneous coronary intervention (PCI), 250
 peri-arrest arrhythmias, 452
 perinatal mortality
 and external cephalic version, 403
 and umbilical cord prolapse, 389–390
 and vaginal breech delivery, 405–406
 perineal trauma assessment of, 424
 levels of classification of, 423–424
 postoperative care, **431**
 repair of first and second degree levels, 424, 426
 peripartum cardiomyopathy, 248 pia mater, **448**

Piper forceps, 367
 placenta, 321, 335–336
 placenta accreta, **329**, 331–334
 placenta increta, 331
 placenta percreta, 331, 345
 placenta praevia, 304, 325, 390
 placental abruption, 304, 311–312
Plasmodium falciparum, 269
 plastic bags, 157–158, **159**
 platelets, 75, **74**
 pocket mask, 99, **99**
 polyhydramnios, 257, 390
 positive end expiratory pressure (PEEP), 158
 postdural puncture headache, 267–268
 posterior axillary traction, 383
 posterior columns, 210
 postnatal depression (PND), 274
 postural supine hypotension, 41
 preconceptual care, 253, 259
 pre-eclampsia algorithm for, **282**
 assessment/management of, 285–288
 definition of, 283
 and headaches, **263**
 and intravenous fluid administration, 80
 and major obstetric haemorrhage, 304
 symptoms of, 285
 pre-eclampsia/eclampsia antepartum/intrapartum
 management, 288–294
 and delivery considerations, 294–297
 health complications from, **285**
 pregnancy
 and arterial blood gas levels, **35**
 and diabetes, 256–261
 and domestic abuse, 172–173
 and emergency monitoring, 20–21
 and increase in blood volume, 41
 and neurological emergencies, 262–268, **263**
 and psychiatric illness, **263**
 and resuscitation considerations, **111**, 110–112,
 116–118
 and seat belt placement, 188
 termination of, 496–497
 and triage assessment, 463–466, **465**
 and use of MEOWS, 26–28
 and use of radiology during, 35–36
 preterm babies, 157–160
 primary apnoea, 141–143, **141**, **142**
 primary survey, 17–18
 prosthetic heart valves, 253
 proteinuria, **286**
 psychiatric illness
 as cause of maternal death, 14–15, 275
 and management of during pregnancy, 276–279
 and neonatal paediatrician needs, 279–280
 in pregnancy, **263**
 rates of pre-and-post pregnancy, 273–274
 puerperal psychosis, 274, **263**
 pulmonary angiography, 133
 pulmonary artery balloon tipped flotation catheters,
 78
 pulmonary contusion, 182 pulmonary embolism

- algorithm for, **128**
 - as cause of maternal death, 129–130
 - clinical presentation of, 131, **131**
 - maintenance treatment of, 135–136
 - management procedures for, 132–134, **133**
 - pulse pressure, 45
 - pulseless electrical activity (PEA), 114
 - pupillary function, 199
 - pyelonephritis, 239
 - pyrexia, 30–31
- Q**
- quadriplegia, 211
- R**
- RADAR, 173 radiology, 35–36
 - red flag symptoms
 - breathlessness, 31
 - headache, 31
 - pyrexia, 30–31
 - sepsis, **56**
 - regional block, 448
 - regurgitant valve disease, 252
 - relative fluid loss, 40–41
 - relative hypovolaemia, 41
 - renal function, 287–288
 - retained placenta, **330**, 335–336
 - reverse Woods’ screw manoeuvre, 382
 - Ringer’s lactate, 71
 - ritodrine, 398
 - Royal College of Obstetricians and Gynaecologists (RCOG), 12
 - Rubin II manoeuvre, 381 rule of nines, **230**
 - ruptured uterus, 345–348, **344**
- S**
- SBAR tool, 28, **29**, **482**
 - scalp wounds, 203
 - second degree burns, 231
 - secondary survey, 18–19
 - seizures, 266–267, 297–298, 452
 - selective serotonin re-uptake inhibitors (SSRIs), 278, 280
 - sepsis
 - algorithm for, **50**
 - and burns, 231
 - causal agents of, **53**, 53–54
 - as cause of headaches, **263**
 - as cause of maternal death, 13, 51, **52**
 - epidemiology of, 54
 - and intravenous fluid administration, 80–81
 - management of, 56–63, **60**
 - pathophysiology of, 54–55
 - symptoms of, **52**, 55–56
 - and use of antibiotics to prevent, **60**, 59–60,
 - septic shock
 - as cause of distributive shock, 43
 - principles of treatment, 47–48
 - symptoms of, **52**
 - septicaemia, **52**
 - septicaemia/intrauterine sepsis, 304
 - Serious Hazards of Transfusion Scheme (SHOTS), 307
 - serious illness
 - and arterial blood gas levels, 34–35
 - clinical and red flag symptoms, 30–33
 - and use of MEOWS, 26–30, **27**
 - and use of radiation, 35–36
 - serious injury, 19–20
 - shock
 - cardiogenic, 42 distributive, 42–44
 - hypovolaemic, 40–42
 - maternal signs of, 305
 - obstructive, 44
 - pathophysiological basis of, 39
 - shoulder dystocia
 - algorithm for, **374**
 - and clinical risks and outcomes, 376–378, **377**
 - definition of, 375
 - prevention of, 378–385
 - and symphysiotomy, 438
 - Sidaway v. Board of Governors (England), 490
 - sigmoid volvulus, 239, 244
 - situational awareness, 483–485, **483**
 - skull fractures, 204
 - socio-economic classification, 10
 - sodium bicarbonate, 115
 - sodium valproate, 279
 - sphincter tear, 368
 - spinal anaesthesia (block)
 - characteristics of, 448–449
 - dosages, 449–450
 - location of injection, **448**, 447
 - management of high and total, 452–454
 - spinal cord, 447–448, **448**
 - spine board, 209
 - spine/spinal cord injuries
 - algorithm for, **206**
 - assessment of, 210–211, **210**, **211**
 - immobilisation techniques, 208–209, **208**, **209**
 - primary survey, 212–213
 - probable causes of, 207
 - spinothalamic tract, 210
 - splenic artery aneurism/rupture, 239
 - St. Vincent Declaration (1991), 258
 - Staphylococcus aureus*, 53–54
 - statins, **251**
 - status epilepticus, 266–267
 - stenotic valve disease, 252
 - steriods, 61
 - Streptococcus*, 53–54
 - stroke volume, 40
 - structured laparotomy, 193
 - subarachnoid block, 448
 - subarachnoid haemorrhage, 203, **263**
 - subdural haemorrhage, 202–203
 - substandard care, 15, 25, 247
 - sucking chest wound, 181
 - suction, 88, 150–151
 - sudden adult death syndrome (SADS), 248
 - sudden unexplained death in epilepsy (SUDEP), 248
 - suicide, 275
 - sulphonamides, 8

508 MANAGING OBSTETRIC EMERGENCIES AND TRAUMA

supraglottic airway devices (SAD), 94–95
 suprapubic pressure, 381, 412
 surfactant deficiency, 158
 surgical cricothyroidotomy, 95, **95**, 101–102
 Surviving Sepsis Campaign, 57, 61–62
 symphysiotomy
 algorithm, **435**
 complications of, 437–438
 and head entrapment, 412
 indications and techniques of, 438–439
 and shoulder dystocia, 380, 383–384
 synthetic colloids, 72
 systemic inflammatory response syndrome (SIRS), **52**

T

T1 injury, 211
 tension pneumothorax, 180
 Term Breech Trial, 405
 terminal apnoea, 141–143, **141**, **142**
 third degree burns, 231
 thoracic injuries, 179–183
 Thorn's manoeuvre, 397
 thromboprophylaxis, 326
 tocolysis, 392, 402, 404
 tone, 146
 total spinal anaesthesia (block), 453–454
 tracheal intubation, 156–157, 160
 tracheobronchial disruption, 183
 transfer
 ACCEPT approach, 467–473
 and pre-eclampsia, 295
 trauma, **166**, 167–169
 Trendelenburg positioning, 391
 triage, 461–466, **462**, **463**, **465**
 triage sieve, **462**
 Twin Birth Study, 417
 twin pregnancy, 412, 417–421, **416**

U

UK Obstetric Surveillance System (UKOSS), 13
 ulcerative colitis, 238
 ultrasound guided access, 69–70
 umbilical cord, 145, **388**, 389–390
 unconscious patient
 algorithm for, **194**

and cerebral perfusion, 196–197
 and primary survey, 197–199
 and secondary survey, 199–201, **200**
 unfractionated heparin, 134–135, **135**
 uterine atony, 304
 uterine evacuation, 118
 uterine inversion, 304, 339–342, **338**
 uterine rupture, 188–189, 304
 uterus
 manual replacement of inverted, 341
 ruptured, 345–348

V

vaginal breech delivery failure to deliver, 410, 412
 and improvement in perinatal mortality, 405–406
 labour and delivery protocols, 406–410, **408**, **409**
 vaginal delivery prerequisites, **350**
 vascular injuries, 223
 vasodilators, 290
 vasopressors, 126, 454
 vena caval occlusion, 116
 venthouse delivery, **351**
 ventilation
 assessment of, 18, 87, 112 and CPR, 112–113
 efficacy of during CPR and pregnancy, 116
 in newborn babies, 140, 147
 and obstetric haemorrhage, 313
 techniques for assisted, 96–97
 ventilation/perfusion scan, 133, **133**
 ventouse cups
 advantages/disadvantages to using, 355
 and determination of appropriate type, 357, 359
 safe use of, **358**
 ventouse delivery, 357–362
 ventricular fibrillation (VF), 113
 vertex presentation, 418

W

Woods' screw manoeuvre, 382
 Wrigley's forceps, 325
 written consent, 490–491

Z

Zavanelli manoeuvre, 380, 383