

Camberwell Assessment of Need for the Elderly

Camberwell Assessment of Need for the Elderly (CANE)

Edited by

Juanita Hoe

City, University of London

Martin Orrell

University of Nottingham



CAMBRIDGE
UNIVERSITY PRESS

Cambridge University Press
978-1-911-62336-6 — Camberwell Assessment of Need for the Elderly (CANE), 2nd Ed.
Edited by Juanita Hoe, Martin Orrell
Frontmatter
[More Information](#)

CAMBRIDGE
UNIVERSITY PRESS

University Printing House, Cambridge CB2 8BS, United Kingdom
One Liberty Plaza, 20th Floor, New York, NY 10006, USA
477 Williamstown Road, Port Melbourne, VIC 3207, Australia
314–321, 3rd Floor, Plot 3, Splendor Forum, Jasola District Centre, New Delhi – 110025, India
79 Anson Road, #06–04/06, Singapore 079906

Cambridge University Press is part of the University of Cambridge.

It furthers the University's mission by disseminating knowledge in the pursuit of education, learning, and research at the highest international levels of excellence.

www.cambridge.org
Information on this title: www.cambridge.org/9781911623366
DOI: 10.1017/9781911623373

© The Royal College of Psychiatrists 2021

This publication is in copyright. Subject to statutory exception and to the provisions of relevant collective licensing agreements, no reproduction of any part may take place without the written permission of Cambridge University Press.

First published 2021

Printed in the United Kingdom by TJ Books Limited, Padstow Cornwall

A catalogue record for this publication is available from the British Library.

ISBN 978-1-911-62336-6 Paperback

Cambridge University Press has no responsibility for the persistence or accuracy of URLs for external or third-party internet websites referred to in this publication and does not guarantee that any content on such websites is, or will remain, accurate or appropriate.

.....

Every effort has been made in preparing this book to provide accurate and up-to-date information that is in accord with accepted standards and practice at the time of publication. Although case histories are drawn from actual cases, every effort has been made to disguise the identities of the individuals involved. Nevertheless, the authors, editors, and publishers can make no warranties that the information contained herein is totally free from error, not least because clinical standards are constantly changing through research and regulation. The authors, editors, and publishers therefore disclaim all liability for direct or consequential damages resulting from the use of material contained in this book. Readers are strongly advised to pay careful attention to information provided by the manufacturer of any drugs or equipment that they plan to use.

Contents

List of Figures vi
List of Tables vii
List of Contributors viii
Acknowledgements x

1	An Introduction to Needs Assessment and Use of the Camberwell Assessment of Need for the Elderly 1	7	Crisis and Assessment of Need in Dementia: Development of a Home Treatment Package 62
	Kunle Ashaye, Dilini Jayalath and Juanita Hoe		Juanita Hoe, Ritchard Ledgerd, Sandeep Toot and Martin Orrell
2	Self-Reported Needs of People with Dementia Living at Home: A Scoping Review 10	8	Needs of People with Young-Onset Dementia 77
	Alžběta Bártová, Iva Holmerová, Vladimíra Dostálová, Hana Bláhová and Michal Štefl		Christian Bakker and Britt Appelhof
3	Needs of Older Primary Care Patients 20	9	Needs of Older People in Long-Term Care Settings 86
	Janine Stein and Steffi Gerlinde Riedel-Heller		Justyna Mazurek, Dorota Szcześniak and Joanna Rymaszewska
4	Unmet Needs of Older People with and Without Depression in Residential Homes 34	10	Needs and Healthcare Costs in Old Age: An Application of the Camberwell Assessment of Need for the Elderly 91
	Hein van Hout, Jannicke M. Iversen and Marijke Boorsma		André Hajek, Janine Stein and Hans-Helmut König
5	Needs of Older People Living Alone: A Critical Review 44	11	The Future of Needs Assessment Research 97
	Raffaella Carvacho and Claudia Miranda-Castillo		Juanita Hoe and Martin Orrell
6	Needs Assessment of People with Dementia and Impact of Caregiver Burden 52	12	Instructions for the CANE 105
	Myonghwa Park, Thi-Thanh-Tinh Giap, Miri Jeong, Younghye Go and Dong Young Lee		
			<i>Index</i> 141

Figures

- 2.1 Search flowchart in accordance with PRISMA guidelines 11
- 3.1 Distribution of unmet needs from the patients' perspective ($n = 231$) 24
- 3.2 Ten most frequently rated unmet needs from the patients' perspective across the four disease-specific subgroups 25
- 3.3 Ten most frequently rated unmet needs from the relatives' perspective across the four disease-specific subgroups 26
- 3.4 Ten most frequently rated unmet needs from the GPs' perspective across the four disease-specific subgroups 27
- 4.1 Sample derivation 35
- 4.2 Percentages of unmet needs in CANE categories among depressed and non-depressed persons 40
- 5.1 Study selection process 46
- 6.1 Unmet needs with highest rate according to burden groups ($n = 320$) 58
- 6.2 Number of unmet needs and burden score according to different groups of number of unmet needs ($n = 320$) 58
- 7.1 Consultation and consensus processes used to develop the home treatment package 66
- 7.2 Home treatment package advisory protocol pathway 71
- 9.1 Profile of an older person staying in a long-term care setting 87

Tables

- | | |
|--|---|
| 1.1 Items Included in the Camberwell Assessment of Need for the Elderly 2 | 6.1 Characteristics of the People with Dementia and Their Family Caregivers 54 |
| 2.1 Summary of Articles Used in Thematic Analysis 12 | 6.2 Needs of People with Dementia and the Source of Help They Received ($n = 320$) 55 |
| 3.1 Socio-demographic Characteristics of Patients 23 | 6.3 Burden on Family Caregivers of People with Dementia ($n = 320$) 57 |
| 4.1 Description of the Total Study Population ($n = 213$) and for Persons Who Are Depressed or Non-depressed 36 | 7.1 Example of Interventions for the Problem Areas Contributing to Crises in Dementia (Behavioural/Psychological Domain) 67 |
| 4.2 Total Needs and Respectively Met and Unmet Needs (CANE) Separately for the Total Group of Persons in Residential Homes ($n = 213$) 37 | 7.2 Categorisation of the CANE Items to the Five Critical Domains that Contribute to Crises 68 |
| 4.3 Needs Measured by CANE Among Persons Who Are Depressed or Non-depressed 39 | 10.1 Sample Characteristics ($n = 1,095$) 93 |
| 4.4 Unmet Needs (CANE) Among Persons Who Are Depressed or Non-depressed 41 | 10.2 Determinants of Total Healthcare Costs (Societal Perspective): Results of Regression Analysis (Generalised Linear Models with Log Link and Gamma Distribution) 94 |
| 5.1 Characteristics of Selected Studies 47 | |

Contributors

Britt Appelhof

Radboud University Medical Centre, Department of Primary and Community Care, Radboudumc Alzheimer Centre, Nijmegen, Netherlands, and Archipel, Centre for Geriatric Care, Eindhoven, Netherlands

Kunle Ashaye

Mental Health Services for Older People, Hertfordshire Partnership University, National Health Service Foundation Trust, London, United Kingdom

Christian Bakker

Radboud University Medical Centre, Department of Primary and Community Care, Radboudumc Alzheimer Centre, Nijmegen, Netherlands; Groenhuysen, Center for Geriatric Care, Roosendaal, Netherlands; and Dutch Young-Onset Dementia Knowledge Centre, Amersfoort, Netherlands

Alžběta Bártová

Department of Longevity Studies, Faculty of Humanities, Charles University, Prague, Czech Republic

Hana Bláhová

Department of Longevity Studies, Faculty of Humanities, Charles University, Prague, Czech Republic

Marijke Boorsma

Amsterdam University Medical Centres, Vrije University, and Department of General Practice, Amsterdam Public Health Institute, Amsterdam, Netherlands

Raffaella Carvacho

Millennium Institute for Research in Depression and Personality and the Pontificia Universidad Católica de Chile, School of Psychology, Santiago, Chile

Vladimíra Dostálová

Department of Longevity Studies, Faculty of Humanities, Charles University, Prague, Czech Republic

Thi-Thanh-Tinh Giap

Research and Education Centre for Evidence-Based Nursing Knowledge, College of Nursing, Chungnam National University, Daejeon, Republic of Korea

Younghye Go

Research and Education Centre for Evidence-Based Nursing Knowledge, College of Nursing, Chungnam National University, Daejeon, Republic of Korea

André Hajek

Department of Health Economics and Health Services Research, University Medical Center Hamburg-Eppendorf, Hamburg Centre for Health Economics, Hamburg, Germany

Juanita Hoe

Division of Nursing, School of Health Sciences, City, University of London, London, United Kingdom

Iva Holmerová

Department of Longevity Studies, Faculty of Humanities, Charles University, Prague, Czech Republic

Jannicke M. Iversen

Amsterdam University Medical Centres, Vrije University, and Department of General Practice, Amsterdam Public Health Institute, Amsterdam, Netherlands

Dilini Jayalath

Mental Health Services for Older People, Hertfordshire Partnership University, National Health Service Foundation Trust, London, United Kingdom

Miri Jeong

Research and Education Center for Evidence-Based Nursing Knowledge, College of Nursing, Chungnam National University, Daejeon, Republic of Korea

Hans-Helmut König

Department of Health Economics and Health Services Research, University Medical Center Hamburg-Eppendorf, Hamburg Centre for Health Economics, Hamburg, Germany

Dong Young Lee

Department of Neuropsychiatry, Seoul National
 University Hospital, Seoul, Republic of Korea

Ritchard Ledgerd

Dementia Care Research Centre, North East London
 National Health Service Foundation Trust, London,
 United Kingdom

Justyna Mazurek

Department and Division of Medical Rehabilitation,
 Wroclaw Medical University, Wroclaw, Poland

Claudia Miranda-Castillo

Faculty of Nursing, Universidad Andres Bello and
 Millennium Institute for Research in Depression and
 Personality, Santiago, Chile

Martin Orrell

Division of Psychiatry and Applied Psychology,
 Faculty of Medicine and Health Sciences, Institute of
 Mental Health, University of Nottingham,
 Nottingham, United Kingdom

Myonghwa Park

Research and Education Center for Evidence-Based
 Nursing Knowledge, College of Nursing, Chungnam
 National University, Daejeon, Republic of Korea

Steffi Gerlinde Riedel-Heller

Medical Faculty, Institute of Social Medicine,
 Occupational Health and Public Health,

University of Leipzig, Leipzig,
 Germany

Joanna Rymaszewska

Department of Psychiatry, Wroclaw Medical
 University, Wroclaw, Poland

Michal Šteffl

Department of Longevity Studies, Faculty of
 Humanities, Charles University, Prague, Czech
 Republic

Janine Stein

Medical Faculty, Institute of Social Medicine,
 Occupational Health and Public Health, University of
 Leipzig, Leipzig, Germany

Dorota Szcześniak

Division of Psychotherapy and Psychosomatic
 Medicine, Department of Psychiatry, Wroclaw
 Medical University, Wroclaw, Poland

Sandeep Toot

Dementia Care Research Centre, North East London
 National Health Service Foundation Trust, London,
 United Kingdom

Hein van Hout

Amsterdam University Medical Centres, Vrije
 University, and Department of General Practice,
 Amsterdam Public Health Institute, Amsterdam,
 Netherlands

Acknowledgements

Firstly, we would like to thank Cambridge University Press for the opportunity to write this second edition of the *Camberwell Assessment of Need for the Elderly (CANE)*. The first paper on the CANE was published in 2000, followed by publication of the first edition of the Camberwell Assessment of Need for the Elderly (CANE), in 2004. Preparing this second edition has given us time to examine 20 years' worth of needs assessment research using the CANE with older people, which has been an enlightening and informative task. The growth in studies using the CANE is encouraging and has helped highlight the potential for future research and use of needs assessment as a health outcome measure. Particularly evident is the need to develop and evaluate interventions that can be used to successfully meet the unmet needs of older people identified within these studies.

We would like to thank all the authors who have contributed chapters to this book and have provided us with new research and insightful perspectives on how the CANE has been or can be used to assess the needs of older people in a variety of contexts. We would also like to extend our appreciation to the

many academics and clinicians who have published research showing the effectiveness of using the CANE within such a diverse range of older populations across the world. We would especially like to thank all the participants and their families for their involvement in the published studies about needs assessment research in older people. Mostly, thank you to all for helping to build the evidence base around needs assessment, which can be used to find ways to improve services and care for older people.

Finally, for help with preparing this second edition for publication, we send a special thanks to Chrissy Bailey, business support officer at the Institute of Mental Health, University of Nottingham, for her proficient administrative support and to Elena Profyri, a researcher at City, University of London, for her excellent editing and proofing skills. We also thank Monica Manela and Lucy Webster, two researchers at University College London who helped with updating the prompts and reviewing the terminology used in the CANE assessment to ensure that these reflect more present-day language and contemporary issues impacting older people's lives. Your contributions have all been invaluable in preparing this work for publication.